

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Warren-Gash, C; (2018) Herpes Zoster: Epidemiological Links With Stroke and Myocardial Infarction. *The Journal of infectious diseases*, 218 (suppl.), S102 – S106. *ISSN*0022 – 1899 *DOI* : <https://doi.org/10.1093/infdis/jiy385>

Downloaded from: <http://researchonline.lshtm.ac.uk/4649478/>

DOI: <https://doi.org/10.1093/infdis/jiy385>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

Herpes zoster: epidemiological links with stroke and myocardial infarction

Running head: Herpes zoster, stroke and MI

Author: Charlotte Warren-Gash¹

Affiliation:

1. Faculty of Epidemiology & Population Health, London School of Hygiene Tropical Medicine, London, UK

Contact details:

Dr Charlotte Warren-Gash, Associate Professor Epidemiology, Room 248c, London School of Hygiene & Tropical Medicine, Keppel Street, London WC1E 7HT

Email: charlotte.warren-gash1@lshtm.ac.uk / Tel: 020 7927 2796

Word count: 1,277

Key words: herpes zoster; cardiovascular disease; acute myocardial infarction; stroke; meta-analysis; vaccination; antiviral treatment.

ABSTRACT

Routine data from electronic health records (EHRs) provide insights into links between herpes zoster (HZ) and cardiovascular complications such as stroke or myocardial infarction (MI) in different populations worldwide. Evidence from large EHR studies using both self-controlled case series and traditional cohort designs suggests that there is a transient increase in the risk of stroke after HZ, which gradually resolves over 6 to 12 months. In these studies, herpes zoster ophthalmicus was associated with a higher risk of stroke than HZ at other sites. A larger effect size was seen in people aged under 40 years. Existing studies also suggest that HZ may have a triggering effect on MI, although fewer studies examine this outcome. Further evidence is needed on the effectiveness and cost-effectiveness of vaccine and antiviral drugs to reduce cardiovascular complications after HZ, in studies which are designed to minimise selection biases and confounding by indication.

Introduction

Acute infections, such as influenza, can trigger cardiovascular events including stroke and myocardial infarction (MI)^{1,2}. Interest is growing in the effect of persistent reactivating viruses such as varicella zoster virus (VZV) on the pathogenesis of cardiovascular diseases, which are the leading global cause of death. There are biologically plausible mechanisms through which VZV could trigger acute vascular events: inflammatory responses following reactivation of VZV from latency to cause herpes zoster (HZ) may lead to endothelial dysfunction and development of a hyper-coagulable state³. Pathological vascular remodelling associated with arterial VZV infection (VZV vasculopathy) may also contribute to vascular risk⁴. As the population ages, the burden of disease due to stroke, including disability, illness and premature death, is projected to double worldwide by 2030⁵. Understanding the current evidence and its limitations is crucial to inform vaccination and treatment strategies to minimise vascular complications of VZV.

Evidence for association between HZ and vascular complications

Historically, the ability to investigate relatively rare complications of infections was hampered by lack of power due to small sample sizes. Now, in the era of “big data”, with routinely collected electronic health records (EHRs) available for large numbers of patients with clinically-diagnosed HZ infections, vascular complications can be investigated in different population subgroups with increasing precision. There remain, however, methodological difficulties with interpreting data robustly from studies with different designs, exposure and outcome definitions, populations and settings. In the last year, several systematic reviews synthesized evidence of the relationship between HZ and stroke from cohort and self-controlled case series studies⁶⁻¹¹. Although the pooled effect estimates presented in these reviews vary due to differing methods and inclusion criteria, the combined evidence suggests the following:

- 1) HZ can trigger stroke.

There is a transient increase in the risk of stroke after HZ, which is highest in the earliest time period after HZ diagnosis – adjusted incidence ratio (IR) for stroke 2.37 (95% CI 2.17-2.59) up to one week after HZ¹² – and gradually diminishes to baseline by around six to twelve months. At four weeks, data from three powerful self-controlled case series studies using primary care EHRs from the US, UK and Germany, show a pooled IR for stroke of 1.55 (95% C.I. 1.46-1.65)¹²⁻¹⁴. These studies have the major benefit of implicitly controlling for fixed between-person confounding effects¹⁵. The findings are corroborated by several prospective cohort studies using data from Asian, European and US populations, which show a similar gradient of stroke risk up to one year¹⁶⁻²¹. Evidence for any longer-term effect of HZ on vascular risk is mixed, with marked heterogeneity between studies. A meta-analysis of four cohort studies using random effects found no association between HZ and stroke after one year: pooled OR 1.20 (95% CI 0.82-1.75)¹¹.

2) Herpes zoster ophthalmicus (HZO) leads to a higher risk of stroke than HZ at other sites.

Point estimates for the effect of HZO on stroke are generally higher than those for HZ at other or unspecified sites. HZO refers to reactivation of VZV in the first division of the trigeminal nerve, and it has been proposed that the proximity of the trigeminal ganglion to cerebral arteries may increase these patients' susceptibility to large vessel stroke.¹¹ However, many EHR studies do not specify zoster site, which is likely to lead to underestimation of effect because some HZ at unclassified sites could actually be HZO. The stronger effect for HZO was noted in two self-controlled case series studies which stratified by HZ site^{12,13} as well as two Asian cohort studies^{20,21}. A further study found no effect of HZ site on long-term stroke risk, although few participants had HZO records²².

3) Stroke risk may be highest for younger people with HZ

Intriguing evidence suggests a more marked effect in younger people. Point estimates for the effect of HZ on stroke are highest for younger people, aged less than 40 years, although the relatively small numbers of strokes in this age group mean that confidence intervals tend to overlap with those for

older ages. Of five cohort studies which stratified by age, four found a greater effect in younger age groups^{16,17,22,23} and one study showed no difference in effect size by age²⁰.

4) HZ can trigger myocardial infarction (MI)

One self-controlled case series study using US Medicare data¹² showed a similar though smaller transient triggering effect of HZ on MI – IR 1.68 (95% CI 1.47 – 1.92) – for week one after HZ. An elevated risk of acute coronary syndrome (ACS) or MI up to 3 months after HZ was also suggested by combined results from two prospective population cohort studies from the US¹⁸ and Taiwan²⁴: pooled OR 1.34 (95% 0.98 to 1.82)¹¹, although this just failed to reach statistical significance in a random effects model. The Taiwanese study also showed a small increase in ACS risk associated with HZ in follow up to 12 years (HR 1.10 (95% CI 1.02 to 1.19)²⁴, which was mirrored in a cohort using UK primary care data with follow up to 24 years²².

Table 1

Effect of HZ prevention on vascular complications

It seems plausible that reducing HZ incidence and severity through vaccination is likely to result in fewer vascular complications. A single dose of the live attenuated zoster vaccine (Zostavax) reduces herpes zoster incidence by around 50%.²⁵ However, at present evidence of direct cardiovascular benefit from the live attenuated zoster vaccine is lacking: in the Shingles Prevention Study, rates of serious adverse events including MI and stroke, were the same in recipients of vaccine and placebo but follow up was short (to 42 days) and the trial was not powered for these endpoints²⁵.

Observational safety studies have also shown no effect of Zostavax on stroke risk in follow up to 5-6 weeks^{26,27}, although longer-term follow up is needed to assess fully the vaccine's effectiveness against vascular outcomes. The recent development of the highly efficacious subunit vaccine HZ/su (Shingrix), which reduces HZ incidence by 90% after two doses in older adults²⁸ has potential for major impact on VZV disease burden, including vascular complications, in settings achieving good

vaccine uptake. A recent modelling study based on RCT data showed that Shingrix, at a price of \$280 for two doses, was more cost effective than Zostavax (current price \$213 per dose).²⁹

Effect of HZ treatment on vascular complications

It remains uncertain whether antiviral agents used to treat HZ alter the risk of vascular events: results from one cohort study¹⁹ and one self-controlled case series study¹³ suggest that patients receiving antivirals may have a lower stroke risk than patients not taking antivirals, especially earlier after HZ diagnosis. Another cohort study, however, found no difference in stroke risk at one year between those who did and did not receive antivirals for HZO²¹, although data on timing and duration of treatment were unavailable in this study and it was difficult to exclude confounding by indication – the situation in which patients who were more unwell and had a higher risk of vascular complications preferentially received antiviral treatment.

Conclusions

In future, our ability to prevent HZ-induced vascular events will depend upon improving understanding of the nature of risk including any effects of sub-clinical VZV reactivation or long-term effects. Large-scale population-based studies to identify the characteristics of affected patients will help to guide intervention targeting. Adequately powered studies of HZ prevention and treatment strategies against specific vascular endpoints are needed, which are carefully designed to avoid selection biases and confounding by indication. As the evidence base develops, it will be important to (re-) assess the cost effectiveness of policy options such as introducing universal varicella vaccination in countries where it is not routine, for example in Northern Europe, or expanding recommendations for zoster vaccination to include younger age groups at high cardiovascular risk.

Funding

CWG is supported by a Wellcome Intermediate Clinical Fellowship (201440/Z/16/Z).

Conflict of interest

None declared.

References

1. Warren-Gash C, Smeeth L, Hayward AC. Influenza as a trigger for acute myocardial infarction or death from cardiovascular disease: a systematic review. *Lancet Infect Dis.* **2009**; 9(10):601–10.
2. Kwok CS, Aslam S, Kontopantelis E, et al. Influenza, influenza-like symptoms and their association with cardiovascular risks: a systematic review and meta-analysis of observational studies. *Int J Clin Pract* **2015**; 69(9):928–37.
3. Corrales-Medina VF, Madjid M, Musher DM. Role of acute infection in triggering acute coronary syndromes. *Lancet Infect Dis.* **2010**; 10(2):83–92.
4. Gilden D, Cohrs RJ, Mahalingam R, Nagel MA. Varicella zoster virus vasculopathies: diverse clinical manifestations, laboratory features, pathogenesis, and treatment. *Lancet Neurol.* **2009**; 8(8):731–740.
5. Feigin VL, Roth GA, Naghavi M, et al. Global burden of stroke and risk factors in 188 countries, during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet Neurol.* **2016**; 15(9):913–24.
6. Yang S-Y, Li H-X, Yi X-H, et al. Risk of Stroke in Patients with Herpes Zoster: A Systematic Review and Meta-Analysis. *J Stroke Cerebrovasc Dis.* **2017**; 26(2):301–7.
7. Liu X, Guan Y, Hou L, et al. The Short- and Long-Term Risk of Stroke after Herpes Zoster: A Meta-Analysis. *PloS One.* **2016**; 11(10):e0165203.
8. Zhang Y, Luo G, Huang Y, Yu Q, Wang L, Li K. Risk of Stroke/Transient Ischemic Attack or Myocardial Infarction with Herpes Zoster: A Systematic Review and Meta-Analysis. *J Stroke Cerebrovasc Dis.* **2017**; 26(8):1807–16.
9. Marra F, Ruckenstein J, Richardson K. A meta-analysis of stroke risk following herpes zoster infection. *BMC Infect Dis.* **2017**; 17(1):198.
10. Lian Y, Zhu Y, Tang F, Yang B, Duan R. Herpes zoster and the risk of ischemic and hemorrhagic stroke: A systematic review and meta-analysis. *PloS One.* **2017**; 12(2):e0171182.
11. Erskine N, Tran H, Levin L, et al. A systematic review and meta-analysis on herpes zoster and the risk of cardiac and cerebrovascular events. *PloS One.* **2017**; 12(7):e0181565.
12. Minassian C, Thomas SL, Smeeth L, Douglas I, Brauer R, Langan SM. Acute Cardiovascular Events after Herpes Zoster: A Self-Controlled Case Series Analysis in Vaccinated and Unvaccinated Older Residents of the United States. *PLoS Med.* **2015**; 12(12):e1001919.
13. Langan SM, Minassian C, Smeeth L, Thomas SL. Risk of Stroke Following Herpes Zoster: A Self-Controlled Case-Series Study. *Clin Infect Dis.* **2014**; 58(11):1497–503.

14. Schink T, Behr S, Thöne K, Bricout H, Garbe E. Risk of Stroke after Herpes Zoster - Evidence from a German Self-Controlled Case-Series Study. *PloS One*. **2016**; 11(11):e0166554.
15. Whitaker HJ, Hocine MN, Farrington CP. The methodology of self-controlled case series studies. *Stat Methods Med Res*. **2009**; 18(1):7–26.
16. Sreenivasan N, Basit S, Wohlfahrt J, et al. The Short-and Long-Term Risk of Stroke after Herpes Zoster-A Nationwide Population-Based Cohort Study. *PloS One*. **2013**; 8(7):e69156.
17. Sundström K, Weibull CE, Söderberg-Löfdal K, Bergström T, Sparén P, Arnheim-Dahlström L. Incidence of herpes zoster and associated events including stroke—a population-based cohort study. *BMC Infect Dis*. **2015**; 15(1):488.
18. Yawn BP, Wollan PC, Nagel MA, Gilden D. Risk of Stroke and Myocardial Infarction After Herpes Zoster in Older Adults in a US Community Population. *Mayo Clin Proc*. **2016**; 91(1):33–44.
19. Calabrese LH, Xie F, Yun H, et al. Herpes Zoster and the Risk of Stroke in Patients With Autoimmune Diseases. *Arthritis Rheumatol Hoboken NJ*. **2017**; 69(2):439–46.
20. Kang J-H, Ho J-D, Chen Y-H, Lin H-C. Increased Risk of Stroke After a Herpes Zoster Attack A Population-Based Follow-Up Study. *Stroke*. **2009**; 40(11):3443–8.
21. Lin H-C, Chien C-W, Ho J-D. Herpes zoster ophthalmicus and the risk of stroke: a population-based follow-up study. *Neurology*. **2010**; 74(10):792–7.
22. Breuer J, Pacou M, Gauthier A, Brown MM. Herpes zoster as a risk factor for stroke and TIA A retrospective cohort study in the UK. *Neurology*. **2014**; 82(3):206–212.
23. Kwon SU, Yun S-C, Kim M-C, et al. Risk of stroke and transient ischaemic attack after herpes zoster. *Clin Microbiol Infect* . **2016**; 22(6):542–8.
24. Wang C-C, Lin C-L, Chang Y-J, Wang G-J, Sung F-C, Kao C-H. Herpes zoster infection associated with acute coronary syndrome: a population-based retrospective cohort study. *Br J Dermatol*. **2014**; 170(5):1122–9.
25. Oxman MN, Levin MJ, Johnson GR, et al. A vaccine to prevent herpes zoster and postherpetic neuralgia in older adults. *N Engl J Med*. **2005**; 352(22):2271–84.
26. Baxter R, Tran TN, Hansen J, et al. Safety of Zostavax™--a cohort study in a managed care organization. *Vaccine*. **2012**; 30(47):6636–41.
27. Tseng HF, Liu A, Sy L, et al. Safety of zoster vaccine in adults from a large managed-care cohort: a Vaccine Safety Datalink study. *J Intern Med*. **2012**; 271(5):510–20.
28. Cunningham AL, Lal H, Kovac M, et al. Efficacy of the Herpes Zoster Subunit Vaccine in Adults 70 Years of Age or Older. *N Engl J Med*. **2016**; 375(11):1019–32.
29. Le P, Rothberg ML. Cost-effectiveness of the adjuvanted herpes zoster subunit vaccine in older adults. *JAMA Intern Med*. **2018**; 178(2):248-258.