



Prochlorococcus as a Possible Source for Transparent Exopolymer Particles (TEP)

Francesca Iuculano¹, Ignacio P. Mazuecos², Isabel Reche² and Susana Agustí^{1,3*}

¹ Department of Global Change, Instituto Mediterráneo de Estudios Avanzados, Consejo Superior de Investigaciones Científicas – Universitat de les Illes Balears, Esporles, Spain, ² Departamento de Ecología, Instituto del Agua, Universidad de Granada, Granada, Spain, ³ Red Sea Research Center, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

OPEN ACCESS

Edited by:

Jonathan P. Zehr,
University of California, Santa Cruz,
USA

Reviewed by:

Lisa Moore,
University of Southern Maine, USA
Paul Berube,
Massachusetts Institute
of Technology, USA

*Correspondence:

Susana Agustí
susana.agusti@kaust.edu.sa

Specialty section:

This article was submitted to
Aquatic Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 19 August 2016

Accepted: 06 April 2017

Published: 26 April 2017

Citation:

Iuculano F, Mazuecos IP, Reche I and
Agustí S (2017) *Prochlorococcus* as
a Possible Source for Transparent
Exopolymer Particles (TEP).
Front. Microbiol. 8:709.
doi: 10.3389/fmicb.2017.00709

Transparent exopolymer particles (TEP), usually associated with phytoplankton blooms, promote the formation of marine aggregates. Their exportation to deep waters is considered a key component of the biological carbon pump. Here, we explored the role of solar radiation and picocyanobacteria in the formation of TEP in oligotrophic surface waters of the Atlantic and Pacific Oceans in ten on-deck incubation experiments during the Malaspina 2010 Expedition. TEP concentrations were low on the ocean's surface although these concentrations were significantly higher on the surface of the Pacific ($24.45 \pm 2.3 \mu\text{g XG Eq. L}^{-1}$) than on the surface of the Atlantic Ocean ($8.18 \pm 4.56 \mu\text{g XG Eq. L}^{-1}$). Solar radiation induced a significant production of TEP in the on-deck experiments from the surface water of the Pacific Ocean, reaching values up to $187.3 \mu\text{g XG Eq. L}^{-1}$ compared with the low production observed in the dark controls. By contrast, TEP production in the Atlantic Ocean experiments was lower, and its formation was not related to the light treatments. *Prochlorococcus* sp. from the surface ocean was very sensitive to solar radiation and experienced a high cell decay in the Pacific Ocean experiments. TEP production in the on-deck incubation experiments was closely related to the observed cell decay rates of *Prochlorococcus* sp., suggesting that this picocyanobacteria genus is a potential source of TEP. The evidence to propose such potential role was derived experimentally, using natural communities including the presence of several species and a variety of processes. Laboratory experiments with cultures of a non-axenic strain of *Prochlorococcus marinus* were then used to test TEP production by this genus. TEP concentrations in the culture increased with increasing cell abundance during the exponential phase, reaching the highest TEP concentration at the beginning of the stationary phase. The average TEP concentration of $1474 \pm 226 \mu\text{g XG Eq. L}^{-1}$ (mean \pm SE) observed at the stationary phase of *P. marinus* cultures is comparable with the values reported in the literature for diatom cultures, also growing in non-axenic as well as axenic cultures. Our results identify *Prochlorococcus* sp. as a possible relevant source of TEP in the oligotrophic ocean.

Keywords: transparent exopolymer particles, *Prochlorococcus*, Pacific Ocean, Atlantic Ocean, UVR, solar radiation

INTRODUCTION

Transparent exopolymer particles (TEP) are organic particles composed mainly of acidic polysaccharides (Passow and Alldredge, 1994), and recognized as a key component of the biological carbon pump in the ocean (Engel et al., 2004b; Riebesell et al., 2007; Passow and Carlson, 2012). TEP form marine aggregates that promote the sedimentation of particles and export carbon to deep waters (Passow, 2002a). Their concentration and distribution in the ocean appear to be related to abiotic aggregation from their dissolved or colloidal exopolymeric precursors (Passow, 2000) and biotic processes via exudation from phytoplankton (Kiørboe and Hansen, 1993) and bacterioplankton (Alldredge et al., 1993; Passow, 2002b; Ortega-Retuerta et al., 2010). Formation of TEP has previously been related to phytoplankton abundance and composition (Passow and Alldredge, 1994), with blooms of diatoms (Mari, 1999) and coccolithophorids (Engel et al., 2004a; Van Oostende et al., 2013) the most studied sources. Cyanobacteria blooms have also been reported as sources of TEP (Berman-Frank et al., 2007). However, the role played by picocyanobacteria, the most abundant and ubiquitous primary producers (Partensky et al., 1999) in the open ocean, as a source for TEP remains unclear.

Nutrient limitation and other stresses have been identified as the main drivers of the exudation of dissolved carbohydrates by primary producers (Berman-Frank et al., 2007). Recently, Ortega-Retuerta et al. (2009) demonstrated that ultraviolet B (UVB) radiation stimulated TEP production in the presence of microorganisms, observing a 17% increase in TEP under +UVB treatment in comparison with dark controls. Orellana and Verdugo (2003) reported results in agreement with those of Ortega-Retuerta et al. (2009). However, there are few estimates of TEP distribution in the oligotrophic ocean, although some studies report low concentrations of TEP (Passow, 2000; Ortega-Retuerta et al., 2010). The oligotrophic ocean is characterized by transparent and relatively warm waters and low nutrient concentrations mostly dominated by picophytoplankton (Agawin et al., 2000), including *Prochlorococcus* sp. and *Synechococcus* sp. (Partensky et al., 1999). Oligotrophic waters are exposed to high solar radiation, because of the high transparency of this water (Tedetti and Sempéré, 2006) as well as their location in most subtropical and tropical regions, which attract maximal incident ultraviolet (UV) radiation. Therefore, in the oligotrophic ocean, photo-processes are likely to significantly influence TEP formation (Ortega-Retuerta et al., 2009). Moreover, it is well documented that solar radiation harms picophytoplanktonic cells by inducing cell mortality (Llabrés and Agustí, 2006; Agustí and Llabrés, 2007). In fact, maximum production of TEP has been related to the decline of phytoplankton blooms and to their senescence phases (Passow, 2002b; Engel et al., 2004a). Berman-Frank et al. (2007) identified autocatalytic programmed cell death in *Trichodesmium* sp. as a process inducing TEP production.

In this study, we experimentally tested the hypotheses that solar radiation enhances the production of TEP in the surface of oceanic and oligotrophic waters. We gathered data to test these hypotheses by performing ten on-deck experiments of oligotrophic surface waters from the North Atlantic and

inter-equatorial Pacific Oceans during the Malaspina 2010 Expedition. We exposed surface waters to different light treatments, including dark controls, full solar radiation, and ultraviolet radiation removed (-UV) to observe differences in TEP production. We simultaneously analyzed the sensitivity of picocyanobacteria and heterotrophic bacteria populations to solar radiation. The results obtained with natural oligotrophic communities suggested a potential role for *Prochlorococcus* as a source of TEP. We tested this potential role in the laboratory using a non-axenic culture of *Prochlorococcus marinus*, where reduced sources of variability relative to field experiments allow test of the inference pointing at *Prochlorococcus* as a source of TEP. We then compared the TEP concentration values observed in the *P. marinus* culture with those described in the literature for phytoplankton species reported to produce TEP growing in non-axenic and axenic cultures.

MATERIALS AND METHODS

Sampling Site and Study Area

We conducted 10 on-deck incubation experiments (Table 1) on board the R/V Hesperides during the Malaspina 2010 Expedition¹. The first set of experiments (#1–4) was performed in the Pacific Ocean during the expedition's leg from Auckland, New Zealand to Honolulu, U.S. during April–May 2011. The second set of experiments (#5–10) was performed in the Atlantic Ocean during the expedition's leg from Cartagena de Indias, Colombia to Cartagena, Spain during June–July 2011 (Figure 1).

Experimental Setup

Water samples were collected at 3 m depth in 30 L Niskin bottles. The water was filtered through a 50 μm mesh to remove the largest fraction of grazers. Water samples were dispensed into duplicate 2 L bottles for each treatment: quartz flasks, which allowed the full solar light spectrum (FULL treatment) to pass through, polycarbonate Nalgene bottles, which removed UVB and most UVA, and between 35% of transmittance at 399 nm and 86% of transmittance at 419 nm, allowing visible wavelengths to pass through (-UV treatment), and black bottles, which blocked all solar radiation (DARK treatment). The bottles were incubated in an open-air tank ($V = 1.9 \times 10^3$ L) and maintained at *in situ* temperature by using a continuous surface water running system. The experiments started after sunrise, at about 07:00 h local time (initial time T_0), and lasted for approximately 24 h (before dawn, final time T_f). The sampling time at T_0 and T_f were used for Chl *a* and TEP analysis (except for Exp 4 during which extra sampling at T_3 in the FULL treatment was also performed). Intermediate samplings coupled with the photoperiod were also performed to study picoplankton abundances: a first sampling (T_1) after 4 h of sunlight exposure, a second sampling (T_2) after 6 to 8 h, a third sampling at sunset (T_3) and a fourth sampling (T_4) 1 h after sunset.

¹<http://www.expedicionmalaspina.es>

TABLE 1 | Biogeographical provinces, geographical coordinates, biogeochemical conditions, and *in situ* incident solar radiation doses for each experiment.

Exp.	Longhurst province (lat-long)	T (°C)	Salinity	Chl a (µg L ⁻¹)	Proch (cells mL ⁻¹)	Syn (cells mL ⁻¹)	HBA (cells mL ⁻¹)	UVB (kJ m ⁻²)	UVA+B (kJ m ⁻²)	Global radiation (kJ m ⁻²)
1	PEQD 7.06° 171.39°W	28.91	35.48	0.14	3.24E+04	4.05E+03	6.84E+05	39	1276	19696
2	PEQD 3.41°S 169.46°W	28.56	35.36	0.23	8.54E+04	2.90E+03	5.54E+05	44.6	1507	16416
3	PEQD 1.60°S 166.85°W	27.65	35.08	0.31	7.30E+04	2.52E+03	1.45E+06	45.1	1569	26591
4	PEQD 6.99°N 164.37°W	27.74	34.81	0.3	6.58E+04	5.36E+03	1.00E+06	46	1539	24639
5	CARB 14.16°N 71.67°W	29.22	35.56	0.13	1.54E+04	5.06E+03	2.86E+05	23.9	1348	21252
6	NATR 17.43°N 59.83°W	29.43	35.52	0.14	3.68E+04	2.07E+03	6.72E+05	24.5	1334	21000
7	NATR 19.00°N 55.15°W	28.51	36.61	0.19	2.49E+04	6.82E+03	2.31E+05	23.7	1311	22558
8	NATR 24.84°N 38.71°W	25.65	37.57	0.07	2.04E+04	4.85E+03	6.11E+05	36.7	1663	27105
9	NASE 27.97°N 29.65°W	23.32	37.32	0.05	1.01E+04	2.14E+03	5.32E+05	34.3	1431	22173
10	NASE 32.08°N 17.27°W	21.81	36.7	0.09	4.29E+04	3.39E+03	7.32E+00	40.4	1660	28909

Exp, number of experiment; T, *in situ* seawater surface temperature (at 3 m depth); Chl a, Chlorophyll a concentration (at initial incubation time). Proch, Syn and HBA represent the cell abundance of *Prochlorococcus* sp., *Synechococcus* sp. and heterotrophic bacteria in seawater, respectively, at the initial incubation time. Incident UV radiation data are expressed as total doses received per day. PEQD, Trades-Pacific Equatorial Divergence; PNAC, Trades-North Pacific Equatorial Countercurrent Province; CARB, Caribbean Province; NATR, Trades-North Atlantic Tropical Gyral Province; NASE, North Atlantic Subtropical Gyral Province (Longhurst, 1998).

In experiments 4, 5, and 6 an additional treatment to control for potential abiotic TEP production, due to spontaneous assemblage from dissolved acidic polysaccharide precursors was included by incubating seawater filtered by gravity through a 0.2 µm cartridge (to eliminate phytoplankton and bacterial cells) into duplicate 2 L quartz bottles.

Solar Radiation

Incident solar radiation (W/m²) was measured every 2 min by a Radiation Sensor (2770; Aanderaa Instruments) installed in the meteorological station of the vessel (SADO) placed over the deck. The sensor is sensitive in the range of 300–2500 nm, but is covered by a glass (borosilicate) dome, filtering out most of the incident UVR. To measure solar radiation in the ultraviolet bands, we used an incident UVA (315–400 nm) and UVB (280–315 nm) radiometer (UVS-AB-T; Kipp & Zonen, Version T2.07.217) also installed in the vessel’s meteorological station. Doses of solar radiation received during the experiments were then calculated by integrating radiation values for the day’s duration and expressed in kJ/m².


Transparent Exopolymer Particles

Transparent Exopolymer Particles were stained with alcian blue, a group of basic, polyvalent and water-soluble dyes, derived from phthalocyanine. In a solution of 3% acetic acid (pH = 2.5) alcian blue stained carboxylated and ester-sulfate groups of acidic mucopolysaccharides. TEP were retained on polycarbonate filters (Poretics) with 0.4 µm pore size and of 25 mm in diameter and stained with 1 mL working solution of alcian blue pre-filtered through a 0.2 µm filter to a final concentration of 0.02%. TEP concentrations were determined spectrophotometrically following the colorimetric method proposed by Passow and Alldredge (1995). For each treatment of the experiments, 400 to 1000 mL volumes of the sample (depending on the saturation of the filter) were collected at the initial (T₀) and final (T_f) times (in Exp 4 also at T₃). Duplicates were filtered under low and constant pressure (150 mmHg). Filters were subsequently stained with 1000 µL of a 0.02% working solution of Alcian blue (pre-filtered through a 0.2 µm filter) in 0.06% acetic acid (pH = 2.5), left for few seconds, filtered again and rinsed twice with MilliQ water to eliminate excess dye. The filters were frozen at –80°C and stored until further processing in the laboratory. The filters were then thawed, soaked in 80% sulfuric acid for 2–3 h, and the absorbances were measured by means of a spectrophotometer (PerkinElmer UV-VIS Lambda 35) at a fixed wavelength of 787 nm. Empty stained filters (stained and frozen in parallel with the samples) were used as blanks.

Alcian blue absorption was calibrated using a Xanthan Gum (XG) solution that was homogenized by a tissue grinder and measured by weight. TEP concentrations were, therefore, expressed in µg of XG equivalents per liter and calculated as follows:

$$TEP (\mu\text{g XG Eq. L}^{-1}) = (a_{\text{sample}} - a_{\text{blank}})V^{-1} \times F, \quad (1)$$

where *a*_{sample} is the absorbance of the sample, *a*_{blank} is the absorbance of the blank, *V* is the filtered volume of the sample



and F is the calibration factor, which is the inverse of the slope between μg of XG and alcian blue absorption at 787 nm. The detection limit of the method was $1 \mu\text{g}$ of XG equivalents L^{-1} and the coefficient of variation was ca. 20%.

The TEP production rate per day (α) was calculated as the natural logarithm of the ratio between the concentrations (μg XG Eq. L^{-1}) observed at the final and initial experimental times ($t = 1$ day):

$$\alpha \text{ (d}^{-1}\text{)} = \ln(\text{TEP } t_f / \text{TEP } t_0) / t, \quad (2)$$

where TEP t_f and t_0 are the TEP concentrations at the final and initial times, respectively.

We also calculated the normalized (Passow, 2012) or net (Egge et al., 2009; MacGilchrist et al., 2014) TEP production (TEP_{PR}) expressed as μg XG Eq. $\text{L}^{-1} \text{d}^{-1}$ and calculated from the difference between the TEP concentration at the end of the experiment (TEP t_f) minus the initial TEP concentration (TEP t_0)

divided by the experimental time:

$$\text{Net TEP}_{\text{PR}} \text{ (}\mu\text{g XG Eq. L}^{-1} \text{d}^{-1}\text{)} = \frac{(\text{TEP } t_f - \text{TEP } t_0)}{(t_f - t_0)}. \quad (3)$$

Chlorophyll a

The concentration of chlorophyll a (Chl a) was quantified by using the fluorimetric method described by Parsons et al. (1984). Seawater samples (250 mL) were collected at the initial (T_0) and final (T_f) time of each experiment and filtered through 25 mm Whatman GF/F filters. Chlorophyll a in the filters was extracted in 90% acetone and left 24 h in the dark at 4°C until processing in a Turner Design fluorimeter, calibrated with pure chlorophyll a .

Picoplankton Abundance

Picoplankton populations were quantified by using flow cytometry.

To quantify the abundance of heterotrophic prokaryotes, we fixed 1.2 mL of each sample treatment from all sampling times ($T_0, T_1, T_2, T_3, T_4, T_f$) with 48 μL of glutaraldehyde (final concentration 1%). The samples were left for 10 min in the dark and then were frozen in liquid nitrogen and stored at -80°C until analysis in a FACScalibur (Becton Dickinson) flow cytometer equipped with a 488 nm argon laser. The samples were defrosted and analyzed following Marie et al. (1997) by staining 400 μL with 4 μL of $10 \times$ SYBR Green I (Molecular probe S-7563) solution (final dilution 1:1000 [vol/vol]) for 10 min and letting the stained samples run through the FACScalibur at low speed (ca. 25 $\mu\text{L min}^{-1}$) and adding 10 μL of a fluorescent 1 μm latex beads solution as an internal standard for cytometric counts. The run terminated when about 10,000 particles had been recorded or after 90 s. The threshold was set in the green channel. Stained bacteria were detected and discriminated from other non-bacterial particles with a

TABLE 2 | Growth rate (d^{-1} , mean \pm SE), maximum and mean (\pm SE) of concentration and specific production rates (d^{-1} , mean \pm SE) of transparent exopolymer particles (TEP) observed in cultures of *Prochlorococcus marinus*.

	<i>P. marinus</i> Culture 1	<i>P. marinus</i> Culture 2
Maximum TEP (μg XG Eq. L^{-1})	2500	1514
<i>Growth phase (days 11–40)</i>		
TEP (μg XG Eq. L^{-1})	295 \pm 39	151 \pm 19
TEP (d^{-1})	0.048 \pm 0.008	0.040 \pm 0.05
Growth rate (d^{-1})	0.4 \pm 0.02	0.23 \pm 0.06
<i>Stationary phase (days 40–56)</i>		
TEP (μg XG Eq. L^{-1})	1647 \pm 333	1185 \pm 208
TEP (d^{-1})	0.055 \pm 0.008	0.038 \pm 0.003
Growth rate (d^{-1})	−0.10 \pm 0.05	−0.07 \pm 0.1

light side scatter (SSC) and green (FL1) and red fluorescence (FL3).

To measure the abundance of phototrophic prokaryotes (*Prochlorococcus* and *Synechococcus*), we ran 400 μL of each sample treatment from all sampling times (T_0 , T_1 , T_2 , T_3 , T_4 , T_f) at medium speed (ca. 55 $\mu\text{L min}^{-1}$) in the flow cytometer. A sample (10 μL) of a calibrated solution of 1- μm diameter yellow-green fluorescent latex beads (Polysciences) was added as an internal standard for the quantification of cell concentrations. Beads concentrations [$9.33 \times 10^5 \text{ mL}^{-1}$ for the Pacific Ocean experiments] [$4.63 \times 10^5 \text{ mL}^{-1}$ for the Atlantic Ocean experiments] in the standard solution were calculated by filtering duplicate aliquots onto black Nucleopore filters and counting under an epifluorescence microscope. All samples were fixed with glutaraldehyde (1% final concentration) and stored at -80°C until their analysis in the laboratory. *Synechococcus* populations were detected by an orange fluorescence dot plot (FL2 = interval between 585 \pm 20 nm of wavelength, corresponding to the fluorescence of the phycoerythrin pigment) versus red fluorescence (FL3 \geq 635 nm, corresponding to the fluorescence of chlorophyll). *Prochlorococcus* had a lower FL3 signal and no FL2 signal. Moreover, the populations were detected in a SSC versus FL3 dot plot (Marie et al., 1997). Decay rates and growth rates (h^{-1}) of *Synechococcus*, *Prochlorococcus*, and heterotrophic bacterial populations were then calculated as the slope of the linear regression between the natural logarithms of the concentrations (cells mL^{-1}) and the experimental time (hours) during each sampling interval.

Prochlorococcus marinus Batch Cultures

To corroborate that *Prochlorococcus* could be a source of TEP we set up two non-axenic cultures of a strain of *P. marinus* (RCC 0156). This strain was grown in the laboratory under one-liter batch aseptic conditions with PCR-11 culture medium at 22°C under low light of 30 $\mu\text{mol Photons m}^{-2} \text{ s}^{-1}$. We did not include any UV light incubation due to the difficulty and limitations of keeping this species growing in the laboratory. The cultures were sampled regularly during 2 months to monitor changes in cell abundance and TEP generation during the exponential and stationary phases. Triplicated volumes of 20 mL for TEP quantification were sampled by syringe, gently filtered under low vacuum pressure and immediately analyzed as previously described (Passow and Alldredge, 1995). *P. marinus* cell abundance (cells mL^{-1}) was quantified in replicated 1 ml fresh samples in a BD FACScalibur flow cytometer.

Statistical Analyses

Statistical analyses of TEP and picoplankton data were carried out using JMP software, Student's *t*-test was used to test for significant differences between oceans, treatments (DARK, FULL, -UV) and other changes in parameters, and linear regression was used for the analysis of the relationship between TEP production and picoplankton cell decay.

RESULTS

Surface Ocean Conditions

Surface temperature and salinity data collected from the stations near the sampling areas indicated that the experiments were conducted in warm waters (Table 1), ranging from 29.4°C in the Caribbean Sea to 21.8°C at the most north-eastern station of the Atlantic Ocean (Figure 1); the most saline waters were located in the North Atlantic Ocean (Table 1).

The chlorophyll *a* concentration was low in all the biogeographic provinces studied, ranging from $0.05 \mu\text{g L}^{-1}$ in the Northern Atlantic to $0.31 \mu\text{g L}^{-1}$ in the equatorial Pacific Ocean.


Initial abundances of picoplankton were generally higher in the waters sampled in the Pacific Ocean than in the waters sampled in the Atlantic Ocean. Heterotrophic bacterial concentrations were also higher in the Pacific Ocean (mean $9.23 \pm 2.00 \times 10^5 \text{ cells mL}^{-1}$) than in the Atlantic Ocean (mean $3.96 \pm 1.39 \times 10^5 \text{ cells mL}^{-1}$), reaching maximum abundance at the Equatorial station ($1.45 \pm 0.43 \times 10^6 \text{ cells mL}^{-1}$). *Prochlorococcus* sp. was more abundant (mean Pacific $6.42 \pm 0.38 \times 10^4 \text{ cells mL}^{-1}$; mean Atlantic $2.51 \pm 0.17 \times 10^4 \text{ cells mL}^{-1}$) than *Synechococcus* sp. (mean Pacific $3.71 \pm 0.50 \times 10^3 \text{ cells mL}^{-1}$; mean Atlantic $4.05 \pm 0.33 \times 10^3 \text{ cells mL}^{-1}$).

UVR conditions measured during the experiments in the Pacific Ocean were slightly higher at the northern latitudes and were the highest during Exp 4 (Table 1). During the Atlantic Ocean experiments, UVR doses were slightly higher at the northern latitudes, in the North Atlantic Gyre, than in the Caribbean experiments but in all the cases lower than during the Pacific Ocean experiments (Table 1). UVB daily doses measured during the Pacific Ocean experiments (mean $43.6 \pm 3.1 \text{ KJ m}^{-2}$) were significantly higher than those measured during the Atlantic Ocean experiments (mean $30.5 \pm 2.5 \text{ KJ m}^{-2}$; $t = 13.09$, $\text{df} = 8$, $p < 0.005$).

Initial TEP concentrations in the Pacific Ocean were significantly higher (mean $24.45 \pm 2.3 \mu\text{g XG Eq. L}^{-1}$) than in the Atlantic Ocean (mean $8.18 \pm 4.56 \mu\text{g XG Eq. L}^{-1}$; *t*-test: $t = 16.26$, $\text{df} = 8$, $p < 0.05$) (Figure 2).

On-deck Solar Radiation Incubation Experiments

In all the experiments performed in the Pacific Ocean, we detected, after the 24 h of incubation, a significant increase in TEP concentrations under full solar radiation, but not in the DARK and -UV treatments (Figure 2A; *t*-test: Full-Initial $t = 81.05$, $\text{df} = 7$, $p < 0.05$). The highest concentration of TEP was consistently observed in the full solar radiation treatments, reaching values up to $187.3 \mu\text{g XG Eq. L}^{-1}$ in the Exp 3 (Figure 2A). The potential that TEP was produced by self-assembly due to abiotic processes was tested in Exp 4. When microorganisms were excluded from the waters, there was no detectable increase in TEP (Figure 2C). An additional sampling time at the end of the day in Exp 4 revealed that the production of TEP during the day increased to $60.2 \pm 8 \text{ XG Eq. L}^{-1}$, although no significantly higher than




the value observed after 24 h (Figure 2A, $47.4 \pm 3.83 \text{ XG Eq. L}^{-1}$).

In the experiments performed in the North Atlantic Ocean (Exps 5–10, Table 1) we also observed an increase in TEP production after the incubation period in most of the experiments, but, in contrast with the Pacific Ocean experiments, this increase in production was not consistently induced by solar radiation exposure (Figure 2B). Maximum concentrations of $26.9 \mu\text{g XG Eq. L}^{-1}$ were produced during Exp 7 (Figure 2B). In the experiments 8, 9, and 10, we observed higher concentrations in the full solar radiation treatments but these increases were not statistically significant (Figure 2B). In the two experiments (#5 and 6) in which an extra treatment excluding the microorganisms was performed, we observed contrasting results (Figure 2C). In Exp 5 significantly more TEP was produced in the $0.2 \mu\text{m}$ -filtrated treatment (Figure 2C) ($\text{ca. } 50 \mu\text{g XG Eq. L}^{-1}$) than in the full treatment including microorganisms (Figure 2B) ($\text{ca. } 10 \mu\text{g XG Eq. L}^{-1}$). This fact suggests the relevance of abiotic self-assembly, although no difference was observed in the case of the Exp 6 (Figure 2C).


Transparent Exopolymer Particles production rates were generally higher in the Pacific Ocean experiments, resulting in a mean net daily TEP production of $44.3 \pm 14.2 \mu\text{g XG Eq. L}^{-1}$, than in the Atlantic Ocean experiments, which averaged $6.6 \pm 1.1 \mu\text{g XG Eq. L}^{-1}$. Furthermore, the TEP production rates (d^{-1}) in the Pacific Ocean experiments were significantly higher under the FULL solar radiation treatment (Figure 3A). By contrast, TEP production in the Atlantic Ocean did not differ significantly among treatments (Figure 3B).

Consistently, in all the experiments performed in the Pacific Ocean, the populations of *Prochlorococcus* sp. were negatively affected by solar radiation ($p < 0.05$), resulting in the decay of the cell population over time (Figure 4), without significant differences in the decay rates between FULL (mean $-0.06 \pm 0.01 \text{ h}^{-1}$) and -UV treatments (mean $-0.07 \pm 0.009 \text{ h}^{-1}$), but with significantly higher cell decay rates in comparison with the DARK treatments (Figure 4; mean $-0.01 \pm 0.002 \text{ h}^{-1}$; t -test: DARK-FULL $t = 0.058$, $\text{df} = 7$, $p < 0.001$; t -test: DARK-UV $t = 0.048$, $\text{df} = 7$, $p < 0.005$). The most pronounced decay in cell concentrations during the sampling time was observed during Exp 2. Dark treatments provided a control for other sources of cell mortality than light, including grazing by small ciliates and protists (passing through a net of $50 \mu\text{m}$ mesh) present in the experimental communities. *Prochlorococcus* decay rates in the dark treatments in the Pacific experiments were low, significantly lower than those in the light treatments. However, in the Atlantic, the differences between dark and light treatments were not significant ($p > 0.05$) and may reflect losses due to other processes. In contrast to *Prochlorococcus*, the *Synechococcus* population exhibited low cell decay rates, which were not significantly different ($p > 0.05$) among treatments (Figure 5). Heterotrophic bacterial cell abundance did not decline significantly across the different treatments and experiments in both Pacific and Atlantic experiments. Overall, picoplankton populations from the Atlantic experiments did not exhibit significant decay rates under light exposure treatments except for Exp. 5, which showed



the highest decay rates (Figure 5). Cell decay of *Prochlorococcus* in the Pacific (mean $-0.047 \pm 0.009 h^{-1}$) was significantly higher than in the Atlantic Ocean (mean $-0.01 \pm 0.005 h^{-1}$; t -test: $t = 0.036$, $df = 19$, $p < 0.001$) and significantly higher than that of *Synechococcus* in the Pacific Ocean (mean $-0.004 \pm 0.002 h^{-1}$; t -test: $t = 0.043$, $df = 23$, $p < 0.001$). However, *Prochlorococcus* and *Synechococcus* decay rates were not significantly different in the Atlantic Ocean (Figure 5).


We did not obtain significant correlations between Chl *a* concentration and bacterioplankton, *Synechococcus* and TEP



concentrations or between Chl *a* concentration and TEP production rates (α). However, the TEP production rate (d^{-1}) increased significantly with increasing *Prochlorococcus* sp. decay rates ($R^2 = 0.41$, $p < 0.0001$). This relationship improved when only the Pacific Ocean experiments were considered (α TEP (d^{-1}) = $0.047 - 18.04 * \text{Prochlorococcus decay } (h^{-1})$, $R^2 = 0.60$, $p < 0.001$) in which the *Prochlorococcus* cell decay accounted for a larger percentage of the variability (Figure 6).

TEP Concentration and Production in *Prochlorococcus* Cultures

Prochlorococcus marinus in the cultures grew at a moderate rate requiring 40 days to reach the stationary phase (Figure 7). TEP concentrations varied highly, increasing with time as




P. marinus cell abundance increased (Figure 7A). The highest concentration of TEP was observed before reaching the stationary phase and remained high during the following days (Figure 7A and Table 2). TEP production rates in the cultures were low (Figure 7B) in comparison with the production rates in the field (Figure 3). There was no net production of TEP during the first 3 weeks of the culture although *P. marinus* was growing actively (Figure 7B). When growth rates of *P. marinus* declined, TEP production still remained at the highest values (Figure 7B). When *P. marinus* experienced negative growth the production of TEP stopped, although the TEP concentration remained high in the culture (Figure 7).


DISCUSSION

Our study showed consistent differences in TEP production on the surfaces of the oligotrophic Atlantic and Pacific Oceans. The results also suggest that the picocyanobacteria *Prochlorococcus* may contribute to the production of TEP in the open ocean, as we observed in the Pacific Ocean experiments under solar radiation stress. This TEP production might affect the exportation of carbon to the deep ocean in oceanic oligotrophic regions.

The values of TEP concentrations from surface waters reported in this study were low, in the range of the lowest reported in the literature for the oligotrophic Mediterranean Sea (Ortega-Retuerta et al., 2010). Passow (2002b) compared values of TEP across different areas and found the values in the North Atlantic Ocean were also lower than those reported in the Pacific Ocean. Wurl et al. (2011) reported higher TEP values for the surface of the North Pacific and tropical North



Pacific than those found in our study, although the chlorophyll *a* concentration was also higher in that study and a bloom of *Trichodesmium* sp. was reported. The main differences between the Pacific and the Atlantic Ocean sampling areas in this study were connected to the incident solar radiation doses in the areas. There were significantly higher UVB radiation doses reaching the Pacific Ocean than the Atlantic Ocean. Also, the phytoplankton biomass and the abundance of *Prochlorococcus* sp. were higher in the Pacific Ocean sampling areas than in the Atlantic Ocean sampling areas. The TEP-to-Chl *a* concentration ratio averaged 357.3 ± 126.6 and 78.6 ± 9.3 in the Pacific and Atlantic Oceans, respectively, and it was indeed significantly higher in the Pacific Ocean. Overall, we found that during the experiments conducted



in the Pacific Ocean, the production of TEP was associated with high-energy solar radiation exposure (both UV and visible wavelengths), suggesting a potential role of light stress in the production of these exopolymers. By contrast, in the experiments # 5, 6, and 7 of the Atlantic Ocean we did not observe significant differences between the dark and the full treatments, but TEP concentrations were higher than the initial conditions. In these experiments, abiotic self-assembling of exopolymers (Verdugo et al., 2004) or likely aggregation mediated by bacteria carbon processing (Stoderegger and Herndl, 1998; Rochelle-Newall et al., 2010) appear to occur independently of light conditions.

Our study is the first to identify *Prochlorococcus* as a possible source of TEP. Callieri et al. (2011) previously suggested this capacity for freshwater *Synechococcus*, resulting in cell aggregation and colonial formation under high ultraviolet radiation exposure. TEP is also produced by marine *Synechococcus* (Deng et al., 2016). The *P. marinus* cultures also suggest that this species is able to generate TEP directly

or indirectly through stimulus of associated bacteria. The TEP concentration increased from the growth phase to the stationary phase, similar to the concentration changes in other phytoplankton species grown in cultures (Passow, 2002a). Most literature data came from non-axenic cultures of phytoplankton rendering our *P. marinus* cultures results comparable. Maximum TEP concentrations were observed at the beginning of the stationary phase of *P. marinus* cultures and were in the same order of magnitude of those reported for other phytoplankton species, such as the diatoms *Chaetoceros affinis* and *Nitzschia angularis* (Passow, 2002b). Although other diatomic species are able to produce higher TEP concentrations (Passow, 2002a,b), other phytoplankton species, such as *Emiliania huxleyi* and *Tetraselmis suecica* produced much lower TEP concentrations, on the order of 740 and 880 $\mu\text{g XG Eq. L}^{-1}$, respectively (Passow, 2002a) than did those found in the *P. marinus* cultures. However, as discussed by Passow (2002b) about other phytoplankton cultures, we can not therefore disregard the role of bacteria in TEP production in our batch cultures. Although phytoplankton is considered the major source for the generation of TEP precursors (Passow, 2002a), heterotrophic bacteria were also identified to have an important role in the formation of TEP in the ocean (Stoderegger and Herndl, 1998; Ortega-Retuerta et al., 2010). Recent works by Gärdes et al. (2011) and Van Oostende et al. (2013) have reported that bacteria attached to diatoms or associated with the coccolithophore *E. huxleyi* are ultimately the promoters of sticky exopolymers that aggregate to larger particles. In fact, bacteria appear to uptake the carbohydrates released by phytoplankton cells producing more refractory and stickier TEP as dissolved organic carbon get older (Rochelle-Newall et al., 2010).

Our results relating TEP production with picocyanobacteria cell decay are in agreement with results from Berman-Frank et al. (2007) who documented that TEP production was associated with cell mortality of the cyanobacteria *Trichodesmium*. In their study, natural populations of *Trichodesmium erythraeum* from New Caledonia were exposed to artificial high irradiance to induce cell mortality (autocatalytic programmed cell death, PCD) and TEP formation was observed. The induced oxidative stress activated PCD and positively correlated with increased TEP production (Berman-Frank et al., 2007). In cultures of the same species, the formation of TEP increased during depleted iron conditions, associated with *T. erythraeum* PCD (Berman-Frank et al., 2007). In the North Adriatic Sea, phytoplankton cell lysis was related to extensive exudation of particulate polysaccharides (Baldi et al., 1997). The production of TEP *in situ* has been related to the decline of phytoplankton blooms and their senescence phases (Passow, 2002b; Engel et al., 2004a). Lysis of *Prochlorococcus* cells dying under high solar radiation may also contribute to the release of intracellular substances to the medium, as observed in oligotrophic systems (Agustí and Duarte, 2013) contributing to TEP production directly or indirectly through bacteria processing. Phosphorus nutrition may influence DOC excretion by *Prochlorococcus* (Bertilsson et al., 2005). It has also been reported that TEP production increased as growth rates decreased at the end of diatom or coccolithophorids blooms,

with consequences on settling cell aggregates and carbon export (Logan et al., 1995; Engel et al., 2004a). High concentrations of TEP have also been monitored during blooms dominated by *Phaeocystis* spp. (Riebesell et al., 1995), dinoflagellates (Alldredge et al., 1998), coccolithophorids (Engel et al., 2004a), and cyanobacteria in lakes (Grossart and Simon, 1997). However, studies addressing the relationship between TEP production and picophytoplankton in the open ocean have been lacking. Our results describe TEP production rates in the experiments in the Atlantic Ocean to be on the order of those reported by Ortega-Retuerta et al. (2010) in the Mediterranean Sea; however, TEP production rates in the experiments in the Pacific Ocean under intense solar radiation were higher. Solar radiation, including both PAR and UVR bands, induced significant mortality in *Prochlorococcus* sp. in the Pacific Ocean and one experiment in the Atlantic Ocean. *Prochlorococcus* sp. has been described as more sensitive to sunlight damage than *Synechococcus* and picoeukaryotes (Llabrés and Agustí, 2006; Agustí and Llabrés, 2007). Even a short exposure of 30 min to high solar radiation was able to induce cell mortality in *Prochlorococcus* sp. indicating its high sensitivity to UV exposure (Agustí and Llabrés, 2007). UVR also induced the lysis of dead *Prochlorococcus* cells (Llabrés et al., 2010). However, a review of available experiments in the Atlantic Ocean revealed large variability of *Prochlorococcus* sp. cell mortality rates under UVR, varying from non-detectable to high cell mortality rates (Agustí and Llabrés, 2007). The cell decay rates were significantly related to the health condition (i.e., percentage of living cells) of the original population (Agustí and Llabrés, 2007). *Prochlorococcus* sp. sensitivity to UVR would depend, moreover on incident UVR doses, on other factors as nutrients conditions, light history, associated bacteria, and differences in the ability of *Prochlorococcus* populations to use mechanisms to reduce oxidative stress or improve photosystem responses (Morris et al., 2011; Mella-Flores et al., 2012). Therefore, the relative importance of *Synechococcus* sp. in comparison to *Prochlorococcus* sp. in the Atlantic Ocean versus the Pacific Ocean (Table 1) could explain, to some extent, the contrasting results observed in the experiments performed in both oceans.

Our study is the first evidence of the association of TEP production with the picocyanobacteria *Prochlorococcus*. In the Pacific Ocean experiments, under high solar radiation, the production of TEP was related with *Prochlorococcus* cell decay. However, this was not a plausible process in the experiments performed in the Atlantic Ocean where both biotic and abiotic

processes contributed to the exopolymer particle dynamics. Our results, verified by direct assays in culture experiments, identify the picocyanobacteria *Prochlorococcus* as a possible source of TEP in the oligotrophic ocean. These new insights, despite its limitations, open new research questions arising from our starting hypothesis. Further field and laboratory experiments are required to deepen our understanding on the effect of high light and UVR, as well as on the relevance of *Prochlorococcus* cell death processes in the production of TEP. Moreover, the effect of other drivers already proven to be relevant to TEP dynamics could not be disregarded. Those drivers include abiotic (e.g., nutrient limitation, temperature, pH, aggregation) and biotic processes (e.g., exudation, viral, and grazing pressure) including the tricky on-going debate on the significance of bacteria in TEP production requiring further analysis of bacteria-*Prochlorococcus* interactions. *Prochlorococcus* is the dominant primary producer in the oligotrophic ocean stressing the need to deepen its potential role in the production of TEP.

AUTHOR CONTRIBUTIONS

SA, FI, and IR designed the study. FI run the experiments on board. FI and IM analyzed the samples. All authors contribute to writing the manuscript.

FUNDING

This research was funded by the Expedición Malaspina 2010 project of the CONSOLIDER program (ref. CSD2008-00077), the project Stress-X (ref. CTM2012-32603) of the Spanish Minister of Economy and Competitiveness, and King Abdullah University of Science and Technology baseline funding to SA. FI was supported by a CSIC JAE-preDOC fellowship.

ACKNOWLEDGMENTS

We are grateful to the chief of the project, C. M. Duarte, and to the Bio-Hesperides crew members and UTM personnel for assistance. We also thank the editor J. P. Zehr and two reviewers for their thoughtful comments and suggestions helping to improve the quality of the manuscript.

REFERENCES

- Agawin, N. S., Duarte, C., and Agustí, S. (2000). Nutrient and temperature control of the contribution of picoplankton to phytoplankton biomass and production. *Limnol. Oceanogr.* 45, 591–600. doi: 10.4319/lo.2000.45.3.0591
- Agustí, S., and Duarte, C. M. (2013). Phytoplankton lysis predicts dissolved organic carbon release in marine plankton communities. *Biogeosciences* 10, 1259–1264. doi: 10.5194/bg-10-1259-2013
- Agustí, S., and Llabrés, M. (2007). Solar radiation-induced mortality of marine pico-phytoplankton in the oligotrophic ocean. *Photochem. Photobiol.* 83, 793–801. doi: 10.1111/j.1751-1097.2007.00144.x
- Alldredge, A. L., Passow, U., and Haddock, H. D. (1998). The characteristics and transparent exopolymer particle (TEP) content of marine snow formed from thecate dinoflagellates. *J. Plankton Res.* 20, 393–406. doi: 10.1093/plankt/20.3.393
- Alldredge, A. L., Passow, U., and Logan, B. E. (1993). The abundance and significance of a class of large, transparent organic particles in the ocean. *Deep Sea Res. I* 40, 1131–1140. doi: 10.1016/0967-0637(93)90129-Q
- Baldi, F., Minacci, A., Saliot, A., Mejanelle, L., Mozetic, P., Turk, V., et al. (1997). Cell lysis and release of particulate polysaccharides in extensive marine mucilage assessed by lipid biomarkers and molecular probes. *Mar. Ecol. Prog. Ser.* 153, 45–57. doi: 10.3354/meps153045
- Berman-Frank, I., Rosenberg, G., Levitan, O., Haramaty, L., and Mari, X. (2007). Coupling between autocatalytic cell death and transparent exopolymeric particle production in the marine cyanobacterium *Trichodesmium*. *Environ. Microbiol.* 9, 1415–1422. doi: 10.1111/j.1462-2920.2007.01257.x

- Bertilsson, S., Berglund, O., Pullin, M. J., and Chisholm, S. (2005). Release of dissolved organic matter by *Prochlorococcus*. *Vie Milieu* 55, 225–232.
- Callieri, C., Lami, A., and Bertoni, R. (2011). Microcolony formation by single-cell *Synechococcus* strains as a fast response to UV radiation. *Appl. Environ. Microbiol.* 77, 7533–7540. doi: 10.1128/AEM.05392-11
- Deng, W., Cruz, B. N., and Neuer, S. (2016). Effects of nutrient limitation on cell growth, TEP production and aggregate formation of marine *Synechococcus*. *Aquat. Microb. Ecol.* 78, 39–49. doi: 10.3354/ame01803
- EGge, J. K., Thingstad, T. F., Larsen, A., Engel, A., Wohlers, J., Bellerby, R. G. J., et al. (2009). Primary production during nutrient-induced blooms at elevated CO₂ concentrations. *Biogeosciences* 6, 877–885. doi: 10.5194/bg-6-8772009
- Engel, A., Delille, B., Jacquet, S., Riebesell, U., Rochelle-Newall, E., Terbrüggen, A., et al. (2004a). Transparent exopolymer particles and dissolved organic carbon production by *Emiliania huxleyi* exposed to different CO₂ concentrations: a mesocosm experiment. *Aquat. Microb. Ecol.* 34, 93–104. doi: 10.3354/ame034093
- Engel, A., Thoms, S., Riebesell, U., Rochelle-Newall, E., and Zondervan, I. (2004b). Polysaccharide aggregation as a potential sink of marine dissolved organic carbon. *Nature* 428, 929–932. doi: 10.1038/nature02453
- Gärdes, A., Iversen, M. H., Grossart, H. P., Passow, U., and Ullrich, M. S. (2011). Diatom-associated bacteria are required for aggregation of *Thalassiosira weissflogii*. *ISME J.* 5, 436–445. doi: 10.1038/ismej.2010.145
- Grossart, H. P., and Simon, M. (1997). Formation of macroscopic organic aggregates (lake snow) in a large lake: the significance of transparent exopolymer particles, phytoplankton, and zooplankton. *Limnol. Oceanogr.* 42, 1651–1659. doi: 10.4319/lo.1997.42.8.1651
- Kiorboe, T., and Hansen, J. L. S. (1993). Phytoplankton aggregate formation: observations of patterns and mechanisms of cell sticking and the significance of exopolymeric material. *J. Plankton Res.* 15, 993–1018. doi: 10.1093/plankt/15.9.993
- Llabrés, M., and Agustí, S. (2006). Picophytoplankton cell death induced by UV radiation: evidence for oceanic Atlantic communities. *Limnol. Oceanogr.* 51, 21–29. doi: 10.4319/lo.2006.51.1.0021
- Llabrés, M., Agustí, S., Alonso-Laita, P., and Herndl, G. J. (2010). *Synechococcus* and *Prochlorococcus* cell death induced by UV radiation and the penetration of lethal UVR in the Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 399, 27–37. doi: 10.3354/meps08332
- Logan, B. E., Passow, U., Alldredge, A. L., Grossart, H. P., and Simon, M. (1995). Rapid formation and sedimentation of large aggregates is predictable from coagulation rates (half-lives) of transparent exopolymer particles (TEP). *Deep Sea Res. II* 42, 203–214. doi: 10.1016/0967-0645(95)00012-F
- Longhurst, A. (1998). *Ecological Geography of the Sea*. San Diego, CA: Academic Press.
- MacGilchrist, G. A., Shi, T., Tyrrell, T., Richier, S., Moore, C. M., Dumousseaud, C., et al. (2014). Effect of enhanced pCO₂ levels on the production of DOC and TEP in short-term bioassay experiments. *Biogeosciences* 11, 3695–3706. doi: 10.5194/bg-11-3695
- Mari, X. (1999). Carbon content and C:N ratio of transparent exopolymeric particles (TEP) produced by bubbling exudates of diatoms. *Mar. Ecol. Prog. Ser.* 183, 59–71. doi: 10.3354/meps183059
- Marie, D., Partensky, F., Jacquet, S., and Vaulot, D. (1997). Enumeration and cell cycle analysis of natural populations of marine picoplankton by flow cytometry using the nucleic acid stain SYBR green I. *Appl. Environ. Microbiol.* 63, 186–193.
- Mella-Flores, D., Six, C., Ratin, M., Partensky, F., Boutte, C., Le Corguillé, G., et al. (2012). *Prochlorococcus* and *Synechococcus* have evolved different adaptive mechanisms to cope with light and UV stress. *Front. Microbiol.* 3:285. doi: 10.3389/fmicb.2012.00285
- Morris, J. J., Johnson, Z. I., Szul, M. J., Keller, M., and Zinser, E. R. (2011). Dependence of the cyanobacterium *Prochlorococcus* on hydrogen peroxide scavenging microbes for growth at the ocean's surface. *PLoS ONE* 6:e16805. doi: 10.1371/journal.pone.0016805
- Orellana, M. V., and Verdugo, P. (2003). Ultraviolet radiation blocks the organic carbon exchange between the dissolved phase and the gel phase in the ocean. *Limnol. Oceanogr.* 48, 1618–1623. doi: 10.4319/lo.2003.48.4.1618
- Ortega-Retuerta, E., Duarte, C. M., and Reche, I. (2010). Significance of bacterial activity for the distribution and dynamics of transparent exopolymer particles in the Mediterranean Sea. *Microb. Ecol.* 59, 808–818. doi: 10.1007/s00248-010-9640-7
- Ortega-Retuerta, E., Passow, U., Duarte, C. M., and Reche, I. (2009). Effects of ultraviolet B radiation on (not so) transparent exopolymer particles. *Biogeosciences* 6, 3071–3080. doi: 0.5194/bg-6-3071-2009
- Parsons, T. R., Maita, Y., and Lalli, C. M. (1984). *A Manual of Chemical and Biological Methods for Seawater Analysis*. Pergamon: Pergamon Press.
- Partensky, F., Hess, W. R., and Vaulot, D. (1999). *Prochlorococcus*, a marine photosynthetic prokaryote of global significance. *Microbiol. Mol. Biol. Rev.* 63, 106–127.
- Passow, U. (2000). Formation of transparent exopolymer particles, TEP, from dissolved precursor material. *Mar. Ecol. Prog. Ser.* 192, 1–11. doi: 10.3354/meps192001
- Passow, U. (2002a). Transparent exopolymer particles (TEP) in aquatic environments. *Prog. Oceanogr.* 55, 287–333. doi: 10.1016/S0079-6611(02)00138-6
- Passow, U. (2002b). Production of transparent exopolymer particles (TEP) by phyto- and bacterioplankton. *Mar. Ecol. Prog. Ser.* 236, 1–12. doi: 10.3354/meps236001
- Passow, U. (2012). The abiotic formation of TEP under different ocean acidification scenario. *Mar. Chem.* 12, 72–80. doi: 10.1016/j.marchem.2011.10.004
- Passow, U., and Alldredge, A. L. (1994). Distribution, size and bacterial colonization of transparent exopolymer particles (TEP) in the ocean. *Mar. Ecol. Prog. Ser.* 113, 185–198. doi: 10.3354/meps113185
- Passow, U., and Alldredge, A. L. (1995). A dye-binding assay for the spectrophotometric measurement of transparent exopolymer particles (TEP). *Limnol. Oceanogr.* 40, 1326–1335. doi: 10.4319/lo.1995.40.7.1326
- Passow, U., and Carlson, C. (2012). The biological pump in a high CO₂ world. *Mar. Ecol. Prog. Ser.* 470, 249–271. doi: 10.3354/meps09985
- Riebesell, U., Reigstad, M., Wassmann, P., Noji, T., and Passow, U. (1995). On the trophic fate of *Phaeocystis pouchetii* (hariat): VI. Significance of *Phaeocystis*-derived mucus for vertical flux. *Neth. J. Sea Res.* 33, 193–203. doi: 10.1016/0077-7579(95)90006-3
- Riebesell, U., Schulz, K. G., Bellerby, R. G. J., Botros, M., Fritsche, P., Meyerhöfer, M., et al. (2007). Enhanced biological carbon consumption in a high CO₂ ocean. *Nature* 450, 545–548. doi: 10.1038/nature06267
- Rochelle-Newall, E. J., Mari, X., and Pringault, O. (2010). Sticking properties of transparent exopolymeric particles (TEP) during aging and biodegradation. *J. Plankton Res.* 32, 1433–1442. doi: 10.1093/plankt/fbq060
- Stodereger, K. E., and Herndl, G. J. (1998). Production of exopolymer particles by marine bacterioplankton under contrasting turbulence conditions. *Mar. Ecol. Prog. Ser.* 189, 9–16. doi: 10.3354/meps189009
- Tedetti, M., and Sempéré, R. (2006). Penetration of ultraviolet radiation in the marine environment. A review. *Photochem. Photobiol.* 82, 389–397. doi: 10.1562/2005-11-09-IR-733
- Van Oostende, N., Moerdijk-Poortvliet, T. C. W., Boschker, H. T. S., Vyverman, W., and Sabbe, K. (2013). Release of dissolved carbohydrates by *Emiliania huxleyi* and formation of transparent exopolymer particles depend on algal life cycle and bacterial activity. *Environ. Microbiol.* 15, 1514–1531. doi: 10.1111/j.1462-2920.2012.02873.x
- Verdugo, P., Alldredge, A. L., Azam, F., Kirchman, D. L., Passow, U., and Santschi, P. H. (2004). The oceanic gel phase: a bridge in the DOM POM continuum. *Mar. Chem.* 92, 67–85. doi: 10.1016/j.marchem.2004.06.017
- Wurl, O., Miller, L., and Vagle, S. (2011). Production and fate of transparent exopolymer particles in the ocean. *J. Geophys. Res.* 116, C00H13. doi: 10.1029/2011JC007342

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Iuculano, Mazuecos, Reche and Agustí. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.