



**UCC Library and UCC researchers have made this item openly available. Please [let us know](#) how this has helped you. Thanks!**

<b>Title</b>	The Preconscious Awareness to Action Framework: an application to promote testicular awareness
<b>Author(s)</b>	Saab, Mohamad M.; Landers, Margaret; Hegarty, Josephine
<b>Publication date</b>	2018-03-01
<b>Original citation</b>	Saab, M. M., Landers, M. and Hegarty, J. (2018) 'The Preconscious Awareness to Action Framework: an application to promote testicular awareness', <i>Nursing Research</i> , 67(2), pp. 169-176. doi: 10.1097/NNR.000000000000268
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="http://dx.doi.org/10.1097/NNR.000000000000268">http://dx.doi.org/10.1097/NNR.000000000000268</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2018, Wolters Kluwer Health, Inc. All rights reserved. This document is the Accepted Manuscript version of a Published Work that appeared in final form in <i>Nursing Research</i> . To access the final edited and published work see <a href="https://dx.doi.org/10.1097/NNR.000000000000268">https://dx.doi.org/10.1097/NNR.000000000000268</a>
<b>Embargo information</b>	Access to this article is restricted until 12 months after publication by request of the publisher.
<b>Embargo lift date</b>	2019-03-01
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/7049">http://hdl.handle.net/10468/7049</a>

Downloaded on 2019-12-02T14:51:07Z

1           **The Pre-Conscious Awareness to Action Framework: An Application to Promote**  
2   **Testicular Awareness**

3  
4           **Running Head:** The Pre-Conscious Awareness to Action Framework

5  
6           Mohamad M. Saab <sup>a,1</sup>, Lecturer, PhD Candidate, MSc, BSc, RN, PGCTLHE

7                           Margaret Landers <sup>a</sup>, Lecturer, PhD, MSc, BSc, RNT, RGN

8                           Josephine Hegarty <sup>a</sup>, Professor, PhD, MSc, BSc, RNT, RGN

9  
10  
11   **<sup>a</sup>Authors' affiliation:** Catherine McAuley School of Nursing and Midwifery, University College  
12   Cork, Brookfield Health Sciences Complex, College Road, Cork, Ireland.

13  
14   **<sup>1</sup>Corresponding author:** Mohamad M. Saab, Catherine McAuley School of Nursing and  
15   Midwifery, University College Cork, Brookfield Health Sciences Complex, College Road, Cork,  
16   Ireland.

17   Tel.: +353 21 490 1518

18   Fax: +353 21 490 1493

19   E-mail addresses: [msaab@ucc.ie](mailto:msaab@ucc.ie) ; [mohamad.m.saab@gmail.com](mailto:mohamad.m.saab@gmail.com)

20  
21   **Conflict of Interest and funding:** None

22  
23



## 1 1. INTRODUCTION

2 Globally, health outcomes among males continue to be markedly poorer than females,  
3 and efforts to promote men's health remain scarce (Baker et al., 2014). This, coupled with the  
4 assumption that men are disinterested in their own health, leads to gender-based health disparity  
5 and discourages men from engaging with health services (Leone & Rovito, 2013).

6 Of the diseases that are seldom discussed in current men's health initiatives, disorders of  
7 the testes can have a major impact on a man's wellbeing. Testicular cancer (TC) is the most  
8 frequently diagnosed solid tumor among men aged 18 to 50 years in the United States (National  
9 Cancer Institute [NCI], 2015). The incidence of TC in developed countries doubled over the past  
10 four decades (Rosen, Jayram, Drazer, & Eggener, 2011) and TC mortality is highest in the  
11 developing world due to late presentation and limited access to care (Znaor, Lortet-Tieulent,  
12 Jemal, & Bray, 2014). TC treatment is associated with chronic complications including fatigue,  
13 neuropathy, metabolic syndromes, and infertility (Saab, Noureddine, Huijjer, & DeJong, 2014).

14 Like TC, a number of benign testicular disorders can be life-threatening if left untreated.  
15 An example is testicular torsion which involves idiopathic twisting of the testes. This condition  
16 is a medical emergency that can lead to testicular ischemia and necrosis (Ringdahl & Teague,  
17 2006). Another example is epididymitis and orchitis (i.e. inflammation of the epididymis and  
18 testes) that are often transmitted sexually among men aged less than 50 and can lead to sepsis  
19 and infertility if not treated promptly (Centers for Disease Control and Prevention [CDC], 2015).

20 Experimental studies aimed at raising men's awareness of benign testicular disorders are  
21 lacking (Saab, Landers, & Hegarty, 2016a), and only half of the studies promoting TC awareness  
22 reviewed by Saab et al. (2016b) were underpinned by theory. Sidani and Fleury (2016) defined

1 interventions as “a set of inter-related activities directed toward attaining common goals” (p.  
2 190). Theory-based interventions help achieve positive health outcomes more than interventions  
3 that lack a theoretical basis (Michie, Johnston, Francis, Hardeman, & Eccles, 2008; Savage,  
4 Farrell, McManus, & Grey, 2010). Moreover, theory development is a key step in the Medical  
5 Research Council framework used to develop and evaluate complex health interventions (Craig  
6 et al., 2013). Therefore, a good theoretical understanding of a phenomenon is key to modelling  
7 an intervention, understanding how an intervention instigates change, and exploring the role of  
8 potential moderating and influencing variables (Michie et al., 2008; Sidani & Fleury, 2016).

9         In nursing, theory serves as a roadmap to design, implement, and evaluate interventions  
10 to promote health, maintain health, manage illness, and provide comfort at end-of-life (Sidani &  
11 Fleury, 2016). The elements of theory building include concepts, statements, and theories  
12 (Walker & Avant, 2011). Concepts are a mental image of a phenomenon and are regarded as the  
13 building blocks of a theory. Statements specify the context of theory building, either by defining  
14 the concepts (i.e. non-relational) or describing the relationship between them (i.e. relational).  
15 Walker and Avant (2011) defined theory as “an internally consistent group of relational  
16 statements that presents a systematic view about a phenomenon and that is useful for description,  
17 explanation, prediction, and prescription or control” (p. 7). It is often used to express a new idea  
18 or insight into the nature of a phenomenon. Theory is built using three key processes; synthesis,  
19 derivation, and analysis. Synthesis involves using information to construct a new concept or  
20 theory, derivation allows researchers to redefine a concept to fit a new context, and analysis  
21 involves clarifying refining, or sharpening concepts or theories (Walker & Avant, 2011).

22         This paper presents a theory-focused analysis of the Pre-Conscious Awareness to Action  
23 Framework (PAAF) developed to enhance testicular awareness and help-seeking intentions and

1 behaviors among men aged 18 to 50 years since those who fall within this age group are at the  
2 highest risk for testicular disorders (CDC, 2015; NCI, 2015; Ringdahl & Teague, 2006).

## 3 **2. METHODS**

4 The PAAF was developed following a synthesis of evidence from the empirical and  
5 theoretical literature on awareness, help-seeking, and symptom appraisal (Table 1).

### 6 **2.1 Reviews of the empirical literature**

7 A systematic review of exploratory studies on men's awareness of TC and self-  
8 examination (n=25 studies) (Saab et al., 2016c), a systematic review of experimental studies  
9 promoting awareness of TC and self-examination (n=11 studies) (Saab et al., 2016b), and an  
10 integrative review of exploratory studies on men's awareness of benign testicular disorders (n=4  
11 studies) (Saab et al., 2016a) were conducted. It was found that men lacked awareness of TC, self-  
12 examination, and benign testicular disorders and many intended to delay help-seeking for  
13 testicular lumpiness, swelling, and pain (Saab et al., 2016a,c). Ten of the 11 reviewed  
14 interventions promoting awareness of TC and self-examination succeeded in doing so.  
15 Moreover, six studies mentioned using theories, yet failed to map them onto the development  
16 and testing of the interventions (Saab et al., 2016b). Of note, none of the reviewed studies  
17 reported on men's preferred learning strategies, explored qualitatively their awareness of benign  
18 testicular diseases, or attempted to raise their awareness of diseases other than TC.

### 19 **2.2 Qualitative Study**

20 Findings and gaps from the reviewed empirical literature informed a qualitative study  
21 aimed at exploring men's (n=29) awareness of benign and malignant testicular disorders, help-

1 seeking intentions for testicular symptoms, and preferred learning strategies in relation to  
2 testicular disorders and symptoms. This study is reported in detail in Saab et al. (2017a,b).

3 Overall, men were unaware of the risk factors, signs and symptoms, and treatment of TC.  
4 Moreover, men's awareness of non-malignant testicular disorders was lacking, which echoes  
5 findings from the empirical literature (Saab et al., 2016a,b,c). A number of men intended to delay  
6 help-seeking for testicular symptoms due to knowledge deficit, symptom misappraisal, fear,  
7 embarrassment, cultural factors, cost and access to care, and inability to differentiate between  
8 normal and abnormal lumps (Saab et al., 2017a) (Figure 1). To help raise awareness, men  
9 recommended using educational interventions that are brief, visually stimulating, novel, and  
10 positively worded, with the use of light and simple language (Saab et al., 2017b).

### 11 **2.3 Review of the theoretical literature**

12 Data from the empirical literature and qualitative study helped plan an interactive  
13 educational intervention aimed at enhancing men's awareness of the normal testes and the most  
14 common testicular symptoms (e.g. lumpiness, swelling, and pain) and disorders (e.g. TC,  
15 epididymitis, orchitis, and testicular torsion). The ultimate goal of this intervention was to enable  
16 men to seek timely medical attention for testicular symptoms.

17 An iterative narrative review process of the theoretical literature on health promotion and  
18 symptom appraisal was undertaken to source and derive an underpinning theory/model.  
19 Derivation involves redefining a concept, statement, or theory from one context to another  
20 (Walker & Avant, 2011).

21 Six categories of multidisciplinary theories and models used in health promotion were  
22 identified as follows: behavioral change theories, intervention-based models, ecological theories

1 and models, planning models, communication theories, and evaluation models (Nutbeam, 2013;  
2 Raingruber, 2014). A behavioral change model, namely the Transtheoretical Model (TTM),  
3 provided insight into the cognitive domains involved in the thinking processes underpinning the  
4 development of awareness and intentions to alter behavior (Prochaska & DiClemente, 1986).  
5 Therefore, the five stages of the TTM (i.e. pre-contemplation, contemplation, preparation, action,  
6 and maintenance) served as a stepping stone to the construction of the PAAF.

7         A side-by-side comparison between the TTM and PAAF is presented in Table 2. The  
8 TTM and PAAF are process models that use predefined and interconnected stages in order to  
9 achieve a certain goal (Prochaska & DiClemente, 1986; Wislon & Schlam, 2004). However,  
10 while the TTM aims to ‘treat’ harmful behaviors, the primary goal of the PAAF is to raise  
11 awareness in order to promote healthy behaviors. Moreover, the TTM is comprised of five  
12 stages, whereas the PAAF is comprised of seven stages (i.e. pre-conscious awareness,  
13 unconscious awareness, conscious awareness, unconscious appraisal, conscious appraisal,  
14 intention, and behavior) derived from the TTM (Prochaska & DiClemente, 1986), and the  
15 literature on neurobehavioral psychology (Baumeister, Masicampo, & Vohs, 2011; Dehane &  
16 Naccahe, 2001) and symptom appraisal (Whitaker, Scott, & Wardle, 2015). Furthermore, while  
17 the TTM addresses the effect of consciousness on behavior, the PAAF acknowledges the impact  
18 of three stages of awareness (i.e. pre-conscious, unconscious, and conscious awareness) on  
19 behavior. Another difference pertains to symptom appraisal which is an integral part of the  
20 PAAF and not the TTM.



### 1 3. RESULTS

2 Movement between the steps of the PAAF is a fluid process (Figure 2). Individuals move  
3 in and out of unconscious and conscious awareness/appraisal as thoughts move in and out of  
4 consciousness and as the individual and the environmental context of thinking changes.

#### 5 3.1 Pre-Conscious Awareness

6 According to Sigmund Freud, pre-consciousness and unconsciousness are not  
7 synonymous (Natsoulas, 1996). While lack of awareness is a possibility, individuals in the pre-  
8 conscious awareness stage can be either under-informed about certain behaviors, have relevant  
9 discrete pieces of unlinked information, or are somewhat informed but not consciously thinking  
10 about the information. For instance, notwithstanding that men in the qualitative study lacked  
11 knowledge of TC risk factors, signs and symptoms, treatment, and screening; almost all of them  
12 reported having heard of TC through the media, school, college, family members, friends, and/or  
13 colleagues (Saab et al., 2017a,b). In other words, men were not oblivious to testicular disorders,  
14 as they were able to recall certain information in relation to these disorders when prompted.

15 There are a number of hypotheses as to why men in the reviewed literature and  
16 qualitative study lacked awareness. One explanation is that men are often preoccupied with life  
17 matters that they may perceive as more important than their own health, such as starting college,  
18 building a career, or establishing a family (Saab et al., 2014). Therefore, conflicting  
19 responsibilities have the potential to mask health awareness or push it to the bottom of their list  
20 of priorities. Furthermore, illness is globally linked to old age, which might cause younger men  
21 to think that they are not prone to getting sick (Saab et al., 2017a). Another explanation could be  
22 what Leone and Rovito (2013) referred to as “social norms and gender scripting” (p. 246) that

1 depict how men should and should not behave. These are thought to negatively affect men's  
2 health awareness and subsequent health outcomes.

### 3 **3.2 Conscious and Unconscious Awareness**

4 If one does not deliberately think about a certain behavior, it does not mean that he or she  
5 is not aware of it, which requires differentiating between conscious and unconscious awareness  
6 (Baumeister et al., 2011). This is key, since the impact both types of awareness have on health  
7 behavior had been overlooked in behavioral change theories, including the TTM.

8 According to Dehane and Naccahe (2001), “durable and explicit information  
9 maintenance, novel combinations of operations, and intentional behavior” (p. 9) are not possible  
10 without conscious awareness which is divided into two subtypes; the phenomenal awareness and  
11 conscious thought (Baumeister et al., 2011). Phenomenal awareness involves subjective  
12 experiences. An example is Armstrong's (1981) absent-minded driver, as driving involves  
13 automaticity and, despite being a learned behavior, is often instinctive rather than a purposefully  
14 planned course of action. In contrast, conscious thought involves reflection and reasoning. An  
15 example is a man trying to make sense of a newly discovered testicular lump. Moreover,  
16 conscious awareness helps combine various mental operations in order to plan, evaluate, and  
17 execute a novel behavior that cannot be attained unconsciously. Of note, certain mental  
18 processes are possible without conscious awareness; nonetheless, they are often short-lived  
19 (Baumeister et al., 2011). Therefore, a long-lasting memory of a recently acquired intentional  
20 behavior cannot be generated in the absence of conscious awareness (Sperling, 1960).

21 Taylor (2001) highlighted a form of long-term memory that is linked to unconscious  
22 awareness, namely the implicit memory which receives, stores, and recovers information outside  
23 one's conscious awareness. It is also thought to affect present behaviors as it involves habits,

1 attitudes, and preferences shaped by past events (Taylor, 2001). An example is the ingrained  
2 gender roles that often impact on men's health-seeking behaviors (Leone & Rovito, 2013).

### 3 **3.2.1 Testicular Awareness**

4 In order to address the lack of awareness of testicular disorders, researchers could adopt a  
5 novel and comprehensive concept namely 'testicular awareness'. To help define this concept, it  
6 is necessary to consult the literature on the concept of 'breast awareness' which was defined by  
7 Thornton and Pillarisetti (2008) as "a woman becoming familiar with her own breasts and the  
8 way they will change throughout her life. It encourages women to know how their own breasts  
9 look and feel normally so that they gain confidence about noticing any change" (p. 2119).

10 In order to achieve awareness, early detection of an abnormality must be followed by a  
11 specific course of action (Scott & Walter, 2010). However, a prerequisite to symptom detection,  
12 is familiarity with the normal state of the body. In the case of testicular awareness, familiarity  
13 would enable a man to establish a baseline of what is normal for him so that he would be able to  
14 recognize changes (e.g. swelling, lumpiness, and pain). This is key since lack of familiarity with  
15 one's own testes has been linked to intentions to delay help-seeking (Saab et al., 2017a).

16 Testicular awareness would also help promote heightened body awareness since it  
17 encourages men to become attentive to a body area that is seldom spoken about (Saab et al.,  
18 2014). Moreover, knowing the risk factors for testicular disorders would alert men to the  
19 aforementioned symptoms and disorders. For instance, men must recognize that having a first  
20 degree relative with TC increases their risk for this malignancy, and that unprotected sex exposes  
21 them to epididymitis and orchitis (CDC, 2015). Therefore, it is presumed that testicular  
22 awareness is a form of conscious rather than unconscious awareness.

1           Of note, testicular awareness does not necessarily involve scheduled self-examination,  
2 especially that the risks and benefits of this practice are debatable (Ilic & Misso, 2011). The U.S.  
3 Preventive Services Task Force (2011) discourages self-examination, mainly due to the potential  
4 harms of false positives, and concomitant anxiety. In contrast, proponents of TC screening argue  
5 that recommendations discouraging this practice are based on speculations rather than empirical  
6 evidence, especially that early diagnosis of TC was found to be more cost-effective than late  
7 diagnosis (Rovito, Manjelievskaia, Leone, Lutz, & Nangia, 2016). A middle ground could be  
8 reached by instructing men to feel their testes in order to establish a baseline of what is normal  
9 for them without necessarily promoting monthly testicular self-examination.

### 10 **3.3 Conscious and Unconscious Appraisal**

11           Symptoms, described as bodily sensation or changes, are subject to complex psychosocial  
12 processes. In a conceptual review of nine symptom appraisal models, Whitaker et al. (2015)  
13 defined the process of symptom appraisal as the “detection of bodily changes, interpretation of  
14 bodily changes, and responses to interpretation” (p. S28).

15           Detection of bodily changes involves recognizing a disturbance that can be general,  
16 localized, visible, palpable, and/or audible. This disturbance can differ in intensity and frequency  
17 (Whitaker et al., 2015). Therefore, for a bodily change to be detected, it must be of a significant  
18 magnitude. For example, a testicular lump must be large enough for a man to be able to detect it.

19           Symptom detection is followed by appraisal which involves “labeling, categorizing, and  
20 evaluating the bodily changes” (Whitaker et al., 2015, p.S28). Like awareness, appraisal is  
21 subject to conscious and unconscious influences as it involves reflection and reasoning in order  
22 to make sense of a symptom (i.e. conscious appraisal), and at the same time it can be impacted  
23 by past attitudes, beliefs, experiences, and/or behaviors (i.e. unconscious appraisal).

1           Symptom appraisal is a ‘delicate’ stage as it is subject to misinterpretation which can  
2 cause people to delay help-seeking; this was the case for a number of men in the qualitative  
3 study (Saab et al., 2017a). In addition, the impact of the external environment on the detection of  
4 bodily changes must not be overlooked (Pennebaker, 1982). For instance, excessive stimulation  
5 from the external environment can shift a person’s attention away from a symptom.

### 6 **3.4 Help-Seeking Intention and Behavior**

7           Following awareness and appraisal, a response is generated, which involves reaching a  
8 decision regarding the action that must be taken in relation to the symptom experienced. In the  
9 case of testicular awareness, a man decides either to seek medical attention or not. Alternately,  
10 he can revert to self-help measures, inform his family and friends, delay help-seeking, or adopt  
11 dysfunctional coping strategies such as denial and avoidance (Saab et al., 2017a).

12           As aforementioned, a number of barriers to help-seeking exist; these including lack of  
13 knowledge, symptom misappraisal, fear, embarrassment, machoism, and conflicting  
14 responsibilities. On the other hand, having the motivation and ability to seek help, and having  
15 access to healthcare settings positively influence one’s intention to seek medical care (Leventhal,  
16 H., Leventhal, E. A., & Contrada, 1998). Social support, disclosure of symptoms to friends,  
17 having an inherent help-seeking drive, and symptom severity also serve as enablers for help-  
18 seeking (Saab et al., 2017a). Finally, one can speculate that, when a man becomes ‘testes aware’,  
19 he is more likely to seek medical attention for testicular symptoms. This also is supported by  
20 evidence from the reviewed literature on men’s awareness of TC and self-examination (Saab et  
21 al., 2016c).

1           Relational statements are key to specifying the relationship between the concepts of the  
2 PAAF (Walker & Avant, 2011). These were simplified and presented in Figure 3, with the “-”  
3 sign depicting a negative relationship and the “+” sign depicting a positive relationship.

#### 4 **4. CONCLUSION**

5           To the best of the authors’ knowledge, the PAAF is the first framework to highlight the  
6 impact of three stages of awareness (i.e. pre-conscious, unconscious, and conscious awareness)  
7 and two stages of symptom appraisal (i.e. unconscious and conscious appraisal) on help-seeking  
8 intentions and behaviors. Moreover, The PAAF was applied to the concept of ‘testicular  
9 awareness’ that was first introduced in this paper. This concept can be instrumental in  
10 familiarizing men with their own testes and enabling them to detect abnormalities. This could  
11 influence their decision to seek timely medical attention, thus preventing potential complications  
12 that are linked to delayed help-seeking such as testicular ischemia, sepsis, and infertility.

13           In order to tailor effective health-promoting messages, researchers are encouraged to  
14 underpin their interventions with behavioral change theories, intervention-based models, and/or  
15 theories of health communication; an example is the PAAF discussed in the present paper. The  
16 authors are currently using the PAAF to build and test an interactive intervention aimed at  
17 raising men’s testicular awareness.

18

19

20

21

1 **REFERENCES**

- 2 Armstrong, D. M. (1981). What is consciousness? In D. M. Armstrong, *The nature of mind*  
3 *and other essays* (pp. 55-67). Ithaca, NY: Cornell University Press.
- 4 Baker, P., Dworkin, S. L., Tong, S., Banks, I., Shand, T., & Yamey, G. (2014). The men's  
5 health gap: Men must be included in the global health equity agenda. *Bulletin of the*  
6 *World Health Organization*, 92, 618-620.  
7 doi:http://dx.doi.org/10.2471/BLT.13.132795
- 8 Baumeister, R. F., Masicampo, E. J., & Vohs, K. D. (2011). Do conscious thoughts cause  
9 behavior? *Annual Review of Psychology*, 62, 331-361.  
10 doi:10.1146/annurev.psych.093008.131126
- 11 Centers for Disease Control and Prevention (2015). *Sexually transmitted diseases treatment*  
12 *guidelines – Epididymitis*. Retrieved from  
13 <http://www.cdc.gov/std/tg2015/epididymitis.htm>
- 14 Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew, M. (2013),  
15 Developing and evaluating complex interventions: The new Medical Research Council  
16 guidance. *International Journal of Nursing Studies*, 50(5), 587-592.  
17 doi:http://dx.doi.org/10.1016/j.ijnurstu.2012.09.010
- 18 Dehane, S. & Naccache, L. (2001). Towards a cognitive neuroscience of consciousness:  
19 Basic evidence and a workspace framework. *Cognition*, 79, 1-37.  
20 [https://doi.org/10.1016/S0010-0277\(00\)00123-2](https://doi.org/10.1016/S0010-0277(00)00123-2)
- 21 Ilic, D. & Misso, M. L. (2011). Screening for testicular cancer. *Cochrane Database of*  
22 *Systematic Reviews*, 16(2), D007853. doi:10.1002/14651858.CD007853.pub2

23

- 1 Leone, J. E. & Rovito, M. J. (2013). "Normative content" and health inequity enculturation a  
2 logic model of men's health advocacy. *American Journal of Men's Health*.  
3 doi:10.1177/1557988312469659
- 4 Leventhal, H., Leventhal, E. A., & Contrada, R. J. (1998). Self-regulation, health, and  
5 behaviour: A perceptual-cognitive approach. *Psychology & Health*, 13(4), 717-733.  
6 <http://dx.doi.org/10.1080/08870449808407425>
- 7 Michie, S., Johnston, M., Francis, J., Hardeman, Q., & Eccles, M. (2008). From theory to  
8 intervention: Mapping theoretically derived behavioural determinants to behaviour  
9 change techniques. *Applied Psychology*, 57(4), 660-680. doi:10.1111/j.1464-  
10 0597.2008.00341.x
- 11 National Cancer Institute (2015). *SEER stat fact sheets: Testis cancer*. Retrieved from  
12 <http://seer.cancer.gov/statfacts/html/testis.html>.
- 13 Natsoulas, T. (1996). Freud and consciousness: IX. James, Freud, and the theory of  
14 consciousness. *Psychoanalysis & Contemporary Thought*, 19(3),461-494.
- 15 Nutbeam, D. (2013). Using theory to guide change at the individual level. In L. Cragg, M.  
16 Davies, & W. Macdowall (Eds.), *Health promotion theory* (pp. 65-78). New York,  
17 NY: Open University Press.
- 18 Pennebaker, J. (1982). *The psychology of physical symptoms*. New York, NY: Springer.
- 19 Prochaska, J. O. & Di Clemente, C. C. (1986). Toward a comprehensive model of change. In  
20 W. R. Miller & N. Heather (Eds.), *Treating addictive behaviors: Processes of change*  
21 (pp. 3-27). New York, NY: Plenum Press.
- 22 Raingruber, B. (2014). Health promotion theories. *Contemporary health promotion in*  
23 *nursing practice* (pp. 53-94). Burlington: Jones & Bartlett Learning.



- 1 Ringdahl, E. & Teague, L. (2006). Testicular torsion. *American Family Physician*, 74, 1739-  
2 1743.
- 3 Rosen, A., Jayram, G., Drazer, M., & Eggener, S. E. (2011). Global trends in testicular  
4 cancer incidence and mortality. *European Urology*, 60, 374-379.  
5 doi:10.1016/j.eururo.2011.05.004
- 6 Rovito, M. J., Manjelievskaia, J., Leone, J. E., Lutz, M. J., & Nangia, A. (2016). From 'D' to  
7 'T': A critique of the current United States preventive services task force  
8 recommendation for testicular cancer screening. *Preventive Medicine Reports*, 3, 361-  
9 366. doi:10.1016/j.pmedr.2016.04.006
- 10 Saab, M. M., Landers, M., & Hegarty, J. (2016a). Males' awareness of benign testicular  
11 disorders: An integrative review. *American Journal of Men's Health*. Advance online  
12 publication. doi:10.1177/1557988315626508
- 13 Saab, M. M., Landers, M., & Hegarty, J. (2016b). Promoting testicular cancer awareness and  
14 screening: A systematic review of interventions. *Cancer Nursing*, 39(6), 473-487.  
15 doi:10.1097/NCC.0000000000000333
- 16 Saab, M. M., Landers, M., & Hegarty, J. (2016c). Testicular cancer awareness and screening  
17 practices: A systematic review. *Oncology Nursing Forum*, 43, E8-E23.  
18 doi:10.1188/16.ONF.E8-E23
- 19 Saab, M. M., Landers, M., & Hegarty, J. (2017a). Exploring awareness and help-seeking  
20 intentions for testicular symptoms among heterosexual, gay, and bisexual men in  
21 Ireland: A qualitative descriptive study. *International Journal of Nursing Studies*, 67,  
22 41-50. doi:10.1016/j.ijnurstu.2016.11.016

- 1 Saab, M. M., Landers, M., & Hegarty, J. (2017b). Exploring men's preferred strategies for  
2 learning about testicular disorders inclusive of testicular cancer: A qualitative  
3 descriptive study. *European Journal of Oncology Nursing*, 26, 27-35.  
4 doi:10.1016/j.ejon.2016.11.001
- 5 Saab, M., Noureddine, S., Huijjer, H. A. S., & DeJong, J. (2014). Surviving testicular cancer:  
6 The Lebanese lived experience. *Nursing Research*, 63, 203-209.  
7 doi:10.1097/NNR.0000000000000033
- 8 Savage, E., Farrel, D., McManus, V., & Grey, M. (2010). The science of intervention  
9 development for type 1 diabetes in childhood: A systematic review. *Journal of*  
10 *Advanced Nursing*, 66, 2604-2619. doi:10.1111/j.1365-2648.2010.05423.x
- 11 Scott, S. & Walter, F. (2010). Studying help-seeking for symptoms: The challenges of  
12 methods and models. *Social and Personality Psychology Compass*, 4, 531-547.  
13 doi:10.1111/j.1751-9004.2010.00287.x
- 14 Sidani, S. & Fleury, J. (2016). Theory-based nursing interventions. In S. J. Henly (Ed.),  
15 *Routledge international handbook of advanced quantitative methods in nursing*  
16 *research* (pp. 189-198). New York, NY: Routledge.
- 17 Sperling, G. (1960). The information available in brief visual presentations. *Psychological*  
18 *Monographs: General and Applied*, 75(11), 1-29.
- 19 Taylor, E. W. (2001). Transformative learning theory: A neurobiological perspective of the  
20 role of emotions and unconscious ways of knowing. *International Journal of Lifelong*  
21 *Education*, 20(3), 218-236. <http://dx.doi.org/10.1080/02601370110036064>

- 1 Thornton, H. & Pillarisetti, R. R. (2008). 'Breast awareness' and 'breast self-examination'  
2 are not the same. What do these terms mean? Why are they confused? What can we  
3 do? *European Journal of Cancer*, 44, 2118-2121. doi:10.1016/j.ejca.2008.08.015
- 4 Trumbo, C. W. (2004). Mass-mediated information effects on testicular self-examination  
5 among college students. *Journal of American College Health*, 52(6), 257-262. doi:  
6 10.3200/JACH.52.6.257-262
- 7 U.S. Preventive Services Task Force (2011). Screening for testicular cancer: U.S. preventive  
8 task force reaffirmation recommendation statement. *Annals of Internal Medicine*,  
9 154(7), 483-486. doi:10.7326/0003-4819-154-7-201104050-00006
- 10 Walker, L. O. & Avant, K. C. (2011). *Strategies for theory construction in nursing*. New  
11 Jersey, NJ: Pearson.
- 12 Wanzer, M. B., Foster, C., Servoss, T., & LaBelle, S. (2014). Educating young men about  
13 testicular cancer: Support for a comprehensive testicular cancer campaign. *Journal of*  
14 *Health Communication*, 19, 303-320. doi:10.1080/10810730.2013.811320
- 15 Whitaker, K. L., Scott, S. E., & Wardle, J. (2015). Applying symptom appraisal models to  
16 understand sociodemographic differences in responses to possible cancer symptoms: A  
17 research agenda. *British Journal of Cancer*, 112, S27-S34. doi:10.1038/bjc.2015.39
- 18 Wilson, G.T. & Schlam, T.R. (2004). The Transtheoretical model and motivational  
19 interviewing in the treatment of eating and weight disorders. *Clinical Psychology*,  
20 24(3), 361-378. doi:10.1016/j.cpr.2004.03.003
- 21 Znaor, A., Lortet-Tieulent, J., Jemal, A., & Bray, F. (2014). International variations and  
22 trends in testicular cancer incidence and mortality. *European Urology*, 65, 1095-1106.  
23 doi:10.1016/j.eururo.2013.11.004

1 **FIGURE LEGENDS**

2 **Figure 1.** Barriers to awareness and appraisal of testicular symptoms (Saab et al., 2017a).

3 **Figure 2.** The PAAF as applied to the concept of testicular awareness (Saab et al., 2016a,b,c;  
4 Whitaker et al., 2015).

5 **Figure 3.** Simplified statements illustrating the relationship between the key concepts of the  
6 PAAF (Walker & Avant, 2011).

7

8

9

10

11

12

13

14

15

16

17

18

19

**Table1.** Synthesis and derivation processes guiding the development of the PAAF

STEPS	METHODS	KEY FINDINGS
Empirical literature reviews	<ul style="list-style-type: none"> <li>• Systematic review of 25 quantitative and qualitative studies exploring men’s knowledge, awareness, and attitudes towards testicular cancer and self-examination (Saab et al., 2016c)</li> </ul>	<ul style="list-style-type: none"> <li>• Men lacked awareness of testicular cancer risk factors, signs and symptoms, and treatment</li> <li>• Few men practiced self-examination with the majority not knowing what to look for</li> <li>• Many reported an intention to delay help-seeking for testicular lumpiness</li> <li>• Men perceived education about testicular cancer and self-examination as important</li> </ul>
	<ul style="list-style-type: none"> <li>• Systematic review of 11 experimental studies promoting men’s awareness of testicular cancer and self-examination (Saab et al., 2016b)</li> </ul>	<ul style="list-style-type: none"> <li>• Ten of the 11 studies successfully raised men’s awareness of testicular cancer and self-examination e.g. a college campaign (Wanzer et al., 2014) and mass media (Trumbo, 2004)</li> <li>• None of the studies reported on men’s preferred learning strategies a priori</li> <li>• Six of the 11 studies were underpinned by theory</li> </ul>
	<ul style="list-style-type: none"> <li>• Integrative review of four quantitative studies exploring men’s awareness of benign testicular disorders (Saab et al., 2016a)</li> </ul>	<ul style="list-style-type: none"> <li>• Men’s awareness of benign testicular disorders was lacking</li> <li>• Many reported an intention to delay help-seeking for testicular lumpiness and pain</li> <li>• No qualitative or experimental studies were identified in the literature search</li> </ul>
Qualitative descriptive study	<ul style="list-style-type: none"> <li>• To explore men’s (n=29) awareness of testicular disorders, help-seeking intentions for testicular symptoms, and preferred strategies for learning about testicular disorders and symptoms (Saab et al., 2017a,b)</li> <li>• 12 semi-structured individual interviews and three focus groups were conducted with men aged 18-47 years.</li> <li>• Data were analyzed using qualitative content analysis</li> <li>• Trustworthiness was enhanced by selecting a heterogeneous sample, cross-checking the analyzed data, member-checks, memoing, and using field notes and audit trails</li> </ul>	<ul style="list-style-type: none"> <li>• Men lacked awareness of testicular cancer; very few reported having heard of benign testicular disorders; and many intended to delay help-seeking for lumpiness, swelling, and/or pain</li> <li>• Barriers to awareness and help-seeking included: lack of prior knowledge, symptom misappraisal, fear, embarrassment, machoism, lack of endorsement by the health system, cost and access to care, and inability to differentiate between normal and abnormal lumps</li> <li>• Facilitators to awareness and help-seeking included: prior knowledge, regular self-examination, clinical testicular examination, personal and/or family history of testicular disease, aging, being gay, access to support, inherent health-seeking drive, and perceived threats to fertility</li> <li>• Men stressed the importance of raising awareness using educational interventions that are brief, simple, visually appealing, novel, and positively worded</li> </ul>
Theoretical literature review	<ul style="list-style-type: none"> <li>• An iterative narrative review process of the theoretical literature on health promotion, symptom appraisal, and neurobehavioral psychology was undertaken.</li> <li>• Six categories of multidisciplinary theories used in health promotion were identified: behavioral change theories, intervention-based models, ecological theories and models, planning models, communication theories, and evaluation models (Nutbeam, 2013; Raingruber, 2014)</li> </ul>	<ul style="list-style-type: none"> <li>• The five stages of the Transtheoretical Model guided the development of the seven stages of the Pre-Conscious Awareness to Action Framework</li> <li>• The literature on neurobehavioral psychology helped derive the concepts of pre-conscious, unconscious, and conscious awareness</li> <li>• The empirical literature, the qualitative study, and the concept of “breast awareness” helped create and define the concept of “testicular awareness” (Thornton &amp; Pillarisetti, 2008)</li> <li>• The literature on symptom appraisal helped derive the concepts of unconscious and conscious appraisal (Whitaker et al., 2015)</li> </ul>

1

2

**Table 2.** Comparison of the key characteristics of the TTM and the PAAF

	<b>TTM<sup>a</sup></b>	<b>PAAF</b>
<b>Discipline</b>	Clinical psychology and psychotherapy	Health promotion and neurobehavioral psychology
<b>Primary goal</b>	To treat and change pre-existing addictive/harmful behaviors (e.g. smoking)	To raise awareness to promote new and healthy behaviors (e.g. early help-seeking)
<b>Development</b>	Comparative analysis of 29 systems of psychotherapy	Iterative process of synthesis, derivation, and analysis
<b>Stages</b>	Five stages: pre-contemplation, contemplation, preparation, action, and maintenance	Seven stages: pre-conscious awareness, unconsciousness awareness, conscious awareness, unconscious appraisal, conscious appraisal, intention, and behavior
<b>Pre-contemplation vs. Pre-conscious awareness</b>	People in the pre-contemplation stage are uninformed or under-informed about the consequences of their behavior	People in the pre-conscious awareness stage are uninformed, under-informed, have relevant discrete pieces of unlinked information, or are somewhat informed but not consciously thinking about the information
<b>Consciousness vs. Conscious awareness</b>	Highlights the role of consciousness raising in promoting intentions to change a behavior	Highlights the role of conscious awareness and memory in retaining new information, combining various mental processes, and understanding and shaping behavior
<b>Symptom appraisal</b>	No symptoms involved as the primary goal is to treat pre-existing addictive behaviors	Involves labeling, categorizing, and evaluating bodily changes
<b>Intention</b>	Intention to change and take action imbedded in the contemplation and preparation stages respectively	A separate stage that is impacted by motivation and ability to seek help and access to healthcare services
<b>Action vs. Behavior</b>	Course of action needed to modify one's lifestyle. It must be maintained to avoid relapse	Ultimate behavior is to seek timely medical attention in response to a symptom
<b>Context</b>	Emphasis on the impact of the harmful behavior on the social environment	Emphasis on the impact of intrinsic and extrinsic factors on the seven stages of the framework
<b>Testing</b>	Tested in various contexts (e.g. smoking cessation, alcohol use, and weight reduction)	Currently being used to build and test an interactive intervention aimed at raising men's testicular awareness


<sup>a</sup>Prochaska and Di Clemente (1986)

1

2

3

4



1

2