

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/75308>

Please be advised that this information was generated on 2018-07-08 and may be subject to change.

## Ground-based observations of the $\beta$ Cephei CoRoT main target HD 180642

M. Briquet<sup>1</sup>, K. Uytterhoeven<sup>2,3</sup>, C. Aerts<sup>1,4</sup>, T. Morel<sup>1,6</sup>, P. De Cat<sup>5</sup>, A. Miglio<sup>6</sup>, P. Mathias<sup>7</sup>, K. Lefever<sup>1</sup>, E. Poretti<sup>2</sup>, S. Martín-Ruiz<sup>8</sup>, M. Páparó<sup>9</sup>, M. Rainer<sup>2</sup>, F. Carrier<sup>1</sup>, J. Gutiérrez-Soto<sup>10</sup>, J.C. Valtier<sup>7</sup>, J.M. Benkő<sup>9</sup>, Zs. Bognár<sup>9</sup>, P.J. Amado<sup>8</sup>, J.C. Suarez<sup>8</sup>, C. Rodríguez-Lopez<sup>8</sup> and R. Garrido<sup>8</sup>

<sup>1</sup>Instituut voor Sterrenkunde,  
 Katholieke Universiteit Leuven, Celestijnenlaan 200 D, B-3001 Leuven, Belgium  
<sup>2</sup>INAF-Osservatorio Astronomico di Brera, Via E. Bianchi 46, I-23807 Merate, Italy  
<sup>3</sup>Instituto de Astrofísica de Canarias,  
 Calle Via Láctea s/n, E-38205 La Laguna, TF, Spain  
<sup>4</sup>Department of Astrophysics,  
 University of Nijmegen, IMAPP, PO Box 9010, 6500 GL Nijmegen, the Netherlands  
<sup>5</sup>Koninklijke Sterrenwacht van België, Ringlaan 3, B-1180 Brussel, Belgium  
<sup>6</sup>Institut d'Astrophysique et de Géophysique de l'Université de Liège,  
 Allée du 6 Août 17, B-4000 Liège, Belgium  
<sup>7</sup>UMR 6525 H. Fizeau, UNS, CNRS, OCA, Campus Valrose,  
 F-06108 Nice Cedex 2, France  
<sup>8</sup>Instituto de Astrofísica de Andalucía, CSIC, Apdo. 3004, 18080 Granada, Spain  
<sup>9</sup>Konkoly Observatory, P.O. Box 67, 1525 Budapest, Hungary  
<sup>10</sup>GEP1, Observatoire de Paris, CNRS, Université Paris Diderot,  
 place Jules Janssen 92195 Meudon Cedex, France

### Abstract

We present the preliminary results of a detailed study of ground-based photometric and spectroscopic observations dedicated to the  $\beta$  Cephei CoRoT main target HD 180642. Besides the non-linear dominant radial mode several low-amplitude modes are detected in both kinds of datasets. Our aim is to derive the wavenumbers ( $\ell, m$ ) of these modes, as additional constraints to the CoRoT pulsation frequencies, for forthcoming asteroseismic modelling of the star.

Individual Objects: HD 180642

### Introduction

The B1.5II-III star HD 180642 (V1449 Aql, HIP 94793,  $m_v = 8.27$ ) is the only known  $\beta$  Cephei star in CoRoT's core programme. It has been observed by the satellite during a long run (some 150 days) from May to October 2007. The star presents a dominant non-linear mode with an amplitude of  $\sim 40$  mmag together with several modes of much lower amplitudes (below 4 mmag). Fourier periodograms of its CoRoT lightcurve are displayed in Michel et al. (2008) and are not repeated here. Numerous frequencies are found in the CoRoT lightcurve, among which harmonics and combination frequencies. Its full analysis will be published elsewhere (Degroote et al., in preparation).

To complement the space white light data, both ground-based photometric and spectroscopic data were collected, to which we added several archival observations. In total, we

Table 1: Logbook of the spectroscopic observations. For each instrument we give the number of spectra  $N$ , the time-span  $\Delta T$  in days, the average S/N-ratio and the resolution of the spectrograph.

Instrument	$N$	$\Delta T$	$\langle S/N \rangle$	resolution
FEROS	223	967.5	186	48 000
SOPHIE	35	10.1	106	40 000
Aurélie	22	8.1	93	25 000

made use of 507 high-quality multi-colour photometric data, 172 white light data, and 280 high-resolution high S/N spectra. In this paper, we describe the first results of the analysis of these datasets.

### Ground photometry

Several telescopes with multi-colour photometric instruments are involved in this project. Strömgren *ubvy* photometry was collected with the 90cm telescope at Sierra Nevada Observatory and with the 1.5m telescope at San Pedro Mártir Observatory; Geneva 7-colour photometry (*UB<sub>1</sub>BB<sub>2</sub>V1VG*) with the 1.2m Mercator telescope at the Observatorio Roque de los Muchachos and with the 70cm Swiss telescope at La Silla Observatory; and Johnson *V* photometry with the 50cm telescope at Konkoly Observatory. Hipparcos measurements are also available. A logbook of the observations is shown in Table 2 of Uytterhoeven et al. (2008).

We combined the datasets for the filters that allow to do so, as follows. First, we merged the Strömgren data of both sites. Second, we constructed ultraviolet, blue, and visual ground-based lightcurves (combined *U, B, V* data hereafter) by merging the Geneva *U* and Strömgren *u, B* and *v*, and *V* and *y* data, respectively. Finally, we also constructed a more extensive visual band lightcurve by adding also the Johnson *V* data and the Hipparcos  $H_p$  data to the Geneva *V* and Strömgren *y* data.

As an illustration of our frequency analysis of the merged datasets, Scargle periodograms calculated from the combined *U* data are shown in Fig. 1. The variations due to the main mode occur with  $f_1 = 5.48694 \text{ d}^{-1}$ ,  $2f_1$  and  $3f_1$ . The corresponding amplitudes in the combined *U* filter are 82.8, 8.3 and 4.3 mmag. Note that HD 180642 has one of the highest amplitudes among the known  $\beta$  Cephei stars (see catalog of Stankov & Handler 2005 for comparison). The dominant mode of HD 180642 was already identified by Aerts (2000) as a radial mode by means of Geneva photometry.

In the combined Strömgren and ultraviolet datasets, we also detect two additional significant frequencies, i.e. their corresponding amplitudes reach at least 4 times the noise level in the periodogram, following the criterion derived empirically by Breger et al. (1993). The results from the CoRoT data (Michel et al. 2008) allow us to pick out the right peak. The second frequency present in our photometric data is  $f_2^p = 0.30818(5) \text{ d}^{-1}$  instead of its one-day-alias whose amplitude is higher in the periodogram (see Fig. 1). Afterwards, we find the frequency  $f_3^p = 7.36673(7) \text{ d}^{-1}$ .

### Spectroscopy

In the framework of the CoRoT ground-based Large Programme (e.g. Uytterhoeven & Poretti 2007; Uytterhoeven et al. 2008), aimed at the follow-up of selected CoRoT targets, HD 180642 was observed with the FEROS spectrograph at the 2.2m telescope at ESO La Silla. The target

was also monitored with the SOPHIE (1.93m telescope) and Aurélie (1.52m telescope) spectrographs, both situated at the Haute-Provence observatory. The logbook of the observations is shown in Table 1.

A phase diagram of the first moment  $\langle v^1 \rangle$  (radial velocity placed at average zero) is displayed in Fig. 2 and clearly shows the non-linear behaviour of the dominant pulsation  $f_1$  in HD 180642. The peak-to-peak amplitude is  $\sim 90 \text{ km s}^{-1}$  and harmonics up to  $5f_1$  are present in  $\langle v^1 \rangle$ . This makes HD 180642 the  $\beta$  Cephei star with the third largest radial-velocity amplitude, BW Vul having a peak-to-peak amplitude of  $\sim 200 \text{ km s}^{-1}$  (Aerts et al. 1995) and  $\sigma$  Sco of  $\sim 110 \text{ km s}^{-1}$  (Mathias et al. 1991).

In the radial velocity measurements, we also clearly detect  $f_2^s = 8.408(1) \text{ d}^{-1}$  as well as  $f_2^p$ , with an amplitude of  $1.5 \text{ km s}^{-1}$  and  $1.1 \text{ km s}^{-1}$ , respectively. In a 2D frequency search across the line profile, we additionally find  $f_3^s = 6.325(1) \text{ d}^{-1}$ . The amplitude and phase distributions for the Si III 4553 Å line for  $f_2^s$  and  $f_3^s$ , also detected in the CoRoT lightcurve, are shown in Fig. 3.

We note that the difference between  $f_2^p$ ,  $f_2^s$  and  $f_3^s$  is  $\sim 1.041 \text{ d}^{-1}$ . The frequency precision of our observations as well as the uninterrupted CoRoT photometry exclude the possibility of dealing with one-day-aliases.

## Conclusions

Besides the dominant radial mode, low-amplitude non-radial modes are detected in our ground-based observations of HD 180642. Our aim is to perform a careful mode identification for them. Because of the highly non-linear behaviour of the radial mode, available mode identification techniques cannot be applied straightforwardly, as they are implemented for a linear pulsation. In particular for spectroscopy, the moment method (Briquet & Aerts 2003) and the FPF method (Zima 2006) need to be adapted. This work is ongoing and the final results will be described in a forthcoming paper.

**Acknowledgments.** MB and FC are Postdoctoral researchers of the Fund for Scientific Research, Flanders. KU acknowledges financial support from a *European Community Marie Curie Intra-European Fellowship*, contract number MEIF-CT-2006-024476. The FEROS data are being obtained as part of the ESO Large Programme: LP178.D-0361 (PI: E. Poretti). This work was supported by the Italian ESS project, contract ASI/INAF I/015/07/0, WP 03170, by the Hungarian ESA PECS project No 98022, and by the European Helio- and Asteroseismology Network (HELAS), a major international collaboration funded by the European Commission's Sixth Framework Programme.

## References

- Aerts, C., Mathias, P., Van Hoolst, T., et al. 1995, *A&A*, 301, 781  
 Aerts, C. 2000, *A&A*, 361, 245  
 Breger, M., Stich, J., Garrido, R., et al. 1993, *A&A*, 271, 482  
 Briquet, M., & Aerts, C. 2003, *A&A*, 398, 687  
 Mathias, P., Gillet, D., & Crowe, R. 1991, *A&A*, 252, 245  
 Michel, E., Baglin, A., Weiss, W., et al. 2008, *CoAst*, 156, 73  
 Stankov, A., & Handler, G. 2005, *ApJS*, 158, Issue 2, 193  
 Uytterhoeven, K., Poretti, E., & the CoRoT SGBOWG 2007, *CoAst*, 150, 371  
 Uytterhoeven, K., Poretti, E., Rainer, M., et al. 2008, in 'HelasII international conference: Helioseismology, Asteroseismology and MHD Connections', *Journal of Physics: Conference Series*, IOP Publishing, in press (astro-ph 0710.4068)  
 Zima, W. 2006, *A&A*, 455, 227




Figure 1: Scargle periodograms calculated from the combined Geneva  $U$  and Strömgren  $u$  data. The uppermost panel shows the spectral window of the data. All subsequent panels show the periodograms at different stages of prewhitening. From top to bottom: periodogram of the observed light variations, of the data prewhitened with  $f_1$  and its two harmonics  $2f_1$  and  $3f_1$ , of the data subsequently prewhitened with  $f_2^p$ , and of the data subsequently prewhitened with  $f_3^p$ . The three frequencies found are indicated by dashed gray lines. The light gray lines indicate the 4 S/N level.




Figure 2: Phase diagram of the first moment computed from the Si III 4553 Å line, for  $f_1 = 5.48695 \text{ d}^{-1}$ .




Figure 3: Amplitude and phase distributions for  $f_2^s = 8.408 \text{ d}^{-1}$  (left) and  $f_3^s = 6.325 \text{ d}^{-1}$  (right) for the Si III 4553 Å line. The amplitudes are expressed in units of continuum and the phases in  $\pi$  radians.