



American Journal of
Medical Genetics
Part B: Neuropsychiatric Genetics

Association of Aggression With a Novel MicroRNA Binding Site Polymorphism In the Wolfram Gene

Journal:	<i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i>
Manuscript ID:	NPG-12-0187.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Kovacs-Nagy, Reka; Semmelweis University, Department of Medical Chemistry Elek, Zsuzsanna; Semmelweis University, Department of Medical Chemistry Szekely, Anna; Eotvos Lorand University, Institute of Psychology Nanasi, Tibor; Semmelweis University, Department of Medical Chemistry Sasvari-Szekely, Maria; Semmelweis University, Department of Medical Chemistry Ronai, Zsolt; Semmelweis University, Department of Medical Chemistry
Keywords:	aggression, microRNA, WFS1, rs1046322, SNP

SCHOLARONE™
Manuscripts

view

Kovacs-Nagy et al. WFS1 and aggression

**ASSOCIATION OF AGGRESSION WITH A NOVEL MICRORNA BINDING
SITE POLYMORPHISM IN THE WOLFRAMIN GENE**

Reka Kovacs-Nagy¹, Zsuzsanna Elek¹, Anna Szekely², Tibor Nanasi¹, Maria Sasvari-
Szekely¹, Zsolt Ronai¹

¹ Institute of Medical Chemistry, Molecular Biology and Pathobiochemistry, Semmelweis
University, Budapest, Hungary

² Institute of Psychology, Eötvös Loránd University, Budapest, Hungary

Corresponding author: Zsolt Ronai

Tuzolto u. 37–47. Budapest, H-1094 Hungary

Tel: +3620-6660135

Fax: +3620-6660104

Email: ronai.zsolt@med.semmelweis-univ.hu

Running Head: WFS1 and aggression

Keywords: aggression, microRNA, WFS1, rs1046322, SNP

Kovacs-Nagy et al. WFS1 and aggression

ABSTRACT

Background

Rare mutations in the WFS1 gene lead to Wolfram syndrome, a severe multisystem disorder with progressive neurodegeneration and diabetes mellitus causing life-threatening complications and premature death. Only a few association studies using small clinical samples tested the possible effects of common WFS1 gene variants on mood disorders and suicide, the non-clinical spectrum has not been studied yet.

Methods

Self-report data on Aggression, Impulsiveness, Anxiety and Depression were collected from a large (N=801) non-psychiatric sample. SNPs were selected to provide an adequate coverage of the entire WFS1 gene, as well as to include putative microRNA binding site polymorphisms. Molecular analysis of the assumed microRNA binding site variant was performed by an *in vitro* reporter-gene assay of the cloned 3' UTR with coexpression of miR-668.

Results

Among the 17 WFS1 SNPs, only the rs1046322, a putative microRNA (miR-668) binding site polymorphism showed significant association with psychological dimensions after correction for multiple testing: those with the homozygous form of the minor allele reported higher aggression on the Buss-Perry Aggression Questionnaire ($p=0.0005$). Functional effect of the same SNP was also demonstrated in a luciferase reporter system: the minor A allele showed lower repression compared to the major G allele, if co-expressed with miR-668.

Conclusions

To our knowledge, this is the first report describing a microRNA binding site polymorphism of the WFS1 gene and its association with human aggression based on a large, non-clinical sample.

Kovacs-Nagy et al. WFS1 and aggression

INTRODUCTION

Wolfram syndrome (WS) is an autosomal recessive multisystem disorder with progressive neurodegeneration and diabetes mellitus [Rigoli and others 2011], also known as DIDMOAD (diabetes insipidus, diabetes mellitus, optic atrophy and deafness). Although the clinical course of WS is highly variable, the prognosis is poor as patients have an impaired quality of life and usually die in their thirties from central respiratory failure as a result of brainstem atrophy [Barrett and others 1995]. Psychiatric manifestations are common in WS patients, including severe depression, psychosis, as well as impulsive verbal and physical aggression [Swift and others 1990].

The mutated gene, WFS1 was identified in 1998 on chromosome 4p16.1 spanning 33.4 kb [Inoue and others 1998]; it has eight exons but the first exon is noncoding. The WFS1 gene encodes a transmembrane protein (wolframin) of the endoplasmatic reticulum (ER), predominantly expressed in the brain and in the pancreatic β cells, but detected nearly in all tissues. Although the molecular function of wolframin is not fully understood to date, current results suggest that wolframin has a critical role in the regulation of ER stress [Fonseca and others 2005; Fonseca and others 2010]. ER homeostasis is based on a sensitive balance of protein synthesis and protein folding capacity of the ER. Accumulation of unfolded proteins induces ER stress, triggering the unfolded protein response (UPR). At the same time it is of vital importance to avoid UPR hyperactivation, which could trigger apoptotic pathways and lead to cell death. According to recent data, wolframin might participate in down regulation of ER stress and thus, in the reestablishment of ER homeostasis [Osowski and Urano 2011]. Further findings pointed out another possible role of wolframin in controlling ER calcium homeostasis [Takei and others 2006]. A recent study demonstrated that wolframin is a calcium-calmodulin binding protein, and three mutations found in WS patients completely abolish this function [Yurimoto and others 2009]. Taken together the findings above about the

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 molecular function of wolframin, this protein is expected to have a vital role in ER
4
5 homeostasis of excretory cells, such as the insulin producing pancreatic β cells or the neurons
6
7 producing synaptosomes for neurotransmission.
8

9
10 Studies on the WFS1 knockout mouse provided further evidence for the significant role of
11
12 wolframin in the brain and in the pancreas. WFS1 deficient mice had a dramatic decrease in
13
14 the mass of the pancreatic islets and a serious impairment of glucose homeostasis [Ishihara
15
16 and others 2004]. Results from behavioral studies of the WFS1 knockout mouse suggested
17
18 that it could be used as an animal model of mood disorder [Kato and others 2008], impaired
19
20 behavioral stress adaptation [Luuk and others 2009] and post-traumatic stress disorder
21
22 [Kesner and others 2009].
23

24
25 Although more than 200 different mutations causing WS have been identified (for a recent
26
27 database see: [M. 2010]), phenotype-genotype relationships are still not clear. Similarly, little
28
29 is known about the clinical and molecular effects of common variants. Several single
30
31 nucleotide polymorphisms (SNPs) have been shown as possible risk factors for type 2
32
33 diabetes mellitus, such as the rs10010131 in intron 4 and rs1046320 in the 3' untranslated
34
35 region (UTR) [Fawcett and others 2010]. Other studies raised the possibility of association
36
37 between several WFS1 SNPs and neuropsychiatric disorders, such as mood disorders
38
39 [Kawamoto and others 2004; Koido and others 2005] and impulsive suicide [Lemondé and
40
41 others 2003; Must and others 2009]. Although the level of significance did not survive
42
43 corrections for multiple testing in most studies, the possible link between WFS1 genetic
44
45 variants and psychiatric disorders is strongly supported by the fact that carriers of various
46
47 WFS1 mutations are 26-fold more likely to be hospitalized in psychiatric institutions than
48
49 noncarriers [Swift and Swift 2000].
50
51

52
53 According to our best knowledge, WFS1 polymorphisms have not been studied yet in relation
54
55 to either aggression or mood characteristics within the non-clinical spectrum. Here we present
56
57
58
59
60

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 an association study of WFS1 SNPs and four psychological dimensions: Aggression,
4 Impulsiveness, Anxiety, and Depression, using self-report data from a large, non-psychiatric
5 sample. Study of these phenotypes was based on the main psychiatric symptoms of WS
6 patients (depression and aggression), as well as on the results of the few psychogenetic
7 association studies related to mood disorders and impulsive suicide. In addition, we
8 characterized the molecular function of rs1046322 which had a significant effect on the
9 measured phenotypes.
10
11
12
13
14
15
16
17

18 **MATERIALS AND METHODS**

19 **Sample**

20
21
22
23
24 Caucasian (Hungarian) participants of the study signed written informed consent, provided
25 buccal samples and filled out questionnaires. They were recruited at different higher
26 education sites on a voluntary basis. The study protocol was approved by the Scientific and
27 Research Ethics Committee of the Medical Research Council (ETT TUKÉB). Valid DNA and
28 questionnaire data were available from 801 subjects (46.2% males and 53.8% females). Mean
29 age was 21.3 years (± 3.3) within the age range of 18–35 years.
30
31
32
33
34
35
36
37
38
39
40

41 **Phenotypic Measures**

42
43 All subjects completed a panel of self-report measures relevant to the study of Aggression,
44 Impulsiveness, Anxiety and Depression. The Buss-Perry Aggression Questionnaire [Buss and
45 Perry 1992] includes 29 items, rated from 1 (extremely uncharacteristic of me) to 5
46 (extremely characteristic of me). The aggression construct was measured with items loading
47 on four distinct factors: Verbal Aggression (5 items), Anger (7 items), Hostility (8 items) and
48 Physical Aggression (9 items). The total score for Aggression was the sum of ratings for all
49 the items, thus possible total scores range from 29 to 145. The 11th version of the Barratt
50
51
52
53
54
55
56
57
58
59
60

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 Impulsiveness Scale was designed to assess general impulsiveness using 30 items scored from
4 1–4 (BIS-11; [Patton and others 1995]). In the present study the total score for Impulsiveness
5 was used, with values ranging from 30–120. The English version of the questionnaires
6 measuring aggression and impulsiveness were translated using the ‘forward-backward’
7 procedure and were pilot tested prior to the present study to enhance translation clarity and
8 applicability. Anxiety and depression was measured by the Hungarian version of the Hospital
9 Anxiety and Depression Scale (HADS; [Muszbek and others 2006]). Items are scored from 0–
10 3 based on the selected response category, possible total scores for both anxiety (7 items) and
11 depression (7 items) range from 0 to 21.
12
13

14 To assess the internal consistency, Cronbach alpha values were calculated for each phenotype
15 construct in the present sample, and were found satisfactory (Aggression Total Score: 0.90;
16 Impulsiveness Total Score: 0.82; Anxiety Scale: 0.77; Depression Scale: 0.74). Internal
17 consistency measures of the four aggression factors were also sufficient (Verbal Aggression:
18 0.64; Anger: 0.84; Hostility: 0.80; Physical Aggression: 0.84).
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 **Marker Selection**

39 Common (higher than 5% minor allele frequency) polymorphic loci were selected from
40 dbSNP database using HapMap information about linkage disequilibrium to provide adequate
41 coverage of the entire WFS1 gene. LD blocks were defined by the “Four Gamete Rule” using
42 Haplowiev 4.2. At least two SNPs have been investigated in each block taking the
43 chromosomal localization also into consideration. Our goal was to generate a balanced
44 “physical coverage” so that no segments longer than 6 kb remained uncovered. Priority was
45 given to those SNPs which were referred in various association studies in connection with
46 diabetes mellitus and psychiatric disorders. SNPs, theoretically influencing microRNA
47 binding, were searched in the polymiRTS and Patrocles databases.
48
49
50
51
52
53
54
55
56
57
58
59
60

Kovacs-Nagy et al. WFS1 and aggression

DNA and genotyping

Genomic DNA was extracted from buccal swabs by Gentra purification kit (Minneapolis, US) according to manufacturer's instructions and quantified by fluorometry employing an intercalation assay (AccuBlue Broad Range dsDNA Quantitation Kit, Biotium, Hayward, CA). Typical DNA concentration was about 50 ng/ μ l (range: 15–200 ng/ μ l).

Genotyping was performed by an Open Array[®] real-time PCR platform of Applied Biosystems (Foster City, CA) based on the 5' nuclease assay with pre-designed and validated primers and allele-specific, fluorescent (TaqMan) probes immobilized to a solid surface. Approximately 100 ng (range: 30–150 ng) DNA was applied for each SNP subarray. The master mix, containing each dNTP and the AmpliTaq Gold DNA-polymerase, was supplied by the manufacturer. The thermocycle was carried out according to the manufacturer's instruction. Allele specific FAM and VIC fluorescent intensities were measured after the PCR amplification, genotypes were called using the TaqMan Genotyper v1.2 software. Call rates for SNPs on the OpenArray system was 88.87% in average (range: 69.37–96.00%). For quality control, three SNPs (rs9457, rs1046320 and rs1046322) were genotyped additionally on a conventional 96 well plate system with the commercially available TaqMan assay kits on a 7300 Real-Time PCR System (Applied Biosystems, Foster City, CA). Re-genotyping proved the reliability of the high-throughput OpenArray platform, as 99.1% agreement was observed comparing the two approaches, moreover increased the call rate of genotypes at these 3 loci. The genotype and allele frequencies obtained as a combination of data determined by the two genotyping platforms for the 801 individuals with valid data of questionnaires were used for association analysis between genotypes and phenotypic measures. Genotype frequency data for each SNP are given in Supplementary Table 1, minor allele frequencies (MAF) are shown in Table 1. The measured genotype frequencies

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 corresponded to the Hardy-Weinberg equilibrium for all SNPs (for p values see:
4
5 Supplementary Table 1). Call rates for each SNP were calculated as the sum of the genotypes
6
7 given in Supplementary Table 1 divided by the total number of participants with valid
8
9 phenotypic data (801).
10

14 **Luciferase constructs of the WFS1 3' UTR**

16 The entire 3' UTR region of the human WFS1 gene was amplified by PCR of a sample with
17
18 known rs1046322 (GG) genotype using the 5' TCG GCG **GAG CTC** GGA TGG TCC GCC
19
20 ACG AGG AGC 3' forward and 5' AAA GGA **AAG CTT** GCG CTG CAG GTT CCA CCA
21
22 GAG G 3' reverse primers. Bold letters indicate the *Sac* I and *Hind* III recognition sites in the
23
24 forward and reverse primer, respectively. PCR product was purified and cloned between the
25
26 *Sac* I and *Hind* III sites downstream to the firefly luciferase gene in pMIR-Report vector
27
28 (Ambion) by standard protocols (pMIR-G). Construct with WFS1 3' UTR possessing the
29
30 rs1046322 A allele (designated pMIR-A) was generated by quick change site directed
31
32 mutagenesis of the cloned pMIR-G, both constructs were confirmed by direct sequencing. A
33
34 control pMIR construct (pMIR-c) was also created by cloning a DNA-fragment with identical
35
36 length, however without the binding site of miR-668 downstream to the luciferase gene.
37
38
39
40
41
42

43 **Cell transfection and luciferase assay**

45 The Human Embryonic Kidney 293 (HEK293) cell line was cultured in DMEM medium
46
47 (Invitrogen–Gibco), supplemented with 10% bovine fetal serum (Lonza) at 37 °C in an
48
49 atmosphere of 5% CO₂. HEK293 cells were transferred into 24-well plate and incubated for
50
51 24 h before transfection. Subsequently, 0.05 µg of the luciferase reporter construct were
52
53 cotransfected with 0.2 µg β-galactosidase vector (Ambion), and 5 pmol miR-668 using
54
55 Lipofectamine™ 2000 (Invitrogen). Cells were collected 48 hours after transfection and
56
57
58
59
60

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 extracted by three consecutive freeze-thaw cycles followed by a centrifugation. Supernatants
4
5 were used for luciferase assay. Luminescence for luciferase as well as optical density for β -
6
7 galactosidase were measured by Varioskan Flash Multimode Reader, luciferase activity data
8
9 were normalized by β -galactosidase activities. All experiments were carried out in triplicates.
10
11 Statistical analysis of normalized reporter data was performed using one-way ANOVA
12
13 followed by Tukey-Kramer test (GraphPad InStat 3.05).
14
15
16
17

18 **Statistical Analysis**

19
20 Statistical analyses were carried out using SPSS 19.0 for Windows. Allele and genotype
21
22 frequency distributions were tested by chi-square analyses. Independent-Samples t-test was
23
24 applied for assessment of gender differences on self-report scales; their correlation with age
25
26 was also calculated. Genetic associations were tested by one way analyses of covariance
27
28 (ANCOVA) assuming a bi-allelic inheritance model and the recessivity of the minor allele.
29
30 Sex and age were used as covariates, since both were possible confounds. Significant
31
32 ($p < 0.001$) gender differences were found in aggression and anxiety (males exposed higher
33
34 aggression and lower anxiety scores). Impulsiveness and depression scores showed no
35
36 significant sex differences. A significant weak negative correlation was found between age
37
38 and aggression $r = -0.073$ ($p = 0.04$). Other measured scales did not correlate with age. In
39
40 association analyses nominal level of significance was corrected to $p < 0.0007$ based on the
41
42 Bonferroni correction for multiple testing: $p = 0.05$ was divided by 68, the number of analyses
43
44 performed (4 phenotypes \times 17 SNPs). Lewontin's D' as well as R^2 values of linkage
45
46 disequilibrium were determined using HaploView 4.2. [Barrett and others 2005].
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Kovacs-Nagy et al. WFS1 and aggression

RESULTS

Higher aggression in AA homozygotes of WFS1 rs1046322

Table 1 summarizes the most important information for the 17 studied WFS1 SNPs. The average distance between the selected SNPs was 1764 bp. Most of them are intronic, but three SNPs in the longest (8th) exon and three SNPs in the 3' UTR region were also investigated. The minor allele frequency (MAF) was calculated from the genotype distribution of each polymorphism (see Supplementary Table 1). As our aim was to study common allelic variants, we selected SNPs with $MAF \geq 0.05$.

Figure 1 shows the pairwise linkage disequilibrium (LD) estimates between the studied SNPs of the WFS1 gene region. R^2 (Figure 1A) as well as D' values (Figure 1B) were assessed, the most apparent differences between the two measures could be observed regarding rs28716718 and rs1801208 polymorphisms. R^2 values suggested practically no LD between these two SNPs and the other sites, whereas D' showed a high level of LD in this region. This finding can be demonstrated in that special form of LD, which can be characterized by the lack of exactly one haplotype, i.e. one allele is freely combined with both alleles of the adjacent site, however the other allele is always present together with one of the two variants of the neighboring polymorphism. The LD structure of WFS1 gene could be characterized by two (D' , block definition according to confidence intervals) or three (R^2 , block definition according to Four Gamete Rule) haploblocks as labeled in Figure 1. All SNPs were used for the subsequent association analysis.

Individual scores for aggression, impulsiveness, anxiety, and depression were derived from self-report questionnaire data. Mean values are listed for all three genotypes in Table 2. Based on the applied recessive model of the minor allele we compared mean values of minor homozygotes (mm) to those carrying the major allele (MM and Mm) for each polymorphism.

Kovacs-Nagy et al. WFS1 and aggression

Results of these one-way ANCOVAs are depicted in Table 2, sex and age were used as covariates.

The WFS1 rs1046322 showed significant association with Aggression [$F(1,797)=12.241$, $p=0.0005$, $\eta^2=0.015$, power=0.938], Impulsiveness [$F(1,791)=6.362$, $p=0.012$, $\eta^2=0.008$, power=0.712], Anxiety [$F(1,797)=7.801$, $p=0.005$, $\eta^2=0.010$, power=0.797] and Depression [$F(1,797)=5.074$, $p=0.025$, $\eta^2=0.006$, power=0.614]. These scales are highly related, correlation values reported below are all positive with a p value lower than 0.0001 Aggression showed a strong correlation with Impulsiveness ($r=0.43$) and Anxiety ($r=0.41$) and a somewhat weaker correlation with Depression ($r=0.37$). Impulsiveness was related to both Anxiety ($r=0.27$) and Depression ($r=0.25$). Correlation was the highest between Anxiety and Depression ($r=0.57$). Participants with the homozygous form of the rs1046322 minor allele (mm) reported significantly higher aggression, they were also more impulsive, more anxious and more depressed than those with genotypes including the major allele (see total scores for each self-report scale in Table 2).

Other WFS1 SNPs did not show significant association with Aggression, Impulsiveness or Anxiety scores. Depression was associated with two other WFS1 SNPs from the 1st haploblock: rs10002743 [$F(1,678)=6.777$, $p=0.009$, $\eta^2=0.01$, power=0.739] and rs6824720 [$F(1,673)=6.789$, $p=0.009$, $\eta^2=0.01$, power=0.740]. These SNPs are in strong linkage disequilibrium with each other, but not with the WFS1 rs1046322 polymorphism (see Figure 1).

Due to performing multiple comparisons, the accepted level of significance was corrected to rule out false positive results. From the nominally significant associations (labeled by bold in Table 2) only effect of the WFS1 rs1046322 on Aggression remained significant after the stringent Bonferroni correction (corrected p value was 0.0007). As highlighted in Table 2, participants with the AA genotype (homozygous for the minor allele) showed the highest

1 Kovacs-Nagy et al. WFS1 and aggression

2 aggression (mean total score = 85.8), whereas subjects with the AG and GG genotypes scored
3 notably lower (their mean total score was 66.3 and 66.4, respectively). It is important to note
4 that only 9 participants out of 801 carried this rare variant (see Supplementary Table 1 for
5 genotype distribution). To test consistency of this genetic effect we calculated mean
6 aggression raw scores for each factor, and compared the average Verbal Aggression, Anger,
7 Hostility and Physical Aggression values for the WFS1 rs1046322 genotypes. Figure 2 shows
8 that aggression level of those with the AA variant are notably higher in all four dimensions as
9 compared to aggression level of carriers of the major allele (AG and GG genotypes). This
10 result indicates that the applied genetic model fits data from the different dimensions of
11 aggression as well as the main construct.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 **Molecular-functional study of rs1046322**

29 The rs1046322 SNP which had a significant effect on aggression, impulsiveness, and
30 depression is located in the 3' UTR of the WFS1 gene (see Table 1) and does not fall into any
31 of the identified haploblocks (see Figure 1). This SNP was selected as a putative microRNA
32 binding site polymorphism (miRSNP). According to the PolymiRTS database, the minor
33 allele (A) generates a mismatch in the target site of the seed region of miR-668. To explore
34 whether the rs1046322 polymorphism alters the miR-668 binding efficiency under
35 experimental conditions, a firefly luciferase reporter vector containing the 3' UTR region of
36 the WFS1 gene (pMIR-G) was constructed, and its minor allelic variant was generated by site
37 directed mutagenesis (pMIR-A). The control construct (pMIR-c) possessed an independent
38 sequence with the same length, lacking any complementary region with miR-668. HEK293
39 cells were cotransfected with miR-668 and with pMIR-G, pMIR-A or pMIR-c, respectively,
40 transfection efficiency was determined by β -galactosidase reporter vector. The average values
41 of the relative luciferase activities are shown in Figure 3. Expression level was decreased by
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Kovacs-Nagy et al. WFS1 and aggression

miR-668 in the presence of both allelic variants (pMIR-A and pMIR-G) compared to the control (pMIR-c). The wild type 3' UTR variant (pMIR-G) inhibited the luciferase expression upto 28% of the control, while pMIR-A resulted only in a 47% silencing effect. ANOVA followed by post hoc analysis showed a statistically significant ($p < 0.05$) difference between the luciferase reporter activity of the pMIR-G and pMIR-A constructs (ANOVA $p = 0.0005$). Co-transfection of a control miRNA lacking a binding site in the WFS1 3' UTR did not result in the decrease of relative luciferase activity.

DISCUSSION

Loss of function mutation of WFS1 gene leads to serious neurodegeneration and early death, suggesting a central role of this protein in the function of the nervous system. Studies on the common WFS1 polymorphisms to date focused primarily on major depression, bipolar depression and suicidality. Kawamoto and colleagues [Kawamoto and others 2004] examined association of mood disorders and certain haplotypes of WFS1. The p.H611R (rs734312) missense exon polymorphism of the WFS1 gene has been linked to mood disorders [Koido and others 2005] in different populations [Zalsman and others 2009]. Suicidal behavior is often a symptom of an underlying psychological disorder, especially mood disorders. Sequeira and colleagues [Sequeira and others 2003] found higher frequency of the WFS1 611R/611R genotype in a small sample of completed suicide victims in the general population. This result, however, has not been confirmed by Zalsman and coworkers [Zalsman and others 2009]. A recent study of 4p chromosomal region identified various risk factors of completed suicide in males in several genes including the p.H611R (rs734312) of WFS1, but these results were not significant after the stringent corrections for multiple testing. Interestingly, the risk allele was A in this recent study in opposite to the results of

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 Sequeira et al. [Sequeira and others 2003] where the G allele emerged as the risk factor of
4
5 completed suicide.

6
7 Based on the above only a handful of association studies investigated the effect of certain
8
9 WFS1 gene variants on mood disorders and suicide using clinical samples. Results are rather
10
11 contradictory, which may be a consequence of the small sample size. It is also possible that
12
13 the most widely investigated WFS1 p.H611R (rs734312) polymorphism is only a marker,
14
15 being in linkage disequilibrium with another functional polymorphism of the WFS1 gene,
16
17 such as the microRNA binding site polymorphism (miRSNP) studied here, which has not
18
19 been assessed previously.
20
21

22
23 Recent investigations suggested that miRNAs are important regulators of protein synthesis
24
25 [Bartel 2004]. There is a short (6–8 basepairs) complementary region between the miRNA
26
27 binding site of the target mRNA and the “seed sequence” of miRNA. A SNP, which
28
29 influences the microRNA binding (miRSNP), probably has a significant effect on the rate of
30
31 target protein synthesis. Therefore, miRSNPs represent a novel class of functional
32
33 polymorphisms, playing an important role in the genetics and pharmacogenetics of complex
34
35 diseases [Bertino and others 2007]. For example, the miRSNP of dihydrofolate reductase gene
36
37 was associated with resistance to methotrexate, an anticancer drug [Mishra and others 2007].
38
39 There are also a few studies emphasizing the role of the miRSNPs in neuropsychiatric
40
41 disorders, such as Parkinson’s disease and Tourette syndrome [Sethupathy and Collins 2008].
42
43 The rs686 of the DRD1 gene, previously associated with nicotine dependence, was shown to
44
45 function as miRSNP [Huang and Li 2009]. Recently, it was demonstrated [Jensen and others
46
47 2009] that the rs13212041 in the serotonin receptor 1B gene, which was associated to
48
49 aggression among college students, disrupts a sequence critical for a miR-96 binding. An
50
51 additional evidence for this miRSNP was also provided by the same group showing an
52
53
54
55
56
57
58
59
60

1 Kovacs-Nagy et al. WFS1 and aggression

2 association of a haplotype involving rs13212041 with self-reported anger and hostility in
3 young men [Conner and others 2010].
4

5 Searching in various databases we found two putative miRSNPs (rs9457, rs1046322) in the
6 WFS1 gene. According to our best knowledge, these common variants have not been tested
7 previously in psychogenetic or diabetes mellitus related association studies. We found that
8 one of these putative miRSNPs, the rs1046322 was associated with aggression and related
9 dimensions. Our *in silico* studies indicated that the A allele weakens the binding of miR-668.
10 We also provided experimental evidence about the effect of this SNP on miR-668 binding:
11 coexpressing the cloned WFS1 3' UTR possessing A or G alleles with miR-668 resulted in
12 changes of luciferase reporter activity. These results suggest that the rs1046322 SNP has an
13 important molecular function in the regulation of WFS1 gene expression and should be
14 included into further association studies of the WFS1 gene. We hypothesize that the negative
15 regulation of wolframin protein synthesis is less pronounced in the presence of the minor
16 allele of the rs1046322 leading to an overexpression of wolframin and an imbalance in ER
17 stress regulation. Limitation of our hypotheses is the lack of *in vitro* or *in vivo* studies that
18 would support the role of miR-668 in the regulation of the wolframin protein synthesis.
19

20 Studies to date did not investigate association of WFS1 polymorphisms with impulsivity,
21 aggression or mood characteristics within the non-clinical spectrum. According to the
22 psychiatric literature, these dimensions are interrelated [Apter and others 1990]. Results of the
23 correlation analysis from our present study also showed that Aggression was in a strong
24 correlation with Impulsiveness ($r=0.43$, $p<0.0001$) and Anxiety ($r=0.41$, $p<0.0001$), although
25 they were assessed with different tools and in populations of different type and size.
26

27 The aim of the present study was to detect possible genetic effects of a comprehensive set of
28 WFS1 polymorphisms, involving putative miRSNPs, on psychological dimensions using self-
29 report data from 801 non-related, healthy Caucasians. Our set of 17 WFS1 SNPs included the
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 p.H611R (rs734312) polymorphism that was previously associated with mood disorders,
4
5 however, we did not find any association with impulsivity, aggression or mood characteristics
6
7 in our large, non-clinical sample. Results from our association analyses revealed a novel risk
8
9 factor of the impulsive-aggressive tendencies within the non-clinical range: The WFS1
10
11 rs1046322 polymorphism, which was identified here as a miRSNP, showed significant
12
13 association with all the measured scales: Aggression ($p=0.0005$), Impulsiveness ($p=0.0119$),
14
15 Anxiety ($p=0.0053$) and Depression ($p=0.0246$). Subjects homozygous for the minor allele
16
17 (AA) showed higher aggression, higher impulsivity, higher anxiety and higher depression as
18
19 compared to those without the rare risk genotype. It is important to note, that when correcting
20
21 the level of significance to rule out false positive results of multiple comparisons only the
22
23 effect of rs1046322 on Aggression remained significant. Although application of the stringent
24
25 Bonferroni correction is an important principle in current association studies, we would like to
26
27 emphasize that most of our nominally significant associations emerged between the WFS1
28
29 rs1046322 and the other three dimensions (Impulsiveness, Anxiety and Depression)
30
31 interrelated with Aggression suggesting a significant and complex effect of this miRSNP on
32
33 the behavior.
34
35
36

37
38 Dimensions of aggression, impulsivity, anxiety, and depression are correlated and are linked
39
40 to certain psychopathologies, but also cause disturbances in the emotions, decision making
41
42 and self-regulation within the non-clinical spectrum. Association studies highlight the role of
43
44 various genetic and environmental predisposing factors in the polygenic model of these
45
46 behaviors and acknowledge the importance of understanding interaction effects between these
47
48 components through the study of psychiatric patients as well as the non-clinical population
49
50 [Courtet and others 2011]. To our knowledge, this is the first association study of a validated
51
52 microRNA binding site polymorphism in the WFS1 gene. This functional polymorphism was
53
54 shown as a genetic factor of individual variations in aggression.
55
56
57
58
59
60

Kovacs-Nagy et al. WFS1 and aggression

1
2
3 Limitation of the present study is the applied self-report method for portraying such a
4
5 complex dimension as aggression, however, this tool was sufficient for large-scale
6
7 assessment. We tested 801 subjects, but there were only 9 subjects (5 males and 4 females)
8
9 homozygous with the risk allele (AA) of the microRNA binding site polymorphism
10
11 (rs1046322) of the WFS1 gene, although the call rate for this SNP was 100% as a result of
12
13 double genotyping (see: Supplementary Table 1). Recessivity of the risk allele might be
14
15 explained by the identified molecular effect of the rs1046322 minor A allele. If microRNA
16
17 serves as a fine tuning of gene expression, the ER balance could be disturbed only in the
18
19 presence of both risk alleles. These subjects showed substantially higher levels of aggression
20
21 (mean value was 86(±22) as compared to the overall aggression average: 67(±16). Due to the
22
23 small number of subjects with the risk genotype replication of these findings would be
24
25 important in an independent non-clinical population. Investigation of this novel functional
26
27 polymorphism in clinical samples with mood disorders and suicide would also be essential.
28
29
30
31
32

33 34 **ACKNOWLEDGEMENTS**

35
36 This work was supported by the Hungarian Scientific Research Funds: OTKA K81466 by
37
38 Maria Sasvari-Szekely, and OTKA K100845 by Anna Szekely and connected to the scientific
39
40 program of the NIH R03 TW007656 Fogarty International Research grant awarded to Maria
41
42 Sasvari-Szekely. This project was also supported by the János Bolyai Research Scholarship
43
44 (BO/00089/10/5) of the Hungarian Academy of Sciences. Corresponding author has reviewed
45
46 publication policy and authors have no conflict of interest to declare.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Kovacs-Nagy et al. WFS1 and aggression

LEGENDS TO THE FIGURES

Figure 1. Linkage disequilibrium plots of the WFS1 SNPs.

A: R^2 (%) measure of linkage disequilibrium between any pair of SNPs, as determined by HaploView, version 4.2. The gray-scale spectrum indicates pairwise R^2 values ranging from black ($R^2=1$) to white ($R^2=0$). B: Pairwise LD values expressed by D' (HaploView v4.2). Red squares indicate 100% LD.

Figure 2. The effect of the rs1046322 WFS1 SNP in the Buss-Perry Aggression Scale.

Mean raw score values with standard errors of mean for the 4 aggression factors are shown.

Figure 3. The silencing effect of the rs1046322

HEK293 cells were cotransfected with miR-668 and with pMIR-G, pMIR-A or pMIR-c construct, respectively. Relative luciferase activities (mean \pm SD) from three independent experiments are shown. ANOVA showed a statistically significant difference between the luciferase activity of pMIR-A and pMIR-G constructs ($p = 0.0005$).

Kovacs-Nagy et al. WFS1 and aggression

REFERENCES

- Apter A, van Praag HM, Plutchik R, Sevy S, Korn M, Brown SL. 1990. Interrelationships among anxiety, aggression, impulsivity, and mood: a serotonergically linked cluster? *Psychiatry Res* 32(2):191-199.
- Barrett JC, Fry B, Maller J, Daly MJ. 2005. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21(2):263-265.
- Barrett TG, Bunday SE, Macleod AF. 1995. Neurodegeneration and diabetes: UK nationwide study of Wolfram (DIDMOAD) syndrome. *Lancet* 346(8988):1458-1463.
- Bartel DP. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116(2):281-297.
- Bertino JR, Banerjee D, Mishra PJ. 2007. Pharmacogenomics of microRNA: a miRSNP towards individualized therapy. *Pharmacogenomics* 8(12):1625-1627.
- Buss AH, Perry M. 1992. The aggression questionnaire. *J Pers Soc Psychol* 63(3):452-459.
- Conner TS, Jensen KP, Tennen H, Furneaux HM, Kranzler HR, Covault J. 2010. Functional polymorphisms in the serotonin 1B receptor gene (HTR1B) predict self-reported anger and hostility among young men. *Am J Med Genet B Neuropsychiatr Genet* 153B(1):67-78.
- Courtet P, Gottesman, II, Jollant F, Gould TD. 2011. The neuroscience of suicidal behaviors: what can we expect from endophenotype strategies? *Transl Psychiatry* 1.
- Fawcett KA, Wheeler E, Morris AP, Ricketts SL, Hallmans G, Rolandsson O, Daly A, Wasson J, Permutt A, Hattersley AT, Glaser B, Franks PW, McCarthy MI, Wareham NJ, Sandhu MS, Barroso I. 2010. Detailed investigation of the role of common and low-frequency WFS1 variants in type 2 diabetes risk. *Diabetes* 59(3):741-746.

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 Fonseca SG, Fukuma M, Lipson KL, Nguyen LX, Allen JR, Oka Y, Urano F. 2005. WFS1 is
4
5 a novel component of the unfolded protein response and maintains homeostasis of the
6
7 endoplasmic reticulum in pancreatic beta-cells. *J Biol Chem* 280(47):39609-39615.

8
9
10 Fonseca SG, Ishigaki S, Oslowski CM, Lu S, Lipson KL, Ghosh R, Hayashi E, Ishihara H,
11
12 Oka Y, Permutt MA, Urano F. 2010. Wolfram syndrome 1 gene negatively regulates
13
14 ER stress signaling in rodent and human cells. *J Clin Invest* 120(3):744-755.

15
16 Huang W, Li MD. 2009. Differential allelic expression of dopamine D1 receptor gene
17
18 (DRD1) is modulated by microRNA miR-504. *Biol Psychiatry* 65(8):702-705.

19
20
21 Inoue H, Tanizawa Y, Wasson J, Behn P, Kalidas K, Bernal-Mizrachi E, Mueckler M,
22
23 Marshall H, Donis-Keller H, Crock P, Rogers D, Mikuni M, Kumashiro H, Higashi K,
24
25 Sobue G, Oka Y, Permutt MA. 1998. A gene encoding a transmembrane protein is
26
27 mutated in patients with diabetes mellitus and optic atrophy (Wolfram syndrome). *Nat*
28
29 *Genet* 20(2):143-148.

30
31
32 Ishihara H, Takeda S, Tamura A, Takahashi R, Yamaguchi S, Takei D, Yamada T, Inoue H,
33
34 Soga H, Katagiri H, Tanizawa Y, Oka Y. 2004. Disruption of the WFS1 gene in mice
35
36 causes progressive beta-cell loss and impaired stimulus-secretion coupling in insulin
37
38 secretion. *Hum Mol Genet* 13(11):1159-1170.

39
40
41 Jensen KP, Covault J, Conner TS, Tennen H, Kranzler HR, Furneaux HM. 2009. A common
42
43 polymorphism in serotonin receptor 1B mRNA moderates regulation by miR-96 and
44
45 associates with aggressive human behaviors. *Mol Psychiatry* 14(4):381-389.

46
47
48 Kato T, Ishiwata M, Yamada K, Kasahara T, Kakiuchi C, Iwamoto K, Kawamura K, Ishihara
49
50 H, Oka Y. 2008. Behavioral and gene expression analyses of Wfs1 knockout mice as a
51
52 possible animal model of mood disorder. *Neurosci Res* 61(2):143-158.

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 Kawamoto T, Horikawa Y, Tanaka T, Kabe N, Takeda J, Mikuni M. 2004. Genetic variations
4
5 in the WFS1 gene in Japanese with type 2 diabetes and bipolar disorder. *Mol Genet*
6
7 *Metab* 82(3):238-245.

8
9
10 Kesner Y, Zohar J, Merenlender A, Gispan I, Shalit F, Yadid G. 2009. WFS1 gene as a
11
12 putative biomarker for development of post-traumatic syndrome in an animal model.
13
14 *Mol Psychiatry* 14(1):86-94.

15
16 Koido K, Koks S, Nikopensus T, Maron E, Altmae S, Heinaste E, Vabrit K, Tammekivi V,
17
18 Hallast P, Kurg A, Shlik J, Vasar V, Metspalu A, Vasar E. 2005. Polymorphisms in
19
20 wolframin (WFS1) gene are possibly related to increased risk for mood disorders. *Int J*
21
22 *Neuropsychopharmacol* 8(2):235-244.

23
24
25 Lemonde S, Turecki G, Bakish D, Du L, Hrdina PD, Bown CD, Sequeira A, Kushwaha N,
26
27 Morris SJ, Basak A, Ou XM, Albert PR. 2003. Impaired repression at a 5-
28
29 hydroxytryptamine 1A receptor gene polymorphism associated with major depression
30
31 and suicide. *J Neurosci* 23(25):8788-8799.

32
33
34 Luuk H, Plaas M, Raud S, Innos J, Sutt S, Lasner H, Abramov U, Kurrikoff K, Koks S, Vasar
35
36 E. 2009. Wfs1-deficient mice display impaired behavioural adaptation in stressful
37
38 environment. *Behav Brain Res* 198(2):334-345.

39
40 M. M, Lesperance. 2010. Human Genetics Laboratory Low-Frequency Hearing Loss WFS1
41
42 Gene Mutation and Polymorphism Database.

43
44
45 http://www.khri.med.umich.edu/research/lesperance_lab/low_freq.php.

46
47 Mishra PJ, Humeniuk R, Longo-Sorbello GS, Banerjee D, Bertino JR. 2007. A miR-24
48
49 microRNA binding-site polymorphism in dihydrofolate reductase gene leads to
50
51 methotrexate resistance. *Proc Natl Acad Sci U S A* 104(33):13513-13518.

52
53
54 Must A, Koks S, Vasar E, Tasa G, Lang A, Maron E, Vali M. 2009. Common variations in 4p
55
56 locus are related to male completed suicide. *Neuromolecular Med* 11(1):13-19.

1 Kovacs-Nagy et al. WFS1 and aggression

2
3 Muszbek K, Szekely A, Balogh EM, Molnar M, Rohanszky M, Ruzsa A, Varga K, Szollosi
4
5 M, Vadasz P. 2006. Validation of the Hungarian translation of Hospital Anxiety and
6
7 Depression Scale. *Qual Life Res* 15(4):761-766.
8

9
10 Osowski CM, Urano F. 2011. The binary switch that controls the life and death decisions of
11
12 ER stressed beta cells. *Current opinion in cell biology* 23(2):207-215.
13

14 Patton JH, Stanford MS, Barratt ES. 1995. Factor structure of the Barratt impulsiveness scale.
15
16 *J Clin Psychol* 51(6):768-774.
17

18 Rigoli L, Lombardo F, Di Bella C. 2011. Wolfram syndrome and WFS1 gene. *Clin Genet*
19
20 79(2):103-117.
21

22
23 Sequeira A, Kim C, Seguin M, Lesage A, Chawky N, Desautels A, Tousignant M, Vanier C,
24
25 Lipp O, Benkelfat C, Rouleau G, Turecki G. 2003. Wolfram syndrome and suicide:
26
27 Evidence for a role of WFS1 in suicidal and impulsive behavior. *Am J Med Genet B*
28
29 *Neuropsychiatr Genet* 119B(1):108-113.
30

31
32 Sethupathy P, Collins FS. 2008. MicroRNA target site polymorphisms and human disease.
33
34 *Trends Genet* 24(10):489-497.
35

36 Swift M, Swift RG. 2000. Psychiatric disorders and mutations at the Wolfram syndrome
37
38 locus. *Biol Psychiatry* 47(9):787-793.
39

40 Swift RG, Sadler DB, Swift M. 1990. Psychiatric findings in Wolfram syndrome
41
42 homozygotes. *Lancet* 336(8716):667-669.
43

44
45 Takei D, Ishihara H, Yamaguchi S, Yamada T, Tamura A, Katagiri H, Maruyama Y, Oka Y.
46
47 2006. WFS1 protein modulates the free Ca(2+) concentration in the endoplasmic
48
49 reticulum. *FEBS Lett* 580(24):5635-5640.
50

51
52 Yurimoto S, Hatano N, Tsuchiya M, Kato K, Fujimoto T, Masaki T, Kobayashi R, Tokumitsu
53
54 H. 2009. Identification and characterization of wolframin, the product of the wolfram
55
56
57
58
59
60


1 Kovacs-Nagy et al. WFS1 and aggression

2
3 syndrome gene (WFS1), as a novel calmodulin-binding protein. *Biochemistry*
4
5 48(18):3946-3955.
6

7 Zalsman G, Mann MJ, Huang YY, Oquendo MA, Brent DA, Burke AK, Ellis SP, Mann JJ.

8
9 2009. Wolframin gene H611R polymorphism: no direct association with suicidal
10
11 behavior but possible link to mood disorders. *Prog Neuropsychopharmacol Biol*
12
13 *Psychiatry* 33(4):707-710.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review




Linkage disequilibrium plots of the WFS1 SNPs.


A: R^2 (%) measure of linkage disequilibrium between any pair of SNPs, as determined by HaploView, version 4.2. The gray-scale spectrum indicates pairwise R^2 values ranging from black ($R^2=1$) to white ($R^2=0$). B: Pairwise LD values expressed by D' (HaploView v4.2). Red squares indicate 100% linkage.

319x475mm (96 x 96 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



The effect of the rs1046322 WFS1 SNP in the Buss-Perry Aggression Scale. Mean raw score values with standard errors of mean for the 4 aggression factors are shown. 210x220mm (96 x 96 DPI)



HEK293 cells were cotransfected with miR-668 and with pMIR-G, pMIR-A or pMIR-c construct, respectively. Relative luciferase activities (mean \pm SD) from three independent experiments are shown. ANOVA showed a statistically significant difference between the luciferase activity of pMIR-A and pMIR-G constructs ($p = 0.0005$).

260x196mm (96 x 96 DPI)

Kovacs-Nagy et al. WFS1 and aggression

Table 1. Main characteristics of the selected WFS1 SNPs.

dbSNP number	Genomic position	Inter-SNP distance (bp)	Type	MAF*
rs10002743	6276581		intron 2	0.191
rs6824720	6278647	2066	intron 2	0.159
rs752854	6281961	3314	intron 3	0.310
rs4689393	6287241	5280	intron 3	0.435
rs10010131	6292915	5674	intron 4	0.391
rs13147655	6293474	559	intron 6	0.384
rs4467645	6294305	831	intron 7	0.394
rs13128674	6294517	212	intron 7	0.409
rs6446482	6295693	1176	intron 7	0.398
rs4689395	6295985	292	intron 7	0.359
rs28716718	6301910	5925	intron 8	0.053
rs1801208	6302889	979	exon 8 (p.R456H)	0.050
rs734312	6303354	465	exon 8 (p.H611R)	0.462
rs1046316	6304087	733	exon 8 (p.S855S)	0.341
rs1046320	6304344	257	3' UTR	0.403
rs1046322	6304448	104	3' UTR	0.099
rs9457	6304799	351	3' UTR	0.433

Kovacs-Nagy et al. WFS1 and aggression

Table 2. Association of WFS1 genotypes with self-report measures

Mean values of the total score are presented for each genotype

*significant with Bonferroni correction

WFS1 SNP	AGGRESSION (BUSS-PERRY)				IMPULSIVITY (BARRAT)				ANXIETY (HADS)				DEPRESSION (HADS)			
	(MM)	(Mm)	(mm)	p	(MM)	(Mm)	(mm)	p	(MM)	(Mm)	(mm)	p	(MM)	(Mm)	(mm)	p
rs10002743	66.2	68.1	66.1	0.8711	58.3	59.8	61.1	0.2133	5.7	5.7	6.5	0.3196	2.5	2.7	3.8	0.0094
rs6824720	66.4	67.1	64.1	0.5821	58.8	59.7	61.6	0.2611	5.7	5.6	6.7	0.3235	2.5	2.7	4.1	0.0094
rs752854	67.5	66.5	66.8	0.8727	58.6	58.8	57.9	0.5744	5.7	5.5	5.3	0.6456	2.7	2.6	2.6	0.9468
rs4689393	66.2	67.5	66.6	0.9975	58.0	59.4	59.4	0.5180	5.5	5.7	5.6	0.9226	2.4	2.7	2.7	0.6312
rs10010131	66.8	66.2	64.8	0.3542	58.1	59.8	58.5	0.6086	5.8	5.9	5.4	0.2070	2.6	2.5	2.7	0.7347
rs13147655	66.6	67.4	66.1	0.7131	57.9	59.4	58.7	0.9993	5.7	5.7	5.5	0.4438	2.6	2.7	2.8	0.7246
rs4467645	66.5	67.2	66.2	0.7985	58.3	59.4	58.9	0.9656	5.6	5.8	5.6	0.6563	2.6	2.6	2.7	0.6657
rs13128674	66.9	67.4	67.2	0.8759	58.1	58.8	59.0	0.6345	5.5	5.6	5.6	0.8510	2.7	2.7	2.9	0.6256
rs6446482	66.3	67.3	67.0	0.8364	57.8	59.4	59.1	0.6555	5.5	5.7	5.6	0.8530	2.5	2.6	2.8	0.4933
rs4689395	66.6	66.6	67.0	0.8654	58.6	59.2	59.0	0.9261	5.7	5.7	5.5	0.7063	2.6	2.6	2.7	0.9011
rs28716718	66.5	66.8	72.2	0.4960	58.8	59.9	57.0	0.6773	5.6	5.7	7.0	0.5337	2.6	2.8	3.8	0.3753
rs1801208	66.6	68.5	72.5	0.5980	59.0	58.0	59.0	0.938	5.6	5.9	7.8	0.1544	2.6	2.7	2.8	0.9045
rs734312	66.0	67.4	66.5	0.9215	58.7	58.8	58.9	0.8644	5.6	5.7	5.7	0.9185	2.6	2.6	2.8	0.4407
rs1046316	67.3	67.5	66.4	0.8790	58.8	58.2	59.7	0.3845	5.5	5.6	5.6	0.9628	2.7	2.7	2.8	0.8360
rs1046320	66.9	66.5	66.3	0.8793	58.1	59.3	58.4	0.6323	5.7	5.8	5.3	0.1697	2.7	2.7	2.5	0.5612
rs1046322	66.4	66.3	85.8	0.0005*	58.6	59.1	66.9	0.0119	5.6	5.8	8.8	0.0053	2.6	2.7	4.7	0.0246
rs9457	66.3	67.3	65.8	0.5462	57.8	59.2	58.2	0.6403	5.4	5.8	5.5	0.6373	2.6	2.8	2.7	0.9249