

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/84242>

Please be advised that this information was generated on 2017-12-06 and may be subject to change.

On Superselection Theory of Quantum Fields in Low Dimensions*

Michael Müger[†]

Institute of Mathematics, Astrophysics and Particle Physics.
Radboud University, Nijmegen, The Netherlands. E-mail: mueger@math.ru.nl

September 14, 2009

Abstract

We discuss finite local extensions of quantum field theories in low space time dimensions in connection with categorical structures and the question of modular invariants in conformal field theory, also touching upon purely mathematical ramifications.

1 Introduction: Superselection theory in Local Quantum Theory

The aim of this contribution is to briefly review some aspects of the ‘quantum symmetry’ question in low dimensional (typically conformal) quantum field theory (QFT), from the point of view of algebraic quantum field theory. The latter (also called ‘local quantum physics’) arguably is the most successful approach to the axiomatic study of QFT in Minkowski space (or on more general pseudo-Riemannian manifolds). Due to space constraints we must refer to [16] or [17] for typical lists of axioms and recall only that a QFT is a map $\mathcal{O} \mapsto \mathcal{A}(\mathcal{O})$ assigning to every decent spacetime region \mathcal{O} an algebra of operators $\mathcal{A}(\mathcal{O})$ satisfying requirements like spacelike commutativity, covariance w.r.t. the group of spacetime symmetries, irreducibility, etc. Under these assumptions, augmented by Haag duality, it was shown in [9] that the category $\text{DHR}(\mathcal{A})$ of ‘localized representations’ (with finite statistical dimension) of a QFT \mathcal{A} in Minkowski space of dimension $\geq 2+1$ is a rigid semisimple symmetric tensor $*$ -category with simple unit. Later it was proven [11] that every such category is the representation category of a unique compact group G . Furthermore [12], there is a (graded local) QFT \mathcal{F} with the following properties: (1) It is acted upon by G and $\mathcal{F}^G \cong \mathcal{A}$. (2) The category $\text{DHR}(\mathcal{F})$ is trivial [4]. (This must be modified suitably when \mathcal{A} has fermionic representations). (3) Upon restriction to $\mathcal{F}^G \cong \mathcal{A}$, the representation space of \mathcal{F} decomposes into a direct sum of irreps, which are precisely the localized representations of \mathcal{A} . (4) Every local extension $\mathcal{B} \supset \mathcal{A}$ is isomorphic to \mathcal{F}^H for some closed subgroup $H \subset G$. As argued in [17, Section 10.6], the above amounts to a very satisfactory Galois theory for QFTs. [17] also provides an alternative and more transparent approach to the construction of \mathcal{F} , based on unpublished work by Roberts and Deligne’s independent approach [7] to proving $\mathcal{C} \simeq \text{Rep } G$ for symmetric categories, cf. also [26].

2 What changes in low dimensional spacetimes?

When one considers QFTs in Minkowski space of dimension $1+1$ or just on the light ray \mathbb{R} , the category $\text{DHR}(\mathcal{A})$ can only be proven to be braided [14, I]. In [19] this was strengthened considerably, in that it was shown that $\text{DHR}(\mathcal{A})$ is maximally non-symmetric, to wit a modular category in the sense of Turaev, for the important class of ‘completely rational’ models (a rigorous axiomatization of rational chiral conformal QFTs). In this situation it is clear that $\text{DHR}(\mathcal{A})$

*Invited lecture at the XVIth International Congress on Mathematical Physics. Prague, August 3-8, 2009.

[†]Partially based on joint work with A. Kitaev and with A. Davydov, D. Nikshych and V. Ostrik.

cannot be equivalent to a representation category of a group. Another advantage of the alternative construction of the field net [17] is that it also applies in low dimensions *provided* a fiber functor from $\text{DHR}(\mathcal{A})$ to the category of Hilbert spaces is given: Roberts' construction of the field net as described in [17] goes through essentially unchanged, and appealing to a version of Woronowicz's Tannakian theorem [29] one obtains a discrete algebraic quantum group acting on \mathcal{F} and an R-matrix describing the spacelike commutation relations of \mathcal{F} (which will be far from local), cf. [28].

However, in general a fiber functor for $\text{DHR}(\mathcal{A})$ will not exist (as can be proven e.g. when the category is finite and contains objects of non-integer dimension). At least in the case of a finite category, one can always get around this problem obtaining [30] a weak Hopf algebra, acting on the reduced field bundle of [14, II], but it is not clear that this is useful: The weak Hopf algebra is not uniquely determined by $\text{DHR}(\mathcal{A})$ so that it does not have an intrinsic physical meaning. (Furthermore, the inclusion $\mathcal{A} \subset \mathcal{F}$ has the undesirable property of being reducible.) For many purposes it seems better to resign oneself to considering the category $\text{DHR}(\mathcal{A})$ itself as the fundamental structure. The rest of this note is an attempt to convince the reader of the feasibility, even elegance, of such an approach. Cf. [28] for details.

3 Categorical analysis of local extensions

The first question we address is the classification of extensions of a QFT, in particular local ones and their representation categories. By an extension $\mathcal{B} \supset \mathcal{A}$ of a QFT we mean an inclusion preserving assignment $\mathcal{O} \mapsto \mathcal{B}(\mathcal{O}) \supset \mathcal{A}(\mathcal{O})$ satisfying covariance and irreducibility, but not necessarily locality. If \mathcal{B} is local we call it a local extension. In [23] it was shown that there is a bijection between (unitary equivalence classes of) extensions $\mathcal{B} \supset \mathcal{A}$ that are finite (i.e. $[\mathcal{B}(\mathcal{O}) : \mathcal{A}(\mathcal{O})]$ is independent of \mathcal{O} and finite) and (equivalence classes of) Q-systems in the category $\text{DHR}(\mathcal{A})$. (A Q-system in a $*$ -category is essentially a monoid/algebra (Γ, m, η) such that $(\Gamma, m, \eta, m^*, \eta^*)$ is a Frobenius algebra.) Furthermore, \mathcal{B} is local iff (Γ, m, η) is commutative, i.e. $m \circ c(\Gamma, \Gamma) = m$, where c is the braiding.¹ When this is the case, it is natural to ask for a determination of the category $\text{DHR}(\mathcal{B})$ in terms of $\text{DHR}(\mathcal{A})$ and (Γ, m, η) . The answer is given by:

3.1 THEOREM [28] *Let $\mathcal{B} \supset \mathcal{A}$ be the extension of the completely rational theory \mathcal{A} corresponding to the commutative algebra (Γ, m, η) in $\text{DHR}(\mathcal{A})$. Then there is an equivalence*

$$\text{DHR}(\mathcal{B}) \simeq \Gamma - \text{Mod}_{\text{DHR}(\mathcal{A})}^0$$

of braided tensor categories. In particular, $\dim(\text{DHR}(\mathcal{B})) = \dim(\text{DHR}(\mathcal{A}))/d(\mathcal{A})^2$.

3.2 REMARK 1. Here, $\Gamma - \text{Mod}_{\text{DHR}(\mathcal{A})}^0$ is the full subcategory of the category $\Gamma - \text{Mod}_{\text{DHR}(\mathcal{A})}$ of left Γ -modules consisting of the objects (X, μ) satisfying $\mu \circ c(X, \Gamma) \circ c(\Gamma, X) = \mu$. These 'dyslexic' or 'local' modules form a braided tensor category in a canonical way, cf. [31]. While the proof of Theorem 3.1 is too long to be exhibited here, it can be given in a few pages including the prerequisites. It uses results of [1] on α -induction to obtain a full and faithful braided tensor functor $\Gamma - \text{Mod}_{\text{DHR}(\mathcal{A})}^0 \rightarrow \text{DHR}(\mathcal{B})$ and [19] together with the result $\dim \Gamma - \text{Mod}_{\mathcal{C}}^0 = \dim \mathcal{C}/d(\Gamma)^2$ from [20] to conclude essential surjectivity.

2. The analogous result in the context of vertex operator algebras appears in [20], but many hard VOA technicalities have not yet been worked out.

3. The above result also holds in $\geq 2 + 1$ dimensions where the representation categories are symmetric and every Γ -module is local. But in this case, $\text{DHR}(\mathcal{A})$ is equivalent to $\text{Rep } G$ and commutative algebras in $\text{Rep } G$ correspond to subgroups of $H \subset G$, cf. [30]. Thus we recover (at least for finite extensions) the Galois correspondence mentioned in the introduction.

4. In view of the above result, almost all questions concerning finite local extensions of QFTs and their representation categories can be reduced to purely categorical, i.e. algebraic considerations. An obvious exception to this is the inverse problem, to wit the question which categories are realized in some QFT model. (The analogous question in the group theoretic situation $d \geq 2 + 1$ was settled in [10].)

¹From now on we will simply write 'commutative algebra' with the understanding that $\dim \text{Hom}(\mathbf{1}, \Gamma) = 1$ and that the algebra is 'strongly separable symmetric Frobenius' or 'étale' [5].

Recall that in $d \geq 2 + 1$ dimensions, the extension \mathcal{F} has trivial representation category. In low dimensions it is *not* true that a local extension with this property always exists, since in view of Theorem 3.1, we have:

3.3 COROLLARY *A completely rational theory \mathcal{A} admits a local extension with trivial representation category iff $\mathcal{C} = \text{DHR}(\mathcal{A})$ contains a commutative algebra Γ such that $\Gamma - \text{Mod}_{\mathcal{C}}^0$ is trivial, equivalently $d(\Gamma)^2 = \dim \mathcal{C}$.*

(Notice that every commutative algebra Γ in a modular category \mathcal{C} satisfies $d(\Gamma)^2 \leq \dim \mathcal{C}$, cf. [20].) The question when such a commutative algebra exists is answered by the following remarkable

3.4 THEOREM *A commutative algebra Γ in a modular category gives rise to a braided equivalence $Z(\Gamma - \text{Mod}_{\mathcal{C}}) \simeq \mathcal{C} \boxtimes \widetilde{\Gamma - \text{Mod}_{\mathcal{C}}^0}$. Thus if $\Gamma - \text{Mod}_{\mathcal{C}}^0$ is trivial, equivalently $d(\Gamma)^2 = \dim \mathcal{C}$, then $\mathcal{C} \simeq Z(\mathcal{D})$, where $\mathcal{D} = \Gamma - \text{Mod}_{\mathcal{C}}$ is a fusion category. Conversely, if \mathcal{D} is a fusion category, then $Z(\mathcal{D})$ contains a commutative algebra Γ with $d(\Gamma)^2 = (\dim \mathcal{D})^2 = \dim Z(\mathcal{D})$.*

Here $Z(\cdot)$ is the categorical center of a tensor category due to Drinfeld, Majid, Joyal and Street which was proven [24] to be modular for every fusion category \mathcal{D} with $\dim \mathcal{D} \neq 0$. Theorem 3.4 was obtained by Kitaev and the author [21]. The first half was discovered independently in [13], and results related to the second half are obtained in [3]. Combining Corollary 3.3 and Theorem 3.4 we have:

3.5 COROLLARY *A completely rational theory \mathcal{A} admits a local extension with trivial representation category iff the modular category $\text{DHR}(\mathcal{A})$ is a Drinfeld center.*

The \Rightarrow direction actually seems to be more useful. E.g., one can use it to prove:

3.6 COROLLARY *Let \mathcal{B} be a completely rational theory with trivial representation category acted upon freely by a finite group G , and let $\mathcal{A} = \mathcal{B}^G$ be the ‘orbifold’ theory. Then there is a unique class $[\omega] \in H^3(G, \mathbb{T})$ such that $\text{DHR}(\mathcal{A}) \simeq D^\omega(G) - \text{Mod}$, where $D^\omega(G)$ is the twisted quantum double of G [8].*

3.7 REMARK A more general analysis of orbifold QFTs $\mathcal{A} = \mathcal{B}^G$ without the triviality assumption on $\text{DHR}(\mathcal{B})$ was given in [25]. However, this did not yield the above corollary since a certain coherence theorem for braided crossed G -categories was missing. But see [27].

Even when \mathcal{A} admits no local extension $\mathcal{B} \supset \mathcal{A}$ with trivial $\text{DHR}(\mathcal{B})$, it follows easily from the above results that maximal local extensions always exist and that every local extension embeds into a maximal one. Contrary to the situation for $d \geq 2 + 1$, it is not true that there is a unique (up to unitary equivalence) maximal local extension. (E.g., if \mathcal{A} is completely rational with $\text{DHR}(\mathcal{A}) \simeq D(G) - \text{Mod}$, then there exist at least two non-isomorphic commutative algebras $\Gamma, \widehat{\Gamma}$ such that $d(\Gamma) = d(\widehat{\Gamma}) = |G|$, giving rise to local extensions that are not unitarily equivalent.)

However, by a result of [5], cf. Section 5, all maximal local extensions have braided equivalent representation categories. In particular, if a local extension with trivial representation category exists, then every local extension embeds into one with trivial representation category.

4 Connections with Rehren’s approach to modular invariants

It is well known that a modular category \mathcal{C} gives rise to a finite dimensional representation of the modular group $SL(2, \mathbb{Z})$, acting on the complexified Grothendieck group of \mathcal{C} . Given two modular categories $\mathcal{C}_L, \mathcal{C}_R$, a *modular invariant* used to be defined as a $\mathbb{Z}_{\geq 0}$ -valued matrix (Z_{ij}) , indexed by the respective sets I_L, I_R of simple objects, that satisfies $Z_{00} = 1$ and intertwines the associated representations of $SL(2, \mathbb{Z})$, i.e. $Z\pi_L(g) = \pi_R(g)Z$. This definition turned out to be insufficient, and from now on ‘modular invariant’ will mean the following for us:

4.1 DEFINITION A modular invariant for a pair $(\mathcal{C}_L, \mathcal{C}_R)$ of modular categories consists of a triple (Γ_L, Γ_R, E) , where Γ_L, Γ_R are commutative algebras in $\mathcal{C}_L, \mathcal{C}_R$, respectively, and E is a braided monoidal equivalence $\Gamma_L - \text{Mod}_{\mathcal{C}_L}^0 \rightarrow \Gamma_R - \text{Mod}_{\mathcal{C}_R}^0$.

The question immediately arises whether two given categories $\mathcal{C}_L, \mathcal{C}_R$ admit a modular invariant. In the left-right symmetric case $\mathcal{C}_L = \mathcal{C}_R = \mathcal{C}$ at least one modular invariant always exists, to wit the triple $(\mathbf{1}, \mathbf{1}, \text{id}_{\mathcal{C}})$ (giving rise to the diagonal matrix $Z = \mathbf{1}$). A very ingenious subfactor theoretic approach to finding non-trivial modular invariants in this left-right symmetric situation was initiated in [1] and pursued further in [2]. This approach turned out [30, 15] to be of essentially categorical nature, revolving around algebras in \mathcal{C} that are not necessarily commutative and thus do not have an interpretation in terms of local extensions. A more physically motivated approach, applicable also when $\mathcal{C}_L \not\cong \mathcal{C}_R$, was proposed in [32], where it was shown that, given two chiral CQFTs A_L, A_R and considering their product $\mathcal{A} = A_L \otimes \overline{A_R}$ as a QFT on $1 + 1$ -dimensional Minkowski space, every finite local extension $\mathcal{B} \supset \mathcal{A}$ gives rise to a matrix Z satisfying $Z_{00} = 1$ and intertwining the T -matrices of the modular categories $\mathcal{C}_L, \mathcal{C}_R$. Furthermore, it was conjectured that $ZS_L = S_RZ$ iff $\text{DHR}(\mathcal{B})$ is trivial. Proofs of this were found by the author and by R. Longo and Y. Kawahigashi (both remained unpublished). In [32] it was also shown that, given such a local extension $\mathcal{B} \supset A_L \otimes \overline{A_R}$, there are chiral local extensions $\widehat{A}_L \supset A_L, \widehat{A}_R \supset A_R$ that are maximal w.r.t. the property $A_L \otimes \overline{A_R} \subset \widehat{A}_L \otimes \overline{\widehat{A}_R} \subset \mathcal{B}$ and have isomorphic fusion rings. In [18] it was even shown that one has an equivalence $E : \text{DHR}(\widehat{A}_L) \rightarrow \text{DHR}(\widehat{A}_R)$ of braided tensor categories. Recalling the correspondence [23] between local extensions $\mathcal{B} \supset \mathcal{A}$ and commutative algebras in $\text{DHR}(\mathcal{A})$ as well as Theorem 3.1, we conclude that a commutative algebra $\Gamma \in \text{DHR}(A_L \otimes \overline{A_R}) \simeq \text{DHR}(A_L) \boxtimes \widetilde{\text{DHR}}(\overline{A_R})$ of maximal dimension gives rise to a modular invariant for $\mathcal{C}_L, \mathcal{C}_R$. The converse follows from the fact, resulting from [23] and [19], that $A \otimes \overline{A}$ admits a local extension with trivial representation category. At this point, two questions arise:

1. What is the significance of the above reasoning for the construction of $d = 2$ CQFTs? Perhaps the requirement that \mathcal{B} have trivial representation category amounts to the absence of an obstruction to existence of a ‘Wick rotated’ Euclidean CQFT that can be defined on arbitrary Riemann surfaces? (The fact that the DHR category has a cohomological interpretation in terms of Roberts’ local cohomology may be taken as further support for this speculation.)
2. Do the above results hold irrespective of whether the categories $\mathcal{C}_{L/R}$ arise from chiral CFTs $A_{L/R}$? The answer is yes:

4.2 THEOREM Given two modular categories $\mathcal{C}_L, \mathcal{C}_R$, a modular invariant (Γ_L, Γ_R, E) gives rise to a commutative algebra $\Gamma \in \mathcal{C}_L \boxtimes \widetilde{\mathcal{C}}_R$ of maximal dimension (i.e. $d(\Gamma) = (\dim \mathcal{C}_L \cdot \dim \mathcal{C}_R)^{1/2}$). Conversely, every such algebra arises from a modular invariant [5], which in fact is unique [6].

(In the case $\mathcal{C}_L = \mathcal{C}_R$ this was proven in [22].) The proof of the first half requires little more than ideas already contained in [23] and [20]. The converse implication follows from facts outlined in the final section.

5 On the classification of modular categories

The circle of ideas of the preceding section has important pure mathematical ramifications, cf. [5]. Namely, given modular categories $\mathcal{C}_L, \mathcal{C}_R$, it is not hard to see that existence of a fusion category \mathcal{D} such that $\mathcal{C}_L \boxtimes \widetilde{\mathcal{C}}_R \simeq Z(\mathcal{D})$ is equivalent to existence of fusion categories $\mathcal{E}_1, \mathcal{E}_2$ such that $\mathcal{C}_L \boxtimes Z(\mathcal{E}_1) \simeq \mathcal{C}_R \boxtimes Z(\mathcal{E}_2)$. In this case we write $\mathcal{C}_L \sim \mathcal{C}_R$, and one easily sees that \sim is an equivalence relation, called Witt equivalence in analogy with the classical theory of quadratic forms. The set W_M of Witt equivalence classes of modular categories acquires an abelian monoid structure by $[\mathcal{C}_1] \cdot [\mathcal{C}_2] := [\mathcal{C}_1 \boxtimes \mathcal{C}_2]$ and $\mathbf{1}_{W_M} = [\text{Vect}]$. In view of $\mathcal{C} \boxtimes \widetilde{\mathcal{C}} \simeq Z(\mathcal{C})$ (by [24]) and $Z(\mathcal{C}) \sim \mathbf{1}$ (by definition), W_M is a group with inverse operation $[\mathcal{C}]^{-1} = [\widetilde{\mathcal{C}}]$.

In view of Theorem 3.4, if Γ is a commutative algebra in \mathcal{C} then $[\mathcal{C}] = [\Gamma - \text{Mod}_{\mathcal{C}}^0]$. (Thus if $A \subset B$ are completely rational CFTs then $[\text{DHR}(A)] = [\text{DHR}(B)]$.) Calling a modular category *completely anisotropic* if every commutative algebra in it is isomorphic to $\mathbf{1}$, one can prove that Witt-equivalent completely anisotropic categories are actually braided equivalent. (In other words: every Witt

equivalence class contains one completely anisotropic category up to braided equivalence). This is the reason for the uniqueness of the representation category of maximal local extensions and of the converse statement in Theorem 4.2: By Theorem 3.4, existence of a maximal commutative algebra in $\mathcal{C}_L \boxtimes \widetilde{\mathcal{C}}_R$ implies $\mathcal{C}_L \boxtimes \widetilde{\mathcal{C}}_R \simeq Z(\mathcal{D})$ and thus $[\mathcal{C}_L] = [\mathcal{C}_R]$. Choosing commutative algebras $\Gamma_{L/R} \in \mathcal{C}_{L/R}$ such that the respective local module categories are completely anisotropic, we have $[\Gamma_L - \text{Mod}_{\mathcal{C}_L}^0] = [\mathcal{C}_L] = [\mathcal{C}_R] = [\Gamma_R - \text{Mod}_{\mathcal{C}_R}^0]$ and thus a braided equivalence $\Gamma_L - \text{Mod}_{\mathcal{C}_L}^0 \simeq \Gamma_R - \text{Mod}_{\mathcal{C}_R}^0$.

Finally, the Witt group holds great promise for the classification of modular categories. The point is that $Z(\mathcal{C})$ is modular for every fusion category, of which there are far too many to hope for a classification. (The fact that inequivalent fusion categories can have equivalent centers [24] does not help much.) Passing to the Witt group not only kills those ‘trivial’ modular categories but has the nice effect of yielding an abelian group W_M . Many generators (from quantum groups at roots of unity) and relations (from conformal extensions and cosets) for W_M are already known, and one may hope that W_M can be determined completely. This would seem to be a rigorous implementation to the idea from CQFT folklore that “all modular categories” arise from the chiral WZW models via local extensions and coset constructions.

References

- [1] J. Böckenhauer, D. E. Evans: Modular invariants, graphs and α -induction for nets of subfactors I-III. *Commun. Math. Phys.* **197**, 361-386 (1998); **200**, 57-103 (1999); **205**, 183-228 (1999).
- [2] J. Böckenhauer, D. E. Evans, Y. Kawahigashi: *Commun. Math. Phys.* **208**, 429-487 (1999); **210**, 733-784 (2000); *Publ. Res. Inst. Math. Sci.* **37**, 1-35 (2001).
- [3] A. Bruguières, A. Virelizier: The Double of a Hopf monad. [arXiv:0812.2443](https://arxiv.org/abs/0812.2443).
- [4] R. Conti, S. Doplicher, J. E. Roberts: Superselection theory for subsystems. *Commun. Math. Phys.* **218**, 263-281 (2001).
- [5] A. Davydov, M. Müger, D. Nikshych, V. Ostrik: In preparation.
- [6] A. Davydov, M. Müger: Work in progress.
- [7] P. Deligne: Catégories tannakiennes. *In: P. Cartier et al. (eds.): Grothendieck Festschrift*, vol. II, pp. 111–195. Birkhäuser Verlag, 1991.
- [8] R. Dijkgraaf, V. Pasquier, P. Roche: Quasi Hopf algebras, group cohomology and orbifold models. *Nucl. Phys. B (Proc. Suppl.)* **18B**, 60-72 (1990).
- [9] S. Doplicher, R. Haag, J. E. Roberts: Local observables and particle statistics I & II. *Commun. Math. Phys.* **23**, 199-230 (1971); **35**, 49-85 (1974).
- [10] S. Doplicher, G. Piacitelli: Any compact group is a gauge group. *Rev. Math. Phys.* **14**, 873-885 (2002).
- [11] S. Doplicher, J. E. Roberts: A new duality theory for compact groups. *Invent. Math.* **98**, 157-218 (1989).
- [12] S. Doplicher, J. E. Roberts: Why there is a field algebra with a compact gauge group describing the superselection structure in particle physics. *Commun. Math. Phys.* **131**, 51-107 (1990).
- [13] V. Drinfeld, S. Gelaki, D. Nikshych, V. Ostrik: Group-theoretical properties of nilpotent modular categories. [math.QA/0704.1095](https://arxiv.org/abs/math/0704.1095).
- [14] K. Fredenhagen, K.-H. Rehren, B. Schroer: Superselection sectors with braid group statistics and exchange algebras I & II. *Commun. Math. Phys.* **125**, 201-226 (1989), *Rev. Math. Phys.*, Special Issue 1992, 113–157.
- [15] J. Fröhlich, J. Fuchs, I. Runkel, C. Schweigert: Correspondences of ribbon categories. *Adv. Math.* **199**, 192-329 (2006).

- [16] R. Haag: *Local Quantum Physics*. 2nd ed., Springer, 1996
- [17] H. Halvorson: Algebraic quantum field theory. *In: J. Butterfield & J. Earman (eds.): Handbook of the Philosophy of Physics*. Elsevier, 2006.
- [18] Y. Kawahigashi, R. Longo: Classification of two-dimensional local conformal nets with $c < 1$ and 2-cohomology vanishing for tensor categories. *Commun. Math. Phys.* **244**, 63-97 (2004).
- [19] Y. Kawahigashi, R. Longo, M. Müger: Multi-interval subfactors and modularity of representations in conformal field theory. *Commun. Math. Phys.* **219**, 631-669 (2001).
- [20] A. Kirillov Jr., V. Ostrik: On q-analog of McKay correspondence and ADE classification of $sl(2)$ conformal field theories. *Adv. Math.* **171**, 183-227 (2002).
- [21] A. Kitaev, M. Müger: Unpublished work 2006-2008.
- [22] L. Kong, I. Runkel: Cardy algebras and sewing constraints I. [arXiv:0807.3356](https://arxiv.org/abs/0807.3356).
- [23] R. Longo, K.-H. Rehren: Nets of subfactors. *Rev. Math. Phys.* **7**, 567-597 (1995).
- [24] M. Müger: From subfactors to categories and topology II. The quantum double of tensor categories and subfactors. *J. Pure Appl. Alg.* **180**, 159-219 (2003).
- [25] M. Müger: Conformal orbifold theories and braided crossed G-categories. *Commun. Math. Phys.* **260**, 727-762 (2005).
- [26] M. Müger: Abstract duality theory for symmetric tensor *-categories. App. to [17].
- [27] M. Müger: On braided crossed G-categories. App. to V. Turaev: *Homotopy Quantum Field Theory*. EMS, to appear.
- [28] M. Müger: Local extensions and modular invariants of quantum fields. In preparation.
- [29] M. Müger, J. E. Roberts, L. Tuset: Representations of algebraic quantum groups and reconstruction theorems for tensor categories. *Alg. Repres. Theor.* **7**, 517-573 (2004).
- [30] V. Ostrik: Module categories, weak Hopf algebras and modular invariants. *Transf. Groups* **8**, 177-206 (2003).
- [31] B. Pareigis: On braiding and dyslexia. *J. Algebra* **171**, 413-425 (1995).
- [32] K.-H. Rehren: Locality and modular invariance in 2D conformal QFT. *Fields Inst. Commun.* **20**, 297-319 (2001).