# Nose to tail – how to allocate the environmental burden of livestock production systems to different meat cuts?



CORE

### Stucki Matthias and Eymann Lea

Zurich University of Applied Sciences Institute of Natural Resource Science Grüental, 8820 Wädenswil Switzerland matthias.stucki@zhaw.ch



**Hirsiger Eva** Eaternity, 8057 Zurich Switzerland ehirsiger@eaternity.ch



# Introduction and Objectives

- **"Nose-to-tail eating**" is a global trend that refers to using every part of an animal instead of valuing only premium cuts such as filet and steak (e.g.[1]).
- It is a matter of allocation, to show how environmentally favourable the use inferior meat cuts is. In LCA studies of agri-food systems economic allocation is frequently used to handle multi-output processes [2]. However, most studies analysing the environmental impact of meat are limited to the

## Method

- In a first step, the global warming potential (GWP) of beef, calf and pork under live weight was calculated applying the IPCC 100a impact assessment method in the SimaPro 8 software to the Agribalyse v1.2 database [7].
- In a second step, the environmental burden of meat and slaughterhouse by-products was determined based on economic allocation using Swiss meat prices [8].
- Subsequently, the environmental impact of meat cuts

For expensive meat cuts like filet, the economic allocation approach results in up to 3.3 times higher environmental impacts than mass allocation. Applying mass allocation, the GWP of beef, veal and pork is 31.8 kg CO<sub>2</sub>-eq/kg, 16.1 kg  $CO_2$ -eq/kg and 3.6 kg  $CO_2$ -eq/kg, respectively. With economic allocation the greenhouse gas emissions of different **beef** cuts range from **12 kg CO<sub>2</sub>-eq/kg** for sausage meat to 105 kg CO<sub>2</sub>-eq/kg for filet (Figure 1). For veal and pork the GWP of meat cuts ranges from 6.0 kg CO<sub>2</sub>-eq/kg (sausage meat) to 38 kg CO<sub>2</sub>-eq/kg (filet) and from 0.4 kg CO<sub>2</sub>-eq/kg (pork rind) to 10.9 kg CO<sub>2</sub>-eq/kg (filet), respectively, if economic allocation is applied. When assessing the environmental impact of meat dishes the choice of the allocation approach has a major influence on the result. For Beef Wellington, for example, the GWP per serving is 6.0 kg CO<sub>2</sub>-eq if mass allocation is used to compute the filet's climate impact, and 18.3 kg  $CO_2$ -eq when economic allocation is applied. In contrast, for pork ragout pie the allocation approach does not have a significant impact on the GWP of the meal.

Sausage

**meat: 12** 

kg CO<sub>2</sub>-

eq./kg

Results

differentiation between meat and slaughter byproducts [3-6].

 This study aims at assessing the environmental impact of individual meat cuts. was calculated using two different allocation approaches (mass and economic allocation).

- Finally, the allocation factors were applied to GWP of beef, calf and pork.
- Furthermore, the impact of meat dishes on climate change was computed using the global warming potential of additional ingredients from the database set up by Eaternity [9].

Figure 1: Carbon footprint of different beef cuts when applying an economic allocation [10]



Minced beef:

18 kg CO<sub>2</sub>-

eq./kg

## **Discussion and Conclusion**

When dealing with the environmental impact of meat, allocation is a crucial issue. Many LCA studies analysing the environmental impact of livestock production systems express the result per of live weight. Using such results to indicate the environmental impact of meat implies that all products, co-products and wastes arising from the slaughterhouse are assigned the same environmental burden, which cannot be justified. Likewise, it is questionable if the same environmental burden should be attributed to sausage meat and filet.

This study proposes the use of economic allocation, since livestock farming is **driven by economic factors** and thus the reason why animals are kept is that eventually a variety of animal products can be sold. The results from the **economic allocation** presented in this study are used to provide climaterelevant information for **restaurants**, consumers and NGOs in collaboration with the ICT-consulting firm Eaternity. The participating restaurants are encouraged to offer **climate-friendly dishes** which can be labelled with the Eaternity Award if their GWP lies below a defined threshold value.

Lean boiled

beef: 27 kg

CO<sub>2</sub>-eq./kg

In line with the "nose-to-tail" principle, an economic allocation approach promotes the use of inferior meat cuts and could thereby contribute to increasing the share of

#### an animal's body used for human nutrition.



- [1] Henderson, F. (2004). *The Whole Beast: Nose to Tail Eating* Bloomsbury Publishing.
- [2] van der Werf, H. M. G., & Nguyen, T. T. H. (2014). Construction cost of plant compounds provides a physical relationship for co-product allocation in life cycle assessment. 9th International Conference LCA of Food San
- տ 🚊 Francisco, USA 8-10 October 2014,

5

Z

🚬 🗒 [3] Blonk, H., & Luske, B. (2008). Greenhouse Gas Emissions of Meat. Methodological issues and establishment of an information infrastructure.

Shoulder: 30

kg CO<sub>2</sub>-eq./kg

- s [4] Desjardins, R. L., Worth, D. E., Vergé, X. P. C., Maxime, D., Dyer, J., & Cerkowniak, D. (2012). Carbon Footprint of Beef Cattle. Sustainability, 4, 3279-3301. doi: 10.3390/su4123279
- Solution issue: Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Processing: the allocation issue. Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Processing: the allocation issue. Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Processing: the allocation issue. Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Processing: the allocation issue. Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Processing: the allocation issue. Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Processing: the allocation issue. Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Processing: the allocation issue. Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Proceedings of the 9th International Conference on Life Cycle
  Solution issue: Proceedings of the 9th International Conference on Life Cycle
  Solution: Proceedings of the 9th International Conference on Life Cycle
  Solution: Proceedings of the 9th International Conference on Life Cycle
  Solution: Proceedings
  Solution: Proc
- Gac, A., Tribot-Laspière, P., Scislowski, V., Lapasin, C., Ponchant, P., Guardia, S., Nassy, G., & Chevillon, P. (2012). Recherche de méthodes d'évaluation de l'expression de l'empreinte carbone des produits viande. No.
  Réf: 00 12 33 023 ISSN 1773-4738
- 🤗 🚊 [7] ADEME (2015). base de données Agribalyse v1.2.
- [8] FOAG. (2014). Consumer prices meat. Excel. Federal Office for Agriculture. BLW. Market Survey Section.
  - [9] Eaternity (2015). CO2-Data by Eaternity, retrieved I January 2015 from the Eaternity Cloud: co2.eaternity.ch [10] Image from insitebrazosvalley.com