

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/92367>

Please be advised that this information was generated on 2017-12-06 and may be subject to change.

STAT genes display differential evolutionary rates that correlate with their roles in the endocrine and immune system

Marnix Gorissen¹, Erik de Vrieze¹, Gert Flik¹ and Mark O Huising^{1,2}

¹Department of Organismal Animal Physiology, Faculty of Science, Radboud University Nijmegen, Heyendaalseweg 135, 6525AJ Nijmegen, The Netherlands

²The Clayton Foundation Laboratories for Peptide Biology, Peptide Biology Laboratories, The Salk Institute for Biological Studies, 10010 North Torrey Pines Road, La Jolla, California 92037, USA

(Correspondence should be addressed to M Gorissen; Email: m.gorissen@science.ru.nl)

Abstract

We identified orthologues of all mammalian Janus kinase (*JAK*) and signal transducer and activator of transcription (*STAT*) genes in teleostean fishes, indicating that these protein families were already largely complete before the teleost tetrapod split, 450 million years ago. In mammals, the *STAT* repertoire consists of seven genes (*STAT1*, -2, -3, -4, -5a, -5b, and -6). Our phylogenetic analyses show that *STAT* proteins that are recruited downstream of endocrine hormones (*STAT3* and *STAT5a* and -5b) show a markedly higher primary sequence conservation compared with *STATs* that convey immune signals (*STAT1-2*, *STAT4*, and *STAT6*).

A similar dichotomy in evolutionary conservation is observed for the *JAK* family of protein kinases, which activate *STATs*. The ligands to activate the *JAK/STAT*-signalling pathway include hormones and cytokines such as GH, prolactin, interleukin 6 (*IL6*) and *IL12*. In this paper, we examine the evolutionary forces that have acted on *JAK/STAT* signalling in the endocrine and immune systems and discuss the reasons why the *JAK/STAT* cascade that conveys classical immune signals has diverged much faster compared with endocrine *JAK/STAT* paralogues.

Journal of Endocrinology (2011) **209**, 175–184

Introduction

Class-I helical cytokines constitute a monophyletic group of proteins consisting of molecules that convey signals of the endocrine system (e.g. GH, prolactin, and erythropoietin (*EPO*)), as well as signalling molecules that coordinate host defence (e.g. interleukins; Huising *et al.* 2006). All class-I helical cytokines fold in a typical four α -helix barrel structure and signal through a group of related receptors (Liongue & Ward 2007). These cytokines activate the Janus kinase/signal transducer and activator of transcription (*JAK/STAT*) pathway, ultimately leading to changes in gene expression. Since its discovery as a regulator of interferon (*IFN*) responses in the immune system (Schindler *et al.* 1992, Darnell *et al.* 1994), *JAK/STAT* molecules were shown to represent a common signalling pathway shared by many cytokines (Shuai & Liu 2003). The binding of a cytokine to its receptor typically leads to dimerisation of the receptor and subsequently to phosphorylation of recruited *JAK* molecules (Chen *et al.* 2004). The phosphorylated *JAKs* in turn phosphorylate several key tyrosines in the intracellular domain of the receptor, which then serve as a docking site for *STAT* proteins (Gadina *et al.* 2001). These *STATs* are phosphorylated on a single tyrosine residue, after which they form homo- or heterodimers with other phosphorylated *STAT* proteins. These dimers detach from the receptor and are then

translocated to the nucleus, where they promote or inhibit gene expression (Darnell 1997, Levy & Darnell 2002, O'Shea *et al.* 2002; Fig. 1).

In mammals, the *JAK* family consists of four distinct genes (*JAK1-3* and *TYK2*), whereas the *STAT* repertoire consists of seven distinct *STAT* genes, (*STAT1*, -2, -3, -4, -5a, -5b, and -6; Darnell 1997). Some promiscuity exists among *STAT* proteins regarding the ligands and cytokine receptors that can activate particular *STAT* members, with sometimes multiple *STATs* that can be activated downstream of the same cytokine receptor (e.g. leptin's main actions are exerted through *STAT3* homodimers, but *STAT1/STAT3* heterodimers also serve in leptin signalling (Bendinelli *et al.* 2000)). Besides the well-established roles of *STAT1-2*, 4, and 6 in the immune system, in recent years, *STAT3* and *STAT5a/b* have emerged as regulators of T regulatory (*Treg*) and T helper 17 (*Th17*) cell development, differentiation, and maintenance (Wei *et al.* 2008). Despite these contributions to vital aspects of the mammalian immune system, a very interesting demarcation exists between *STAT3* and *STAT5* and the other *STAT* family members, as most of the classical hormones within the class-I helical cytokine family, such as GH, PRL, and *EPO*, signal predominantly via *STAT3* and *STAT5*, whereas the other members of the *STAT* family serve predominantly in the immune response (Horvath *et al.* 1995). A similar differentiation exists between *JAK1* and *JAK2*, which serve both in the




Figure 1 JAK–STAT signalling. (A) The cytokine binds to a cytokine-specific receptor α -chain. (B) The receptor then dimerises with a β -chain shared by multiple cytokine receptor complexes. After dimerisation, the β -chain is phosphorylated. (C) Members of the JAK family can now bind to the receptor, are activated by transphosphorylation and in turn transphosphorylate a second site on the receptor. (D) Members of the STAT family can now bind to the receptor complex at this second site, and are phosphorylated by JAK. Phosphorylated STAT molecules dimerise and translocate to the nucleus to alter gene transcription.

signalling of immune cytokines and endocrine cytokines, and JAK3 and Tyk2, which serve in immunology only.

All STATs share a highly conserved domain structure, including an SRC2 homology (SH2) domain – involved in the formation of STAT dimers (Shuai *et al.* 1994), a DNA-binding domain (Horvath *et al.* 1995), and a transactivation domain (TAD; Shuai *et al.* 1993). The latter domain shows the highest degree of variability among STATs at the primary sequence level and the gene structure (Supplementary Figure 1, see section on supplementary data given at the end of this article) and enables them to interact with different cofactors required for activation of a STAT-selective transcriptional profile (Levy & Darnell 2002).

Almost all of our knowledge on intracellular signalling of class-I helical cytokines is based on rodent and primate models. In recent years, genomes of sufficient and sufficiently diverse vertebrate species have been elucidated to initiate a comprehensive study on the phylogeny and evolution of this key family of proteins. In the present study, we compare the *JAK* and *STAT* repertoires of mammals with those of key distantly related vertebrate species, including teleostean fishes.

This approach gives us unique tools to reconstruct an evolutionary history, which is surprisingly dynamic and features multiple gene duplications and subsequent deletions. Moreover, our phylogenetic analyses reveal differential evolutionary rates for the immune and endocrine members of the JAK and STAT protein family.

Materials and Methods

Identification of *JAK* and *STAT* orthologues from databases

We retrieved JAK and STAT sequences from the NCBI protein (www.ncbi.nlm.nih.gov/protein) and swissprot (www.expasy.org/sprot/) databases. To complete the JAK/STAT repertoire of key vertebrate species, we conducted an extensive BLAST (Altschul *et al.* 1997) search in the publicly available genome databases (www.ensembl.org; Hubbard *et al.* 2007). Because of the (as of yet) incomplete annotation of several genomes, it is inevitable that some BLAST searches yielded JAK/STAT orthologues that were overtly incomplete. These incomplete annotations were corrected manually, by

searching for the correct intron–exon splice sites and coding sequences in the genome. In our phylogenetic analysis, only complete coding sequences of *JAK* and *STAT* genes were used.

Reconstruction of phylogeny

Multiple sequence alignments were constructed with ClustalW (www.ebi.ac.uk/Tools/clustalw2/index.html) and uploaded into MEGA 3.0 (Kumar *et al.* 2004). Phylogeny was constructed on the basis of amino acid differences (*P* distance) using the neighbour-joining algorithm. Phylogenetic trees were constructed using both pairwise and complete deletion parameters, which rendered trees with similar topology to one another. Only the phylogenetic analyses using pairwise deletion are shown. Reliability of the trees was assessed by bootstrapping (1000 replications).

Characterisation of the nature of selective force: pN/pS ratios

We calculated the ratio between the proportion of non-synonymous (pN) and synonymous (pS) substitutions for all *stats*. In order to do this, the coding region of each *stat* paralogue of zebrafish and pufferfish was aligned pairwise to its human orthologue. We corrected these nucleotide alignments manually for overt mismatches, guided by the corresponding amino acid alignments. Then, the number of (non)synonymous sites and (non)synonymous substitutions were determined using MEGA 3.0, according to the Nei & Gojobori (1986) method. To test if the level of purifying selection (i.e. pN/pS < 1) is statistically different from neutral selection (i.e. pN/pS = 1), we conducted a Z-test on the pN/pS ratios.

Results

Phylogeny of vertebrate *JAK/STATs*

By scrutinizing available genomes and protein databases, we identified teleostean orthologues for all mammalian *JAK* and *STAT* family members. This indicates that the contemporary *JAK/STAT* repertoire was already complete before the teleost–tetrapod split (~450 Mya), with the exception of the mammalian and teleostean *STAT5* paralogues (as discussed later in this paper). Although *STATs* have been found in several invertebrates, a repertoire of *STAT* proteins reminiscent of the vertebrate *STAT* family has not been identified in any non-vertebrate. In the sea squirt (*Ciona intestinalis*), a key species in chordate evolution as it represents one of the closest non-vertebrate relatives to the vertebrate subphylum, only one *jak* and two *stat* genes (*stat-a* and *stat-b*) have been identified (Hino *et al.* 2003). This suggests that both the *JAK* and the *STAT* repertoire radiated early during vertebrate evolution, after the urochordate–vertebrate bifurcation but before the teleost–tetrapod split.

Our phylogenetic analyses show that *JAK1* and *JAK2* (Fig. 2), which act downstream of immune and endocrine signals, display a noticeably higher primary sequence conservation than *JAK3* and *Tyk2*, which are restricted to signalling in immune pathways, as they cluster more compactly. Similarly, *STAT3* and *STAT5* (Fig. 3), which both convey endocrine signals, display noticeably higher primary sequence conservation than the *STAT* proteins that serve in immunity. To quantify this observation, we calculated the ratio of non-synonymous to synonymous substitutions (pN/pS ratios) of the *stat* repertoire of zebrafish (*Danio rerio*) and Japanese pufferfish (*Takifugu rubripes*) in comparison to the human repertoire of *STAT* genes (Table 1). The ratio between the pN and the pS substitutions provides us with insight into the type and strength of the selective pressure that has acted on a protein sequence in a given evolutionary time frame. In addition, pS values provide information regarding divergence time between two sequences. As synonymous substitutions generally experience no selection, orthologous or paralogous genes that have separated earlier in evolution, in general will have acquired more synonymous substitutions – and therefore carry a higher pS value – than genes that separated more recently, and therefore the pS value is an indicator of the divergence time between two sequences. By definition, pN/pS values < 1 indicate purifying selection, which is aimed at maintaining an amino acid sequence constant. The counterpart of purifying selection is positive selection, which favours amino acid changes and is characterised by a pN/pS ratio > 1. Neutral selection is assumed if neither purifying nor positive selection is demonstrated.

For all *stat* family members, pN/pS ratios < 1 are observed, which indicates that all *stats* have been subjected to some degree of purifying selection over the examined time frame. *Stat3* and *stat5* show markedly lower pN/pS values compared with the other *stat* genes, and this corroborates our earlier observation concerning the noticeably more compact clustering of these *stats* in phylogenetic analysis (Fig. 2) and indicates that *stat3* and *stat5* experienced stronger purifying selection over the course of vertebrate evolution which led to better conservation of their primary amino acid sequences in comparison to the other *stat* family members. In addition to the pN value for each protein, we also examined the distribution of the non-synonymous substitutions (between the *STAT* repertoire of zebrafish and that of human) within each member of the *STAT* family (Fig. 4). As already indicated by the phylogenetic analysis and pN/pS values, *STAT1-2*, 4, and 6 show markedly more non-synonymous substitutions. Interestingly, the domain that displays most non-synonymous substitutions is the TAD. The complete lack of studies addressing the properties of teleostean TADs precludes speculation on the consequences of the widely variable and poorly conserved TAD domains; from mammalian studies, we know that the TAD is involved in the binding of different co-factors required for *STAT*-induced transcription of target genes.




Figure 2 Neighbour-joining phylogenetic analysis of vertebrate Janus kinase (JAK) proteins, performed under pairwise deletion and *P* distance conditions in MEGA3 (Kumar *et al.* 2004). For mammals (red), marsupials (purple), birds (yellow), amphibians (green), and teleostean fishes (blue), several key species are included in this phylogenetic analysis. Bootstrap values of all main branches are indicated by the size of the dots. JAK proteins involved in immunology only (viz. JAK3 and Tyk2) show longer branch lengths than JAKs downstream of both immune and endocrine signalling molecules.

A model for the genesis of contemporary vertebrate STAT repertoires

Alternating views exist on early key formative events that shape contemporary vertebrate genomes. The 2R hypothesis postulates that two successive rounds of whole genome duplication occurred before the teleost–tetrapod split, accounting for the presence of many genes, or gene clusters found on four paralogous loci (Sharman & Holland 1996, Sidow 1996, Meyer & Van de Peer 2005). Others have pointed out that series of tandem duplications of large genomic segments could also account for the distribution of ancestral genes across paralogous loci (Hughes & Friedman 2003, 2004). Regardless, both hypotheses agree on the occurrence of large-scale genomic duplication events in the formative stages of the ancestral vertebrate genome. Following these large-scale rearrangement events, many of the newly acquired duplicates were lost in order to arrive at the present-day genomic distribution of many gene families, including the STATs.

In mammals, *STAT* genes are distributed over three independent chromosomal regions (Fig. 5), with each locus carrying two genes, with the exception of the region that contains *STAT3* and both *STAT5* paralogues. This suggests that the ancestral *stat* gene was duplicated by tandem duplication. Indeed, the *stat* repertoire of *C. intestinalis*, the sea squirt, consists of two *stat* genes (*stat-a* and *stat-b*) that reside on separate loci (Hino *et al.* 2003). One of these loci may be the representative of the ancestral *stat* gene that gave rise to the contemporary vertebrate *stat* repertoire, while the other sea squirt *stat* gene may have originated independently of the mechanisms that gave rise to the vertebrate *stat* family, or was lost in the course of vertebrate evolution. After the first tandem duplication, the ancestral locus, carrying two tandem copies of *stat*, was subsequently distributed over three independent loci by two large-scale genome duplication events, possibly involving the genome duplications that constitute the 2R hypothesis (Copeland *et al.* 1995). The mammalian *STAT5a/5b* duplication is the result of a much more recent tandem duplication event that took place




Figure 3 Neighbour-joining phylogenetic analysis of vertebrate STATs, performed under pairwise deletion and *P* distance conditions in MEGA3 (Kumar *et al.* 2004). For mammals (red), marsupials (purple), birds (yellow), amphibians (green), and teleostean fish (blue), key species are included in the phylogenetic analysis. Bootstrap values of all main branches are indicated by the size of the dots. Protein accession codes are listed in Supplementary Table 1, see section on supplementary data given at the end of this article.

Table 1 pN/pS ratios for human versus teleostean fish STATs and representatives of the signals they convey. pN and pS are calculated with MEGA3 software (Kumar *et al.* 2004). For zebrafish, values for the duplicated *stat5* genes are indicated with 5.1 and 5.2 in brackets

Human	Zebrafish	S.E.M.	Tiger pufferfish	S.E.M.	Main cytokine ligand
<i>STAT1</i> (immune)	0.326	0.021	0.268	0.019	IFN α/β , IFN γ
<i>STAT2</i> (immune)	0.537	0.026	0.615	0.037	IFN α/β
<i>STAT3</i> (endocrine/immune)	0.088	0.010	0.086	0.010	G-CSF, IL6, leptin
<i>STAT4</i> (immune)	0.281	0.018	0.314	0.020	IL12
<i>STAT5a</i> (endocrine)	0.179 (5.1) 0.216 (5.2)	0.015 0.016	0.171 –	0.014 –	PRL, EPO, TPO
<i>STAT5b</i> (endocrine)	0.167 (5.1) 0.221 (5.2)	0.013 0.016	0.239 –	0.018 –	GH
<i>STAT6</i> (immune)	0.576	0.034	0.655	0.038	IL4, IL13

We employed a Z-test based on the pN/pS ratios; all pN/pS ratios were proven significant with $P < 0.001$.

in the mammalian lineage after the teleost–tetrapod split, as is discussed in the next section.

In contrast to mammalian *STATs* that are nicely arranged in tandem repeats, teleostean *stat* genes (with the exception of the *stat3/stat5.1* pair) are no longer distributed in tandem pairs. Although it is clear, based on our phylogenetic analysis, that teleostean *stats* are orthologues of the mammalian *STATs*, the genes that encode them have somehow been scattered over their genome. To understand the events that underlie this distribution, we compared the synteny of the teleostean and mammalian *stat* genes to arrive at the following scenario for the distribution of *stat* genes in teleostean fish.

Early in the teleostean lineage, an additional large-scale gene duplication occurred (Wittbrodt *et al.* 1998, Jaillon *et al.* 2004), also known as the ‘fish-specific genome duplication’ or ‘3R hypothesis’ and it appears that all teleostean loci that carry *stat* genes were duplicated in this event. In order for both duplicate copies to be maintained, each paralogue must acquire a distinct function that is subject to selection (in some cases, gene dosage may result in the maintenance of both paralogues (Kondrashov *et al.* 2002). However, as both paralogues will initially act fully redundantly, a failure to acquire distinct function, spatial or temporal expression patterns will usually lead to one member of each pair disappearing through genetic drift. In general, it appears that the majority of these duplicated genes in the 3R event was subsequently lost, as the total estimated gene number of teleostean species does not greatly exceed the number of genes in the human or mouse genome (Aparicio *et al.* 2002). This is also true for the *stat* gene family. After duplication, one of the duplicated genes was subsequently lost in a manner that left the contemporary teleostean *stats* apparently isolated at their respective loci (Fig. 5). Human *STAT1* and *STAT4*, for example, are located on chromosome 2. In zebrafish, *stat4* is located on chromosome 9, but *stat1* is positioned on chromosome 22 (Fig. 4A). However, both zebrafish chromosomes carry neighbouring genes that are orthologous to the neighbouring genes on the human *STAT1/STAT4* locus; zebrafish chromosome 9 and 22 each shares three genes in synteny with the human 2q32.2/3 locus that carries human *STAT1* and *STAT4*. Interestingly, on zebrafish chromosome 9, which carries zebrafish *stat4*, a remnant of

another *stat-like* gene can be found (Stein *et al.* 2007). It consists only the first 16 of the 25 exons that typically encode a full *STAT* protein, and is present in the zebrafish EST databases (EH485578.1), indicating that this truncated *stat-like* gene is expressed in zebrafish and therefore does not constitute a pseudogene. This *STAT-like* protein does not have a TAD and may possess regulatory properties such as the mammalian truncated *STAT3 β* , which is considered a dominant negative regulator of *STAT* signalling (Maritano *et al.* 2004). More importantly, this observation underpins our hypothesis that *stat* genes in the teleostean ancestor resided in tandem pairs prior to their duplication. One of the tandem copies on most loci disappeared subsequently; an hypothesis further strengthened by the fact that, in zebrafish, *stat3* and *stat5.1* are located in tandem, whereas *stat5.2* is on a separate locus.




Figure 4 Localisation of non-synonymous substitutions between zebrafish and human *STAT* genes. For each domain of the *STAT* proteins, the primary sequence similarity is indicated as a percentage, i.e. the lower the percentage, the higher the amount of non-synonymous substitutions.




Figure 5 Conserved synteny of human and zebrafish *stat1/stat4* (A), *stat3/stat5* (B), and *stat2/stat6* (C) loci. The partial zebrafish *stat1* orthologue is named (*p*)*stat1*. In Fig. 3C, we omitted several genes that are not of interest regarding the conserved synteny of the *stat* loci.

The same phenomenon can be witnessed for the other zebrafish chromosomes containing *stat* genes (Fig. 5): neighbouring genes of the zebrafish *stat* paralogues have maintained synteny with their human orthologues. Some duplicate genes are conserved (i.e. neither of the two paralogues is discarded), and are present in synteny on both zebrafish loci. It is apparent that the teleost’s additional genome duplication resulted in the scattering of *stat* genes over more loci than in tetrapods (Fig. 6).

Stat5 underwent two independent tandem duplications during vertebrate evolution

Although the framework for the STAT protein family was largely complete before the teleost–tetrapod split, two additional gene duplications have occurred thereafter. In both mammals and teleostean fish, but not in birds and amphibians, duplicate *stat5* genes are found. Where mammals have *STAT5a* and *STAT5b* paralogues, the teleostean duplicate *stat5* genes have been named *stat5.1* and *stat5.2* (Lewis & Ward 2004). Teleostean *stat5* duplicates are present in both zebrafish (*D. rerio*) and Japanese medaka (*Oryzias latipes*), but appear to be absent in pufferfishes (*Tetraodon nigroviridis*, *Takifugu rubripes*). The pN/pS ratios for *stat5.1* and *stat5.2*, compared with either *STAT5a* or *-5b*, are similar and lower than the ratios for the ‘immune’ *stats* (Table 1),

indicating that relatively strong purifying selection has acted on both *stat5* duplicates in teleosts and tetrapods alike.

The presence of *stat5.1* and *stat5.2* genes in zebrafish and medaka indicates that these two genes have arisen before the estimated divergence time of these species (~300 Mya), early in teleostean evolution. The pufferfish lineage arose ~180 Mya (Muffato & Roest-Crollius 2008) and would therefore be expected to have duplicate *stat5* paralogues as well. However, the genomic landscape of the puffers is substantially different from most vertebrate genomes as it is very condense and contains relatively little non-protein-coding DNA (Aparicio *et al.* 2002). In light of these profound changes in genetic makeup that were experienced in the pufferfish lineage, it is plausible that pufferfishes lost one of their *stat5* paralogues after the teleostean genome duplication in the course of evolution, although we cannot conclusively rule out that we are unable to retrieve a second *stat5* gene in both pufferfish species, as their respective genomes may not be entirely covered. The mammalian and teleostean *stat5* duplicates do not form a uniform clade in our phylogenetic tree, as teleostean *stat5* paralogues form a clade with other teleostean *stat5s*. This further supports our assertion that the teleostean *stat5* duplication occurred independently from the mammalian duplication. The pS values calculated for both human and zebrafish *stat5* paralogues provide an estimate of when the duplications of teleostean and mammalian *stat5*




Figure 6 A proposed mechanism for STAT evolution: including 2R hypothesis as a basis of the radiation of the *stat* repertoire, the fish-specific whole genome duplication (3R), and subsequent (partial) loss of paralogous *stat* genes. Animal diagrams from left to right: sea squirt (*Ciona intestinalis*), zebrafish (*Danio rerio*), Japanese medaka (*Oryzias latipes*), opossum (*Monodelphis domestica*), and mouse (*Mus musculus*).

paralogues may have occurred. Synonymous mutations occur and are fixed in a population at a relatively constant rate, since there is generally no selective pressure acting on these nucleotide positions. Instead, selective pressure acts on amino acid sequences, and those are not affected by synonymous substitutions. The pS value for *STAT5a* versus *STAT5b* is 0.377, whereas for the teleostean *stat5.1* versus *stat5.2* paralogues, the pS value is 0.766. Under the assumption of constant nucleotide substitution rate that is equal in both lineages, these numbers indicate that the teleostean paralogues arose independently from mammalian *STAT5a* and *STAT5b* and approximately twice as early in evolution. As the *STAT5* duplication events in mammals and teleostean fishes occurred independently, the fact that the duplicated *stat5s* still exist in contemporary mammals and fish suggests that the presence of two *stat5* genes presented evolutionary advantages to mammals and bony fish that has led to their maintenance in both lineages (with the aforementioned exception of the pufferfishes).

We know that the mammalian *STAT5* paralogues, while highly similar in primary sequence, acquired partially independent functions: *STAT5a* serves in prolactin signalling, *STAT5b* acts downstream of GH (Schindler 2002). This is illustrated by the observations from genetic models, which

revealed that *Stat5a* knock-out mice are deficient in prolactin signalling, while *Stat5b* knock-out mice display sexually dimorphic growth retardation. Although both single knock-outs are viable, mice that lack functional copies of both *Stat5a* and *-5b* die a few weeks after birth (O'Shea *et al.* 2002), suggesting that some redundancy still exists between these paralogues. For the teleostean *stat5* paralogues, it is not known if they exert identical functions (and thus act fully redundantly) or if they have acquired different functions during the course of teleostean evolution. Nevertheless, given the fact that the primary sequences of *stat5.1* and *-5.2* share less identity with each other than mammalian *STAT5a* and *STAT5b*, it is tempting to speculate that the teleostean *stat5* genes too have adopted at least partially unique functions.

Discussion

We have seen a dynamic evolution of the STAT transcription factors family. Just as for the class-I helical cytokines (Huising *et al.* 2006) and JAKs (Fig. 2), a differential primary sequence conservation for the endocrine and immune STATs is observed, with the endocrine signals being better conserved

than the immune signals. In mammals, it is now clear that STAT3 and STAT5a/b serve in the balance of Treg and Th17 cells (Wei *et al.* 2008). Our understanding of the early vertebrate immune system is not sufficient enough to proclaim that Treg and/or Th17 cells are common aspects of vertebrate immunity or constitute evolutionary recent additions to the mammalian immune systems. Regardless, the strong (endocrine driven) purifying selection that acted on STAT3 and STAT5 may mask the additional weak, immune-driven, purifying selection, resulting from additional roles of STAT3 and STAT5 in immunity.

The continuous threat of invasion by a large array of potential pathogens may have stimulated the evolutionary rate of immune signalling cascades. For example, members of the paramyxovirus family target STAT1 and STAT2 proteins in an attempt to evade the immune response. Some viruses prevent STAT tyrosine phosphorylation, and thus activation; others express STAT ubiquitin ligases, which results in the degradation of STAT proteins (Horvath 2004). STAT1 and STAT2, involved in the anti-viral response downstream of IFNs, display higher pN/pS ratios, indicating a faster rate of evolution, compared with STAT3 and STAT5a/5b. As can be seen in Fig. 2, the family of JAK proteins displays a similar dichotomy in primary sequence conservation. JAK1 and JAK2 serve in the immune system and in the endocrine system, whereas JAK3 and Tyk2 are restricted to immune system. Indeed, JAK1 and JAK2 display shorter branch lengths in phylogenetic analysis, reflecting higher primary sequence conservation.

As the challenges for the immune system are ever-changing, molecular adaptation may provide an answer to these threats. This loop of continuous adaptations of virus and host finds a remarkable homology in the 'Red Queen hypothesis' (Van Valen 1973). This hypothesis states that prey and predator co-evolve as one adapts to the changes of the other in a continuous loop. The endocrine system evolved under relatively constant conditions, as the communication principles in the endocrine system changed relatively little over time. On the other hand, vertebrates are under continuous threat of invasion by a large number of different and continuously evolving potential pathogens, and this may have culminated in an evolutionary arms race between the immune system and the plethora of ever-changing pathogens. Under these conditions, it may have proven advantageous for the vertebrate hosts to relax the constraints of purifying selection sufficiently to enable those STATs that concern themselves with host defence to adapt to the constantly changing playing field of pathogenic insults and thereby contribute to lasting homeostatic equilibrium and reproductive success.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/JOE-11-0033>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Acknowledgments

The authors want to thank Prof. Dr Tom Gerats for constructive comments on the manuscript.

References

- Altschul SF, Madden TL, Schaffer AA, Zhang JH, Zhang Z, Miller W & Lipman DJ 1997 Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* **25** 3389–3402. (doi:10.1093/nar/25.17.3389)
- Aparicio S, Chapman J, Stupka E, Putnam N, Chia JM, Dehal P, Christoffels A, Rash S, Hoon S, Smit A *et al.* 2002 Whole-genome shotgun assembly and analysis of the genome of *Fugu rubripes*. *Science* **297** 1301–1310. (doi:10.1126/science.1072104)
- Bendinelli P, Maroni P, Pecori G, Giraldi F & Piccoletti R 2000 Leptin activates Stat3, Stat1 and AP-1 in mouse adipose tissue. *Molecular and Cellular Endocrinology* **168** 11–20. (doi:10.1016/S0303-7207(00)00313-0)
- Chen W, Daines MO & Khurana Hershey GK 2004 Turning off signal transducer and activator of transcription (STAT): the negative regulation of STAT signaling. *Journal of Allergy and Clinical Immunology* **114** 476–489 (quiz 490). (doi:10.1016/j.jaci.2004.06.042)
- Copeland NG, Gilbert DJ, Schindler C, Zhong Z, Wen Z, Darnell JE, Mui ALF, Miyajima A, Quelle FW, Ihle JN *et al.* 1995 Distribution of the mammalian Stat gene family in mouse chromosomes. *Genomics* **29** 225–228. (doi:10.1006/geno.1995.1235)
- Darnell JE Jr 1997 STATs and gene regulation. *Science* **277** 1630–1635. (doi:10.1126/science.277.5332.1630)
- Darnell JE Jr, Kerr IM & Stark GR 1994 Jak–STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science* **264** 1415–1421. (doi:10.1126/science.8197455)
- Gadina M, Hilton D, Johnston JA, Morinobu A, Lighvani A, Zhou YJ, Visconti R & O'Shea JJ 2001 Signaling by type I and II cytokine receptors: ten years after. *Current Opinion in Immunology* **13** 363–373. (doi:10.1016/S0952-7915(00)00228-4)
- Hino K, Satou Y, Yagi K & Satoh N 2003 A genomewide survey of developmentally relevant genes in *Ciona intestinalis*. VI. Genes for Wnt, TGF β , Hedgehog and JAK/STAT signaling pathways. *Development Genes and Evolution* **213** 264–272. (doi:10.1007/s00427-003-0318-8)
- Horvath MC 2004 Weapons of STAT destruction. *European Journal of Biochemistry* **271** 4621–4628. (doi:10.1111/j.1432-1033.2004.04425.x)
- Horvath CM, Wen Z & Darnell JE Jr 1995 A STAT protein domain that determines DNA sequence recognition suggests a novel DNA-binding domain. *Genes and Development* **9** 984–994. (doi:10.1101/gad.9.8.984)
- Hubbard TJP, Aken BL, Beal K, Ballester B, Caccamo M, Chen Y, Clarke L, Coates G, Cunningham F, Cutts T *et al.* 2007 Ensembl 2007. *Nucleic Acids Research* **35** D610–D617. (doi:10.1093/nar/gkl996)
- Hughes AL & Friedman R 2003 2R or not 2R: testing hypotheses of genome duplication in early vertebrates. *Journal of Structural and Functional Genomics* **3** 85–93. (doi:10.1023/A:1022681600462)
- Hughes AL & Friedman R 2004 Pattern of divergence of amino acid sequences encoded by paralogous genes in human and pufferfish. *Molecular Phylogenetics and Evolution* **32** 337–343. (doi:10.1016/j.ympev.2003.12.007)

- Huising MO, Kruiswijk CP & Flik G 2006 Phylogeny and evolution of class-I helical cytokines. *Journal of Endocrinology* **189** 1–25. (doi:10.1677/joe.1.06591)
- Jaillon O, Aury JM, Brunet F, Petit JL, Stange-Thomann N, Mauceli E, Bouneau L, Fischer C, Ozouf-Costaz C, Bernot A *et al.* 2004 Genome duplication in the teleost fish *Tetraodon nigroviridis* reveals the early vertebrate proto-karyotype. *Nature* **431** 946–957. (doi:10.1038/nature03025)
- Kondrashov FA, Rogozin IB, Wolf YI & Koonin EV 2002 Selection in the evolution of gene duplications. *Genome Biology* **3** RESEARCH0008. (doi:10.1186/gb-2002-3-2-research0008)
- Kumar S, Tamura K & Nei M 2004 MEGA 3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Briefings in Bioinformatics* **5** 150–163. (doi:10.1093/bib/5.2.150)
- Levy DE & Darnell JE Jr 2002 Stats: transcriptional control and biological impact. *Nature Reviews. Molecular Cell Biology* **3** 651–662. (doi:10.1038/nrm909)
- Lewis RS & Ward AC 2004 Conservation, duplication and divergence of the zebrafish *stat5* genes. *Gene* **338** 65–74. (doi:10.1016/j.gene.2004.05.012)
- Liongue C & Ward AC 2007 Evolution of class I cytokine receptors. *BMC Evolutionary Biology* **7** 120. (doi:10.1186/1471-2148-7-120)
- Maritano D, Sugrue ML, Tininini S, Dewilde S, Strobl B, Fu X, Murray-Tait V, Chiarle R & Poli V 2004 The STAT3 isoforms α and β have unique and specific functions. *Nature Immunology* **5** 401–409. (doi:10.1038/ni1052)
- Meyer A & Van de Peer Y 2005 From 2R to 3R: evidence for a fish-specific genome duplication (FSGD). *BioEssays* **27** 937–945. (doi:10.1002/bies.20293)
- Muffato M & Roest-Crollius H 2008 Paleogenomics in vertebrates, or the recovery of lost genomes from the mist of time. *BioEssays* **30** 122–134. (doi:10.1002/bies.20707)
- Nei M & Gojobori T 1986 Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Molecular Biology and Evolution* **3** 418–426.
- O'Shea JJ, Gadina M & Schreiber RD 2002 Cytokine signaling in 2002: new surprises in the Jak/Stat pathway. *Cell* **109** 121–131. (doi:10.1016/S0092-8674(02)00701-8)
- Schindler CW 2002 JAK–STAT signaling in human disease. *Journal of Clinical Investigation* **109** 1133–1137. (doi:10.1172/JCI15644)
- Schindler C, Shuai K, Prezioso VR & Darnell JE Jr 1992 Interferon-dependent tyrosine phosphorylation of a latent cytoplasmic transcription factor. *Science* **257** 809–813. (doi:10.1126/science.1496401)
- Sharman AC & Holland PWH 1996 Conservation, duplication, and divergence of developmental genes during chordate evolution. *Netherlands Journal of Zoology* **46** 47–67. (doi:10.1163/156854295X00050)
- Shuai K & Liu B 2003 Regulation of JAK–STAT signalling in the immune system. *Nature Reviews. Immunology* **3** 900–911. (doi:10.1038/nri1226)
- Shuai K, Stark GR, Kerr IM & Darnell JE Jr 1993 A single phosphotyrosine residue of Stat91 required for gene activation by interferon- γ . *Science* **261** 1744–1746. (doi:10.1126/science.7690989)
- Shuai K, Horvath CM, Tsai Huang LH, Qureshi SA, Cowburn D & Darnell JE Jr 1994 Interferon activation of the transcription factor Stat91 involves dimerization through SH2–phosphotyrosyl peptide interactions. *Cell* **76** 821–828. (doi:10.1016/0092-8674(94)90357-3)
- Sidow A 1996 Gen(om)e duplications in the evolution of early vertebrates. *Current Opinion in Genetics & Development* **6** 715–722. (doi:10.1016/S0959-437X(96)80026-8)
- Stein C, Caccamo M, Laird G & Leptin M 2007 Conservation and divergence of gene families encoding components of innate immune response systems in zebrafish. *Genome Biology* **8** R251. (doi:10.1186/gb-2007-8-11-r251)
- Van Valen L 1973 A new evolutionary law. *Evolutionary Theory* **1** 1–30.
- Wei L, Laurence A & O'Shea JJ 2008 New insights into the roles of Stat5a/b and Stat3 in T cell development and differentiation. *Seminars in Cell and Developmental Biology* **19** 394–400. (doi:10.1016/j.semcdb.2008.07.011)
- Wittbrodt J, Meyer A & Schartl M 1998 More genes in fish? *BioEssays* **20** 511–515. (doi:10.1002/(SICI)1521-1878(199806)20:6<511::AID-BIES10>3.0.CO;2-3)

Received in final form 24 January 2011

Accepted 17 February 2011

Made available online as an Accepted Preprint

17 February 2011