



## Toxic effect of herbicides used for water hyacinth control on two insects released for its biological control in South Africa

Martin P. Hill , Julie A. Coetzee & Claudia Ueckermann

To cite this article: Martin P. Hill , Julie A. Coetzee & Claudia Ueckermann (2012) Toxic effect of herbicides used for water hyacinth control on two insects released for its biological control in South Africa, *Biocontrol Science and Technology*, 22:11, 1321-1333, DOI: [10.1080/09583157.2012.725825](https://doi.org/10.1080/09583157.2012.725825)

To link to this article: <https://doi.org/10.1080/09583157.2012.725825>



Accepted author version posted online: 31 Aug 2012.  
Published online: 31 Aug 2012.



Submit your article to this journal [↗](#)



Article views: 214



Citing articles: 8 View citing articles [↗](#)

## RESEARCH ARTICLE

### Toxic effect of herbicides used for water hyacinth control on two insects released for its biological control in South Africa

Martin P. Hill<sup>a\*</sup>, Julie A. Coetzee<sup>a</sup> and Claudia Ueckermann<sup>b</sup>

<sup>a</sup>Department of Zoology and Entomology, Rhodes University, Grahamstown, South Africa;

<sup>b</sup>Department of Botany, University of Pretoria, Pretoria, South Africa

(Received 20 July 2012; returned 20 August 2012; accepted 28 August 2012)

The integrated control of water hyacinth, *Eichhornia crassipes* (Martius) Solms-Laubach (Pontederiaceae) has become necessary in South Africa, as biological control alone is perceived to be too slow in controlling the weed. In total, seven insect biological control agents have been released on water hyacinth in South Africa. At the same time, herbicides are applied by the water authorities in areas where the weed continues to be troublesome. This study investigated the assumption that the two control methods are compatible by testing the direct toxicity of a range of herbicide formulations and surfactants on two of the biological control agents released against water hyacinth, the weevil, *Neochetina eichhorniae* Warner (Coleoptera: Curculionidae) and the water hyacinth mirid, *Eccritotarsus catarinensis* (Carvalho) (Hemiptera: Miridae). A number of the formulations used resulted in significant mortality of the mirid and the weevil. Products containing 2,4-D amine and diquat as active ingredients caused higher mortality of both agents (up to 80% for the mirid) than formulations containing glyphosate. Furthermore, when surfactants were added to enhance herbicide efficiency, it resulted in increased toxicity to the insects. We recommend that glyphosate formulations should be used in integrated control programmes, and that surfactants be avoided in order to reduce the toxic nature of spray formulations to the insect biological control agents released against water hyacinth.

**Keywords:** *Eccritotarsus catarinensis*; *Eichhornia crassipes*; integrated control; *Neochetina eichhorniae*

#### Introduction

Water hyacinth, *Eichhornia crassipes* (Martius) Solms-Laubach (Pontederiaceae), is a perennial, herbaceous, free-floating, aquatic plant native to the Amazon Basin of South America (Center 1994). This species is now widespread throughout the tropical and subtropical areas of the world (Holm, Plucknett, Pancho, and Herurger 1977), and is invasive in South Africa where dense mats of the weed degrade aquatic ecosystems (Midgley, Hill, and Villet 2006) and severely limit their use (Hill 2003). Biological control of water hyacinth has been highly successful in some areas of South Africa, where a suite of seven biological control agents have been used to reduce populations of the weed (Coetzee, Hill, Byrne, and Bownes 2011). However, in colder areas of the country (Hill and Olckers 2001) and in eutrophic waters, biological control has been less successful (Coetzee, Byrne, and Hill 2007; Coetzee

---

\*Corresponding author. Email: m.hill@ru.ac.za

and Hill 2012). In these areas, biological control is considered to be inadequate, or too slow acting and attempts have been made to integrate different control methods to achieve the benefits of both short-term and sustainable, long-term control (Cilliers, Campbell, Naude, and Nesor 1996; Coetzee and Hill 2012). The integration of biological and herbicide control is currently the most widely advocated method. However, this approach relies on the assumption that these two methods are compatible and especially that the herbicides themselves are not toxic to the biological control agents (Hill and Olckers 2001).

There have been a number of studies investigating the direct effect of herbicides on some of the biological control agents released on water hyacinth (e.g. Roorda, Schulten, and Pieterse 1978; Center, Stewart, and Bruner 1982; Haag 1986a,b; Haag, Glenn, and Jordan 1988; Wright and Skilling 1987; Grodowitz and Pellessier 1990; Jianqing, Wang, Zhiquan, and Weidong 1999; Jadhav, Hill, and Byrne 2008). Generally, these studies found that the herbicide increased mortality of the biological control species. It was also found that it was not always the active ingredient in the herbicide that made it toxic to the agents, but rather the surfactants and other additives (e.g. drift retardants) that caused mortality (Affeld, Hill, Smith, and Syrett 2004). In addition, the rapid loss of habitat through the sinking of the water hyacinth mat after herbicide application decimated biological control agent populations, and required re-inoculation once the mat had regenerated through seedling recruitment (Center, Dray, Jubinsky, and Grodowitz 1999).

In an attempt to improve integrating herbicides with water hyacinth biological control agents, the toxicity of a number of herbicides registered for use on the weed in South Africa and elsewhere in the world was tested on two of the species released as biological control agents. The two agents chosen were the water hyacinth weevil, *Neochetina eichhorniae* Warner (Coleoptera: Curculionidae), which has been released in at least 30 countries around the world (Julien and Griffiths 1998) and the leaf-sucking mirid, *Eccritotarsus catarinensis* (Carvalho) (Heteroptera: Miridae), which was released on water hyacinth in South Africa in 1996 (Hill, Cilliers, and Nesor 1999). These two agents differ in that the adult weevils are nocturnal, hiding amongst the youngest, rolled water hyacinth leaves during the day and the larvae feed within the plant tissue (DeLoach and Cordo 1976), whereas *E. catarinensis* nymphs and adults feed externally on the leaf surface during the day (Hill et al. 1999).

The approach to integrated control of water hyacinth in South Africa employed in the larger systems, such as the Vaal River, is to spray 75% of the weed infestation with herbicides and leave a 25% 'reserve' of the weed onto which the biological control agents can move (Cilliers et al. 1996). This approach has also been suggested in the USA (Haag et al. 1988) and has the benefit of providing both short- and long-term control. However, this method relies on the assumptions that the biological control agents can and do disperse and not sink with the sprayed plants. Grodowitz, Center, and Freedman (1997) showed that adult *N. eichhorniae* and its congener *Neochetina bruchi* (Hustache) frequently had reduced wing musculature and would thus not be capable of flight, and the sessile stages of all of the agents would sink with the mat. Secondly, the method assumes that the natural enemy populations in the 'reserve' are large enough to curb the spread of the weed, and the reserve itself will not become the source of re-infestation. Lastly, it assumes that the herbicides themselves are not directly toxic to the biological control agents. It is the last assumption that was tested in this study.

## Materials and methods

### *Herbicide selection*

In this study, a series of herbicide formulations (Table 1) was applied directly to the two insect test species. Diquat, glyphosate and glyphosate-trimesium are registered for use on water hyacinth in South Africa (Vermeulen, Grobler, and Van Zyl 1998), and although 2,4-D amine is not, it was included as it is popular in neighbouring countries due to its effectiveness and affordability. For each product, a range of concentrations was tested, from below to far above the manufacturers' recommended rates for water hyacinth control. In addition to these herbicides, two surfactants were also tested; Agral<sup>®</sup>, which is recommended for use with Midstream<sup>®</sup> (diquat) and Add-2<sup>®</sup>, which is recommended for use with Touchdown<sup>®</sup> (glyphosate-trimesium). Combinations of Midstream<sup>®</sup> with Agral<sup>®</sup>, and Touchdown<sup>®</sup> with Add-2<sup>®</sup>, were tested to determine whether adding surfactants to the herbicides affected toxicity.

### *Test organisms*

#### *Eccritotarsus catarinensis*

Insects were collected approximately 2 days before treatment from ponds of water hyacinth at the Plant Protection Research Institute (PPRI) in Pretoria, South Africa, and kept in tubs with recently severed water hyacinth leaves from plants maintained at the Institute. Insects were sorted into groups of eight adults (four males and four females) and placed in small Petri-dishes (65 mm diameter) lined with moist filter paper, 1 day before treatment in order to acclimatise. Fresh dilutions of test chemicals were mixed less than 1 hour before application. A 20- $\mu$ l droplet of chemical was applied dorsally to each insect. A chemical-free control also was set up, whereby a 20- $\mu$ l droplet of distilled water was applied to each insect. Insect mortality was monitored every 24 hours for up to 120 hours after application of the chemical. Water hyacinth leaf pieces (20  $\times$  30 mm) were provided as food every 24 hours and the humidity in the Petri-dish was maintained by moistening the filter paper with water. Petri-dishes were exposed to a light regime of 16 hours light, 8 hours darkness and temperatures varied between 22 and 28°C. Three replicates of eight insects were used for each of the chemical treatments and the control.

#### *Neochetina eichhorniae*

Adult weevils were collected from a field site (Bon Accord Dam 25°38'15.4"S/28°11'01.6"W), just north of Pretoria, approximately 2 weeks before initiation of the experiment and kept in plastic tubs with recently severed water hyacinth leaves, from plants maintained in pools at PPRI. Six weevils (three male and three female) were placed in 500 ml plastic containers lined with moistened filter paper and aerated through small holes in the top, at least 48 hours before commencement of the experiment. Insects were exposed to different chemical concentrations by applying a 20- $\mu$ l droplet dorsally to the elytra of the insect. Again, a distilled water control was set up. Food was provided daily in the form of a severed water hyacinth leaf. The experiment was conducted under 16 hours light, 8 hours darkness and temperatures varied between 22 and 28°C. Mortality was noted every 24 hours for up to 120 hours

Table 1. Specifications of the herbicides and surfactants selected for testing (formulation composition and recommended dosages), and dosages tested on two agents released for the biological control of water hyacinth in South Africa.

Brand name	Manufacturer	Content	g a.e/l	Recommended dosages (% vol/vol)	Dosages tested (% vol/vol)	
2,4-D amine	Sanachem	2,4-Dichlorophenoxyacetic acid 480 amine	480	2.00–6.00	1, 3, 9, 18	
Mamba 360 SL <sup>a</sup>	Sanachem	Glyphosate SL (IPA salt)	360	2.00–6.00	3, 4, 12, 24	
Midstream <sup>a</sup>	Zeneca	Diquat (dibromide salt)	373.5	3.75–5.00	1, 2, 3.75, 4, 5	
Mon 52276	Monsanto	Glyphosate (IPA salt)	360	2.00–6.00	3, 4, 12, 24	
Muster <sup>a</sup>	Zeneca	Glyphosate (Trimethyl sulfonium salt)	330	2.00–3.00	2, 3, 12, 24	
Rodeo <sup>a</sup>	Monsanto	Glyphosate (IPA salt)	480	4.50–9.00	1.5, 3, 4, 9, 18	
Roundup <sup>a</sup>	Monsanto	Glyphosate (IPA salt)	360	2.00–6.00	1, 2, 3, 4, 12, 24	
Roundup Ultra <sup>a</sup>	Monsanto	Glyphosate	360	2.00–6.00	3, 4, 12, 24	
Touchdown <sup>a</sup>	Zeneca	Glyphosate (Trimethyl sulfonium salt)	480	2.00	2, 4, 8, 20	
Tumbleweed <sup>a</sup>	Enviro Weed Control	Glyphosate (IPA salt)	240	2.00–6.00	1, 4, 12, 24, 50	
Surfactants		Description				
Add-2 <sup>a</sup>	Zeneca	Spreader adjuvant (with Touchdown)	Polysaccharide (hexitan ester)	600	0.20–0.30	0.2, 0.3, 0.6, 1.0
Agral <sup>a</sup>	Zeneca	Wetting and sticking adjuvant (with Midstream)	Alkylated phenol-ethylene oxide concentrate	940	0.75	0.05, 0.75, 1

g a.e/l, gram acid equivalent per litre.

<sup>a</sup>Registered under Act 36 (1947) for control of water hyacinth.

after chemical application. Three replicates of six insects each were used for each treatment and the control.

### *Statistical analyses*

A lethal concentration ( $LC_{50}$ ) value for each herbicide and/or adjuvant was obtained for each insect species, after 24 hours and 120 hours exposure, by deriving regressions of insect mortality against chemical concentration, using Probit regression analysis in Statistica v10 (StatSoft, Inc. 2011). The safety of each chemical was then rated based on its toxicity to the biocontrol agents, and recommended dosages used in practice.

### **Results**

The two biological control agents responded differently to the chemical applications: the mirid was far more susceptible to the chemicals than the weevil. Generally, higher mortality of the mirid was recorded at higher concentrations, which increased over time until 72 hours, after which mortality was constant (Figure 1). The herbicides Rodeo® and Roundup® were the only two that did not cause 100% mortality of the mirid at any of the concentrations tested. Midstream® caused high mortality at all concentrations over the entire time period, while the addition of the surfactant, Agral®, caused 100% mortality at both concentrations tested. All of the other herbicides tested induced high mortality, particularly at higher dosages. The surfactant Agral® used alone was toxic to the mirid at all concentrations tested, but mortality of the mirid exposed to Add-2®, the other surfactant tested, was less than 60% at all of the dosages, except the highest dosage used, where mean mortality was  $91 \pm 0.93\%$  across the time period. Mirid mortality remained between 8 and 17% across the time period for the control treatment.

The  $LC_{50}$  values were generally higher after 24 hours exposure than 120 hours exposure to the chemicals tested, implying that toxicity increased with time (Table 2). In certain cases (Midstream®, Muster®, Rodeo®, Roundup® and Touchdown®), the  $LC_{50}$ s could not be calculated for either or both the 24 and 120 hour periods, because they fell outside of the range of concentrations tested.

Based on the  $LC_{50}$  values and the recommended doses of each chemical, all the herbicides that were rated as safe for use at recommended dosages with *E. catarinensis* were glyphosate-based, namely Mamba®, Rodeo® and Roundup®. The surfactant Add-2® also was rated safe, whereas the two glyphosate-trimesium products, Muster® and Touchdown®, and 2,4-D amine, were both rated toxic. Applying Add-2® (41.7% mortality) with Touchdown® (4.2% mortality) increased the toxicity of Touchdown® (33.3% mortality) (Figure 1). The diquat-based herbicide Midstream® was the most toxic herbicide tested against *E. catarinensis*, 100% mortality was recorded for all concentrations of Midstream® at 96 hours after treatment, confirming its toxic nature (Figure 1). The surfactant Agral® was also considered toxic on its own (62.5% mortality), and when combined with Midstream® (93.3% mortality; Table 2, Figure 1).

The weevil *N. eichhorniae* was less susceptible to the toxic effects of the herbicides and/or adjuvants used than *E. catarinensis*. Weevil mortality was low overall and only Agral®, Muster® and Roundup Ultra® resulted in high mortality (Figure 2).




Figure 1. Mean percentage mortality of *Eccritotarsus catarinensis* as a result of direct treatment with 12 selected herbicides and herbicide/adjuvant combinations, at various concentrations over five time intervals. For comparison, mirid mortality in the control treatment was low: 24 hours – 8%; 48 hours – 10%; 72 hours – 15%; 96 and 120 hours – 17%. Legends represent % concentration (%vol/vol) of herbicide applied. Error bars represent 95% CI.




Figure 1. Continued



Table 2. LC<sub>50</sub> values, after 24 and 120 hours of exposure to herbicides and surfactants registered for use in the control of water hyacinth, and the rating of each according to the effect that they had on the control agents, *Eccritotarsus catarinensis* and *Neochetina eichhorniae*, during acute toxicity tests.

Herbicide <sup>a</sup>	Dosages (%) used in practice <sup>b</sup>	<i>E. catarinensis</i>			<i>N. eichhorniae</i>		
		LC <sub>50</sub> ± SE (χ <sup>2</sup> value, *P < 0.05)		Rating <sup>c</sup>	LC <sub>50</sub> ± SE (χ <sup>2</sup> value, *P < 0.05)		Rating <sup>c</sup>
		24 h	120 h		24 h	120 h	
2,4-D amine	2.0–6.0	4.3 ± 0.02 (χ <sup>2</sup> = 192.0*)	0.7 ± 0.04 (χ <sup>2</sup> = 120.9*)	Toxic			Safe
Add-2	0.2–0.3	0.81 ± 0.11 (χ <sup>2</sup> = 135.3*)	0.38 ± 0.11 (χ <sup>2</sup> = 62.4*)	Safe			Safe
Agral	0.8	0.3 ± 0.18 (χ <sup>2</sup> = 11.2*)	< 0.01 (χ <sup>2</sup> = 3.18)	Toxic	0.69 ± 0.66 (χ <sup>2</sup> = 119.4*)	0.63 ± 0.58 (χ <sup>2</sup> = 138.2*)	Toxic
Mamba	2.0–6.0	10.1 ± 0.01 (χ <sup>2</sup> = 242.8*)	7.8 ± 0.02 (χ <sup>2</sup> = 246.8*)	Safe			Safe
Midstream	3.8–5.0	< 0.05 (χ <sup>2</sup> = 18.3*)	NA (100% mortality)	Toxic			Safe
Mon 52276	2.0–6.0	8.56 ± 0.01 (χ <sup>2</sup> = 144.4*)	4.2 ± 0.02 (χ <sup>2</sup> = 142.3*)	Hazardous			Safe
Muster	2.0–3.0	< 0.01 (χ <sup>2</sup> = 91.2*)	1.46 ± 0.20 (χ <sup>2</sup> = 83.9*)	Toxic	10.73 ± 0.02 (χ <sup>2</sup> = 420.0*)	8.54 ± 0.03 (χ <sup>2</sup> = 516.2*)	Safe
Rodeo	4.5–9.0	> 18 (χ <sup>2</sup> = 44.2*)	> 18 (χ <sup>2</sup> = 15.7*)	Safe			Safe
Roundup	2.0–6.0	> 26 (χ <sup>2</sup> = 23.9*)	> 26 (χ <sup>2</sup> = 0.9)	Safe			Safe
Roundup ultra	2.0–6.0	4.38 ± 0.02 (χ <sup>2</sup> = 172.3*)	< 2 (χ <sup>2</sup> = 93.8*)	Toxic	> 26 (χ <sup>2</sup> = 22.2*)	18.89 ± 0.01 (χ <sup>2</sup> = 59.4*)	Safe
Touchdown	2.0	> 20 (χ <sup>2</sup> = 11.34*)	< 0.01 (χ <sup>2</sup> = 0.28)	Toxic <sup>d</sup>			Safe
Tumbleweed	2.0–6.0	10.48 ± 0.01 (χ <sup>2</sup> = 214.1*)	3.61 ± 0.01 (χ <sup>2</sup> = 170.5*)	Hazardous			Safe

<sup>a</sup>Formulation or surfactant.

<sup>b</sup>Refer to product labels for specified dosages and application instructions.

<sup>c</sup>Safe, LC<sub>50</sub> never fell within recommended dosage range; Hazardous, LC<sub>50</sub> fell within or below the recommended dosage range at any time during 120 hours; Toxic, LC<sub>50</sub> fell within or below the recommended dosage range from 24 hours.

<sup>d</sup>Considered toxic as it caused high mortality from 72 hours onwards.




Figure 2. Mean percentage mortality of *Neochetina eichhorniae* as a result of direct treatment with four selected herbicides and herbicide/adjuvant combinations, at various concentrations over five time intervals. Control mortality of the weevils was 0% at all of the time intervals. Legends represent % concentration (%vol.vol) of herbicide applied. Error bars represent 95% CI.

Zero mortality was experienced for the control treatment. The surfactant Agra1<sup>®</sup> was the only chemical rated toxic to the weevil and all the other chemicals were found to be safe at recommended dosages (Table 2).

## Discussion

This study showed that many of the herbicides/surfactants used in the control of water hyacinth resulted in significant mortality of two of the agents released as biological control agents. The mirid *E. catarinensis* was more susceptible to the toxic effects of the chemicals than the weevil *N. eichhorniae*. Herbicides containing glyphosate as an active ingredient were less toxic, whereas formulations with surfactants were more toxic. The study also showed that toxicity increased at higher concentrations. These results are consistent with other studies (e.g. Roorda et al. 1978; Center et al. 1982; Haag 1986a; Wright and Skilling 1987; Grodowitz and Pellessier 1990; Jianqing et al. 1999) that showed some level of toxicity of herbicide formulations on water hyacinth biological control agents.

Most herbicides are thought to have low toxicity to animals because the active ingredients have been developed to act on plant pathways such as the shikimic acid pathway (Franz, Mao, and Sikorski 1997). However, the surfactants added to herbicide formulations are designed to breakdown the surface tension of the herbicide and increase coverage. They also play a role in dissolving the waxy cuticle of plants, increasing the uptake of herbicide (Affeld et al. 2004). Therefore, the surfactants also may be harmful to insects by destroying the waxy cuticle of the exoskeleton. Wax removal would lead to water loss and dehydration, or possibly liquid flooding into the trachea via the spiracles, impeding gas exchange and, although not tested in this study, direct toxicity could occur due to ingestion of herbicide contaminated food (Ainsworth 2003).

Herbicide application causes biochemical changes that might alter the palatability of water hyacinth plants, making them either more, or less attractive to the biological control agents. Wright and Bourne (1990) showed that the application of 2,4-D amine onto water hyacinth decreased the laminar hardness and increased the nitrogen content of the plants, thereby improving plant quality for *N. eichhorniae*, *N. bruchi* and the moth *Niphograptia albiguttalis* Warren [= *Sameodes albiguttalis* (Warren)] (Lepidoptera: Crambidae), whereas applications of glyphosate-based products do not appear to change the plant quality (Jadhav et al. 2008).

Extrapolation of these results from the laboratory to the field must take the behaviour of the insect species into consideration. The weevil is nocturnal, hiding in the base of the petioles during the day when herbicide application occurs, feeding on the leaves at night. The weevil is, therefore, highly unlikely to come into direct contact with the herbicide, and if it does, this study has shown that it will not be greatly affected. The mirid, however, feeds on the leaves during the day (Hill et al. 1999) and is, therefore, likely to come into contact with droplets of the herbicide. Thus, mirid populations will be negatively impacted by herbicide application, not only through the loss of habitat, but also acute toxicity.

Populations of all of the agents thus far released on water hyacinth are negatively affected by herbicide interventions due to mortality of sessile stages that are unable to disperse from a sinking mat. In an attempt to better integrate herbicide and biological control, Haag (1986b) suggested that water hyacinth mats should be

sprayed in strips leaving unsprayed sections of the mat for the weevils to move on to, but this still resulted in mortality of the eggs, larvae and pupae in the sprayed sections. Van and Center (1994) showed that the growth retardant, paclobutrazol, acted synergistically with *N. eichhorniae* to control water hyacinth, although this was never implemented in the field. In a more recent study, Jadhav et al. (2008) showed in the laboratory that by spraying water hyacinth with a sub-lethal dose of glyphosate (0.8%) as opposed to the recommended dose of 3%, the plants did not die, but did stop growing, flowering and producing daughter plants. The low dose preserved the habitat for the immature stages of the *Neochetina* weevils, allowing them to build up high populations, but would still result in high mortalities of the more susceptible species such as *E. catarinensis*.

For the integrated management of water hyacinth to succeed in South Africa, site specific plans need to be developed that take into consideration compatibility of the insects with the herbicides to be used. Some herbicide formulations are better suited to integrated control where insects are already established. For example, in systems where *E. catarinensis* populations are to be maintained, herbicides with glyphosate as an active ingredient, a low surfactant content and preferably at low concentrations should be used. In systems where *N. eichhorniae* is present, the choice of herbicide formulation is less vital. Timing and mode of application also are critical to successful integrated control of water hyacinth. In South Africa, the weevils, *N. eichhorniae* and *N. bruchi* overwinter as third instars in the crown of the plant (Byrne et al. 2010), thus herbicide applications in early spring (September and October) should be avoided as these insect stages are unable to move onto unsprayed plants. The timing of releases of agents could also be manipulated where susceptible agents such as the mirid could be released after herbicide application onto mats left unsprayed. Most of the agents used as biological control agents against water hyacinth in the world can easily be mass-reared (Julien, Griffiths, and Wright 1999; Julien, Griffiths, and Stanley 2001) and could be used in this way. Although the direct application of herbicides to several of the biological control agents released against *E. crassipes* has been shown to be toxic, a thorough understanding of the ecology of the weed and the arthropods used in its control, and the formulations of herbicides has shown that chemical and biological control of this weed can be integrated in a compatible manner.

### Acknowledgements

We would like to thank the late Mr. I.G. Oberholzer of the Plant Protection Research Institute for his technical assistance, and the University of Pretoria for the use of their facilities. This work was funded by the Water Research Commission of South Africa (Project 915/1/01). The sponsorship of chemicals and advice received from Enviro Weed Control, Monsanto, Sanachem and Zeneca are greatly appreciated.

### References

- Affeld, K., Hill, K., Smith, L.A., and Syrett, P. (2004), 'Toxicity of Herbicides and Surfactants to Three Insect Biological Control Agents for *Cytisus scoparius* (Scotch broom)', in *Proceedings of the XI International Symposium on Biological Control of Weeds*, eds. J.M. Cullen, D.T. Briese, D.J. Kriticos, W.M. Lonsdale, L. Morin and J.K. Scott, April 27–May 2, 2003, Canberra, Australia, pp. 375–380.

- Ainsworth, N. (2003), 'Integration of Herbicides with Arthropod Biocontrol Agents for Weed Control', *Biocontrol Science and Technology*, 13, 547–570.
- Byrne, M., Hill, M.P., Robertson, M.P., King, A., Jadhav, A., Katembo, N., Wilson, J.R., Brudvig, R., and Fisher, J. (2010), 'Integrated Management of Water Hyacinth in South Africa: Development of an Integrated Management Plan for Water Hyacinth Control, Combining Biological Control, Herbicide Control and Nutrient Control, Tailored to the Climatic Regions of South Africa', WRC Report No. TT 454/10, p. 285.
- Center, T.D. (1994), 'Biological Control of Weeds: Waterhyacinth and Waterlettuce', in *Pest Management in the Tropics. Biological Control – A Florida Perspective*, eds. D. Rosen, F.D. Bennett and J.L. Capinera, Andover: Intercept Ltd., pp. 481–521.
- Center, T.D., Dray, F.A., Jubinsky, G.P., and Grodowitz, M.J. (1999), 'Biological Control of Water Hyacinth under Conditions of Maintenance Control: Can Herbicides and Insects be Integrated?', *Environmental Management*, 23, 241–256.
- Center, T.D., Stewart, K.K., and Bruner, M.C. (1982), 'Control of Waterhyacinth (*Eichhornia crassipes*) with *Neochetina eichhorniae* (Coleoptera: Curculionidae) and a Growth Retardant', *Weed Science*, 30, 453–457.
- Cilliers, C.J., Campbell, P.L., Naude, D., and Naser, S. (1996), 'An Integrated Water Hyacinth Control Programme on the Vaal River, in a Cool, High Altitude Area in South Africa', in *Strategies for Water Hyacinth Control. Report of a Panel of Experts Meeting*, ed. R. Charudattan, 11–14 September 1995, Fort Lauderdale, Florida, USA., pp. 87–103.
- Coetzee, J.A., Byrne, M.J., and Hill, M.P. (2007), 'Impact of Nutrients and Herbivory by *Eccritotarsus catarinensis* on the Biological Control of Water Hyacinth, *Eichhornia crassipes*', *Aquatic Botany*, 87, 179–186.
- Coetzee, J.A., and Hill, M.P. (2012), 'The Role of Eutrophication in the Biological Control of Water Hyacinth, *Eichhornia crassipes*, in South Africa', *BioControl*, 57, 247–261.
- Coetzee, J.A., Hill, M.P., Byrne, M.J., and Bownes, A.B. (2011), 'A Review of the Biological Control Programmes on *Eichhornia crassipes* (C. Mart.) Solms (Pontederiaceae), *Salvinia molesta* D.S. Mitch. (Salviniaceae), *Pistia stratiotes* L. (Araceae), *Myriophyllum aquaticum* (Vell.) Verdc. (Haloragaceae) and *Azolla filiculoides* Lam. (Azollaceae) in South Africa Since 1999', *African Entomology*, 19, 451–468.
- DeLoach, C.J., and Cordo, H.A. (1976), 'Life Cycle and Biology of *Neochetina bruchi*, a Weevil Attacking Waterhyacinth in Argentina, with Notes on *N. eichhorniae*', *Annals of the Entomological Society of America*, 69, 643–652.
- Franz, J.E., Mao, M.K., and Sikorski, J.A. (1997), '*Glyphosate: A Unique Global Herbicide*', Chemical Society Monograph 189. Washington DC, USA: American Chemical Society, p. 653.
- Grodowitz, M.J., Center, T.D., and Freedman, J.E. (1997), 'A Physiological Age – Grading system for *Neochetina eichhorniae* (Warner) (Coleoptera: Curculionidae), a Biological Control Agent of Water Hyacinth, *Eichhornia crassipes* (Mart.) Solms', *Biological Control*, 9, 89–105.
- Grodowitz, M.J., and Pellessier, D.C. (1990), '*Effects of Chemical Applications on the Biological Control Agents of Water Hyacinth*', US Army of Engineers Unpublished Report.
- Haag, K.H. (1986a), 'Effects of Herbicidal Application on Mortality and Dispersive Behaviour of the Water Hyacinth Weevils, *Neochetina eichhorniae* and *Neochetina bruchi* (Coleoptera: Curculionidae)', *Environmental Entomology*, 15, 1193–1198.
- Haag, K.H. (1986b), 'Effective Control of Water Hyacinth Using *Neochetina* and Limited Herbicide Application', *Journal of Aquatic Plant Management*, 15, 70–75.
- Haag, K.H., Glenn, M.S., and Jordan, J.C. (1988), 'Selective Patterns of Herbicide Application for Improved Biological Control of Waterhyacinth', *Journal of Aquatic Plant Management*, 26, 17–19.
- Hill, M.P. (2003), 'The Impact and Control of Alien Aquatic Vegetation in South African Aquatic Ecosystems', *African Journal of Aquatic Science*, 28, 19–24.
- Hill, M.P., Cilliers, C.J., and Naser, S. (1999), 'Life History and Laboratory Host Range of *Eccritotarsus catarinensis* (Carvalho) (Heteroptera: Miridae) a New Natural Enemy Released on Water Hyacinth (*Eichhornia crassipes* (Mart.) Solms-Laub.) (Pontederiaceae) in South Africa', *Biological Control*, 14, 127–133.

- Hill, M.P., and Olckers, T. (2001), 'Biological Control Initiatives Against Water Hyacinth in South Africa: Constraining Factors, Success and New Courses of Action', in *Proceedings of the Second Global Working Group Meeting for the Biological and Integrated Control of Water Hyacinth*, Eichhornia crassipes, eds. M.H. Julien, M.P. Hill, T.D. Center and D. Jianqing, October 9–12, 2000, Beijing, China. ACIAR Proceedings No. 102, pp. 33–38.
- Holm, L.G., Plucknett, D.L., Pancho, J.V., and Herurger, J.P. (1977), *The World's Worst Weeds, Distribution and Biology*, Honolulu, Hawaii: The University Press of Hawaii.
- Jadhav, A., Hill, M.P., and Byrne, M. (2008), 'Identification of a Retardant Dose of Glyphosate with Potential for Integrated Control of Water Hyacinth, *Eichhornia crassipes* (Mart.) Solms-Laubach', *Biological Control*, 47, 154–158.
- Jianqing, D., Wang, R., Zhiquan, C., and Weidong, F. (1999), 'Towards Integrated Management of Water Hyacinth with Insects and Herbicides in Southern China', in *Proceedings of the First IOBC Global Working Group Meeting for the Biological and Integrated Control of Water Hyacinth*, eds. M.P. Hill, M.H. Julien, and T.D. Center, November 16–19, 1998, Zimbabwe. Pretoria, South Africa: Plant Protection Research Institute, pp. 42–147.
- Julien, M.H., and Griffiths, M.W. (1998), *Biological Control of Weeds. A World Catalogue of Agents and their Target Weeds*, (4th ed.), New York: CABI Publishing.
- Julien, M.H., Griffiths, M.W., and Wright, A.D. (1999), 'Biological Control of Water Hyacinth. The Weevils, *Neochetina bruchi* and *N. eichhorniae*: Biologies, Host Ranges, and Rearing, Releasing and Monitoring Techniques for Biological Control of *Eichhornia crassipes*', *ACIAR Monograph No. 60*, 87.
- Julien, M.H., Griffiths, M.W., and Stanley, J.N. (2001), 'Biological Control of Water Hyacinth 2. The Moths, *Niphograptia albiguttalis* and *Xubida infusellus*: Biologies, Host Ranges, and Rearing, Releasing and Monitoring Techniques for Biological Control of *Eichhornia crassipes*', *ACIAR Monograph No. 79*, 91.
- Midgley, J.M., Hill, M.P., and Villet, M.H. (2006), 'The Effect of Water Hyacinth, *Eichhornia crassipes* (Martius) Solms-Laubach (Pontederiaceae), on Benthic Biodiversity in Two Impoundments on the New Year's River, South Africa', *African Journal of Aquatic Science*, 31, 25–30.
- Roorda, F.A., Schulten, G.G.M., and Pieterse, A.H. (1978), 'The Susceptibility of *Orthogalumna terebrantis* Wallwork (Acarina: Galumnidae) to Various Pesticides', in *Proceedings of the European Weeds Research Society 5th Symposium on Aquatic Weeds*, pp. 375–381.
- StatSoft, Inc. (2011), 'STATISTICA (Data Analysis Software System), Version 10'. www.statsoft.com.
- Van, T.K., and Center, T.D. (1994), 'Effect of Paclobutrazol and Water Hyacinth Weevil (*Neochetina eichhorniae*) on Plant Growth and Leaf Dynamics of Water Hyacinth (*Eichhornia crassipes*)', *Weed Science*, 42, 665–672.
- Vermeulen, J.B., Grobler, H., and Van Zyl, K. (1998), *A Guide to the Use of Herbicides*, (16th ed.), Pretoria: Plant Protection Research Institute.
- Wright, A.D., and Bourne, A.S. (1990), 'Effect of 2,4 D on the Quality of Water Hyacinth as Food for Insects', *Plant Protection Quarterly*, 5, 139–141.
- Wright, A.D., and Skilling, L. (1987), 'Herbicide Toxicity and Biological Control Agents', in *Proceedings of the Eighth Australian Weeds Conference*, eds. D. Lemerle and A.R. Leys, New South Wales, Sydney, pp. 93–96.