



Received on 05 January, 2018; received in revised form, 01 May, 2018; accepted, 13 May, 2018; published 01 September, 2018

EFFECTIVE MEDICINAL PLANTS IN TREATING HEPATITIS B

Zeinab Nazarian Samani¹ and Mahmoud Rafieian Kopaei^{*2}

School of Veterinary Medicine¹, Shahrekord University, Iran.

Medical Plants Research Center², Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran.

Keywords:

Liver, Hepatitis B virus,
Medical plants, Viral disease

Correspondence to Author:

Prof. Mahmoud Rafieian-Kopaei

Head of Medical Plants Research
Center, Basic Health Sciences
Institute, Shahrekord University of
Medical Sciences, Shahrekord, Iran.

E-mail: rafieian@skums.ac.ir

ABSTRACT: Medicinal plants and natural products have many applications in the treatment of viral diseases, due to much lower side effects. Different types of hepatitis, especially hepatitis B, are of the viral diseases whose treatment using herbal medicines is currently of great interest due to the limited availability of effective chemical drugs and having numerous side effects. Due to the attention paid to herbal drugs and their importance for the treatment of hepatitis B, we conducted this review to take a look at the evidence regarding the action mechanisms of viruses, antiviral herbal remedies and various herbal anti-hepatitis drugs and their therapeutic mechanism published until 2017. To provide data for conducting this review, the most up-to-date electronic journals including those indexed in the Pubmed, Elsevier, Institute for Scientific Information, Google Scholar and Scientific Information Database databases and various books, has been used. There are several plants for the treatment of viral diseases, including hepatitis B, which prevent or decrease infection via various mechanisms. Medicinal plants counteract viruses by various mechanisms, but most of them including *Ganoderma lucidum* and *Oenanthe javanica* inhibit the transcription of hepatitis B virus in hepatocytes. Most of action mechanism of medical plants exert an antiviral effect through inhibiting the transcription of HBV in hepatocytes; the need to study effective medicinal plants for treating different types of hepatitis, including hepatitis B and their action mechanisms have been intensified more than ever before, which makes this subject even more important.

INTRODUCTION: The liver is a vital organ of the body, which is also considered to be the largest gland. The liver plays a major role in the physiological processes of the body, which includes metabolism, secretion and storage. Detoxification of various drugs and xenobiotics is performed in the liver. Bile secretion is another task of the liver that plays a very important role in digestion^{1,2}.

Many toxic chemicals (certain antibiotics, chemotherapeutics, peroxidised oil, aflatoxin, carbon tetrachloride, acetaminophen, chlorinated hydrocarbons, etc.), food, alcohol, infections such as parasites, viruses, fungi or bacteria and also autoimmune disorders can cause liver diseases such as hepatitis, inflammatory liver disease, jaundice, hepatitis (non-inflammatory liver disease, cirrhosis (a digestive disorder that is the result of liver fibrosis), liver cancer, etc.^{1,3}

It is estimated that approximately 90% of hepatitis is of viral type, including hepatitis B, A, C, D (delta agents), E and G. Of these, hepatitis B often causes chronic liver disease and cirrhosis of the liver³. Hepatitis B virus (HBV) is a virus from the Hepadnaviridae family⁴.

QUICK RESPONSE CODE 	DOI: 10.13040/IJPSR.0975-8232.9(9).3589-96
	Article can be accessed online on: www.ijpsr.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.9(9).3589-96	

HBV is the cause of hepatitis B, which is transmitted to the virus through infected blood and fluids of the body. Although acute hepatitis is spontaneously healed, treatment for chronic cases is recommended because of the likelihood of cirrhosis of the liver. The hepatitis B vaccination program is being implemented in endemic countries, which reduces the cases of infection with this virus⁵. First, human beings were required to prepare the necessary medicines to cure their suffering and diseases from nature, especially plants. Chemical drugs are supplied worldwide for about 150 years.

A comparison of the use of chemical and herbal drugs shows that chemical drugs, while having good effects, have side effects some of which last until the end of life, and even in some cases, are transmitted to the next generation, while the side effects of herbal drugs are comparatively lower and in many cases, they lead to very few or no complications.

In the past decade, due to the problems that chemical drugs have caused for many people, the tendency to using and paying attention to plant-derived medicinal products have increased. Another important issue is that for certain diseases, herbal remedies have better effects, and for some others, only herbal remedies are available⁶.

Due to the diversity of liver diseases, accurate diagnosis is very difficult and therefore a physician cannot do the exact and effective treatment for the disease. In most cases, treatment is symptomatic and supportive. On the other hand, current and modern drugs have high toxicity and it is therefore needed to replace these medications with high-value drugs with comparatively lower toxicity^{1,2}.

Medicinal plants play an important role in maintaining health and survival of humans and animals. On the other hand, they contribute to the treatment of nontoxic liver disease^{7, 8, 9, 10}. The purpose of this article is to investigate and identify effective medicinal plants in the treatment of viral diseases, especially hepatitis B, and the involved therapeutic mechanisms.

1. Medicinal Plants: Recently, a large proportion of the world population has turned to natural products to treat and prevent diseases. Influenced

by this issue, pharmaceutical companies are developing new antimicrobial formulations that are derived from medicinal plants.

Currently, phytotherapy is widely used across the world, especially in developed countries including some European countries and the United States¹¹. Approximately 45% of commercially available therapeutic products have been derived from medicinal plants or their derivatives¹². Recently, a worldwide upsurge in the preparation of medicinal plants and the isolation of their active compounds has emerged in health care¹³. Many medicinal plants that are traditionally used have been reported to have antiviral properties, some of which are used to treat animals and people who suffer from viral infections^{14, 15, 16}.

In a research on post - World War II antiviral agents in 1952 in Europe, a pharmaceutical company in England tested the antiviral activity of 288 medicinal plants against influenza A virus in the egg embryo, showing that 12 of these plants prevented viral amplification¹⁷. Over the past 40 years, widespread programs around the world have begun to evaluate antiviral activity of medicinal plants *in-vitro* and *in-vivo*. Canadian researchers reported antiviral activity of grapes, apples, strawberries and other fruit juices against herpes simplex virus (HSV), poliovirus type 1, coxsackie virus B5 and Echovirus in the 1970s^{18, 19, 20, 21}.

Antiviral activity of 100 British Columbian medicinal plants against seven viruses was investigated²². At the tested concentrations of extract, 12 of the medicinal plants were found to have antiviral properties. The extracts of *Rosa nutkana* and *Amelanchier alnifolia* were very potent against enteric corona virus. *Potentilla arguta* root extract and *Sambucus racemosa* extract completely inhibit respiratory syncytial virus (RSV). It has been shown that the effect of *Ipomopsis aggregata* extract against the influenza type 3 virus is potent.

Lomatium dissectum root extract completely inhibits the cytopathic effects of rotavirus. In addition, the extracts of *Cardamine angulata*, *Conocephalum conicum*, *Lysichiton americanus*, *Polypodium glycyrrhiza* and *Verbascum thapsus* have antiviral activity against herpes virus type 1.

The extracts of 40 different plant species used in the traditional medicine have antiviral properties against human cytomegalovirus (HCMV), Ross River virus (RRV), two RNA viruses, DNA virus and poliovirus type 123. The human rotavirus (HRV), RSV and influenza A virus are susceptible to *Eleutherococcus senticosus* root extract, while

DNA viruses, HSV type 1 virus (HSV-1) and adenoviruses are not inhibited by its extract²⁴. Crude *Sanicula europaea* extract has antiviral properties against human parainfluenza virus type 2, whereas its ethanol extract does not have such property²⁵. There are many medicinal plants with antiviral properties **Table 1**.

TABLE 1: ANTIVIRAL MEDICINAL PLANTS

Medicinal plant name	Product type	Viruses
<i>Melaleuca alternifolia</i> ²⁶	essential oils	HSV-1(herpes simplex virus-1), HSV-2(herpes simplex virus-2)
<i>Santolina insularis</i> ²⁷	essential oils	HSV-1 , HSV-2, herpes types
<i>Santalum album</i> ²⁸	essential oils	HSV-1
<i>Sanicula europaea</i> (L.) ²⁹	water-soluble extract	influenza
<i>Nigella sativa</i> ³⁰	black seed oil	murine cytomegalovirus (MCMV)
<i>Eleutherococcus senticosus</i> ³¹	liquid extract	DNA viruses respiratory syncytial virus (RSV) adenovirus
<i>Rosa nutkana</i> ²²	extract	enteric corona virus
<i>Amelanchier alnifolia</i> ²²	extract	enteric corona virus
<i>Ipomopsis aggregate</i> ²²	extract	parainfluenza virus type 3
<i>Lomatium dissectum</i> ²²	root extract	rotavirus
<i>Potentilla arguta</i> ²²	root extract	respiratory syncytial virus (RSV)
<i>Sambucus racemosa</i> ²²	branch tip extract	respiratory syncytial virus (RSV)
<i>Cardamine angulata</i> ²²	extract	herpes virus type 1
<i>Polypodium glycyrrhiza</i> ²²	extract	herpes virus type 1
<i>Verbascum Thapsus</i> ²²	extract	herpes virus type 1
<i>Conocephalum conicum</i> ²²	extract	herpes virus type 1
<i>Lysichiton americanum</i> ²²	extract	herpes virus type 1
<i>Dianella longifolia</i> var. <i>grandis</i> (Liliaceae) ²³	root extract	poliovirus type 1
<i>Pterocaulon sphacelatum</i> (Asteraceae) ²³	extract	poliovirus type 1
<i>Euphorbia australis</i> (Euphorbiaceae) ²³	extract	human cytomegalovirus (HCMV)
<i>Scaevola spinescens</i> (Goodeniaceae) ²³	extract	human cytomegalovirus (HCMV)
<i>Eremophila latrobei</i> subsp. <i>glabra</i> (Myoporaceae) ²³	extract	Ross River virus (RRV)
<i>Pittosporum phylliraeoides</i> var. <i>Microcarpa</i> (Pittosporaceae) ²³	extract	Ross River virus (RRV)
<i>Sanicula europaea</i> ²⁵	crude extract	Parainfluenza virus type 2
<i>Myrcianthes cispalatensis</i> ²⁵	extract	RSV, adenovirus serotype7, HSV-1
<i>Azadirachta indica</i> ³²	leaf extract	Smallpox, chicken pox, poxvirus, herpes viruses, poliomyelitis
<i>Opuntia streptacantha</i> ³³	extract	HSV, equine herpes virus, pseudorabies virus, influenza virus
<i>Bergenia ligulata</i> ³⁴	extract	influenza virus, HSV
<i>Nerium indicum</i> ³⁴	extract	influenza virus, HSV
<i>Holoptelia integrifolia</i> ³⁴	extract	influenza virus, HSV

2. Hepatitis B Virus: Hepatitis B is one of the most important global health issues that lead to high morbidity and mortality, and two factors have contributed to the epidemiology of this disease; vaccination programs and migration. Nearly 240 million people worldwide have acquired chronic infection with this virus^{35, 36, 37, 38}. HBV-related mortality rates are more likely to be due to chronic infection, including liver cirrhosis and hepatocellular carcinoma (HCC).

HCC is the most common cancer in the world, with chronic HBV infection responsible for 50 - 90% of cases of this cancer³⁹. HBV is a viral prototype of the Hepadnaviridae family that is double-stranded DNA genome⁴ and is transcribed by the reverse transcription of the RNA pregenome^{4, 40}. HBV had eight genotypes from A to H with different virological characteristics and geographical distributions^{40, 41}.

More than 8% of the nucleotide sequences of each genotype are different from each other. Considering these conditions, the likelihood of recombination between genotypes, especially B/C and A/D, increases. Depending on the genotype of the virus, the DNA genome is composed of 3182-3248 nucleotides⁴². HBV is the cause of hepatitis B that is transmitted through the blood or the body fluids (secretions) that contain this virus. Although there is self-healing in acute hepatitis, treatment of chronic cases is recommended because of the possibility of cirrhosis of the liver.

The hepatitis B vaccine program is implemented in endemic countries, which reduces the incidence of this virus⁵. HBV primarily infects humans, but there are also reports regarding the detection of this infection in chimpanzee, Chacma baboons and tree shrews contamination⁴³. Most of the target cells of this virus are hepatocytes, but this virus is also found in other cells, such as those in the pancreas and kidney, bile duct lining cells and peripheral blood mononuclear cells⁴³.

HBV is an enveloped virus with a diameter of 42 - 47 nm. The envelope consists of a small amount of lipid and three surface proteins including: Small (SHB), medium (MHB) and large (LHB), which comprise disulfide-linked hetero and homodimers. In the serum of the people with the infection, two types of particle are found: Small spherical particles with approximately 20 nm diameter and filamentous particles with approximately 20 nm diameter⁴². The nucleocapsid is made up of nuclear protein that is composed of 185 - 188 amino acids depending on the genome. The N-terminal of 149 - 151 amino acids is responsible for the self assembly of nucleocapsids. This assembly has certain steps. The first step is the formation of homodimer links by disulfide bridges. The nucleocapsid contains poles that allow the nucleotide to bind during DNA synthesis. C-terminal amino acids in this core protein play the role of genome packet⁴⁴.

In the bloodstream of people with acute and chronic infections, high levels of HBV virions can be detected. Hepatocytes, which are the main target of this virus, are removed from the bloodstream by endothelial cells and kupffer cells, which are in direct contact with the sinusoids. Liver sinusoidal

endothelial cells have a large cytoplasmic content of 50 - 100 nm in diameter. The virions of the bloodstream in the sinuses reach Dis space (the space between the endothelial cystic fibrosis and hepatocyte membrane).

Virions bind hepatocyte surface by the LHB protein, or perhaps by a lipid envelope^{42, 43}. The nucleocapsid then releases into the cytoplasm and moves through the microtubules to the Microtubules - Organizing Center (MTOC) near the nucleus. Access to nucleus through nuclear pores is accomplished through the mediation of polymerase and heat shock proteins⁴². Then, transcription, translation steps and other steps are carried out.

3. Anti - Hepatitis Medicinal Plants: Due to the diversity of liver diseases, accurate diagnosis is very difficult, so a physician cannot do the exact and effective treatment of the disease. On the other hand, the only drugs used to treat chronic infections are lamivudine, interferon α , HBV, and adefovir dipivoxil. Most of these available drugs inhibit reverse transcriptase activity. Modern and currently used drugs have high toxicity. In addition, the emergence of resistant mutants during long-term treatments is another major problem with modern medicines.

It is, therefore, necessary to replace these medications with high-value ones with comparatively lower toxicity^{1, 2}. Many medicinal plants that are traditionally used have antiviral properties, and some of them are used to treat animals and people suffering from viral diseases^{45, 46}. Medicinal plants have several phytochemicals that have potent antioxidant properties including alkaloids, carotenoids, saponins, flavonoids (isoflavones, flavonones, anthocyanins, catechins, flavones, isocatechins and quercetin), terpenoids, polyphenols (ellagic acid, gallic acid and tannins), vitamins (A, C, E and K), carotenoids, minerals (manganese, selenium, copper, chromium, zinc and iodine), polysaccharides, enzymes (superoxide dismutase, catalase and glutathione peroxidase), lignins, saponins, xanthonenes and pigments^{47, 48}.

Antioxidants treat various diseases by protecting cells against free radical-induced damage⁴⁷. Several medicinal plants are used to treat hepatitis or hepatotoxicity, and some of them have anti-HBV

activity. For example, *Curcuma longa* Linn. has antiviral properties against HBV, which inhibits the production of the constituents of HBV and HBV RNA in hepatocytes. The extract of this plant is indeed an inhibitor of transcription of HBV.

In addition, studies have shown that the antiviral activity of this plant is associated with an increase

in the intracellular accumulation of the P53 protein, which is due to increased transcription of the P53 protein gene⁴⁹. Another plant is *Ganoderma lucidum*. Studies have shown that ganoderic acid isolated from this medicinal plant inhibits HBV transcription in the HepG2215 hepatocytes, resulting in an antiviral activity against HBV⁵⁰.

TABLE 2: ANTIHEPATITIS MEDICAL PLANTS

Family	Medicinal plant name	Hepatitis
Zingiberaceae	<i>Curcuma longa</i> Linn.	Hepatitis B Virus ⁴⁹
Ganodermataceae	<i>Ganoderma lucidum</i>	Hepatitis B Virus ⁵⁰
Euphorbiaceae	<i>Phyllanthus amarus</i> or <i>Phyllanthus niruri</i>	Hepatitis B Virus ^{54, 55, 56, 57}
Euphorbiaceae	<i>Phyllanthusanus</i>	Hepatitis B Virus ^{55, 56, 57, 58}
Acanthaceae	<i>Acanthus ilicifolius</i> L	Hepatitis B Virus ^{52, 53}
Apiaceae	<i>Oenanthe javanica</i>	Hepatitis B Virus ^{59, 60}
Gentianaceae	<i>Swertia patens</i>	Hepatitis B Virus ^{61, 62}
Gentianaceae	<i>Swertia chirayita</i>	Hepatitis B Virus ^{61, 62}
Urticaceae	<i>Boehmeria nivea</i>	Hepatitis B Virus ⁶³
Rutaceae	<i>Citrus Sinensis</i>	Hepatitis C Virus ⁶⁵
Crassulaceae	<i>Rhodiola kirilowii</i>	Hepatitis C Virus ⁶⁶
Mimosoideae	<i>Acacia nilotica</i>	Hepatitis C Virus ⁶⁴
Burseraceae	<i>Boswellia carterii</i>	Hepatitis C Virus ⁶⁴
Myrsinaceae	<i>Embelia schimperi</i>	Hepatitis C Virus ⁶⁴
Fagaceae	<i>Quercus infectoria</i>	Hepatitis C Virus ⁶⁴
Apiaceae	<i>Trachyspermum ammi</i>	Hepatitis C Virus ⁶⁴
Piperaceae	<i>Piper cubeba</i>	Hepatitis C Virus ⁶⁴
Myrtaceae	<i>Syzygium aromaticum</i>	Hepatitis C Virus ⁶⁴
Zingiberaceae	<i>Zingiber officinale</i>	Hepatitis C Virus ⁶⁷
Asteraceae	<i>Silybum marianum</i>	Hepatitis C Virus ⁶⁷
Saxifragaceae	<i>Saxifraga melanocentra</i>	Hepatitis C Virus ⁶⁸
Lamiaceae	<i>Mentha longifolia</i>	Hepatitis A Virus ⁶⁹
Lamiaceae	<i>Ocimum basilicum</i>	Hepatitis A Virus ⁶⁹
Asteraceae	<i>Taraxacum Officinale</i>	Hepatitis ⁷⁰
Brassicaceae	<i>Lepidium sativum</i>	Hepatitis ⁷¹
Fabaceae	<i>Trigonella foenum graecum</i>	Hepatitis ⁷²
Meliaceae	<i>Azadirachta indica</i>	Hepatitis ⁷³
Fabaceae	<i>Glycyrrhiza glabra</i>	Hepatitis ⁷⁴
Euphorbiaceae	<i>Jatropha curcas</i> Linnaeus	Hepatitis ⁷⁵
Asteraceae	<i>Cynara scolymus</i>	Hepatitis ⁷⁶
Asteraceae	<i>Matricaria chamomilla</i>	Hepatitis ⁷⁷
Fabaceae	<i>Cassia fistula</i>	Hepatitis ⁷⁸
Marchantiaceae	<i>Marchantia thallus</i>	Hepatitis ⁷⁹
Asteraceae	<i>Silybum marianum</i>	Hepatitis ⁸⁰
Fabaceae	<i>Sophora flavescens</i>	Hepatitis ⁸¹
Salicaceae	<i>Flacourtia indica</i> Governor's Plum	Hepatitis ⁸²
Rubiaceae	<i>Morinda citrifolia</i>	Hepatitis ⁸³

Clinical studies have indicated that phyllanthus species have positive effects on HBsAg in the clearance of serum from HBV carriers⁵¹. For example, *P. amarus* (L.) is an inhibitor of HBV polymerase activity and mRNA transcription⁴⁷. Another plant that has antiviral activity against hepatitis is *Oenanthe javanica*. Studies have shown that this plant is helpful to treat HBV-induced infections by inhibiting transcription of HBV in

hepatocytes⁴⁹. *Acanthus ilicifolius* L. reduces HBV-induced liver damage by reducing transaminase^{52, 53}. Other plants also have anti-hepatitis properties. In general, there is not enough information on the action mechanisms of medicinal plants on HBV, although there are effective natural products against HBV and the resulting infections **Table 2.**

CONCLUSION: Recently, due to the many side effects and the presence of medicinal residues of the chemical drugs for the treatment of viral diseases, especially hepatitis, attention has been paid to medicinal plant-derived products has increased.

It is, therefore, necessary, more than ever before, to address and study effective medicinal plants to treat different types of hepatitis, including hepatitis B, and their action mechanisms, which makes this subject even more important.

The action mechanism of a number of anti-hepatitis B drugs has been understood. Most of them exert an antiviral effect through inhibiting the transcription of HBV in hepatocytes; however, the action mechanism of many plants remains unknown and needs to be further studied. It should be noted that usually phenolic compounds in medicinal plants possess antimicrobial activities. Hence, other plants which have these compounds^{84, 85} may possess anti-hepatitis activities, too, which worth examining.

ACKNOWLEDGEMENT: Nil

CONFLICT OF INTEREST: Nil

REFERENCES:

- Govind P and Pandey SP: Phytochemical and toxicity study of *Embllica officinalis* (Amla). *Int Res J Pharm* 2011; 2(3): 270-272.
- Thomas D, Leoutsakas D, Zabransky T and Kumar M: Hepatitis C in HIV infected individuals: cure and control, right now. *Journal of the International AIDS Society* 2011; 14: 1-8.
- Kumar CH, Ramesh A, Kumar SJN and Ishaq MB: A review on hepatoprotective activity of medicinal plants. *Int J Pharmaceu Sci Res* 2011; 2(3): 501-515.
- Liang TJ: Hepatitis B. The virus and disease. *Hepatology* 2009; 49(5S): S13-21.
- Ni YH and Chen DS: Hepatitis B vaccination in children. The Taiwan experience. *Pathologiebiologie* 2010; 58: 296-300.
- Rabiei Z, Bigdeli MR and Lorigooini Z: A review of medicinal herbs with antioxidant properties in the treatment of cerebral ischemia and reperfusion. *Journal of Babol University of Medical Sciences* 2015; 17(12): 47-56.
- Madhuri S: Studies on oestrogen induced uterine and ovarian carcinogenesis and effect of ProImmu in rats. PhD thesis. Jabalpur, MP, India: RDVV 2008.
- Madhuri S and Pandey G: Some anticancer medicinal plants of foreign origin. *Curr Sci* 2009; 96(6): 779-783.
- Pandey G and Madhuri S: Some anticancer agents from plant origin. *PI Arch* 2008; 8(2): 527-532.
- Pandey G and Madhuri S: Significance of fruits and vegetables in malnutrition cancer. *PI Arch* 2010; 10(2): 517-522.
- Solati K, Heidari-Soureshjani S, Luther T and Asadi-Samani M: Iranian medicinal plants effective on sexual disorders: A systematic review. *International journal of pharmaceutical sciences and research* 2017; 8(6): 2415-20.
- Lahlou M: The success of natural products in drug discovery. *Pharmacol Pharm* 2013; 4: 17-31.
- Jassim SAA and Najji MA: Novel antiviral agents. a medicinal plant perspective. *Journal of Applied Microbiology* 2003; 95: 412-427.
- Solati K, Heidari-Soureshjani S and Pocock L: Effects and mechanisms of medicinal plants on stress hormone (cortisol): A systematic review. *World family medicine/middle east journal of family medicine* 2017; 15(9): 117-23.
- Venkateswaran PS, Millman I and Blumberg BS: Effects of an extract from *Phyllanthus niruri* on hepatitis B and Woodchuck hepatitis viruses: *in vitro* and *in vivo* studies. *Proceedings of the National Academy of Sciences of the USA* 1987; 84: 274-278.
- Thyagarajan SP, Jayaram S and Valliammai T: *Phyllanthus amarus* and hepatitis B. *Lancet* 1990; 336: 49-50.
- Chantrill BH, Coulthard CE, Dickinson L, Inkley GW, Morris W and Pyle AH: The action of plant extracts on a bacteriophage of *Pseudomonas pyocyanea* and on influenza A virus. *Journal of General Microbiology* 1952; 6: 74-84.
- Konowalchuk J and Speirs JJ: Virus inactivation by grapes and wines. *Applied and Environmental Microbiology* 1976; 32: 757-763.
- Konowalchuk J and Speirs JJ: Antiviral activity of fruit extracts. *Journal of Food Science* 1976b; 41: 1013-1017.
- Konowalchuk J and Speirs JJ: Antiviral effect of apple beverages. *Applied and Environmental Microbiology* 1978a; 36: 798-801.
- Konowalchuk J and Speirs JJ: Antiviral effect of commercial juices and beverages. *Applied and Environmental Microbiology* 1978b; 35: 1219-1220.
- McCutcheon AR, Roberts TE, Gibbons E, Ellis SM, Babiuk LA, Hancock RE and Towers GH: Antiviral screening of British Columbian medicinal plants. *Journal of Ethnopharmacology* 1995; 49: 101-110.
- Semple SJ, Reynolds GD, O'Leary MC and Flower RL: Screening of Australian medicinal plants for antiviral activity. *Journal of Ethnopharmacology* 1998; 60: 163-172.
- Glatthaar-Saalmuller B, Sacher F and Esperester A: Antiviral activity of an extract derived from roots of *Eleutherococcus senticosus*. *Antiviral Research* 2001; 50: 223-228.
- Karagoz A, Arda N, Goren N, Nagata K. and Kuru A: Antiviral activity of *Sanicula europaea* L. extracts on multiplication of human parainfluenza virus type 2. *Phytotherapy Research* 1999; 13: 436-438.
- Schnitzler P, Schon K and Reichling J: Antiviral activity of Australian tea tree oil and eucalyptus oil against herpes simplex virus in cell culture. *Die Pharmazie* 2001; 56: 343-347.
- De Logu A, Loy G, Pellerano ML, Bonsignore L and Schivo ML: Inactivation of HSV-1 and HSV-2 and prevention of cell-to-cell virus spread by *Santolina insularis* essential oil. *Antiviral Research* 2000; 48: 177-185.

28. Benencia F and Courreges MC: Antiviral activity of sandalwood oil against herpes simplex viruses-1 and -2. *Phytomedicine* 1999; 6: 119-123.
29. Turan K, Nagata K and Kuru A: Antiviral effect of *Sanicula europaea* L. leaves extract on influenza virus-infected cells. *Biochemical and Biophysical Research Communications* 1996; 225: 22-26.
30. Salem ML and Hossain MS: Protective effect of black seed oil from *Nigella sativa* against murine cytomegalovirus infection. *International Journal of Immunopharmacology* 2000; 22: 729-740.
31. Glatthaar-Saalmuller B, Sacher F and Esperester A: Antiviral activity of an extract derived from roots of *Eleutherococcus senticosus*. *Antiviral Research* 2001; 50: 223-228.
32. Rao AR, Kumar SSV, Paramasivam TB, Kamalakshi S, Parashuraman AR and Shanta B: Study of antiviral activities of leaves of margosa tree on vaccinia and variola virus. A preliminary report. *Indian Journal Medical Research* 1969; 57: 495-502.
33. Ahmad A, Davies J, Randall S and Skinner GR: Antiviral properties of extract of *Opuntia streptacantha*. *Antiviral Research* 1996; 30: 75-85.
34. Rajbhandari M, Wegner U, Julich M, Schopke T and Mentel R: Screening of Nepalese medicinal plants for antiviral activity. *Journal of Ethnopharmacology* 2001; 74: 251-255.
35. EASL: European Association for the Study of the Liver. clinical practice guidelines: Management of chronic hepatitis B virus infection. *J Hepatol* 2012; 57: 167-185.
36. Schweitzer A, Horn J, Mikolajczyk RT, Krause G and Ott JJ: Estimations of worldwide prevalence of chronic hepatitis B virus infection: A systematic review of data published between 1965 and 2013. *Lancet* 2015; 386: 1546-1555.
37. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K and Aboyans V: Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380: 2095-2128.
38. Ott JJ, Stevens GA, Groeger J and Wiersma ST: Global epidemiology of hepatitis B virus infection: New estimates of age-specific HBsAg seroprevalence and endemicity. *Vaccine* 2012; 30: 2212-2219.
39. Stanaway JD, Flaxman AD, Naghavi M, Fitzmaurice C, Vos T and Abubakar I: The global burden of viral hepatitis from 1990 to 2013: findings from the global burden disease study 2013. *Lancet* 2016; 388: 1081-1088.
40. Schaeffer DJ and Krylov VS: Anti-HIV activity of extracts and compounds from algae and cyanobacteria. *Ecotoxicology and Environmental Safety* 2000; 45: 208-227.
41. Kramvis A and Kew MC: Epidemiology of hepatitis B virus in Africa, its genotypes and clinical association of genotypes. *Hepatol Res* 2007; 37: s1S9-S19.
42. Kann M: Structural and molecular virology. In: *Hepatitis B Virus Guide*. International medicine press 2002; 9-22.
43. Coa J, Yang EB, Su JJ, Li Y and Chow P: The tree shrews. Adjunct and alternative to primates as models for biomedical research. *J Med Primatol* 2003; 32: 123-130.
44. Bruss V: Hepatitis B Virus morphogenesis. *World J gastroenterol* 2007; 13: 65-73.
45. McCutcheon AR, Roberts TE, Gibbons E, Ellis SM, Babiuik LA and Hancock REW: Antiviral screening of British Columbian medicinal plants. *Journal of Ethnopharmacology* 1995; 49(2): 101-110.
46. Rahimi-Madiseh M, Lorigoini Z, Zamani-gharaghoshi H and Rafieian-kopaei M: *Berberis vulgaris*: specifications and traditional uses. *Iranian Journal of Basic Medical Sciences* 2017; 20(5): 569.
47. Govind P: Medicinal plants against liver diseases. *International Research Journal of Pharmacy* 2011; 2(5): 115-121.
48. Ghasemi S and Lorigoini Z: A review of significant molecular mechanisms of flavonoids in prevention of prostate cancer. *Journal of Chemical and Pharmaceutical Sciences*. 2016; 9: 3388-94.
49. Kim HJ, Yoo HS, Kim JC, Park CS, Choi, MS and Kim M: Antiviral effect of *Curcuma longa* Linn. extract against hepatitis B virus replication, *Journal of ethnopharmacology* 2009; 124(2): 189-96.
50. Li YQ and Wang SF: Anti-hepatitis B activities of ganoderic acid from *Ganoderma lucidum*. *Biotechnol Lett* 2006; 28(11): 837-841.
51. Liu J, Lin H and McIntosh H: Genus *Phyllanthus* for chronic hepatitis B virus infection: a systematic review. *J Viral Hepatitis* 2001; 8: 358-366.
52. Wu YH: Naturally derived anti-hepatitis B virus agents and their mechanism of action. *World J Gastroenterol* 2016; 22: 188-204.
53. Wei PH, Wu SZ, Mu XM, Xu B, Su QJ, Wei JL and Yang Y: Effect of alcohol extract of *Acanthus ilicifolius* L. on anti-duck hepatitis B virus and protection of liver. *J Ethnopharmacol* 2015; 160: 1-5.
54. Lee CD, Ott M, Thyagarajan SP, Shafritz DA, Burk RD and Gupta S: *Phyllanthus amarus* down-regulates hepatitis B virus mRNA transcription and replication. *Eur J Clin Invest* 1996; 26: 1069-1076.
55. Ott M, Thyagarajan SP and Gupta S: *Phyllanthus amarus* suppresses hepatitis B virus by interrupting interactions between HBV enhancer I and cellular transcription factors. *Eur J Clin Invest* 1997; 27: 908-915.
56. Patel JR, Tripathi P, Sharma V, Chauhan NS, and Dixit VK: *Phyllanthus amarus*: Ethnomedicinal uses, phytochemistry and pharmacology: A review. *J. Ethnopharmacol* 2011; 138(2): 286-313.
57. Kumar A: A review on hepatoprotective herbal drugs. *Int. J. Res. Pharm. Chem.* 2012; 2(1).
58. Lam WY, Leung KT, Law PT, Lee SM, Chan HL and Fung KP: Antiviral effect of *Phyllanthus nanus* ethanolic extract against hepatitis B virus (HBV) by expression microarray analysis. *J Cell Biochem* 2006; 97: 795-812.
59. Huang ZM, Yang XB and Cao WB: Effects of Qin ling ke li in the treatment of 90 patients with chronic hepatitis B. *Pharm J Chinese People's Lib Army* 2001; 17: 41-44.
60. Jacob JR, Korba BE, You JE, Tennant BC and Kim YH: Korean medicinal plant extracts exhibit antiviral potency against viral hepatitis. *J Altern Complement Med* 2004; 10: 1019-1026.
61. He K, Geng CA, Cao TW, Wang HL, Ma YB and Zhang XM: Two new secoiridoids and other anti-hepatitis B virus active constituents from *Swertia patens*. *J Asian Nat Prod Res.* 2016; 18(6): 528-534.
62. Zhou NJ, Geng CA, Huang XY, Ma YB, Zhang XM, Wang JL and Chen JJ: Anti-hepatitis B virus active constituents from *Swertia chirayita*. *Fitoterapia* 2015; 100: 27-34.
63. Chang JM, Huang KL, Yuan TTT, Lai YK and Hung LM: The anti-hepatitis B virus activity of *Boehmeria nivea* extract in HBV-viremia SCID mice. *Evid Based Complement Alternat Med* 2010; 7: 189-195.
64. Hussein G, Miyashiro H and Nakamura N: Inhibitory effects of Sudanese medicinal plant extracts on hepatitis C virus (HCV) protease. *Phytother Res* 2000; 14: 510-516.

65. Nahmias Y, Goldwasser J, Casali M, Poll DV, Wakita T, Chung RT and Yarmush ML: Apolipoprotein β -dependent hepatitis C virus secretion is inhibited by the grapefruit flavonoid naringenin, *Hepatology*, American Association for the study of liver diseases, pp. Apolipoprotein β -dependent hepatitis C virus secretion is inhibited by the grapefruit flavonoid naringenin 2008; 1437-1443.
66. Khare P, Singh L and Yadav G: A review on novel therapies to combat hepatitis C, *African Journal of Pharmacy and Pharmacology* 2015; 9(3): 37-42.
67. El-Adawi H, El-Demellawy M, El-Wahab AA: Some medicinal plant extracts exhibit potency against viral hepatitis C. *J Biosci Tech* 2011; 2: 223-231.
68. Zuo GY, Li ZQ, Chen LR and Xu XJ: *In vitro* anti-HCV activities of *Saxifraga melanocentra* and its related polyphenolic compounds. *Antivir Chem Chemother* 2005; 16: 393-398.
69. Al-Ali KH and El-Badry AA: Anti-viral activity of two labiatae plants [naana (hassoi, habak) and basil (rahan)] of Al-Madiah Al-Munawarah. *J Med Biomed Sci* 2010; 2: 67-73.
70. Al-Malki AL, Abo-Golayel MK, Abo-Elnaga G and Al-Beshri H: Hepatoprotective effect of dandelion (*Taraxacum officinale*) against induced chronic liver cirrhosis, *Journal of Medicinal Plants Research* 2013; 7(20): 1494-1505.
71. Al-Asmaria AK, Athar T, Al-Shahranib HM, Al-Dakheel SI and Al-Ghamdic MA: Efficacy of *Lepidium sativum* against carbon tetrachloride induced hepatotoxicity and determination of its bioactive compounds by GC-MS, *Toxicology Reports* 2015; 2: 1319-1326.
72. Al-Sultan SI and El-Bahr SM. Effect of Aqueous Extract of Fenugreek (*Trigonella foenum-graecum* L.) On selected biochemical and oxidative stress biomarkers in rats intoxicated with carbon tetrachloride. *International Journal of Pharmacology* 2015; 11: 43-49.
73. Lakshmi T, Krishnan V, Rajendran R and Madhusudhanan N: *Azadirachta indica*: A herbal panacea in dentistry - An update. *Pharmacognosy Reviews* 2015; 9(17): 41-44.
74. Lakshmi and Geetha: *Glycyrrhiza glabra* Linn. commonly known as licorice: A therapeutic review. *International Journal of Pharmacy and Pharmaceutical Sciences* 2011; 3(4): 20-25.
75. Punjaji SA: Traditional Hepatopathic Treatments in Newasa Tahasil of Ahmednagar District (M.S.) India, *International Journal of Pharmaceutical and Phytopharmacological Research* 2012; 1(6): 354-356.
76. Hamza RG, El shahat AN and Mekawey HMS: Protective Effect of γ -Irradiated Dried Powder of Artichoke Leaves against CCl_4 Oxidative Stress Induced in Rat Liver, *Journal of Radiation Research and Applied Sciences* 2012; 5(3): 549-564.
77. Mannaa FA, Ibrahim NA, Ibrahim SS, Abdel-Wahhab KG, Hassan NS and Mohammed SG: Preventive role of chamomile flowers and fennel seeds extracts against liver injury and oxidative stress induced by an immunosuppressant drug in rats, *Hepa. Res.* 2015; 1(3): 125-135.
78. Dawada S, Zade V, Dabhadkar D and Pare S: Hepatoprotective activity of *Cassia fistula* root against Carbon tetrachloride-Induced Hepatic Injury in rats (Wistar), *International Journal of Pharma Sciences and Research (IJPSR)* 2012; 3(4): 368-373.
79. Gupta SK, Sharma A and Moktan S: A review on some species of Marchantia with reference to distribution, characterization and importance, *World journal of pharmacy and pharmaceutical sciences* 2015; 4(4): 1576-1584.
80. Rambaldi A, Jacobs BP and Gluud C: No evidence supporting or refuting milk thistle for alcoholic and/or hepatitis B or C virus liver diseases 2005; 100(11): 2583-2591.
81. Wang L, Li F, Lu J, Li G, Li D, Zhong X, Guo GL and Ma X: The Chinese Herbal Medicine *Sophora flavescens* activates Pregnane X Receptor, *Drug Metabolism and Disposition* 2010; 38(12): 2226-2231.
82. Pandey A: Herbal remedies used in the treatment of liver disease and development of those remedies by the help of micro-propagation: A review, *International Journal of Pharmaceutical and Medicinal Research* 2014; 2(1): 8-14.
83. Kumar A: A review on hepatoprotective herbal drugs. *Int. J. Res. Pharm. Chem* 2012; 2(1): 2231-2781.

How to cite this article:

Samani ZN and Kopaei MR: Effective medicinal plants in treating Hepatitis B. *Int J Pharm Sci & Res* 2018; 9(9): 3589-96. doi: 10.13040/IJPSR.0975-8232.9(9).3589-96.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)