

1 **Title:** Association between gentamicin resistance and stress tolerance in water isolates of  
2 *Ralstonia pickettii* and *R. mannitolilytica*

3

4 **Authors:** Pompeyo Ferro, Ivone Vaz-Moreira, Célia M. Manaia\*

5

6 Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina –  
7 Laboratório Associado, Escola Superior de Biotecnologia, Rua Arquiteto Lobão Vital, 172,  
8 4200-374 Porto, Portugal

9

10 **Author for correspondence (\*):**

11 C.M. Manaia, PhD

12 Escola Superior de Biotecnologia, Universidade Católica Portuguesa, 4200-374 Porto  
13 Portugal

14 Tel: +351 22 5580059

15 Fax: +351 22 5090351

16 e-mail: [cmanaia@porto.ucp.pt](mailto:cmanaia@porto.ucp.pt)

17

18 **ORCID:**

19 Pompeyo Ferro: 0000-0003-2986-2774

20 Ivone Vaz-Moreira: 0000-0001-8912-9836

21 Célia M. Manaia: 0000-0002-3674-1789

22

23 **ACKNOWLEDGEMENTS**

24 This work was financially supported by National Funds from FCT - Fundação para a  
25 Ciência e a Tecnologia through projects UID/Multi/50016/2013 and NanoDiaBac  
26 ENMed/0001/2014, and IVM grant (SFRH/BPD/87360/2012); by FCT through the project  
27 WaterJPI/0001/2013 - STARE (Stopping Antibiotic Resistance Evolution); and by Erasmus  
28 Mundus Project SUD-UE (EMA2-STRAND 1, LOT 14) through PF grant (2013-2589).

29 **ABSTRACT**

30 Members of the species *Ralstonia pickettii* and *R. mannitolilytica*, although ubiquitous and  
31 lacking major virulence factors, have been associated with nosocomial outbreaks.  
32 Tolerance to metals, antibiotics and disinfectants may represent an advantage for their  
33 ubiquity and opportunistic pathogenic potential. In this study we compared five strains that  
34 differed on the origin (hospital effluent, tap water, mineral water) and in the susceptibility  
35 to aminoglycosides, regarding their tolerance to metals and disinfection. The growth  
36 kinetics and biofilm formation capacity were tested in four *R. pickettii* strains and one *R.*  
37 *mannitolilytica* at sub-inhibitory concentrations of aminoglycosides or arsenite. The  
38 survival to UV radiation, chlorine or hydrogen peroxide was also compared in  
39 aminoglycoside resistant and susceptible strains. Aminoglycoside resistant strains presented  
40 a higher tolerance to arsenite than the susceptible ones and either aminoglycosides or  
41 arsenite were observed to stimulate the biofilm formation. Sub-inhibitory concentrations of  
42 the aminoglycoside gentamicin or arsenite significantly decreased the growth rate and  
43 yield, but only arsenite caused a significant increase of the lag phase. Hydrogen peroxide  
44 presented higher disinfection effectiveness against aminoglycoside susceptible than against  
45 resistant strains, an effect that was not observed for UV or chlorine. Although this  
46 conclusion needs validation based on a larger number of isolates, including clinical, the  
47 results suggest that aminoglycoside resistance may be associated with traits that influence  
48 *Ralstonia* spp. fitness in the environment.

49

50

51 **KEYWORDS**

52 aminoglycosides resistance; biofilm; disinfectants; kinetics; sub-inhibitory concentrations

53

54

55

## 56 INTRODUCTION

57 Members of the phylum *Proteobacteria*, mainly some families and genera of the classes  
58 *Alpha-*, *Beta-* and *Gammaproteobacteria* are amongst the most prevalent bacteria in water  
59 habitats (Vaz-Moreira et al. 2014; Vaz-Moreira et al. 2017). Some species of the genus  
60 *Ralstonia* within the class *Betaproteobacteria*, in particular the species *Ralstonia pickettii*  
61 and *R. mannitolilytica*, are frequently observed in aquatic habitats, specifically in  
62 wastewater, potable water, surface water and mineral water (Becerra-Castro et al. 2015;  
63 Falcone-Dias et al. 2012; Ryan et al. 2011; Vaz-Moreira et al. 2017). These *Ralstonia*  
64 species are comprised of ubiquitous bacteria that have been found in a wide variety of  
65 environments, such as in plastic (Poly Vinyl Chloride) pipes forming biofilms structures,  
66 aerospace samples, purified water, saline solutions, skin disinfectants, biological samples,  
67 caps of blood culture bottles or even in human patients in the respiratory tract or the oral  
68 cavity (Adley et al. 2005; Anderson et al. 1990; Boutros et al. 2002; Coenye et al. 2002;  
69 Coman et al. 2017; Daxboeck et al. 2005; Koenig and Pierson 1997; Kulakov et al. 2002;  
70 Labarca et al. 1999; Maroye et al. 2000; McNeil et al. 1985; Mijndonckx et al. 2013;  
71 Riley and Weaver 1975; Ryan et al. 2006; Stelzmueller et al. 2006; Verschraegen et al.  
72 1985). The ubiquitous character of these bacteria is related to the requirement for minimal  
73 nutrient resources and explains the transmission from various sources to humans  
74 (Daxboeck et al. 2005). The capacity to grow in moist environments and to form biofilm  
75 has also been proposed as a reason for the *Ralstonia* spp. persistence in some environments  
76 (Adley et al. 2005). Motility, a characteristic of *Ralstonia* spp., is also sometimes  
77 associated with the increased capacity to form biofilm (Guttenplan and Kearns 2013;  
78 O'Toole and Kolter 1998). Although the decrease of motility after prolonged preservation

79 and sub-culture has been reported in *Ralstonia* spp. (Ryan 2009; Vaneechoutte et al. 2001),  
80 the association between motility and biofilm formation have been proposed in this genus.  
81 In particular, in some *Ralstonia* spp. aerotaxia was observed to regulate the biofilm  
82 formation (Yao and Allen 2007).

83 *R. pickettii* and *R. mannitolilytica* are not considered primary pathogens and, hence, are not  
84 screened in routine monitoring analyses in hospitals (Coenye et al. 2002; Orme et al. 2015;  
85 Waugh et al. 2010). Nevertheless, it has been argued that the low frequency of infection  
86 episodes attributed to *Ralstonia* spp. may be a consequence of misidentifications of these  
87 bacteria, being suggested that members of this group may be more widespread, invasive  
88 and severe than previously thought (Coman et al. 2017; Daxboeck et al. 2005; Ryan et al.  
89 2006). A recent study reported an association between the presence of intestinal *Ralstonia*  
90 *pickettii* and an augmented glucose intolerance in obesity (Udayappan et al. 2017).

91 Although members of *R. pickettii* and *R. mannitolilytica* lack major virulence factors, and  
92 rarely are reported as causing infection, members of these species have been considered the  
93 most pathogenic species of the genus (Vaneechoutte et al. 2004). Indeed, nosocomial  
94 outbreaks attributed to *R. pickettii* or *R. mannitolilytica* have been reported regularly over  
95 the last 30 years (Coman et al. 2017; De Baere et al. 2001; Fernandez et al. 1996; Khajuria  
96 et al. 2014; Labarca et al. 1999; Riley and Weaver 1975; Ryan et al. 2006; Vaneechoutte et  
97 al. 2001; Verschraegen et al. 1985).

98 In part, the ubiquitous character and potential to infect humans may be associated with the  
99 capability to stand environmental stresses observed in *R. pickettii* and *R. mannitolilytica*.

100 For instance, the capacity to survive in hospital disinfectants, as chlorhexidine and  
101 ethacridine lactate (acrinol) (Ryan et al. 2006) or to participate in bioremediation processes

102 through the breakdown of xenobiotic compounds was described in *R. pickettii* (Ryan et al.  
103 2007). Also metal tolerance seems to be a relevant property of these bacteria illustrated by  
104 the fact that the microbiota enriched from a hospital effluent in copper led to the isolation  
105 of monospecies cultures of metal resistant *Ralstonia* spp., in spite of the complex microbial  
106 community of such an effluent (Becerra-Castro et al. 2015). The array of antibiotic  
107 resistance phenotypes to gentamicin, chloramphenicol, colistin, tobramycin, polymyxin B,  
108 and many others observed in members of these species are probably also related with their  
109 ubiquity (Daxboeck et al. 2005; Pan et al. 2011; Ryan et al. 2009; Stelzmueller et al. 2006).

110 Given this background information, the occurrence of *R. pickettii* and *R. mannitolilytica*  
111 throughout the urban water cycle (wastewater and tap water) and in pristine water sources  
112 (mineral water) (Becerra-Castro et al. 2015; Falcone-Dias et al. 2012; Kulakov et al. 2002;  
113 Ryan et al. 2006; Vaz-Moreira et al. 2017), raises interest on their ecology. Able to  
114 withstand the drinking water treatment, bacteria of these species can reach humans through  
115 the water consumption. This work was based on the hypothesis that traits, such as tolerance  
116 to metals and antibiotics, to disinfection or the capacity to produce biofilm in the presence  
117 of antibiotics, may differ among *Ralstonia* spp. strains and hence affect their response to  
118 stress and environmental fitness. As part of the experimental design, a set of five *Ralstonia*  
119 spp. strains, isolated from hospital wastewater, mineral water, and tap water, were tested for  
120 their response under the abovementioned stress types.

121

## 122 **MATERIALS AND METHODS**

123 *Bacterial strains*

124 Five *Ralstonia* spp. isolates were selected for this study, four *Ralstonia pickettii*, two from  
125 hospital wastewater, one from mineral water and one from tap water, and one *Ralstonia*  
126 *mannitolilytica* from tap water (Table 1). Cultures were maintained and preserved in  
127 nutritive (Luria-Bertani) broth supplemented with 15% (v/v) glycerol.

128 The identification of the strains was made based on the 16S rRNA gene sequence analysis  
129 using the primers 27F and 1492R as previously described (Ferreira da Silva et al. 2007).

130 The sequences were compared with the public database EzBioCloud (Yoon et al. 2017).

131 The five strains were characterized based on selected biochemical tests using the  
132 commercial kits API 20E, API 20NE, and API ZYM (bioMérieux) following the  
133 manufacturer's instructions. Capsule presence was tested by negative staining (McKinney  
134 1953) in cultures grown in the absence and presence of sub-inhibitory concentrations of  
135 gentamicin. These additional characterizations were done in an attempt to find some traits  
136 that could be associated with the antibiotic and metals resistance phenotypes.

137 The 16S rRNA gene sequences of the studied strains were compared with other good  
138 quality (>1000 bp) sequences of *R. pickettii* and *R. mannitolilytica* strains of different  
139 origins available in the GenBank (<http://www.ncbi.nlm.nih.gov/>). The nucleotide sequence  
140 analysis was performed using the MEGA6 software (Tamura et al. 2013), based on the  
141 model of Jukes and Cantor (Jukes and Cantor 1969), and a dendrogram was created using  
142 the neighbour-joining method. The iTol software v3.2.4 (Letunic and Bork 2016) was used  
143 to represent the isolates source in the dendrogram.

#### 144 *Determination of antibiotic and metal resistance phenotypes*

145 The antibiotic resistance phenotypes were determined by disk diffusion method as  
146 recommended by the Clinical Laboratory Standards Institute (CLSI 2015), for 12



147 antibiotics: nalidixic acid (NA, 30 µg); ciprofloxacin (CIP, 5 µg); streptomycin (STR, 10  
148 µg); gentamicin (GEN, 10 µg); tetracycline (TET, 30 µg); cephalothin (CP, 30 µg);  
149 meropenem (MER, 10 µg); ceftazidime (CEF, 30 µg); ticarcillin (TIC, 75 µg); colistin  
150 sulphate (CT, 50 µg); sulfamethoxazole (SUL, 25 µg) and sulfamethoxazole/trimethoprim  
151 (SXT, 23.75/1.25 µg). The interpretation criteria (R, resistance; S, susceptible) based on  
152 inhibition zone diameters were as follows (mm): NA30: R≤13, S≥19; CIP5: R≤15, S≥21;  
153 STR10: R≤11, S≥15; GEN10: R≤12, S≥15; TET30: R≤11, S≥15; CP30: R≤14, S≥18;  
154 MER10: R≤15, S≥19; CEF30: R≤14, S≥18; TIC75: R≤15, S≥24; CT50: R≤10, S≥11;  
155 SUL25: R≤12, S≥17; SXT25: R≤10, S≥16. In each assay, the reference strain *Pseudomonas*  
156 *aeruginosa* DSM 1117 was used for quality control.

#### 157 *Determination of Minimum Inhibitory Concentrations (MICs)*

158 The minimum inhibitory concentrations (MICs) were determined using the Etest or the  
159 microdilution method at 30 °C. The Etest (BioMérieux, France) or MICE (OXOID, United  
160 Kingdom) were used for the antibiotics gentamicin (CN 256-0.015 µg/mL, OXOID,  
161 MA0116F), streptomycin (SM 0.064-1024 µg/mL, BioMérieux, 526800), ceftazidime (TZ  
162 0.016-256 µg/mL, BioMérieux, 412293), meropenem (MEM 32-0.002 µg/mL, OXOID,  
163 MA0121F), and sulfamethoxazole (SX 0.064-1024 µg/mL, BioMérieux, 412458). The  
164 microdilution method was used for tetracycline and metals (Andrews 2001), using bacterial  
165 suspensions of absorbance 0.08-0.1 at 625 nm in Mueller-Hinton broth supplemented with  
166 0.1 - 32 mg/L of tetracycline, 0.001 - 2 mmol/L of NaAsO<sub>2</sub>, 0.01 - 10 mmol/L of  
167 NiCl<sub>2</sub>·6H<sub>2</sub>O or 1 - 14 mmol/L of CuSO<sub>4</sub>·5H<sub>2</sub>O. For concentrations of CuSO<sub>4</sub>·5H<sub>2</sub>O above  
168 5 mmol/L the MICs were tested in Tris-buffered Mueller-Hinton broth. The MICs were

169 determined as the minimum concentration that inhibited visible bacterial growth after 24 h  
170 of incubation.

171 Based on preliminary distinctive results between the tested strains, the aminoglycoside  
172 gentamicin, and the metal arsenite were selected to assess their effects as stressors and will  
173 be from this point forward designated as stressors. Each of the five strains was assayed in  
174 stressor-free (SF) culture medium and in the presence of gentamicin or arsenite at  
175 concentrations close to the MIC value.

#### 176 *Stressors and growth kinetics*

177 Cultures were assayed in Mueller-Hinton broth or in this culture medium supplemented  
178 with adequate concentrations and volume of stressor solution. Therefore, strains H2Cu2,  
179 T6BT1 and L1PA1 were assayed in 125 mg/L gentamicin or 1.1 mmol/L As<sup>3+</sup>; strain  
180 H2Cu5 was assayed in 6 mg/L gentamicin or 0.01 mmol/L As<sup>3+</sup>; and strain T6BT10 was  
181 assayed in 0.4 mg/L gentamicin or 0.01 mmol/L As<sup>3+</sup>. Bacterial suspensions with an initial  
182 absorbance of 0.05 at 610 nm (A<sub>610</sub>) were incubated at 30 °C with orbital shaking (~70  
183 rpm) and were monitored every hour until reached the stationary phase (~24 h). Growth  
184 curves and kinetic parameters (growth rate, lag phase, and yield) were determined in  
185 triplicate in independent assays. Growth curves were represented as log values of A<sub>610</sub> in  
186 function of time. The lag phase was the period of time necessary to start the exponential  
187 phase. The growth rate ( $\mu$ ) was determined based on the slope of the curve during the  
188 exponential growth phase, according to the equation  $\ln N_t - \ln N_0 = \mu(t - t_0)$ , where N is  
189 the number of cells at time t. The growth yield corresponded to the maximum A<sub>610</sub>  
190 reached.

191 *Stressors and biofilm formation*

192 The capacity of each strain to form biofilm was tested in modified Luria-Bertani broth  
193 (mLB) (tryptone 5 g/L, yeast extract 2.5 g/L and sodium chloride 1 g/L) over a range of  
194 different stressor concentrations: 0.01, 0.05, 0.5 and 1.1 mmol/L As<sup>3+</sup>; 25, 75 and 125 mg/L  
195 of GEN; and 125, 250, 500 and 750 mg/L of STR, concentrations below the MICs for the  
196 strains H2Cu2, L1PA1 and T6BT1; and of 0.01 mmol/L of As<sup>3+</sup>; 0.4 and 6 mg/L of GEN;  
197 and 50 mg/L of STR for the strains H2Cu5 and T6BT10. The assays were performed in  
198 clear flat bottom 96-well polystyrene microtiter plates (Orange Scientific, Belgium) as  
199 described by Simões *et al.* (2007). Briefly, the microtiter wells were filled with 200 µL of  
200 bacterial suspension (A<sub>610</sub> = 0.1; prepared from overnight cultures in mLB at 30 °C) in  
201 mLB or in mLB supplemented with one of the stressors, incubated for 48 h at 30 °C and  
202 measured the absorbance at 620 nm (A<sub>620</sub>) in a microplate reader (FLUOstar optima,  
203 BMG Labtech, Germany). After that, the plates were washed with phosphate buffer and air-  
204 dried for 30 min. To assess and compare the biofilm formation, the biomass was fixed with  
205 methanol, left to dry, stained with crystal violet, washed again and the dye resuspended  
206 with glacial acetic acid prior to measuring the absorbance at 570 nm (A<sub>570</sub>). A negative  
207 control consisting of non-inoculated culture medium and a reference culture (*Pseudomonas*  
208 *aeruginosa* DSM 1117) were included in each assay. Each experiment was performed at  
209 least six times for each strain. The quantification of the biofilm formation was performed as  
210 described by Rode *et al.* (Rode *et al.* 2007), through the calculation of a ratio A<sub>570</sub>/A<sub>620</sub>,  
211 referring to absorbance at 570 nm (to measure the biofilm formation) and absorbance at 620  
212 nm (to measure the bacterial growth). The absorbance values were corrected by the  
213 subtraction of the respective absorbance measured in the negative control (non-inoculated

214 culture medium). With the procedure used, the possible contribution of the growth yield for  
215 the capacity to form biofilm was normalized by the use of the ratio  $A_{570}/A_{620}$ , referring to  
216 absorbance at 570 nm (measure of the biofilm formation) and absorbance at 620 nm  
217 (measure of the bacterial growth).

### 218 *Disinfectants and inactivation*

219 The effectiveness of the germicide UV radiation, chlorine or hydrogen peroxide, was tested  
220 in saline solution (0.85% (w/v) NaCl) bacterial suspensions of  $A_{610} = 0.1$ . Suspensions  
221 were prepared from 24 h Plate Count Agar (PCA) cultures. Samples collected at the  
222 beginning and over the assay were cultivated for enumeration on PCA and incubated at 30  
223 °C for 24-48 h.

224 For UV disinfection was used a germicide UV lamp with a wavelength of 254 nm, under  
225 which were exposed PCA plates onto which were spread 100  $\mu$ L of a bacterial suspension  
226 with about 10 to 300 CFU/mL. Exposure times were of periods of 0, 15, 30, 45, 60, 90 and  
227 150 seconds.

228 To test the effect of chlorine was used a solution of 10 mg/L sodium hypochlorite prepared  
229 from commercial bleach with a concentration of sodium hypochlorite equivalent to 5% (50  
230 g/L). Bacterial suspensions ( $A_{610} = 0.1$ ) prepared in saline solution were exposed to  
231 sodium hypochlorite at a final concentration of 5 mg/L. A solution of 1.5% (w/v) sodium  
232 thiosulfate was used to neutralize the effect of chlorine at different exposure times of: 0, 2,  
233 7, 12, 17, 25 and 60 minutes. Cultures were plated immediately after the addition of the  
234 neutralizing agent.

235 The effect of hydrogen peroxide was tested using a 0.1% solution prepared from a 30%  
236 stock (Carlo Erba Reagents, Italy). Bacterial suspensions ( $A_{610} = 0.1$ ) were exposed to  
237 hydrogen peroxide at a final concentration of 0.05% (v/v). A freshly prepared solution of  
238 bovine liver catalase (0.1 g/L) was used in a ratio 0.1/5 (v/v) to eliminate residual hydrogen  
239 peroxide (Fiorentino et al. 2015) after exposure times of: 0, 2, 7, 12, 17, 25 and 60 minutes.  
240 Cultures were plated after catalase addition.

#### 241 *Statistical analyses*

242 The effect of different stressors and the behavior of different strains was compared based  
243 on the parametric test one way ANOVA or the nonparametric tests Kruskal-Wallis and  
244 Mann-Whitney, depending if the results followed or not a normal distribution. The capacity  
245 to form biofilm in the presence and absence of stressors was compared based on the  
246 nonparametric test Mann-Whitney. The effect of disinfectants on cells inactivation was  
247 compared based on parametric one way ANOVA test with post-hoc test Tuckey. All the  
248 statistical analyzes were performed with the SPSS software package, version 23.0 (IBM  
249 SPSS software, Chicago, IL).

250

## 251 **RESULTS**

### 252 *Ralstonia* spp. tolerance to antibiotics and heavy metals

253 Based on the 16S rRNA gene sequence analyses, the *R. pickettii* and *R. mannitolilytica*  
254 strains studied clustered together with others from sources such as plant/animal,  
255 clinical/human, water and soil, or other environments (e.g. air) (**Fig. 1**). Although not

256 related to the isolation origin, three phylogenetic subgroups could be distinguished, one that  
257 included strains H2Cu2, L1PA1 and T6BT1, sharing a 16S rRNA gene sequence identity of  
258 99.7 - 99.9%, other including strain H2Cu5, with a 16S rRNA identity with first group of  
259 99.0 - 99.2% and another one of *R. mannitolilytica*, which, non-surprisingly included the  
260 strain T6BT10 with a 16S rRNA gene sequence identity of 97.8 - 98.2% with the *R.*  
261 *pickettii* isolates tested. These differences were not confirmed at the biochemical phenotype  
262 for which the five strains displayed a similar profile (data not shown).

263 All strains were observed to be resistant to colistin and ticarcillin and susceptible to the  
264 quinolones (nalidixic acid, ciprofloxacin), sulfonamides (sulfamethoxazole and  
265 sulfamethoxazole/trimethoprim), beta-lactams (cephalothin and ceftazidime) and  
266 tetracycline. Variable phenotypes were observed for meropenem and aminoglycosides  
267 susceptibility. Strains *R. pickettii* H2Cu5 and *R. mannitolilytica* T6BT10, both susceptible  
268 to gentamicin, differed on the susceptibility to streptomycin observed only in *R.*  
269 *mannitolilytica* T6BT10. Hence, gentamicin resistance was observed in the group of  
270 phylogenetically closely related strains H2Cu2, L1PA1 and T6BT1, with MIC-gentamicin  
271 values >256 mg/L, while strains H2Cu5 and T6BT10, in distinct phylogenetic subgroups,  
272 presented lower MIC values for both gentamicin and streptomycin (Table 2). MIC values  
273 for As<sup>3+</sup> were about 30 times higher in gentamicin resistant than in gentamicin susceptible  
274 isolates, while no differences among strains were observed for Ni<sup>2+</sup> or Cu<sup>2+</sup>. This finding  
275 suggested that aminoglycoside and arsenite resistance mechanisms might be associated.

276 Stressors and growth kinetic

277 Based on the hypothesis that a common mechanism of resistance could be used by these  
278 strains for gentamicin and arsenite, growth kinetic parameters were determined in the  
279 absence and in the presence of each of those stressors (Table 3). In the absence of any  
280 stressor, the growth rates for the five strains were similar ( $\sim 0.4 \text{ h}^{-1}$ ) (Table 3). Either  
281 gentamicin or arsenite led to significant ( $p < 0.05$ ) reductions in the growth rate, being the  
282 highest reductions observed in the presence of sub-inhibitory concentrations of gentamicin  
283 (Table 3). The lag phases in the absence of stressor ranged 0.7-0.9 h. In the presence of  
284 arsenite, but not in the presence of gentamicin, these values significantly ( $p < 0.05$ )  
285 increased (to 2.4-3.4 h) in the strains with highest MIC-As<sup>3+</sup> values (Table 3). In absence of  
286 stressors, growth yield ranged 2.4-2.9. These values that were significantly ( $p < 0.05$ )  
287 reduced in the presence of arsenite for strain H2Cu2 (to 2.2) or in the presence of  
288 gentamicin for strains H2Cu2, H2Cu5, L1PA1, and T6BT1 (to 0.5-0.9). In general, the  
289 reduction of growth yield was more pronounced in the presence of gentamicin than of  
290 arsenite. These differences in the growth parameters in the presence of gentamicin or  
291 arsenite suggest that even if a common resistance mechanism is used to grow in the  
292 presence of each of those stressors, probably distinct functions are targeted in the cell by  
293 the antibiotic or the metal.

#### 294 Effect of stressors in the capacity of biofilm formation

295 The capacity to form biofilm may be an advantage in *Ralstonia* spp. to face adverse  
296 conditions (Adley et al. 2005; Anderson et al. 1990; Di Domenico et al. 2016; Ryan et al.  
297 2011). Hence, it was hypothesized that the stressors aminoglycosides and arsenite could  
298 stimulate the capacity to produce biofilm (**Fig. 2**). The low concentrations of stressor  
299 tolerated by the aminoglycoside susceptible strains H2Cu5 and T6BT10 were not observed

300 to induce in those strains an increased capacity to form biofilm. In contrast, the strains  
301 resistant to the aminoglycosides gentamicin and streptomycin (H2Cu2, T6BT1 and L1PA1)  
302 presented significant increases in the capacity to form biofilm, with increases of 2-4 times  
303 for the lower concentrations and 5-11 times for the highest concentrations of  
304 aminoglycosides tested, in comparison with the non-stressor assays (**Fig. 2**). In the same  
305 way, 1.1 mmol/L arsenite a significant, although lower (1.5-1.9 times), increase of biofilm  
306 formation. The capacity to form biofilm can be associated with the production of  
307 polysaccharide capsules that facilitate the adherence to surfaces and the formation of  
308 biofilms (Moxon and Kroll 1990). It was thus hypothesized that the increased capacity to  
309 form biofilm could be due to an observable overproduction of capsule polysaccharides in  
310 the presence of sub-inhibitory concentration of gentamicin. However, this hypothesis was  
311 not proved, eventually because the method used to observe capsules was not sufficiently  
312 sensitive.

### 313 Disinfectants and inactivation

314 The hypothesis beyond these assays was that aminoglycoside and arsenite resistant strains  
315 would present a higher resilience against the different types of disinfectant - UV radiation,  
316 chlorine, and peroxide disinfection (**Fig. 3**). However, it was observed that only peroxide  
317 disinfection supported that hypothesis. Neither UV radiation nor chlorine were observed to  
318 produce a distinct effect on the gentamicin resistant or susceptible strains (**Fig. 3**). The UV  
319 radiation promoted a reduction of 1-2 log at each 15 min of exposure till the maximum  
320 period tested of 45 min (**Fig. 3A**). In the presence of 5 mg/L chlorine it was observed a  
321 sharp culture inactivation (2 min), to reach after 7 min of exposure, counts < 10 CFU/mL  
322 (Log 1) (**Fig. 3B**). In contrast to the other two disinfectants, hydrogen peroxide revealed



323 higher antibacterial effectiveness against the gentamicin and arsenite susceptible strains  
324 than against the resistant. Susceptible strains decreased to counts below the quantification  
325 limit (one log-unit) after 7 min of exposure, in contrast to the resistant strains that required  
326 12 min to reach < one log-unit (**Fig. 3C**).

327

## 328 **DISCUSSION**

329 *Ralstonia* spp. are ubiquitous, mainly in aquatic environments, including drinking water  
330 (Vaz-Moreira et al. 2017), and have been reported as contaminants of clinical sterile  
331 solutions or materials (Boutros et al. 2002; Labarca et al. 1999; McNeil et al. 1985) or as  
332 the prevalent species in hemodialysis water samples (Vincenti et al. 2014). *Ralstonia*  
333 *pickettii* strains from different clinical and environmental origins were reported as being  
334 resistant to gentamicin, ticarcillin and meropenem, although highly susceptible to  
335 ciprofloxacin, tetracycline and sulfamethoxazole/trimethoprim (Ryan and Adley 2013),  
336 confirming the phenotypes of the strains examined in this study. The observation that  
337 phylogenetically and phenotypically close *Ralstonia* strains, isolated from distinct aquatic  
338 environments, differed on the susceptibility to gentamicin (Table 1 and 2), suggested that it  
339 might be due to gene acquisition. Indeed, in a parallel study, whole genome sequencing  
340 from strains H2Cu2 and H2Cu5 showed that only the aminoglycoside resistant isolate  
341 contains genes associated with resistance to arsenic, and toxic compounds, encoding  
342 lysozyme inhibitors, or phages/prophages receptors (Vaz-Moreira et al. 2016).  
343 The correlation observed between the resistance to arsenite and gentamicin, may also be an  
344 indication of a possible mechanism of co-resistance (genetic linkage between two or more

345 resistance genes) or cross-resistance (same genetic determinant confers resistance to both  
346 antibiotics and metals), frequently reported for heavy metals and antibiotics (Baker-Austin  
347 et al. 2006; Dib et al. 2008; Seiler and Berendonk 2012; Zhou et al. 2015) specially in  
348 contaminated environments (Ahemad and Malik 2013).

349 Supposedly the physiological response to arsenite and aminoglycosides involves diverse  
350 mechanisms, as the response for biofilm formation and growth kinetic did not respond in  
351 the same mode for both antimicrobials (Table 2). However, the stress imposed by sub-  
352 inhibitory concentrations of aminoglycosides or of metals increased the capacity to form  
353 biofilms of the aminoglycoside resistant isolates (**Fig. 2**). This can be due to an unspecific  
354 stress response, not related with the resistance mechanism, in particular enhanced  
355 production of extracellular polymeric substances (EPS) which lead to cell adhesion (Baker-  
356 Austin et al. 2006; Donlan 2002; Donlan and Costerton 2002; Lindsay and von Holy 2006),  
357 and biofilm formation (Balaban 2008; Donlan and Costerton 2002; Lindsay and von Holy  
358 2006). Similar results were observed for *Pseudomonas aeruginosa* and *Escherichia coli*  
359 isolates, increasing their capacity to form biofilm in the presence of aminoglycosides (Aka  
360 and Haji 2015; Hoffman et al. 2005). However, Paul *et al.* (2014) observed a negative  
361 influence of 0.01 mmol/L arsenite on the capacity to form biofilm of *Pseudomonas* spp. or  
362 *Rhizobium* spp. strains. Some authors refer to the importance of the bacteria motility on the  
363 capacity to form biofilm (Guttenplan and Kearns 2013; O'Toole and Kolter 1998; Yao and  
364 Allen 2007). Indeed, *Ralstonia* spp. are motile and this property might influence the  
365 variable capacity to form biofilm observed in the present study. However, it was not  
366 possible to assess differences in motility capabilities in the tested strains or to investigate if  
367 stressors interfere with flagella and therefor with biofilm formation. Considering the history

368 of *Ralstonia* spp. as important colonizers of highly oligotrophic environments, other  
369 conditions to explore could be the effect of nutritional stress.

370 In contrast to what was observed for hydrogen peroxide, the survival to UV radiation or  
371 chlorine disinfection did not differ in aminoglycoside resistant or susceptible strains (**Fig.**  
372 **3**). This observation is probably related with the inactivation mechanisms involved,  
373 DNA/RNA damage for UV, cell metabolism for chlorine and oxidative stress for hydrogen  
374 peroxide (Estrela et al. 2002; Hijnen et al. 2006; McDonnell and Russell 1999). These  
375 results suggest that the mechanisms involved in the aminoglycoside resistance are probably  
376 not directly associated with the mechanisms of survival to UV radiation or chlorination.

377 Studies performed with a higher number of isolates, from different origins may give more  
378 consistency to these results. The capability of *Ralstonia* spp. to survive in some  
379 environments subjected to disinfection processes, as for example the wastewater or  
380 drinking water systems, is probably determinant for their capacity to spread or be  
381 transmitted to humans. It is curious to note that aminoglycoside resistant strains have  
382 increased capacity to form biofilm in the presence of some environmental stressors, since  
383 this may be a relevant factor to facilitate the survival and spread of aminoglycoside  
384 resistant *Ralstonia* spp. strains in the environments subjected to stress conditions, as  
385 antimicrobial challenges. These characteristics combined with the oligotrophic character  
386 may contribute for the *Ralstonia* ubiquity in aquatic habitats.

387

## 388 **CONCLUSIONS**

389 The aminoglycosides resistance was associated with the highest tolerance to arsenite.

390 Sub-inhibitory concentrations of gentamicin or arsenite significantly decreased the growth  
391 rate and yield, while arsenite but not gentamicin caused a significant increase of the lag  
392 phase. The biofilm formation was stimulated in the presence of aminoglycosides or  
393 arsenite, in the aminoglycoside resistant but not in the susceptible strains.

394 Disinfection with UV or chlorine presented identical effectiveness in aminoglycoside  
395 resistant or susceptible strains. In contrast, hydrogen peroxide presented higher  
396 effectiveness against aminoglycoside susceptible than resistant strains.

397 The results support the hypothesis that antibiotic resistance is associated with improved  
398 tolerance to stress.

399

#### 400 **ACKNOWLEDGEMENTS**

401 This work was financially supported by National Funds from FCT - Fundação para a  
402 Ciência e a Tecnologia through projects UID/Multi/50016/2013 and NanoDiaBac  
403 ENMed/0001/2014, and IVM grant (SFRH/BPD/87360/2012); by FCT through the project  
404 WaterJPI/0001/2013 - STARE (Stopping Antibiotic Resistance Evolution); and by Erasmus  
405 Mundus Project SUD-UE (EMA2-STRAND 1, LOT 14) through PF grant (2013-2589).

406

407

#### 408 **REFERENCES**

409 Adley C, Ryan M, Pembroke J, Saieb F (2005) *Ralstonia pickettii*: biofilm formation in  
410 high-purity water. *Biofilms: persistence and ubiquity*:261-271

411 Ahemad M, Malik A (2013) Prevalence of heavy metal and antibiotic resistance in bacterial  
412 isolates from metal polluted soils. Microbiol J. DOI:10.3923/mj.2013

413 Aka ST, Haji SH (2015) Sub-MIC of antibiotics induced biofilm formation of  
414 *Pseudomonas aeruginosa* in the presence of chlorhexidine. Braz J Microbiol  
415 46:149-154

416 Anderson RL, Holland BW, Carr JK, Bond WW, Favero MS (1990) Effect of disinfectants  
417 on pseudomonads colonized on the interior surface of PVC pipes. Am J Public  
418 Health 80:17-21

419 Andrews JM (2001) Determination of minimum inhibitory concentrations. J Antimicrob  
420 Chemother 48 Suppl 1:5-16

421 Baker-Austin C, Wright MS, Stepanauskas R, McArthur JV (2006) Co-selection of  
422 antibiotic and metal resistance. Trends Microbiol 14:176-182

423 Balaban N (2008) Control of biofilm infections by signal manipulation vol 2. Springer,

424 Becerra-Castro C, Machado RA, Vaz-Moreira I, Manaia CM (2015) Assessment of copper  
425 and zinc salts as selectors of antibiotic resistance in Gram-negative bacteria. Sci  
426 Total Environ 530-531:367-372

427 Boutros N, Gonullu N, Casetta A, Guibert M, Ingrand D, Lebrun L (2002) *Ralstonia*  
428 *pickettii* traced in blood culture bottles. J Clin Microbiol 40:2666-2667

429 CLSI (2015) Performance Standards for Antimicrobial Susceptibility Testing; Twenty-fifth  
430 Informational Supplement M100-S25. vol 35. Clinical and Laboratory Standards  
431 Institute, USA

432 Coenye T, Vandamme P, LiPuma JJ (2002) Infection by *Ralstonia* species in cystic fibrosis  
433 patients: identification of *R. pickettii* and *R. mannitolilytica* by polymerase chain  
434 reaction. Emerg Infect Dis 8:692-696

435 Coman I, Bilodeau L, Lavoie A, Carricart M, Tremblay F, Zlosnik JE, Berthiaume Y  
436 (2017) *Ralstonia mannitolilytica* in cystic fibrosis: A new predictor of worse  
437 outcomes. *Respir Med Case Rep* 20:48-50

438 Daxboeck F, Stadler M, Assadian O, Marko E, Hirschl AM, Koller W (2005)  
439 Characterization of clinically isolated *Ralstonia mannitolilytica* strains using  
440 random amplification of polymorphic DNA (RAPD) typing and antimicrobial  
441 sensitivity, and comparison of the classification efficacy of phenotypic and  
442 genotypic assays. *J Med Microbiol* 54:55-61

443 De Baere T et al. (2001) Classification of *Ralstonia pickettii* biovar 3/'thomasii' strains  
444 (Pickett 1994) and of new isolates related to nosocomial recurrent meningitis as  
445 *Ralstonia mannitolilytica* sp. nov. *Int J Syst Evol Microbiol* 51:547-558

446 Di Domenico EG et al. (2016) Development of an in vitro assay, based on the biofilm ring  
447 test®, for rapid profiling of biofilm-growing bacteria. *Frontiers in microbiology*  
448 7:1429

449 Dib J, Motok J, Zenoff VF, Ordonez O, Farias ME (2008) Occurrence of resistance to  
450 antibiotics, UV-B, and arsenic in bacteria isolated from extreme environments in  
451 high-altitude (above 4400 m) Andean wetlands. *Curr Microbiol* 56:510-517

452 Donlan RM (2002) Biofilms: microbial life on surfaces. *Emerg Infect Dis* 8:881-890

453 Donlan RM, Costerton JW (2002) Biofilms: survival mechanisms of clinically relevant  
454 microorganisms. *Clin Microbiol Rev* 15:167-193

455 Estrela C, Estrela CR, Barbin EL, Spano JC, Marchesan MA, Pecora JD (2002) Mechanism  
456 of action of sodium hypochlorite. *Braz Dent J* 13:113-117

457 Falcone-Dias M, Vaz-Moreira I, Manaia CM (2012) Bottled mineral water as a potential  
458 source of antibiotic resistant bacteria. *Water Res* 46:3612-3622

459 Fernandez C et al. (1996) Nosocomial outbreak of *Burkholderia pickettii* infection due to a  
460 manufactured intravenous product used in three hospitals. Clin Infect Dis 22:1092-  
461 1095

462 Ferreira da Silva M, Vaz-Moreira I, Gonzalez-Pajuelo M, Nunes OC, Manaia CM (2007)  
463 Antimicrobial resistance patterns in *Enterobacteriaceae* isolated from an urban  
464 wastewater treatment plant. FEMS Microbiol Ecol 60:166-176

465 Fiorentino A, Ferro G, Alferez MC, Polo-Lopez MI, Fernandez-Ibanez P, Rizzo L (2015)  
466 Inactivation and regrowth of multidrug resistant bacteria in urban wastewater after  
467 disinfection by solar-driven and chlorination processes. J Photochem Photobiol B  
468 148:43-50

469 Guttenplan SB, Kearns DB (2013) Regulation of flagellar motility during biofilm  
470 formation. FEMS Microbiol Rev 37:849-871

471 Hijnen WA, Beerendonk EF, Medema GJ (2006) Inactivation credit of UV radiation for  
472 viruses, bacteria and protozoan (oo)cysts in water: a review. Water Res 40:3-22

473 Hoffman LR, D'Argenio DA, MacCoss MJ, Zhang Z, Jones RA, Miller SI (2005)  
474 Aminoglycoside antibiotics induce bacterial biofilm formation. Nature 436:1171-  
475 1175

476 Jukes TH, Cantor CR (1969) Evolution of protein molecules. In: Munro (ed) Mammalian  
477 Protein Metabolism, vol 3. Academic Press, New York,

478 Khajuria A, Praharaj AK, Grover N, Kumar M (2014) Emergence of VIM-2 metallo-beta-  
479 lactamase producing *Ralstonia pickettii* clinical isolate in India. Indian J Med  
480 Microbiol 32:191-193

481 Koenig DW, Pierson DL (1997) Microbiology of the Space Shuttle water system. Water  
482 Sci Technol 35:59-64

483 Kulakov LA, McAlister MB, Ogden KL, Larkin MJ, O'Hanlon JF (2002) Analysis of  
484 bacteria contaminating ultrapure water in industrial systems. *Appl Environ*  
485 *Microbiol* 68:1548-1555

486 Labarca JA et al. (1999) A multistate nosocomial outbreak of *Ralstonia pickettii*  
487 colonization associated with an intrinsically contaminated respiratory care solution.  
488 *Clin Infect Dis* 29:1281-1286

489 Letunic I, Bork P (2016) Interactive tree of life (iTOL) v3: an online tool for the display  
490 and annotation of phylogenetic and other trees. *Nucleic Acids Res* 44:W242-245

491 Lindsay D, von Holy A (2006) Bacterial biofilms within the clinical setting: what  
492 healthcare professionals should know. *J Hosp Infect* 64:313-325

493 Maroye P, Doermann HP, Rogues AM, Gachie JP, Megraud F (2000) Investigation of an  
494 outbreak of *Ralstonia pickettii* in a paediatric hospital by RAPD. *J Hosp Infect*  
495 44:267-272

496 McDonnell G, Russell AD (1999) Antiseptics and disinfectants: activity, action, and  
497 resistance. *Clin Microbiol Rev* 12:147-179

498 McKinney RE (1953) Staining bacterial polysaccharides. *J Bacteriol* 66:453

499 McNeil MM et al. (1985) Nosocomial *Pseudomonas pickettii* colonization associated with a  
500 contaminated respiratory therapy solution in a special care nursery. *J Clin Microbiol*  
501 22:903-907

502 Mijnenonckx K, Provoost A, Ott C, Venkateswaran K, Mahillon J, Leys N, Van Houdt R  
503 (2013) Characterization of the survival ability of *Cupriavidus metallidurans* and  
504 *Ralstonia pickettii* from space-related environments. *Microb Ecol* 65:347-360

505 Moxon ER, Kroll JS The role of bacterial polysaccharide capsules as virulence factors. In,  
506 Berlin, Heidelberg, 1990. *Bacterial Capsules*. Springer Berlin Heidelberg, pp 65-85



507 O'Toole GA, Kolter R (1998) Flagellar and twitching motility are necessary for  
508 *Pseudomonas aeruginosa* biofilm development. Mol Microbiol 30:295-304

509 Orme J, Rivera-Bonilla T, Loli A, Blattman NN (2015) Native valve endocarditis due to  
510 *Ralstonia pickettii*: a case report and literature review. Case Rep Infect Dis  
511 2015:324675

512 Pan W, Zhao Z, Dong M (2011) Lobar pneumonia caused by *Ralstonia pickettii* in a sixty-  
513 five-year-old Han Chinese man: a case report. J Med Case Rep 5:377

514 Paul D, Poddar S, Sar P (2014) Characterization of arsenite-oxidizing bacteria isolated from  
515 arsenic-contaminated groundwater of West Bengal. J Environ Sci Health A Tox  
516 Hazard Subst Environ Eng 49:1481-1492

517 Riley PS, Weaver RE (1975) Recognition of *Pseudomonas pickettii* in the clinical  
518 laboratory: biochemical characterization of 62 strains. J Clin Microbiol 1:61-64

519 Rode TM, Langsrud S, Holck A, Moretro T (2007) Different patterns of biofilm formation  
520 in *Staphylococcus aureus* under food-related stress conditions. Int J Food Microbiol  
521 116:372-383

522 Ryan MP (2009) Genotypic and phenotypic analysis of *Ralstonia pickettii* high purity water  
523 isolates. University of Limerick

524 Ryan MP, Adley CC (2013) The antibiotic susceptibility of water-based bacteria *Ralstonia*  
525 *pickettii* and *Ralstonia insidiosa*. J Med Microbiol 62:1025-1031

526 Ryan MP, Pembroke JT, Adley CC (2006) *Ralstonia pickettii*: a persistent Gram-negative  
527 nosocomial infectious organism. J Hosp Infect 62:278-284

528 Ryan MP, Pembroke JT, Adley CC (2007) *Ralstonia pickettii* in environmental  
529 biotechnology: potential and applications. J Appl Microbiol 103:754-764

530 Ryan MP, Pembroke JT, Adley CC (2009) Novel Tn4371-ICE like element in *Ralstonia*  
531 *pickettii* and genome mining for comparative elements. BMC Microbiol 9:242

532 Ryan MP, Pembroke JT, Adley CC (2011) Genotypic and phenotypic diversity of *Ralstonia*  
533 *pickettii* and *Ralstonia insidiosa* isolates from clinical and environmental sources  
534 including High-purity Water. Diversity in *Ralstonia pickettii*. BMC Microbiol  
535 11:194

536 Seiler C, Berendonk TU (2012) Heavy metal driven co-selection of antibiotic resistance in  
537 soil and water bodies impacted by agriculture and aquaculture. Front Microbiol  
538 3:399

539 Simões LC, Simões M, Vieira MJ (2007) Biofilm interactions between distinct bacterial  
540 genera isolated from drinking water. Appl Environ Microbiol 73:6192-6200

541 Stelzmueller I et al. (2006) *Ralstonia pickettii*-innocent bystander or a potential threat? Clin  
542 Microbiol Infect 12:99-101

543 Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular  
544 Evolutionary Genetics Analysis version 6.0. Mol Biol Evol 30:2725-2729

545 Udayappan SD et al. (2017) Intestinal *Ralstonia pickettii* augments glucose intolerance in  
546 obesity. PloS one 12:e0181693

547 Vaneechoutte M, De Baere T, Wauters G, Steyaert S, Claeys G, Vogelaers D,  
548 Verschraegen G (2001) One case each of recurrent meningitis and hemoperitoneum  
549 infection with *Ralstonia mannitolilytica*. J Clin Microbiol 39:4588-4590

550 Vaneechoutte M, Kampfer P, De Baere T, Falsen E, Verschraegen G (2004) *Wautersia* gen.  
551 nov., a novel genus accommodating the phylogenetic lineage including *Ralstonia*  
552 *eutropha* and related species, and proposal of *Ralstonia* [*Pseudomonas*] *syzygii*  
553 (Roberts et al. 1990) comb. nov. Int J Syst Evol Microbiol 54:317-327

554 Vaz-Moreira I, Nunes OC, Manaia CM (2014) Bacterial diversity and antibiotic resistance  
555 in water habitats: searching the links with the human microbiome. FEMS Microbiol  
556 Rev 38:761–778

557 Vaz-Moreira I, Nunes OC, Manaia CM (2017) Ubiquitous and persistent *Proteobacteria*  
558 and other Gram-negative bacteria in drinking water. Sci Total Environ 586:1141-  
559 1149

560 Vaz-Moreira I, Tamames J, Martinez JL, Manaia CM (2016) Draft genome sequences of  
561 two *Ralstonia pickettii* strains with different aminoglycoside resistance phenotypes.  
562 Genome Announc 4

563 Verschraegen G, Claeys G, Meeus G, Delanghe M (1985) *Pseudomonas pickettii* as a cause  
564 of pseudobacteremia. J Clin Microbiol 21:278-279

565 Vincenti S et al. (2014) Non-fermentative gram-negative bacteria in hospital tap water and  
566 water used for haemodialysis and bronchoscope flushing: prevalence and  
567 distribution of antibiotic resistant strains. Sci Tot Environ 499:47-54

568 Waugh JB, Granger WM, Gaggar A (2010) Incidence, relevance and response for *Ralstonia*  
569 respiratory infections. Clin Lab Sci 23:99

570 Yao J, Allen C (2007) The plant pathogen *Ralstonia solanacearum* needs aerotaxis for  
571 normal biofilm formation and interactions with its tomato host. J Bacteriol  
572 189:6415-6424

573 Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J (2017) Introducing EzBioCloud:  
574 a taxonomically united database of 16S rRNA gene sequences and whole-genome  
575 assemblies. Int J Syst Evol Microbiol 67:1613-1617

576 Zhou Y, Xu YB, Xu JX, Zhang XH, Xu SH, Du QP (2015) Combined toxic effects of  
577 heavy metals and antibiotics on a *Pseudomonas fluorescens* strain ZY2 isolated  
578 from swine wastewater. Int J Mol Sci 16:2839-2850  
579

580 **TABLES:**

581 Table 1. Group of isolates tested in this study

582

<b>Strain</b>	<b>Species</b>	<b>Source of isolation</b>	<b>Isolation medium and conditions</b>	<b>Abundance in the source (order of magnitude, CFU's/mL)</b>	<b>Reference</b>
<b>H2Cu2</b>	<i>R. pickettii</i>	Hospital wastewater	Culture enrichment in modified Luria-Bertani broth with Cu <sup>2+</sup> (2.5 mmol/L)	10 <sup>3</sup>	Becerra-Castro <i>et al.</i> 2015
<b>H2Cu5</b>	<i>R. pickettii</i>	Hospital wastewater	Culture enrichment in modified Luria-Bertani broth with Cu <sup>2+</sup> (2.5 mmol/L)	10 <sup>3</sup>	Becerra-Castro <i>et al.</i> 2015
<b>L1PA1</b>	<i>R. pickettii</i>	Mineral water	Pseudomonas isolation agar with 32mg/L amoxicillin	10 <sup>1</sup>	Falcone-Dias <i>et al.</i> 2012
<b>T6BT1</b>	<i>R. pickettii</i>	Tap water	Tergitol 7-agar	10 <sup>-1</sup>	Vaz-Moreira <i>et al.</i> 2013
<b>T6BT10</b>	<i>R. mannitolilytica</i>	Tap water	Tergitol 7-agar	10 <sup>-1</sup>	Vaz-Moreira <i>et al.</i> 2013

583 Table 2. Minimum Inhibitory Concentrations (MICs) for antibiotics and metals determined  
 584 for the *Ralstonia* spp. strains under study.

Strain	MICs								
	GEN	STR	TET	MER	CEF	SUL	As <sup>3+</sup>	Ni <sup>2+</sup>	Cu <sup>2+</sup>
	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mmol/L)	(mmol/L)	(mmol/L)
<b>H2Cu2</b>	> 256	>1024	1	>32	6	24	1.4	4	12
<b>H2Cu5</b>	6	56	0.25	6	6	4	0.05	4	12
<b>L1PA1</b>	> 256	>1024	1	>32	6	24	1.4	4	12
<b>T6BT1</b>	> 256	>1024	1	16	8	24	1.4	4	12
<b>T6BT10</b>	0.5	4	8	>32	4	4	0.05	4	12

585 GEN, gentamicin; TET, tetracycline; MER, meropenem; CEF, ceftazidime; SUL, sulfamethoxazole; STR,  
 586 streptomycin and metal salts of As<sup>3+</sup>, Ni<sup>2+</sup> and Cu<sup>2+</sup>

587 Table 3. Variations on the bacterial growth parameters growth rate, phase lag and yield, under sub-inhibitory concentrations of arsenite  
 588 ( $\text{As}^{3+}$ ) or gentamicin (GEN) or control conditions (stressor free, SF).

Strain (stressor concentration)	Growth rate (per hour)			Phase Lag (hours)			Yield (A610)					
	SF	$\text{As}^{3+}$	GEN	SF	$\text{As}^{3+}$	GEN	SF	$\text{As}^{3+}$	GEN			
H2Cu2 125 mg/L GEN or 1.1 mmol/L $\text{As}^{3+}$	0.4±0.03	1;a,b 0.3±0.02	2;a 0.1±0.01	3;a	0.9±0.2	1;a 3.4±0.5	2;a 2.0±0.7	1,2;a	2.9±0.1	1;a 2.2±0.2	2;a 0.5±0.1	3;a
H2Cu5 6 mg/L GEN or 0.01 mmol/L $\text{As}^{3+}$	0.3±0.01	1;a 0.3±0.04	1;a 0.1±0.01	2;b	0.7±0.4	1;a 1.0±0.3	1;b 2.2±2.5	1;a	2.5±0.3	1;a 2.0±0.3	1;a 0.9±0.2	2;a
L1PA1 125 mg/L GEN or 1.1 mmol/L $\text{As}^{3+}$	0.4±0.04	1;a,b 0.3±0.04	1;a,b 0.2±0.01	2;c	0.7±0.4	1;a 3.0±0.9	2;a 0.8±0.1	1;a	2.9±0.4	1;a 2.5±0.5	1;a 0.8±0.1	2;a
T6BT1 125 mg/L GEN or 1.1 mmol/L $\text{As}^{3+}$	0.4±0.02	1;a,b 0.3±0.01	2;a 0.2±0.01	3;b	0.9±0.3	1;a 2.4±0.3	2;a 1.4±0.4	1;a	2.7±0.1	1;a 2.4±0.3	1;a 0.7±0.1	2;a
T6BT10 0.4 mg/L GEN or 0.01 mmol/L $\text{As}^{3+}$	0.4±0.01	1;b 0.4±0.01	1,2;b 0.4±0.01	2;d	0.8±0.1	1;a 0.8±0.1	1;b 0.7±0.1	1;a	2.4±0.5	1;a 2.1±0.4	1;a 1.9±0.6	1;b

589 A610, bacterial suspension absorbance at 610 nm;

590 Statistically significant differences between stress conditions (SF,  $\text{As}^{3+}$  and GEN) are indicated by the numbers: 1, 2, 3; and significant differences between

591 strains are indicated by the letters: a, b, c, d.

592

593 **FIGURES:**

594

595 **Fig. 1** Environmental distribution of *Ralstonia pickettii* and *Ralstonia mannitolilytica*,  
596 including the strains studied (in red).

597

598

599 **Fig. 2** Influence of stressors on the ability of biofilm formation, for the strains H2Cu2 (A),  
600 T6BT1 (B) and L1PA1 (C). The quantification of the biofilm formation was performed  
601 through the calculation of a ratio  $A_{570}/A_{620}$ , referring to absorbance at 570 nm (measure of  
602 the biofilm formation) and absorbance at 620 nm (measure of the bacterial growth). The  
603 non-inoculated control presented a ratio  $A_{570}/A_{620}$  of  $1.0\pm 0.1$ ; and the *P. aeruginosa*  
604 presented ratios  $A_{570}/A_{620}$  of:  $3.1\pm 0.7$  for stressor-free (SF);  $2.2\pm 0.9$  for gentamicin (GEN)  
605 6 mg/L;  $2.8\pm 0.7$  for streptomycin (STR) 50 mg/L;  $0.2\pm 0.09$  for meropenem 4 mg/L;  
606  $1.9\pm 0.9$  for arsenite (As) 0.01 mmol/L;  $0.3\pm 0.05$  for copper 6.0 mmol/L; and  $0.7\pm 0.2$  for  
607 nickel 2.5 mmol/L.

608

609

610 **Fig. 3** Bacterial inactivation with A) UV radiation, B) chlorine (5 mg/L), and C) hydrogen  
611 peroxide (0.05%).