



The influence of alkalosis on repeated high-intensity exercise performance and acid–base balance recovery in acute moderate hypoxic conditions

Lewis Anthony Gough^{1,2} · Danny Brown² · Sanjoy K. Deb² · S. Andy Sparks² · Lars R. McNaughton^{2,3}Received: 12 June 2018 / Accepted: 24 August 2018 / Published online: 8 September 2018
© The Author(s) 2018

Abstract

Purpose Exacerbated hydrogen cation (H^+) production is suggested to be a key determinant of fatigue in acute hypoxic conditions. This study, therefore, investigated the effects of $NaHCO_3$ ingestion on repeated 4 km TT cycling performance and post-exercise acid–base balance recovery in acute moderate hypoxic conditions.

Methods Ten male trained cyclists completed four repeats of 2×4 km cycling time trials (TT_1 and TT_2) with 40 min passive recovery, each on different days. Each TT series was preceded by supplementation of one of the 0.2 g kg^{-1} BM $NaHCO_3$ (SBC2), 0.3 g kg^{-1} BM $NaHCO_3$ (SBC3), or a taste-matched placebo (0.07 g kg^{-1} BM sodium chloride; PLA), administered in a randomized order. Supplements were administered at a pre-determined individual time to peak capillary blood bicarbonate concentration ($[HCO_3^-]$). Each TT series was also completed in a normobaric hypoxic chamber set at $14.5\% FiO_2$ ($\sim 3000 \text{ m}$).

Results Performance was improved following SBC3 in both TT_1 (400.2 ± 24.1 vs. $405.9 \pm 26.0 \text{ s}$; $p = 0.03$) and TT_2 (407.2 ± 29.2 vs. $413.2 \pm 30.8 \text{ s}$; $p = 0.01$) compared to PLA, displaying a very likely benefit in each bout. Compared to SBC2, a likely and possible benefit was also observed following SBC3 in TT_1 ($402.3 \pm 26.5 \text{ s}$; $p = 0.15$) and TT_2 ($410.3 \pm 30.8 \text{ s}$; $p = 0.44$), respectively. One participant displayed an ergolytic effect following SBC3, likely because of severe gastrointestinal discomfort, as SBC2 still provided ergogenic effects.

Conclusion $NaHCO_3$ ingestion improves repeated exercise performance in acute hypoxic conditions, although the optimal dose is likely to be 0.3 g kg^{-1} BM.

Keywords Buffers · Alkalosis · Individual pursuit · Personalised nutrition · Track cycling

Abbreviations

Ca^{2+}	Calcium
Cl^-	Chloride
CI	Confidence intervals

FiO_2	Fraction of inspired oxygen
GI	Gastrointestinal
HCO_3^-	Bicarbonate anion concentrations
H^+	Hydrogen cation
K^+	Potassium
SID	Strong ion difference
VO_{2max}	Maximal rate of oxygen uptake
Na^+	Sodium
O_2	Oxygen
SpO_2	Haemoglobin saturation of oxygen
$NaHCO_3$	Sodium bicarbonate
$NaCl$	Sodium chloride
TT	Time trial

Communicated by Michael Lindinger.

✉ Lewis Anthony Gough
lewis.gough@bcu.ac.uk

¹ Sport and Physical Activity Department, Faculty of Health and Life Sciences, Birmingham City University, Birmingham B13 3TN, UK

² Sports Nutrition and Performance Group, Department of Sport and Physical Activity, Edge Hill University, Ormskirk, Lancashire L39 4QP, UK

³ Department of Sport and Movement Studies, Faculty of Health Science, University of Johannesburg, Johannesburg, South Africa

Introduction

Repeated bouts of high-intensity exercise are a frequent feature of training and competition in athletes (Monedero and Donne 2000; Barnett 2006). The recovery between these exercise bouts is an essential component for determining the effectiveness of the subsequent bout. Enhanced recovery can allow athletes to tolerate greater training loads in the subsequent bout, potentially enhancing the post-training adaptation as a result (Barnett 2006). Whereas, in competition, enhancing recovery is an important component to sustain performance within the subsequent bout. This is applicable to sports such as track cycling, swimming, or a rowing regatta series which involve heats, semi-finals, and finals within a short amount of time (Al-Nawaiseh et al. 2016; Monedero and Donne 2000). Specifically, the gap between the men's team pursuit first round and the final at the Rio 2016 Olympics was separated by just 60 min. Considering that most national and Olympic records are achieved within the preliminary rounds of these events (Al-Nawaiseh et al. 2016), this suggests that full recovery is not always possible during these time frames or that current recovery practices are not optimal. Therefore, interventions to improve recovery and sustain subsequent exercise performance are important.

A major factor that may hamper post-training recovery and the subsequent bout of exercise is the metabolic disturbance that occurs following an initial high-intensity exercise bout (Barnett 2006). Ward et al. (2016) reported that the decline in pH and HCO_3^- following a 4 km cycling time trial (TT; team pursuit distance) was substantial, and reflective of metabolic acidosis (pH 7.16 ± 0.08 , HCO_3^- 11.9 ± 2.3 mmol l^{-1}), which was measured using capillary blood samples. Full recovery of such disturbances is likely to take over 75 min, as Callahan et al. (2017) reported HCO_3^- was 6.2 mmol l^{-1} below baseline at this point (19.5 ± 1.4 vs. 25.7 ± 1.0 mmol l^{-1}). If only 60 min is available for recovery, therefore, such as that during track cycling events, an existing acid–base balance perturbation will be evident. Although contentious (Westerblad 2016), critical rises in hydrogen cation (H^+) accumulation are linked to a reduction in both the release and uptake of calcium ions (Ca^{2+}) from the sarcoplasmic reticulum (Allen et al. 2008), disruption of key enzymes of the glycolytic pathway (Hollidge-Horvat et al. 1999), and a reduction in muscle excitability and action potentials by reducing the strong ion difference (SID) (Cairns and Lindinger 2008). In turn, this may hamper subsequent performance by reducing the capability for muscle force production (Cairns 2006; Fitts 2016). It is intuitive to suggest, therefore, that interventions to accelerate post-exercise

recovery of acid–base balance could be beneficial for a subsequent bout of exercise.

The ingestion of sodium bicarbonate (NaHCO_3) has been shown to accelerate post-exercise acid–base balance recovery and subsequent exercise performance (Pruscino et al. 2008; Zabala et al. 2008, 2011; Gough et al. 2017a). Pruscino et al. (2008) reported a 'trivial' to 'moderate' benefit in the second bout of a 2×200 m freestyle swim (interspersed with a 30 min recovery) following pre-exercise ingestion of NaHCO_3 . It was then later reported that NaHCO_3 ingestion 30 min into a 90 min post-exercise recovery significantly improved subsequent cycling capacity at 100% peak mean minute power by 16.6% (Gough et al. 2017a). Both studies reported that capillary blood pH and HCO_3^- recovery were accelerated and above baseline at the end of the recovery period, whereas the placebo condition failed to fully recover. This time frame between bouts might explain why ergogenic effects were observed, as this allowed pH and HCO_3^- to recover sufficiently. Conversely, no effect of NaHCO_3 ingestion has been reported on three repeated Wingate tests separated shorter recovery time frames of between 15 and 30 min (Zabala et al. 2008, 2011), or during three repeated high-intensity swimming bouts separated by 20 min (Pierce et al. 1992). Zabala et al. (2008) showed that neither pH, nor HCO_3^- recovered back to baseline levels between the three Wingate tests, suggesting that the recovery of pH and HCO_3^- to this level may be important to produce ergogenic effects in the subsequent exercise bouts.

Despite the promising effects of NaHCO_3 ingestion to improve repeated bouts of exercise in normoxia, this strategy has yet to be applied to acute hypoxia. Nonetheless, Robergs et al. (2005) reported the use of NaHCO_3 combined with sodium citrate lead to post-exercise recovery of pH and HCO_3^- to baseline in approximately 50 min at 1570 m terrestrial altitude, whereas the placebo condition failed to recover to baseline within the 80 min sampling period. Robergs et al. (2005) featured no subsequent bout of exercise, however, and it is also unclear if the participant cohort completed an acclimatisation period in the terrestrial altitude location. This enhanced post-exercise recovery displayed by Robergs et al. (2005) is potentially important, nonetheless, as athletes may complete multiple bouts of high-intensity exercise to maximise the adaptation from hypoxic training schedules, therefore, highlighting the need for optimal recovery. The use of pre-exercise NaHCO_3 ingestion may, in turn, result in a blunting of the initial stress of acid–base balance during an initial bout of exercise, but also improve recovery, leading to improved subsequent exercise performance. This may have a cumulative effect in sustaining training volume and intensity during hypoxic training schedules, particularly considering chronic NaHCO_3 ingestion has been shown to be effective at improving performance following training schedules at sea level (Egger et al. 2014; Durkalec-Michalski et al.

2018). The aim of this study, therefore, was to investigate the effects of both 0.2 g kg^{-1} BM and 0.3 g kg^{-1} BM NaHCO_3 on repeated bouts of 4 km TT cycling performance in acute moderate hypoxic conditions. The hypothesis of this study was that both doses of NaHCO_3 would improve both bouts of exercise compared to the placebo; however, 0.3 g kg^{-1} BM NaHCO_3 would improve performance to the greatest extent.

Methods

Participants and compliance with ethical standards

Ten trained male cyclists (age 27 ± 8 years, body mass 82 ± 9 kg, hypoxic maximal rate of oxygen uptake ($\text{VO}_{2\text{max}}$) $48.5 \pm 5.6 \text{ ml kg min}^{-1}$, and hypoxic peak power output $331 \pm 40 \text{ W}$) volunteered for this study. All participants' training load was reflective of a 'trained' cyclist (De Pauw et al. 2013). Ethical approval was granted from the Universities Research Ethics Committee (URESC16-LG01; Edge Hill University), and all participants provided written informed consent.

Experimental overview

Using methods previously described (Gough et al. 2017b), an initial maximal rate of oxygen uptake ($\text{VO}_{2\text{max}}$) test was conducted in a normobaric hypoxic chamber set at a fraction of inspired oxygen (FiO_2) of 14.5% ($\sim 3000 \text{ m}$). Participants then visited the laboratory on a further six separate occasions in a block randomized, crossover, and double-blind designed study ($2 \times$ identification of peak blood HCO_3^- , and $4 \times$ cycling TTs). Individual time to peak HCO_3^- was determined prior to the cycling time trials, using a previously described method (Gough et al. 2017c). This entailed participants ingesting either 0.2 g kg^{-1} BM NaHCO_3 (SBC2) or 0.3 g kg^{-1} BM NaHCO_3 (SBC3) on separate occasions, followed by a quiet rest for 180 min. Finger prick capillary blood samples were taken every 10 min, and the highest value of HCO_3^- was then used as the individual time to peak. Twenty-four hours prior to each cycling TT, participants refrained from consumption of alcohol and caffeine, any strenuous activity, and maintenance of nutritional intake, which was confirmed via use of written nutrition/training diaries. Finally, participants were verbally screened to ensure avoidance of beta alanine ingestion had not occurred prior to enrolment onto the study, to account for the long washout of carnosine (Baguet et al. 2009).

Time trial protocol, supplementation of sodium bicarbonate, and blood measures

Participants completed $2 \times 4 \text{ km}$ TTs (TT_1 and TT_2) interspersed with a 40 min recovery. The protocols for the TT, including the self-selected warm-up, were identical to those detailed in the previous research (Gough et al. 2017c), whereby only cadence and gear was displayed to the participant. Each TT series was preceded by supplementation of one of the SBC2, SBC3, or a taste-matched placebo (0.07 g kg^{-1} BM sodium chloride; PLA) administered in a block-randomized order. The randomisation of supplements was carried out by an individual who was not involved in the research, and performance times remained double-blind until completion of the study. Participants remained seated until their respective pre-determined time to peak HCO_3^- in normoxic conditions. Once reached, participants then entered the normobaric hypoxic chamber (FiO_2 14.5%) for 10 min, prior to beginning the TT_1 warm-up. Following TT_1 , participants completed a passive recovery entailing a quiet seated rest for 40 min within the hypoxic environment. Finger prick capillary blood samples for acid–base balance (pH and HCO_3^-), electrolytes (K^+ , Na^+ , Ca^{2+} , and Cl^-), lactate, and haemoglobin saturation with oxygen (SpO_2) were recorded pre-exercise, at time to peak HCO_3^- , immediately post-exercise and at 10 min intervals during the 40 min passive recovery. These blood samples were immediately analysed using a reliable blood gas analyser (ABL800BASIC, Radiometer Medical Ltd., Denmark), apart from lactate, which was analysed using a reliable and accurate lactate pro 2 analyser (Arkray, Japan) (Pyne et al. 2000; Bonaventura et al. 2015). To maintain the double-blind nature of the study, the screen on the blood gas analyser was covered, and an individual who was not part of the research stored the data until completion of the study. Calculation of the apparent SID was conducted using the formula ($[\text{K}^+] + [\text{Na}^+] + [\text{Ca}^{2+}] + [\text{Na}^+] - [\text{Cl}^-] - [\text{Lac}^-]$) (Lloyd 2004). Afterwards, participants completed the TT_2 warm-up, followed by TT_2 . Additional blood samples were obtained following the warm-up in TT_2 and immediately post TT_2 for the same measures previously described.

Perceptual measures

Rating of perceived exertion for overall body exertion (RPE_O) and the legs (RPE_L) was recorded during TT_1 and TT_2 every 1 km, whilst HR and SPO_2 were recorded during both TT_1 and TT_2 at every 500 m split and at 10 min intervals during recovery. Gastrointestinal (GI) discomfort was recorded at 10 min intervals after ingestion of the supplement (i.e., SBC2, SBC3, or PLA) up to the

individual time to peak HCO_3^- , and at 10 min intervals during the 40 min recovery. At individual time to peak HCO_3^- , participants were asked if they could determine which supplement they had ingested using a supplement belief questionnaire.

Statistical analysis

No evidence of a violation for normality and sphericity was evident in any assessed variable, and therefore, the appropriate parametric statistical tests were employed. A paired t test was conducted for the following: both the time to peak and absolute change in pH and HCO_3^- , and both the severity and aggregated score for GI discomfort following SBC treatments. Performance data (time to TT completion and mean power) and blood parameters (change in pH and HCO_3^- during TT_1 , recovery, and TT_2) were analysed using a repeated measures ANOVA. In addition, magnitude-based inferences (MBI) with 90% confidence intervals (CI) were calculated for performance data and interpreted using an adapted method from a freely available spreadsheet (Batterham and Hopkins 2006). The thresholds to depict a benefit or harm were set as the typical error of the 4 km TT when converted from a percentage to an absolute value. This was completed by calculating the difference score in each individual, then calculating the standard deviation of the difference scores, and finally by dividing this by $\sqrt{2}$ (Swinton et al. 2018). This was used as in many cases, the 0.2 small effect size of Cohen d (Cohen 1988) is less than the typical error and, therefore, produces inflated positive results. Otherwise, a two-way [treatment \times time] repeated measures ANOVA was conducted with a Bonferroni correction. Effect size for interactions is reported as partial eta squared ($P\eta^2$) and where appropriate, between treatment Hedge's g effect sizes (g) are reported and interpreted as per conventional thresholds (Cohen 1988). Significant effects are displayed with 95% CI where appropriate. Reproducibility of the absolute changes in pH and HCO_3^- in the preliminary trial and the subsequent cycling trials was assessed using intraclass correlation coefficients (ICC). Data are reported as mean \pm standard deviation (SD) and statistical significance was set at $p < 0.05$. Data were analysed using a statistical software package, SPSS (V.22, IBM Inc., Chicago, IL, USA).

Results

Preliminary trials to determine time to peak blood bicarbonate

Time to peak pH ranged between 30 and 100 min in SBC2 (mean 66 ± 22 min; median 60 min; CV 34%) and between 40 and 120 min in SBC3 (mean 76 ± 21 min; median 75 min;

CV 27%; $p = 0.04$). The absolute change from baseline to peak pH was similar in SBC2 and SBC3 (0.08 ± 0.02 vs. 0.09 ± 0.02 ; $p = 0.27$). In the subsequent cycling trials, the reproducibility of the absolute change in pH was fair in SBC2 ($r = 0.50$, $p = 0.09$) and good in SBC3 ($r = 0.60$, $p = 0.06$). Time to peak HCO_3^- was achieved between 30 and 110 min in SBC2 (mean 67 ± 21 min; median 60 min; CV 31%) compared to between 50 and 100 min in SBC3 (mean 77 ± 17 min; median 75 min; CV 22%; $p = 0.20$). The absolute change from baseline was greater ($+1 \text{ mmol l}^{-1}$) in SBC3 compared to SBC2 (7.1 ± 1.2 vs. $6.0 \pm 0.9 \text{ mmol l}^{-1}$; $p = 0.04$; $g = 1.0$). In the subsequent cycling trials, the reproducibility of the change from baseline to peak HCO_3^- was good in SBC2 ($r = 0.70$, $p = 0.04$) and excellent in SBC3 ($r = 0.77$, $p = 0.02$).

Performance

The decline in performance from TT_1 to TT_2 was similar in all treatments (SBC2 8.0 ± 6.8 vs. SBC3 7.0 ± 6.3 vs. PLA 7.3 ± 6.4 s; $p > 0.05$). In TT_1 , SBC3 improved performance compared to PLA by $1.4 \pm 1.5\%$ (400.2 ± 24.1 vs. 405.9 ± 26.0 s; $p = 0.03$; CI = 10.6, 0.8; $g = 0.2$; Fig. 1), which was determined as a very likely benefit in MBI analysis. Meanwhile, SBC2 displayed a likely benefit compared to PLA, improving performance by $0.9 \pm 1.1\%$ (402.3 ± 26.5 s; $p = 0.14$; $g = 0.1$; Fig. 1). A likely benefit was also observed in SBC3 vs. SBC2 ($p = 0.15$; $g = 0.1$). Findings were similar in TT_2 , where SBC3 again displayed the fastest completion times by $1.4 \pm 1.1\%$ compared to PLA (407.2 ± 29.2 vs. 413.2 ± 30.8 s; $p = 0.01$; CI = 10.5, 1.5; $g = 0.2$), which MBI analysis determined this as a very likely effect. Whereas, SBC2 improved performance by $0.7 \pm 1.2\%$ compared to PLA (410.3 ± 30.8 s) and this was determined as a likely benefit ($p = 0.35$; $g = 0.1$; Fig. 2). A possible benefit was determined for SBC3 compared to SBC2 for TT_2 completion time ($p = 0.44$; $g = 0.1$).




Fig. 1 Mean (\pm SD) and individual (horizontal lines) time to complete time trial 1 (TT_1) following SBC2, SBC3, and PLA. Asterisk denotes significantly improved compared to PLA ($p < 0.05$)




Fig. 2 Mean (\pm SD) and individual (horizontal lines) time to complete time trial 2 (TT₂) following SBC and PLA treatments. Asterisk denotes significantly improved compared to PLA ($p < 0.05$)

Mean power in TT₁ was 4.1% greater in SBC3 compared to PLA (247 ± 41 vs. 258 ± 41 ; $p = 0.03$; CI = 1.7, 19.6; $g = 0.3$), showing a very likely improvement. Meanwhile, SBC2 improved mean power by 2.5% compared to PLA, also revealing a very likely benefit (vs. 254 ± 43 W; $p = 0.68$; $g = 0.2$). A likely benefit was determined for SBC3 vs. SBC2 ($p = 0.39$; $g = 0.1$). Mean power in TT₂ was improved by 3.8% in SBC3 compared to PLA (247 ± 46 vs. 237 ± 47 W; $p = 0.005$; CI = 3.0, 15.5; $g = 0.2$), and demonstrated a most likely benefit. Whereas, a likely benefit was determined in SBC2 (vs. 242 ± 47 W; $p = 0.34$; $g = 0.1$). A likely benefit was determined for SBC3 vs. SBC2 ($p = 0.48$; $g = 0.1$).

Blood responses

A [treatment \times time] interaction was observed for HCO_3^- ($P\eta^2 = 0.65$, $p < 0.001$), as HCO_3^- was greater post-supplementation of NaHCO_3 in SBC3 compared to both SBC2 ($p = 0.02$; CI = 0.3, 2.5, $g = 1.5$) and PLA ($p < 0.001$; CI = 6.3, 7.9; $g = 8.4$; Fig. 3). Whereas, SBC2 was greater than PLA only ($p < 0.001$; CI = 4.4, 7.1; $g = 5.7$). Post TT₁, HCO_3^- was greater in both SBC2 and SBC3 compared to PLA (both $p < 0.001$), with no differences between SBC conditions ($p = 0.38$). There was a [treatment] effect for HCO_3^- change during TT₁ ($P\eta^2 = 0.69$, $p < 0.001$), whereby both SBC2 and SBC3 were greater than PLA ($p < 0.005$), with a small effect size between SBC treatments (10.6 ± 3.4 vs. 11.5 ± 3.2 mmol l⁻¹; $p = 0.63$; $g = 0.26$). A significant [treatment \times time] interaction was observed for pH ($P\eta^2 = 0.36$, $p = 0.002$), as pH was greater post-supplementation and post-TT₁ warm-up in SBC3 compared to both SBC2 and PLA (both $p < 0.01$), whilst SBC2 was greater compared to PLA ($p < 0.001$). Blood lactate was greater post-TT₁ in both SBC treatments compared to PLA ($p < 0.005$), with no differences observed otherwise (all $p > 0.05$; Fig. 3).

Both SBC treatments elicited reductions in K^+ , Ca^{2+} , and Cl^- , and increases in Na^+ compared to PLA (Fig. 4). The




Fig. 3 Mean (\pm SD) responses for blood **a** pH, **b** bicarbonate (HCO_3^-), and **c** lactate following NaHCO_3 across time. SBC3 (asterisk) and SBC2 (hash symbol) significantly greater than PLA. SBC3 (dagger symbol) significantly greater than SBC2 ($p < 0.05$). R recovery, PW post warm-up

SID at post-TT₁ warm-up and post-TT₁ was greater in both SBC treatments compared to PLA ($p < 0.005$). However, the SID in SBC3 was greater than SBC2 post- NaHCO_3 supplementation ($p = 0.005$; CI = 0.7, 3.5; $g = 1.0$) and post-TT₁ warm-up ($p = 0.049$; CI = 0.01, 3.6; $g = 0.8$; Fig. 5). During recovery, HCO_3^- was greater following SBC3 compared to PLA at all recovery timepoints ($p < 0.01$), however, only greater at 20 (23.5 ± 1.7 mmol l⁻¹; 21.1 ± 2.7 mmol l⁻¹; $p = 0.04$; CI = 0.2, 4.7; $g = 1.0$) and 40 min (27.7 ± 0.8 vs. 25.7 ± 1.3 mmol l⁻¹; $p = 0.006$; CI = 0.6, 3.3; $g = 1.8$) compared to SBC2. Similarly, SBC2 was greater than PLA at all recovery timepoints (all $p < 0.01$). The absolute change in HCO_3^- from post TT₁ to 40 min recovery was significantly greater compared to PLA (5.0 ± 1.5 mmol l⁻¹) in both SBC2




Fig. 4 Mean (\pm SD) potassium (a), sodium (b), calcium (c), and chloride (d) responses over time following SBC treatments. SBC3 (asterisk) and SBC2 (hash symbol) significantly different compared

to PLA. SBC3 (dagger symbol) significantly different compared to SBC2 ($p < 0.05$). *R* recovery, *PW* post warm-up




Fig. 5 Mean (\pm SD) strong ion difference (SID) responses over time following SBC treatments. SBC3 (asterisk) and SBC2 (hash symbol) significantly greater than PLA. SBC3 (dagger symbol) significantly greater than SBC2 ($p < 0.05$). *R* recovery, *PW* post warm-up

($10.1 \pm 1.4 \text{ mmol l}^{-1}$, $p < 0.001$; $CI = 3.2, 7.1$; $g = 3.4$) and SBC3 ($11.1 \pm 2.5 \text{ mmol l}^{-1}$, $p < 0.001$; $CI = 4.2, 8.1$; $g = 2.8$), with a small effect size between SBC treatments ($p = 0.45$; $g = 0.3$). The absolute change in the SID from post-TT₁ to 40 min recovery was only significantly greater for SBC3 compared to PLA ($p = 0.05$; $CI = 0.01, 9.0$; $g = 1.2$).

Post warm-up in TT₂, HCO_3^- was greater in both SBC2 and SBC3 compared to PLA ($p < 0.001$); however, SBC3 was greater than any treatment ($p < 0.001$). There was a [treatment] effect for HCO_3^- change during TT₂ ($P\eta^2 = 0.71$, $p < 0.001$), whereby both SBC treatments were greater than

PLA ($p < 0.01$); however, SBC3 was greater compared to SBC2 (10.7 ± 2.9 vs. $9.2 \pm 2.7 \text{ mmol l}^{-1}$; $p = 0.02$; $CI = 0.3, 2.6$; $g = 0.5$; Fig. 3). Post-TT₂ warm-up, and post-TT₂, pH in both SBC2 and SBC3 were greater than PLA ($p < 0.001$), although no differences between SBC treatments were observed ($p > 0.05$). Blood lactate was greater post-TT₂ in both SBC2 (18.0 ± 4.2 vs. $14.4 \pm 3.3 \text{ mmol l}^{-1}$; $p = 0.05$; $CI = -0.01, 7.2$; $g = 0.9$) and SBC3 ($18.6 \pm 3.5 \text{ mmol l}^{-1}$; $p = 0.009$; $CI = 1.1, 7.2$; $g = 0.9$) compared to PLA, with no difference between SBC treatments ($p = 0.424$; $g = 1.2$). Post-TT₂ warm-up, the SID was greater for both SBC2 and SBC3 compared to PLA ($p < 0.001$); however, no difference was observed between SBC treatments (SBC2 35 ± 3 vs. SBC3 $37 \pm 3 \text{ meq/L}$; $p = 0.20$; $g = 0.6$).

Perceptual responses

During TT₁, NaHCO_3 did not affect RPE_O ($P\eta^2 = 0.24$, $p = 0.07$) or RPE_L ($P\eta^2 = 0.10$, $p = 0.38$), HR ($P\eta^2 = 0.07$, $p = 0.63$), or SPO_2 ($P\eta^2 = 0.18$, $p = 0.16$). Similarly, in TT₂, no changes in RPE_O ($P\eta^2 = 0.17$, $p = 0.12$), RPE_L ($P\eta^2 = 0.11$, $p = 0.35$), HR ($P\eta^2 = 0.07$, $p = 0.78$), or SPO_2 ($P\eta^2 = 0.02$, $p = 0.83$) were observed.

Gastrointestinal (GI) discomfort

In total, 6/10 (60%) participants suffered from GI discomfort in SBC2, whereas 9/10 (90%) suffered from GI discomfort




Fig. 6 Individual aggregated score and severity of gastrointestinal (GI) responses following SBC treatments. Asterisk denotes SBC3 significantly greater than SBC2 ($p < 0.05$)

in SBC3. The most common GI discomfort symptom was belching (2/10) in SBC2, whilst in SBC3, diarrhoea, bowel urgency, and feeling of vomiting were most common (5/10). Both the aggregated GI discomfort and the severity of the most severe GI discomfort symptom suffered were greater in SBC3 compared to SBC2 (both $p < 0.05$, Fig. 6). On 3/30 (10%) occasions, the supplement was correctly identified by the participant.

Discussion

This study investigated the effects of NaHCO_3 ingestion on post-exercise acid–base balance recovery and repeated 4 km TT performance in moderate acute hypoxic conditions. Both SBC2 and SBC3 improved TT_1 and TT_2 performance compared to PLA, displaying ‘likely’ and ‘very likely’ beneficial effects in magnitude-based inferences analysis, respectively. The current study findings suggest that this occurred due to the greater magnitude of acid–base balance recovery between TT_1 and TT_2 or that the initial acid–base balance stress during TT_1 was blunted by NaHCO_3 ingestion. A greater magnitude of performance improvement was observed in SBC3, however, showing ‘likely’ and ‘possibly’ beneficial effects in TT_1 and TT_2 compared to SBC2, respectively. As such, SBC3 is the most optimal to improve repeated efforts of high-intensity exercise in acute moderate hypoxic conditions. One individual did display an ergolytic effect following SBC3, however, despite still gaining ergogenic effects from SBC2 vs. PLA, possibly caused by the onset of severe GI discomfort (Saunders et al. 2014). Individuals who display similar responses may, therefore, wish to select SBC2.

The current study findings suggest the enhanced exercise performance in both bouts of exercise following NaHCO_3 ingestion occurred due to the combination of the greater alkalotic state of the acid–base balance prior to TT_1 and the greater magnitude of acid–base balance recovery prior

to TT_2 . The change in HCO_3^- during TT_1 was increased following NaHCO_3 (SBC2 + 29%, SBC3 + 34%), whilst post TT_1 , blood lactate was also greater compared to PLA (SBC2 + 21%, SBC3 + 23%). These changes suggest an increased H^+ buffering from intramuscular to extracellular compartments, which may lead to an increased anaerobic energy provision and glycogen utilization, as intramuscular pH is better protected (Lopes-Silva et al. 2018; Percival et al. 2015). It is argued that these indirect biomarkers of upregulated glycolytic flux in the present study instead show a reduction in lactate by inactive tissue however, as the samples in the present study do provide data on use/uptake (Granier et al. 1996). Likewise, it is strongly argued that acidosis does not inhibit contractile machinery and that alkalization may cause instability of the buffering systems, and thus, the benefit to the ATP-generating process is either negligible, or actually harmful (Korzeniewski and Zoladz 2002; Sahlin et al. 1998). A recent study by Lopes-Silva et al. (2018), nonetheless, reported similar post-exercise HCO_3^- and lactate responses to the current study following NaHCO_3 ingestion, yet also reported both a 34% greater estimated glycolytic energy contribution to exercise and an improved performance. It is alternately conceivable that the increases in HCO_3^- and CO_2 following NaHCO_3 ingestion are part of a causal sequence of the primary mechanism of performance, by accelerating VO_2 kinetics at the onset of high-power outputs (Zoladz et al. 2005). Unfortunately, VO_2 was not measured during the TTs in the study, so this cannot be concluded and thus warrants further research. The current study findings, nonetheless, further support past research (Fitts 2016) that acid–base balance disturbances are an important determinant of fatigue during high-intensity exercise.

The current study adds that NaHCO_3 ingestion increased the SID prior to each TT bout, which may explain the improved performance. The SID has been suggested to be an important component of action potentials by increasing muscle excitability (Allen et al. 2008). Notable changes

NaHCO₃ ingestion elicits include a lowering of extracellular K⁺ and Cl⁻, whilst increasing Na⁺, all of which have collectively been identified to be important for muscle contraction during high-intensity exercise previously (Cairns and Lindinger 2008). Here, the current study findings expand upon those of Sostaric et al. (2006) that reported an improvement in finger flexion to exhaustion combined with an increased SID following NaHCO₃ ingestion, by showing a similar effect but during dynamic whole-body exercise. These findings also provide an alternative mechanism to the pH and HCO₃⁻-mediated mechanisms often criticised in the literature (Westerblad 2016). It is worth noting that the highlighted changes in the SID may also have further reaching benefits to health-related outcomes. Mild metabolic acidosis has been linked with multiple health-related outcomes including protein metabolism, by decreasing synthesis through increases of non-enzymatic proteolysis (Wiederkehr and Krapf 2001). Likewise, regular consumption of alkaline rich supplements has been shown to improve mineral balance and rate of bone formation (Sebastian et al. 1994). Therefore, more research is required investigating the use of chronic NaHCO₃ ingestion in sporting populations that could also be prone to such issues, particularly endurance runners and adolescent athletes (Tenforde et al. 2017; Scofield and Hecht 2012).

Both pH and HCO₃⁻ following SBC treatments were increased prior to TT₂, such that the absolute change in HCO₃⁻ from TT₁ to 40 min recovery was over twofold greater compared to PLA (SBC2 + 51%, SBC3 + 55%), whilst the change in the SID was also more superior (SBC2 + 29%, SBC3 + 31%). This suggests a greater amount of H⁺ buffering occurred during this time, which subsequently facilitated a more substantial recovery of acid–base balance compared to PLA, in support of previous research (Pruscino et al. 2008; Gough et al. 2017a). Alternatively, the improvement in TT₂ may have been due to the stress on acid–base balance being blunted by NaHCO₃ during TT₁, as pH, HCO₃⁻, and the SID were all greater immediately post TT₁ compared to PLA. Based on data from Gough et al. (2017a), however, it is more likely that the enhancement of the acid–base balance state between TT₁ and TT₂ explains the improvement in the current study, as the authors reported NaHCO₃ ingestion improved a subsequent exercise even when supplementation was after an initial bout of exercise. Moreover, SBC3 elicited a significantly greater magnitude of acid–base balance recovery prior to TT₂, and change in HCO₃⁻ during TT₂ compared to SBC2. Correspondingly, SBC3 improved performance within half of the sample compared to SBC2, compared to only two displaying greater improvements in SBC2 vs. SBC3 when using the 3.1 s TE of the test. Combined, these greater increases in acid–base balance blood analytes may explain the greater magnitude of improvement in TT₂ produced by SBC3.

At 40 min recovery pH, HCO₃⁻ and the SID were still rising following NaHCO₃ ingestion, such that pH at 40 min recovery was 7.49 ± 0.02 in SBC3, which was identical to the increase prior to TT₁ following the same dose (7.49 ± 0.01). This is in agreement with the previous research in normoxia showing similar increases at the end of a recovery period typically seen with pre-exercise NaHCO₃ ingestion (Callaghan et al. 2017; Pruscino et al. 2008). Equally, in the current study, and others (Callaghan et al. 2017; Pruscino et al. 2008), acid–base balance status was still significantly rising to a more alkalotic state. It is plausible to suggest, therefore, if a longer period of recovery was employed, a more pronounced performance effect may have been observed compared to PLA. Moreover, this also suggests that re-dosing NaHCO₃ following an initial fatiguing bout is not required, as acid–base balance increased well above baseline, despite no re-dosing of NaHCO₃. These findings may be of importance to individuals who suffer from GI discomfort, as no instances were reported during recovery in the current study. Future research should, therefore, investigate the performance responses on repeated exercise following NaHCO₃ ingestion with a longer period of recovery.

One participant presented an ergolytic effect in both TT₁ and TT₂ after ingestion of SBC3. This was likely due to the occurrence of severe GI discomfort (diarrhoea = 10; aggregate score = 63), as this participant still improved their performance in SBC2. These findings support the previous research, whereby ergolytic effects were observed in participants who suffered from severe GI discomfort following 0.3 g kg⁻¹ BM NaHCO₃ (Saunders et al. 2014; Froio de Araujo Dias et al. 2015). It is therefore important to monitor the GI discomfort responses following NaHCO₃ ingestion on an individual basis, as those who display severe symptoms following SBC3 may instead benefit from ingesting SBC2. The use of this smaller dose, however, will be dependent on an improvement in performance still being observed compared to a placebo.

Conclusion

This study investigated the effects of NaHCO₃ ingestion on repeated 4 km TT performance and acid–base balance recovery in acute moderate hypoxic conditions. Both amounts of NaHCO₃ employed in this study ensured recovery of acid–base balance back to baseline or above within 20–40 min, whereas this was not achieved for PLA. For the first time, blunting of the acid–base balance stress during the initial bout of exercise, or a greater magnitude of acid–base balance recovery, has translated into improved subsequent high-intensity exercise performance following

NaHCO₃ ingestion in acute hypoxic conditions. The performance improvement was greater in SBC3, which is likely due to the greater alkalotic status of acid–base balance both prior to TT₁, and during recovery compared to SBC2 and PLA. The onset of GI discomfort was an issue with SBC3, however, and one participant displayed an ergolytic effect on performance following this dose. Individuals should, therefore, employ SBC3 to improve performance in acute hypoxic conditions, only if severe GI discomfort does not occur.

Author contributions Conception or design of the work: the original conception of the work was conducted by LG. Data collection: data were collected by LG. Data analysis and interpretation: completed by LG. Drafting the article: completed by LG. Critical revision of the article: all authors (DB, SD, SAS, and LRM) on the manuscript contributed. Final approval of the version to be published: this was sought from all authors (DB, SD, SAS, and LRM).

Funding No sources of funding were used to conduct this study or prepare this manuscript.

Compliance with ethical standards

Conflict of interest LA Gough, SK Deb, D Brown, SA Sparks, and LR McNaughton have no conflicts of interest that are directly relevant to the content of this article.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Allen DG, Lamb GD, Westerblad H (2008) Impaired calcium release during fatigue. *J Appl Physiol* 104(1):296–305
- Al-Nawaiseh AM, Pritchett RC, Bishop PA (2016) Enhancing short-term recovery after high-intensity anaerobic exercise. *J Strength Cond Res* 30(2):320–325
- Baguet A, Harmen R, Andries P, Inge E, Stefanie C, Eric A, Wim D (2009) Carnosine loading and washout in human skeletal muscles. *J Appl Physiol* 106(3):837–842
- Barnett A (2006) Using recovery modalities between training sessions in elite athletes does it help? *Sports Med* 36(9):781–786
- Batterham AM, Hopkins WG (2006) Making meaningful inferences about magnitudes. *Int J Sports Physiol Perform* 1(1):50–57
- Bonaventura JM, Sharpe K, Knight E, Fuller KL, Tanner RK, Gore CJ (2015) Reliability and accuracy of six hand-held blood lactate analysers. *J Sports Sci Med* 14(1):203
- Cairns SP (2006) Lactic acid and exercise performance: culprit or friend? *Sports Med* 36(4):279–291
- Cairns SP, Lindinger MI (2008) Do multiple ionic interactions contribute to skeletal muscle fatigue? *J Physiol* 586(17):4039–4054
- Callahan MJ, Parr EB, Hawley JA, Burke LM (2017) Single and combined effects of beetroot crystals and sodium bicarbonate on 4-km cycling time trial performance. *Int J Sport Nutr Exerc Metab* 27(3):271–278
- Cohen J (1988) *Statistical power analysis for the behavioural sciences*. Lawrence Erlbaum Associates, Hillsdale
- De Pauw K, Roelands B, Cheung SS, De Geus B, Rietjens G, Meeusen R (2013) Guidelines to classify subject groups in sport-science research. *Int J Sports Physiol Perform* 8(2):111–122
- Durkalec-Michalski K, Zawieja EE, Podgórski T, Łoniewski I, Zawieja BE, Warzybok M, Jeszka J (2018) The effect of chronic progressive-dose sodium bicarbonate ingestion on CrossFit-like performance: a double-blind, randomized cross-over trial. *PLoS One* 13(5):e0197480
- Egger F, Meyer T, Such U, Hecksteden A (2014) Effects of sodium bicarbonate on high-intensity endurance performance in cyclists: a double-blind, randomized cross-over trial. *PLoS One* 9(12):e114729
- Fitts RH (2016) The role of acidosis in fatigue. *Med Sci Sports Exerc* 48(11):2335–2338
- Froio de Araujo Dias G, Da Eira Silva V, De Salles Painelli V, Sale C, Giannini Artioli G, Gualano B, Saunders B (2015) Consistencies in responses to sodium bicarbonate supplementation: a randomised, repeated measures, counterbalanced and double-blind study. *PLoS One* 10(11):e0143086
- Gough LA, Rimmer S, Osler CJ, Higgins MF (2017a) Ingestion of sodium bicarbonate (NaHCO₃) following a fatiguing bout of exercise accelerates postexercise acid-base balance recovery and improves subsequent high-intensity cycling time to exhaustion. *Int J Sport Nutr Exerc Metab* 27(5):429–438
- Gough LA, Deb SK, Sparks SA, McNaughton LR (2017b) The reproducibility of blood acid base responses in male collegiate athletes following individualised doses of sodium bicarbonate: a randomised controlled crossover study. *Sports Med* 47(10):2117–2127
- Gough LA, Sanjoy K, Deb S, Sparks SA, McNaughton LR (2017c) Sodium bicarbonate improves 4 km time trial cycling performance when individualised to time to peak blood bicarbonate in trained male cyclists. *J Sports Sci* 36(15):1705–1712
- Granier P, Dubouchaud H, Mercier B, Mercier J, Ahmaidi S, Préfaut C (1996) Lactate uptake by forearm skeletal muscles during repeated periods of short-term intense leg exercise in humans. *Eur J Appl Physiol Occup Physiol* 72(3):209–214
- Hollidge-Horvat MG, Parolin ML, Wong D, Jones NL, Heigenhauser GJ (1999) Effect of induced metabolic acidosis on human skeletal muscle metabolism during exercise. *Am J Physiol* 277(4):647–658
- Korzeniewski B, Zoladz JA (2002) Influence of rapid changes in cytosolic pH on oxidative phosphorylation in skeletal muscle: theoretical studies. *Biochem J* 365(1):249–258
- Lloyd P (2004) Strong ion calculator—a practical bedside application of modern quantitative acid-base physiology. *Crit Care Resusc* 6(4):285–294
- Lopes-Silva JP, Da Silva Santos JF, Artioli GG, Loturco I, Abbiss C, Franchini E (2018) Sodium bicarbonate ingestion increases glycolytic contribution and improves performance during simulated taekwondo combat. *Eur J Sport Sci* 18(3):1–10
- Monedero J, Donne (2000) Effect of recovery interventions on lactate removal and subsequent performance. *Int J Sports Med* 21(8):593–597
- Percival ME, Martin BJ, Gillen JB, Skelly LE, Macinnis MJ, Green AE, Tarnopolsky MA, Gibala MJ (2015) Sodium bicarbonate ingestion augments the increase in PGC-1 α mRNA expression during recovery from intense interval exercise in human skeletal muscle. *J Appl Physiol* 119(11):1303–1312
- Pierce EF, Eastman NW, Hammer WH, Lynn TD (1992) Effect of induced alkalosis on swimming time trials. *J Sports Sci* 10(3):255–259
- Pruscino CL, Ross MLR, Gregory JR, Savage B, Flanagan TR (2008) Effects of sodium bicarbonate, caffeine, and their combination

- on repeated 200-m freestyle performance. *Int J Sport Nutr Exerc Metab* 18(2):116–130
- Pyne DB, Boston T, Martin DT, Logan A (2000) Evaluation of the Lactate Pro blood lactate analyser. *Eur J Appl Physiol* 82(1–2):112–116
- Robergs R, Hutchinson K, Hendee S, Madden S, Siegler J (2005) Influence of pre-exercise acidosis and alkalosis on the kinetics of acid-base recovery following intense exercise. *Int J Sport Nutr Exerc Metab* 15(1):59–74
- Sahlin K, Tonkonogi M, Söderlund K (1998) Energy supply and muscle fatigue in humans. *Acta Physiol Scand* 162(3):261–266
- Saunders B, Sale C, Harris RC, Sunderland C (2014) Sodium bicarbonate and high-intensity-cycling capacity: variability in responses. *Int J Sports Physiol Perform* 9(4):627–632
- Scofield KL, Hecht S (2012) Bone health in endurance athletes: runners, cyclists, and swimmers. *Curr Sports Med Rep* 11(6):328–334
- Sebastian A, Harris ST, Ottaway JH, Todd KM, Morris Jr RC (1994) Improved mineral balance and skeletal metabolism in postmenopausal women treated with potassium bicarbonate. *N Engl J Med* 330(25):1776–1781
- Sostaric SM, Skinner SL, Brown MJ, Sangkabutra T, Medved I, Medley T, McKenna MJ (2006) Alkalosis increases muscle K^+ release, but lowers plasma $[K^+]$ and delays fatigue during dynamic forearm exercise. *J Physiol* 570(1):185–205
- Swinton PA, Hemingway BS, Saunders B, Gualano B, Dolan E (2018) A statistical framework to interpret individual response to intervention: paving the way for personalized nutrition and exercise prescription. *Front Nutr* 5:41
- Tenforde AS, Carlson JL, Chang A, Sainani KL, Shultz R, Kim JH, Fredericson M (2017) Association of the female athlete triad risk assessment stratification to the development of bone stress injuries in collegiate athletes. *Am J Sports Med* 45(2):302–310
- Ward R, Bridge CA, McNaughton LR, Sparks SA (2016) The effect of acute taurine ingestion on 4-km time trial performance in trained cyclists. *Amino Acids* 48(11):2581–2587
- Westerblad H (2016) Acidosis is not a significant cause of skeletal muscle fatigue. *Med Sci Sports Exerc* 48(11):2339–2342
- Wiederkehr M, Krapf R (2001) Metabolic and endocrine effects of metabolic acidosis in humans. *Swiss Med Wkly* 131(9–10):127–132
- Zabala M, Requena B, Sánchez-muñoz C, González-Badillo JJ, García I, Ööpik V, Pääsuke M (2008) Effects of sodium bicarbonate ingestion on performance and perceptual responses in a laboratory-simulated bmx cycling qualification series. *J Strength Cond Res* 22(5):1645–1653
- Zabala M, Peinado AB, Calderón FJ, Sampedro J, Castillo MJ, Benito PJ (2011) Bicarbonate ingestion has no ergogenic effect on consecutive all out sprint tests in BMX elite cyclists. *Eur J Appl Physiol* 111(12):3127–3134
- Zoladz JA, Szkutnik Z, Duda K, Majerczak J, Korzeniewski B (2005) Preexercise metabolic alkalosis induced via bicarbonate ingestion accelerates VO_2 kinetics at the onset of a high-power-output exercise in humans. *J Appl Physiol* 98(3):895–904