

International Journal of Horticultural Science 2014, 20 (3–4): 7–13.

Agroinform Publishing House, Budapest, Printed in Hungary

ISSN 1585-0404

The uses of wolfberry (*Lycium barbarum* L.) as a fruit in an international breadth of view

Sóspataki, R., Józsa, M. & Simon, G.

¹Corvinus University of Budapest Faculty of Horticulture, Department of Pomology

Summary: *Lycium barbarum* is known exclusively as an ornamental plant in Hungary, and is planted so, as popular belief deem it a toxic plant. The plant's fruit receives great respect in countries abroad, thus, due to its favourable content values the *Lycium* has achieved the title of 'biological gold mine'. The last couple of years has seen the import of *Lycium* shrubs and its corresponding products, dried goods or in the form of various processed products which have been marketed and sold at extreme prices (under name of Goji, Wolfberry, Lifeberry). Our goal was to examine the similarities of the content values of wolfberry found in Hungary and those cultivated abroad. Along with Miklós Józsa the domestic *Lycium* population was surveyed based on foreign examples, between the years 2009 and 2011. Those defined sweet and large-fruited were selected for further investigations to be set into a clone repository. This clone repository – which contains 67 different clones from a number of regions of Hungary – was established in the nursery of Dr. Miklós Józsa, located in the city of Szombathely. The phenological and morphological characteristics and the fruit ripening and quality indicators of plants in the clone repository were investigated. Six 'best' clones – selected based on flavour, disease-resistance and vegetative characteristics (plant size, fruit set, yield and fruit size) were analysed based on their content values. The control plant was a cultivar imported by a delivery service, found also in growing. In addition to the results of the selection, the results of the content values of the six selected 'best' clones (total soluble sugar content, glucose-fructose ratio, carotene content, FRAP value) is documented in this paper. Further, based on the resulting information the possibilities of the fruit's utilization are suggested. Significant differences were measured in the vegetative characteristics of the clones (plant size, fruit set, yield and fruit size) and of those of its inner content values. The resulting properties are considerable in regards of the plant's introduction into cultivation and also utilization. Based on growing characteristics, more favourable clones were selected than the foreign varieties already in growing. Those clones selected by us have had similar or better content values than foreign breeds. The investigation of these content values is still in progress.

Keywords: *Lycium barbarum*, Goji berry, Wolfberry, Antioxidant, HPLC, FRAP

Introduction

Wolfberry (*Lycium barbarum*) is related to the shrub family of solanaceous (*Solanaceae*). The vegetative organs of the plant (belonging to this family) are rich in alkaloid and many of them are ornamental plants (Vancsura, 1992). The *Lycium* genus is made up of approximately 80-90 species (Krüssmann, 1977). There are 15 native species in Europe, most can be found in the Mediterranean region (Vancsura, 1992). 7 *Lycium* species, 3 varieties and 35 various cultivated plant type can be found in China. The most respected type is the *Lycium barbarum* thanks to its curative influence (Young et. al., 2006).

It is spread widely: South and Central Europe, North Africa, West Asia, North America. It is an adventive plant to most regions, originating most likely from Central China. The occurrence in Hungary is frequent, however the plant is absent in some regions (ie: northern part of the Great Plain) (Bartha, 1999).

Wolfberries (syn. name Goji) are able to grow in all soil types and exposure, and tolerate the heavily saline soil as well (Schmidt, 1996).

Lycium barbarum most probably got its name from the province of Lycia, located in Asia Minor. Chinese call the

plant itself 'gouqi', and the fruit 'gouqizi', 'zi' meaning small fruit. This is where the name 'Goji' has evolved from (Young et. al., 2006).

Wolfberries are long-living and fast growing 2-3 m high deciduous plants (Schmidt & Tóth, 2006). The plant can be characterized by shoots and branches, with significant number of thin branches emerging from the rootstock (Bartha, 1999). Young shoots are upright, and will take curved shapes later. (Vancsura, 1992). Wolfberry leaves can take many forms, may have lanceolate spear point or elliptic, mid-widening shape (Bartha, 1997). The flowers grow segregated in the leaf axils of the long shoot, and in bundles of two to five of the short shoot (Bartha, 1999). The flowering often starts in May and may continue until September or even October, which makes the wolfberry a valued apiculture plant (Halmágyi, 1991). Fruits are in clusters of one to four, round to egg-shaped, green colour at first (Bartha, 1997). Ripe berries turn bright red, are 1.5-2 cm long, and contain small sized seeds numbering 10-15 (Vancsura, 1992). The seeds are 2.2-2.8 mm long, wide oval shaped, are laterally compressed from its sides, have fine degree of spherical surface and are orange-red coloured (Bartha, 1997).

There is no information on consumption of crops in Hungary. This can be justified as authoritative publications

consider the fruits toxic, and thus do not recommend consumption under any circumstances (Pap, 2010).

Wolfberry can be easily vegetative propagated by green, semi-hardwood and hardwood cuttings. Cuttings are able to get good root even among inappropriate conditions. Generative propagation is also known by seeding from September through October (Schmidt, 1996).

Lycium genus is infected mainly by the powdery mildew (*Arthrocladiella mougeotii* – previously *Erysiphe mougeotii*), which is the only known species of the *Arthrocladiella* genus. Infected parts of the plant are slightly deformed, in case of serious infections the plant ornamental values may reduce (www.cabi.org, 2011). Its most common disease is the lineate patterns of the leaves and the yellow mosaic disease, however infected shrubs did not initiate remarkable growth reduction or weakening (Salamon, 2002).

Chinese have been consuming the wolfberry for 5000 years. Wolfberry has been cultivated in northern China and Tibet for a long time, nowadays primarily in the regions of Ningxia, Hei Bei, Xin Jiang and Inner Mongolia.

The crops are marketed as dried fruit, but numerous other product types are also produced. Wolfberry fruits have several wide-ranging, positive effects. More than 15% of its weight is composed of proteins, wolfberry contains more than 21 essential minerals, 18 amino acids and a considerable amount of chromium. Its vitamin content (vitamins B₁, B₃, C) is excessively (*Table 1.*). It has a strong supporting effect for the immune system by its outstanding high antioxidant content (Young et. al., 2006), (*Table 2.*).

Zeaxanthin and its esters are major contributors to the carotenoids in *L. barbarum* (Li et. al., 1998), (Weller & Breithaupt, 2003). The content of zeaxanthin esters in ripening goji fruits can reach as high as >77.5% of total carotenoids (Li et. al., 1998), (Peng et. al., 1988), (*Table 2.*). Zeaxanthin and other carotenoids are also potent antioxidants, which contribute to the health effects of goji berry against oxidative stress-mediated diseases.

Table 1. Nutrient data for dried goji berries compared with dried apricots (Source: Paul M. Gross, 2006. Tables Ciquel (AFFSA))

Nutrients	Dried Goji berries	Dried Apricot
	Value: 100 g	
Energy (kcal)	370	226
Protein (g)	11.7	2.5
Sugar (g)	67.7	46.3
Total lipid (Fat) (g)	8.2	0.28
Fiber (g)	10	13
Vitamin C (mg)	29	3.8
Calcium (mg)	112	55
Potassium (mg)	1132	1520
Beta-carotene (microgram)	7400	2280

Table 2. Contents of zeaxanthin dipalmitate and total carotenoid in Fructus Lycii from different species of the genus *Lycium* in China (Source: Yong et. al., 2005.)

Sample no.	Species	Contents of zeaxanthin dipalmitate (%)	Contents of total carotenoid (%)	Contents of zeaxanthin dipalmitate in total carotenoids (%)
1	<i>L. dasystemum</i> Pojark	0.146	0.269	54.28
2	<i>L. barbarum</i> L.	0.214	0.386	55.44
3	<i>L. chinense</i> Mill	0.219	0.444	49.32
4	<i>L. truncatum</i> Y.C. Wang	0.2	0.383	52.22
5	<i>L. cylindricum</i> Kuang et A.M. Lu	0.174	0.345	50.43
6	<i>L. ruthenicum</i> Murr	0.031	0.084	36.9
7	<i>L. chinense</i> Mill. var. <i>potaninii</i> A.M. Lu	0.149	0.306	48.69
8	<i>L. dasystemum</i> Pojark var. <i>rubricaulium</i> A.M. Lu	0.262	0.473	55.39
9	<i>L. barbarum</i> L. var. <i>auranticarpum</i> K.F. Ching	0.011	0.035	31.43

Table 3. Oxygen radical absorbance capacity (ORAC) of berries and other fruits (adapted from USDA 2010)

Fruit	ORAC (μmol TE/100 g)
Acai berry	102.700
Black currant	7.957
Black raspberry	19.220
Blackberry	5.905
Blueberry	4.669
Cranberry	9.090
Elderberry	14.697
Goji	3.290
Gooseberry	1.700
Grape	1.837
Pomegranate	4.479
Red currant	3.387
Red raspberry	5.065
Strawberry	4.302
Apple	2.500
Banana	800
Orange	2.000

Material and methods

Gathering of *Lycium* genotypes in Hungary started in the summer of 2009. It was found 91-94% of *Lycium* population has no valuable crop (fruits). During gathering sweet, thornless, less susceptible or resistant to disease examples were given priority. Various flowering, ripening times and different berry shapes and sizes were found from the beginning of the summer to the end of October. Most of the

tasty *Lycium* were detected in the triangle of Székesfehérvár – Dunaújváros – Zalaegerszeg.

Hardwood cuttings were cut between 10 and 20 October 2009. Cuttings were set into seed trays with 66-144 cells. Porous, not coarse propagation media with efficient drainage property was applied. The soil mixture was the following: 2/3 part of peat, 1/3 part of perlite including 12-14 months old Osmocote. The sprout cuttings were grown in a plastic tunnel under appropriate conditions. The clones were set between 10 and 20 of October 2009, and three plants per clone were planted between 20 and 27 of April 2010 into the clone repository.

Overall, 67 clones were planted in the clone repository, from which 4 were candidate varieties originating from various European nurseries. These clones were set into the clone repository for comparison.

Shrubs of the *Lycium* clones were examined in terms of various characteristics (shrub height, width and habitat). Clone 29 (as control) was compared with the examination results of the six 'best' clones. Characteristics (shrub height and diameter) of two year-old plants on 3 shrubs from each clones were measured by tape measure in October 2011. Habits of the shrubs were characterised based on the shrub height and width ratio.

Flower set and crop load were estimated between 2011.04.30 and 2011.08.10 on the second years old plants. The blooming estimation categories: No flower = 0; Only few flowers = 0,5; Medium flower set = 1; Good flower set = 2; Very good flower set = 3

Crop load categories: No berries = 0; Only few berries = 0,5; Light crop load = 1; Medium yielding = 2; Good yielding = 3.

The fruit size parameters were measured in different time in the ripening period according to ripening status of berries.

3x5 (15) fruits were used from each genotypes as a sample for measuring of the typical characteristics of berries. The fruit weight of *Lycium* berries were measured with a digital balance. The length of the crops and his width a tenth accuracy I measured it with a digital calliper. The formal index of fruits was calculated from fruit length and the width.

The water-soluble sugar content (Brix%), was measured with ATAGO PR 101-digital refractometer from smashed, and homogenised mash of 150 pieces frozen *Lycium* fruits.

HPLC analysis for nutrient content of the *Lycium* fruits:

Sample preparation. After thawing, fruit samples were placed in 1.5 ml Eppendorf tubes and centrifuged (Hettich 23R) for 5 min at 15.000 rpm. One ml supernatant was then pipetted off. D-glucose, D-fructose and L-malic acid, analytical standards from Sigma-Aldrich Chemical Co. (3050 Spruce Street, St. Louis, MO 63103, USA), were used as HPLC standards. The standards were first dissolved in double distilled water at a concentration of 0.01 g 50 ml⁻¹, and a 1:50 dilution of these solutions was used for the HPLC analysis. Each measurement was repeated three times.

The HPLC equipment. The HPLC equipment (Waters Corporation, 34 Maple Street, Milford, MA 01757, USA) consisted of the following hardware: 2487 Dual Wavelength

Absorbance Detector (for the determination of organic acids), 2414 Refractive Index Detector (for the determination of sugars), 1525 Binary HPLC Pump, Colonna thermostat, 717 plus automatic injector. The equipment was controlled by the EmPower™ 2 software programme.

Determination of sugars. The sugars were separated on a Sugar-Pak™ column placed in a thermostat at 90°C. The mobile phase was a 0.0001 M aqueous solution of Ca-EDTA [304695-78-1]. The flow rate was 0.5 ml/min, leading to a pressure of 450 ± 10 psi on the column. Detection was continued for 30 min. The injected sample quantity was 20 ml. The retention time was 10.8 min for glucose. 11.77 min for fructose.

The β-carotene: The definition of carotene content was based on the method drawn up by Konzerv Kutató. (KPKI method collection. 1990,2/4. method.)

The antioxidant capacity his definition (FRAP), Benzie and Stran happened with a modified method.

Results and discussion

Collected clones from the different areas of Hungary were planted in a trial orchard under uniform ecological (soil, moisture, precipitation, temperature) and growing conditions, and were compared objectively.

From among the 67 clones which can be found in the clone collection based on different viewpoints, selected the six best clones. Selected clones were compared to the control which is a cultivar of a foreign country delivery service.

Flowering and yielding were observed from spring to late autumn. From 67 collected genotypes 6 were selected as the best by the bush features, crop quantity, fruit size and it was not based on their flavour.

It was considered, that on the different clones when can be found most crops simultaneously, or which one bringing yielding towed. It was also considered the proportional fruit quantity to the bush size. It was also important in terms of the crop picking how soft their berry skin is, or, that how long they are storable without storage disorders. It was selected clones with no tasteless and without bitter aftertaste.

Vegetative characteristics, bush sizes

The measurement of vegetative parameters started in a two-year-old *Lycium* trial orchard in 2011, October. Height of bushes and the bush diameter were measured by tape measure. The ratio of bush height and diameter was calculated, giving information about the bush shape. By the statistical analysis of the data it was found significant differences between the height of bush and bush diameter of selected clones (*Table 4.*)

It was found that Hungarian selected genotypes (excepted clone No. 12) had less bush size parameters than the commercial cultivar (clone No. 29.) The shrub (plant) size has major effect on the spacing and growing conditions of the *Lycium* orchard. Of course the less plant size is not

evaluated as a negative vegetative character, because by the compact growth of shrubs can be planted more intensive and more easily cultivated and harvested orchard. It is a question whether the moderate growth vigour interferes in the fertility.

Table 4. Bush size parameters and the calculated ratio of bush height and diameter in a two-year-old *Lycium* orchard, in 2011.

Clones' number	Height of the bush	Bush diameter	The ratio of bush height and diameter
55.	73,33	143,67	0,51
47.	85,33	151,00	0,57
21.	86,67	154,00	0,56
30.	91,67	137,33	0,67
67.	112,67	168,67	0,67
29.	137,33	237,33	0,58
12.	141,00	200,00	0,71

Generative parameters, flower set and crop load

Table 5. contains blooming and yield estimation data, which were collected in different times during the growing season. The clone No. 29. – used as the control in the field trial – showed very low blooming and yielding characteristic in the two-year-old plantation. Genotypes (clones) selected by us in Hungary showed more favourable values in both parameters.

Table 5. Flower set and crop load data based on estimation method in two-year-old *Lycium* orchard in 2011.

Clones' number	Flower set	Crop load/ Yielding
12.	2	2
21.	2	3
29.	0,5	0,5
30.	2	2
47.	0,5	2
55.	1	2
67.	3	2
Comment	No flower = 0 Only few flowers = 0,5 Medium flower set = 1 Good flower set = 2 Very good flower set = 3	No yield = 0 Only few berries = 0,5 Light crop load = 1 Medium yielding = 2 Good yielding = 3

In the flower set the clone No. 67. was outstanding, but clones No. 12, 21 and 30 showed also good flower set. The clone No. 21. had the highest crop load and all the other clones selected by us resulted medium yielding capacity.

The crop size parameters

From among the fruit parameters, the size is the most considerable characteristic. Commonly measured fruit size

parameters are the followings: the average fruit weight, longitude and width of fruits. In the average berry weight parameter data significant difference was found between genotypes. The clone No. 67 had the significantly smallest average berry weight (Table 6.). Clones No. 47. and 55. had the significantly largest berry weight. It was also significant difference between clones No. 67. 30. and 21., but they did not differ significantly from clones No. 12. and 29.

Table 6. The average berry size parameter in two-year-old *Lycium* orchard in 2011.

Clones' number	Average berry weight (g)	Homogeneous group	Compared to the control cultivar (%)
67	0.28	a	45.66
30	0.49	b	80.26
21	0.50	b	81.45
12	0.57	bc	93.06
29*	0.61	bc	100.00
47	0.64	c	103.69
55	0.66	c	108.24
SzD _{5%} = 0.13			

* Clone No. 29. (a commercial cultivar) was applied as a control

The clones can be divided into three groups by the fruit size parameter (Figure 1.):

Small fruit size (avg. berry weight < 0,3 g): clone No. 67.;

Medium fruit size (0,3 g < avg. berry weight < 0,5g): clones No. 30. and 21

Large fruit size (0,5 g < avg. berry weight < 0,7): clones No. 55, 47, 29 and 12.

The clone No. 55 significantly did not differ from the 29 clone applied as the control, but it had higher fruit weight by 8%.



Figure 1. Comparison of *Lycium* fruit (berry) size, 2011.

Table 7. The formal index of *Lycium* clone's fruit, 2011.

Clones' number	Formal index of clone's fruit	Homogeneous group	Compared to the control cultivar (%)
12.	1.80	a	98.50
29.*	1.82	a	100.00
55.	1.91	ab	104.83
21.	2.00	ab	109.58
30.	2.20	bc	120.72
67.	2.25	bc	123.45
47.	2.83	c	155.05
SzD _{5%} = 0.93			

* Clone No. 29. (a commercial cultivar) was applied as a control

From the longitudinal and latitudinal data of fruits the formal index data of the crops was calculated.

This numerical value is the quotient of the longitude of the fruits and his width. The berry shape is more longish the value formal index is higher (Table 7.).

- Longish a little bit (1.8 > formal index > 2): clones No. 12, 29 and 55;
- Medium longish (2 > formal index > 2.5) clones No. 21, 30 and 67
- Very longish (2.5 > formal index) clone No. 47.

It can be seen clearly that there are clones selected by us with more favourable fruit size and growing parameters than the control (clone No. 29) which is already grown in abroad.

For different consumptions (dried fruits, dried fruits for tea or muesli) where the fruit size is determining it is offered to use the large fruited genotypes (clones No. 47 and 55).

For processing (jam, fruit juice) where the fruit size is not important, but the nutrition and inner content values are determining those clones are offered to use which have small fruits with higher nutrition and inner content values.

Inner contents values of the tested Wolfberry genotypes

For the processing industry the crops nutrition components (sugar, acids, antioxidant effect compounds and alkaloids) are defining the aims of suitability.

Table 8. Water-soluble total sugar content (Brix%) of *Lycium* genotypes in 2011

Clones' number	Total sugar content (Brix%)	Homogeneous group	Compared to the control cultivar (%)
29.*	12.37	a	100.00
55.	14.63	b	118.33
47.	15.03	c	121.56
12.	15.43	d	124.80
67.	15.53	d	125.61
30.	16.07	e	129.92
21.	16.13	e	130.46
SzD _{5%} = 0.33			

* Clone No. 29. (a commercial cultivar) was applied as a control

The sample fruit harvesting period was free of bigger amount of precipitation, so it means the weather conditions did not have effect on the sugar content of berries. Significant differences appeared between the observed clones in the total sugar content (Table 9.). All the clones selected by us showed higher total sugar content than the control clone No. 29. The significant highest total sugar content gave clones No. 30. and 21., which had higher total sugar content by 30% than the control No. 29. clone.

Table 9. Glucose – fructose ratio of *Lycium* genotypes in 2011.

Clones' number	Glucose / fructose ratio
47	0,988
67	1,017
21	1,018
55	1,025
30	1,041
12	1,042
29*	1,063

* Clone No. 29. (a commercial cultivar) was applied as a control

The quantity and proportion of different sugar content (glucose, fructose, sorbitol etc.) is considerable character inside the total sugar content quantities. Glucose (retentions time: 10,3) and fructose (retention time: 12,3) quantitative analysis was made by HPLC equipment. From quantity of glucose and fructose it was calculated glucose – fructose ratio of *Lycium* berries. With one exception in the case of all observed Wolfberry genotypes the glucose content was higher than fructose content. Clone No. 47 was the exception, where the fructose showed higher level than glucose (Table 9.). Our results harmonize partly with result of Huang et. al., (1998), the glucose was measured as outstanding sugar component of *Lycium barbarum*, but the fructose was not mentioned.

Table 10. The average carotene content (mg/kg) of *Lycium* berries, in 2011

Clones' number	Carotene content (mg/kg)	Homogeneous group	Compared to the control cultivar (%)
67.	181.75	a	78.81
55.	198.12	ab	85.91
12.	221.05	b	95.85
30.	223.30	b	96.83
29.	230.61	b	100.00
47.	271.36	c	117.67
21.	305.23	c	132.36
SzD _{5%} = 37.08			

The carotene content of Wolfberry fruit pulp was determined by spectrophotometric analysis. It was experienced significant differences between the clones in the carotene content (Table 10.). The clone No. 67. was found as containing the significantly lowest carotene content.

Clones No. 47. and 21. showed the significant highest carotene values in their fruits. The clone No. 29 (used as the control) finished in the middle of field. The results showed that carotene content difference between observed genotypes reached even 50% value.

This difference can influence the consumption aims of the selected clones. Instrumental colour measurement of fruit pulp is planned in the future for verification of spectrophotometric analysis results. Based on our results fruits of clones No. 47. and 21. can be suitable for the making of natural fruit colouring matter.

The carotene antioxidant dissolving in fat effect compound that is why the high carotene content fruits of observed clones can be used as of functional foods. In our trial the carotene content of Wolf berries was determined because the carotene compounds are proportional, with the capacity of an antioxidant.

The FRAP method (Ferric Reducing Antioxidant Power) the water-soluble antioxidant is suitable for the definition of antioxidant capacity. The measured results showed significant differences between the observed *Lycium* genotypes. All the clones selected by us in Hungary showed higher FRAP value than the No. 29. clone used as the control (Table 11.).

Examined *Lycium* clones can be divided to four groups by the FRAP value their fruit pulp.

- Clones with the same FRAP value (2,5-2,6 mM/l) as No. 29. clone (control): clone No. 12.;
- Clones with slightly higher FRAP value (2,6-3 mM/l) than clone No. 29: clone No. 21.;
- Clones with significantly higher FRAP value (3-3,6 mM/l) than the control No. 29. clone: clones No. 30, 47. and 67.;
- Clones with outstanding FRAP value (3,6 mM/l and higher) compared to the control No. 29 clone: clone No. 55.

Table 11. *Lycium* clones' average FRAP values (mM/l), 2011

Clones' number	FRAP value (mM/l)	Homogeneous group	Compared to the control cultivar (%)
29.	2.51	a	100.00
12.	2.53	a	101.05
21.	2.87	b	114.59
30.	3.27	c	130.47
47.	3.40	c	135.58
67.	3.52	c	140.62
55.	4.08	d	162.97
SzD _{5%} = 0.28			

Similarly to the sand thorn and briar rose (Kovács et al., 2010) from which fruit juices, pulps, syrups are prepared for tumorous and cancerous patients, *Lycium* it would be possible to use for this aim especially fruits of selected clones No. 67 and 55.

By the literature it is well known, that fruits with high colour matter content (sour cherry, small-berry fruits) usually

have high water-soluble antioxidant capacity. The water soluble antioxidant capacity of measured *Lycium* clones is well commensurable to the water soluble antioxidant capacity of Hungarian sour cherry cultivars (Érdi bőtermő – FRAP 2,45-2,75 mM/l -, Kántorjánosi – FRAP 2,55-4,05 mM/l, IV-3/48 -4,47-7,50 mM/l; Ficzek, 2012; Ficzek et. al., 2009; Ficzek et al., 2010).

It was interesting that those clones which had higher FRAP values (clone No. 67 and 55) the content of the carotene was lower. Clones (No. 29, 21. and 12.) with higher carotene content had lower water-soluble antioxidant values (Figure 2.). To check this correlation, FRAP and TEAC analysis are planned in the future.




Figure 2. Carotene content and water-soluble antioxidant capacity (FRAP) of *Lycium* berries, in 2011.

Presented results justify that domestic *Lycium* population has favourable bush and fruit characteristics from the fruit growing aspect. Data above was based one-year laboratory measurements, so it is necessary repeat them in different years under different ecological and weather conditions in field trials. Examinations in the future should be included determinations of alkaloids (atropine and solanine), acids with the HPLC analytic method.

References

- Bartha D. (1997):** Fa- és cserjehatározó. Mezőgazda Kiadó. Budapest. 36: 112, 182, 238, 282 p.
- Bartha D. (1999):** Magyarország Fa és Cserjefajai. Mezőgazda Kiadó. Budapest. 138 p.
- Ficzek G., Bujdosó G., Tóth M., Stégerné-Máté M., Nótin B., Kállay E., Szügyi S. (2009):** Changes in the Antioxidant Components in Hungarian Bred Sour Cherry Cultivars during the Ripening Period. In: **B.Patil, O. van Kooten, M-J. Amiot-Carlin.** Acta Horticulturae 1040, Proceedings of the Third International Symposium on Human Health Effects of Fruit and Vegetables, Avignon, France, 84-85 p.
- Ficzek G., Stégerné-Máté M., Nótin B., Kállay E., Szügyi S., Bujdosó G. and Tóth M. (2010):** Inner content and processing industrial characteristics of new Hungarian bred sour cherry cultivar candidate. Review of Faculti of Engineering. Analecta Technica Szegediensia. Innovatív Nyomdaipari Kft., Szeged. 68-73. p.

- Ficzek G. (2012):** A hazai alma és meggyfajták egészségvédő és felhasználhatósági értékei gyümölcsanalízis alapján. Doktori értekezés, Budapesti Corvinus Egyetem, Budapest.
- Halmágyi L. (1991):** Erdei és díszfák, cserjék. [In: Halmágyi L. & Keresztes B. (szerk.): A Méhlegelő.] Akadémia Kiadó, Budapest. 176 p.
- Huang, L., Lin, Y., Tian, G. (1998):** Isolation, purification and physicochemical properties of immunoactive glycoconjugates from fruit of *Lycium barbarum* L. *Yaoxue Xuebao* 33: 512-516 p.
- Kovács, S., Udvardy, L. and Tóth, M. (2010):** Breeding Rosa taxa native to the Carpathian Basin for fruit purposes – fruit quality. *Acta Agronomica Hungarica*, 58 (3): 273–281. p.
- Krüssmann, G. (1977):** Lycium In: Manual of Cultivated Broad – Leaved Trees § Shrubs, Volume II, E-PRO, Timber Press, Oregon. 259-261 p.
- Li, Z., Peng, G. & Zhang, S. (1998):** Separation and determination of carotenoids in *Fructus lycii* by isocratic non-aqueous reversed-phase liquid chromatography. *Chinese Journal of Chromatography*, 16: 341–343 p.
- Pap E. (2010):** Szuperélelmiszer az Ördögcérna. *Kertészet és Szőlészet*, 21: 20 p.
- Paul M. Gross (2006):** Wolfberry Nature's bounty of nutrition and health, Tables Ciquel (AFFSA)
- Peng, G. H., Li, Z. & Zhang, S. H. (1998):** Separation and identification of carotenoids in *Fructus lycii* by thin-layer chromatography. *Acta Nutrimenta Sinica*, 20: 76–78 p.
- Salamon P. (2002):** Az Ördögcérna, mint vírusforrás. *Kertészet és Szőlészet*. 27: 15-16 p.
- Schmidt G. & Tóth I. (2006):** A díszfák és díszcserjék ismerete [In: Schmidt G. és Tóth I., (szerk.): *Kertészeti Dendrológia.*] Mezőgazda Kiadó, Budapest, 275 p.
- Schmidt G. (1996):** A legfontosabb díszfák, díszcserjék szaporítása, nevelése és növényvédelme. [In: Schmidt G. & Tóth I. (szerk.): *Díszfaiskola.*] Mezőgazda Kiadó, Budapest, 567 p.
- USDA (2010):** U.S. Department of agriculture, agricultural research service oxygen radical absorbance capacity (ORAC) of selected foods, release 2. Nutrient Data Laboratory
- Vancsura R. (1992):** A fás növények fejlődéstani és rendszertani vonatkozásai. [In: Gencsi L. & Vancsura R., (szerk.): *Dendrológia. Erdészeti növénytan II.*] Mezőgazda Kiadó, Budapest, 689-700 p.
- Weller, P. & Breithaupt, E. (2003):** Identification and quantification of zeaxanthin esters in plants using liquid chromatography–mass spectrometry. *Journal of Agricultural and Food Chemistry*, 51: 7044–7049 p.
- Yong P., Chen M., Yawell L., Kelvin SZE-Yin L., Zhi-Hong J. & Zhongzhen Z. (2005):** Quantification of Zeaxanthin Dipalmitate and Total Carotenoids in *Lycium* Fruits (*Fructus Lycii*). *Plant Foods for Human Nutrition* 60: 161–164 p.
- Young, G., Roland, L., Schreuder, M. (2006):** Ningxia Wolfberry: the Ultimate Superfood. Essential Science Publishing.