



Swansea University  
Prifysgol Abertawe



## Cronfa - Swansea University Open Access Repository

---

This is an author produced version of a paper published in:  
*Ecological Indicators*

Cronfa URL for this paper:  
<http://cronfa.swan.ac.uk/Record/cronfa40894>

---

### Paper:

Roces-Díaz, J., Vayreda, J., Banqué-Casanovas, M., Cusó, M., Anton, M., Bonet, J., Brotons, L., De Cáceres, M., Herrando, S., et. al. (2018). Assessing the distribution of forest ecosystem services in a highly populated Mediterranean region. *Ecological Indicators*, 93, 986-997.  
<http://dx.doi.org/10.1016/j.ecolind.2018.05.076>

Released under the terms of a Creative Commons Attribution Non-Commercial No Derivatives License (CC-BY-NC-ND).

---

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>

# Assessing the distribution of forest ecosystem services in a highly populated Mediterranean region

Jose V. Rocés-Díaz<sup>1,2\*</sup>, Jordi Vayreda<sup>1</sup>, Mireia Banqué-Casanovas<sup>1</sup>, Martí Cusó<sup>1</sup>, Marc Anton<sup>3</sup>, José A. Bonet<sup>4,5</sup>, Lluís Brotons<sup>1,4,6</sup>, Miquel De Cáceres<sup>1,4</sup>, Sergi Herrando<sup>1,3</sup>, Juan Martínez de Aragón<sup>4</sup>, Sergio de-Miguel<sup>5</sup>, Jordi Martínez-Vilalta<sup>1,7</sup>

1: CREA, E08193 Bellaterra (Cerdanyola del Vallès), Catalonia, Spain

2: Department of Geography, Swansea University, Singleton Park, Swansea SA2 8PP, United Kingdom

3: Catalan Ornithological Institute, Natural History Museum of Barcelona, 08019 Barcelona, Catalonia (Spain)

4: Forest Sciences Centre of Catalonia (CEMFOR-CTFC), 25280 Solsona, Catalonia (Spain)

5: Department of Crop and Forest Sciences, Universitat de Lleida-Agrotecnio Center (UdL-Agrotecnio), Av. Rovira Roure, 191, E-25198 Lleida, Catalonia (Spain)

6: Consejo Superior de Investigaciones Científicas (CSIC), 08193 Cerdanyola del Valles, Catalonia (Spain)

7: Universitat Autònoma de Barcelona, E08193 Bellaterra (Cerdanyola del Vallès), Catalonia, Spain

\*corresponding author José V. Rocés-Díaz ([jvroces@gmail.com](mailto:jvroces@gmail.com)) CREA, +34935814850 Autonomous University of Barcelona, E-08193 Cerdanyola del Vallès, Barcelona (Spain)

## Abstract

Forest ecosystems provide a wide range of goods and services to society and host high levels of biodiversity. Nevertheless, forest ecosystem services (ES) are often quantified and assessed using simplified methodologies (e.g., proxy methods based exclusively on Land Use Land Cover maps) that introduce substantial uncertainty in the analysis by ignoring, for instance, the species composition and spatial configuration of the ecosystems studied. In this work we defined and calculated a set of 12 indicators of several ES for the forests of the highly populated region of Catalonia (North-eastern Iberian Peninsula). The indicators combined different sources of information such as forest surveys, ecological model predictions and official statistics, but also included additional land cover information. All ES indicators were aggregated at the municipality level to compare their values and distribution patterns. We assessed spatial trade-offs and synergies among ES, as well as their relationships with a set of socioeconomic, climatic and biodiversity variables using correlation analyses and mixed-effects models. The results suggest a clustering of provisioning and regulating ES in mountainous zones towards the North of the study area. These two types of services showed a high degree of spatial similarity and presented high positive correlations. In contrast, cultural ES showed a more scattered pattern, which included lower elevation areas in the South of the study region. Climatic conditions were the main determinants of the spatial variability in the supply of the different ES, with most indicators being positively associated with precipitation and negatively associated with temperature. In addition, biodiversity (particularly woody species richness) showed positive relations with most of these ES, while socioeconomic indicators (such as

population density and the percentage employment in agriculture) showed negative associations with most of them. The combination of information from different data sources (including primary data) allowed for a detailed analysis of forest ES, likely removing some of the problems derived from approaches based only on proxy methods. In addition, the use of municipalities as study unit makes results directly relevant to management and planning strategies operating at this scale (e.g., forest management and planning).

**Keywords:** Mediterranean forests; Catalonia; ecosystem services indicators; ecosystem services mapping; Trade-offs and synergies; Hotspots

### **Highlights**

A set of indicators was developed to assess the spatial distribution of a wide range of forest ecosystem services (ES).

Provisioning and regulating ES showed an aggregated spatial pattern, whereas cultural were more scattered.

Most ES were highly associated with climate and species richness variables.

Cultural ES were more associated with socioeconomic variables.

## **1. Introduction**

Forest ecosystems are key elements for the maintenance of global biodiversity (Brooks et al., 2006). They support a range of ecosystem functions and provide multiple and essential ecosystem services (ES) to society (MEA, 2005). Some of the main forest ES can be classified as regulating services: climate and water regulation, erosion and flood control, etc. (Miura et al., 2015). However, materials and energy provision and cultural services are also relevant in forests (MEA, 2005). Forest ecosystems have been strongly disturbed and modified by the human use of the landscapes, although the intensity of historical disturbances and the current condition of these ecosystems are highly heterogeneous in space (FAO, 2014; Trumbore et al., 2015).

Several authors have highlighted the relevance of the biodiversity contained in Mediterranean landscapes (Brooks et al., 2006) and in particular in the Mediterranean Basin (Medail and Quezel, 1999; Hampe and Petit, 2005), which is considered a biodiversity hotspot of global relevance (Myers et al., 2000). The forests of this region have been managed and modified for millennia due to the historical use of natural resources by human societies (Underwood et al., 2009). In the context of global change, the development of effective management and conservation strategies is key for the maintenance of their diversity and ecosystem functions (Costanza et al., 1997). A series of drivers have been identified as having potential effects on

forest ecosystems and their supply of ES (EME, 2011; Thom and Seidl, 2015), including land-use changes, wildfires, climate change, alien species, pests and pathogens (Vila et al., 2010; Doblas-Miranda et al., 2015).

Methodological factors may have a large impact on the quantification of ES (Eigenbrod et al., 2010; Van der Biest et al., 2015) and are important sources of uncertainty in ES assessments (Hou et al., 2013). Land Use/Land Cover (LULC) information often constitutes the basis for ES assessments (Hou et al., 2013). However, the use of proxy-based methods relying only on LULC data assumes that if one class (an ecosystem type) provides a specific ES, the level of supply is constant in space, neglecting the importance of other ES drivers not represented by land use categories. This leads to a potentially large generalization error in ES assessments (Plummer, 2009). Notably, these proxy-based approaches often hide large differences in the composition and structure of the forests that drive ecosystem functioning (Vila et al., 2007; Ruiz-Benito et al., 2014) and ES supply (Alamgir et al., 2016; Sutherland et al., 2016). Recent studies overcome some of the limitations of proxy-based methods by defining specific bio-physical indicators (e.g., Garcia-Nieto et al., 2013) or by using specific information about the structure and the composition of these ecosystems (Roces-Díaz et al., 2017). Finally, accurate assessments of ES should include the analysis of ES spatial patterns and their spatial associations (Andrew et al., 2015), including synergies and trade-offs as well the identification of areas with particularly high levels of overall supply (hotspots; Mouchet et al., 2014; Schröter and Remme, 2016).

Forest planning and management strategies are beginning to include forest ES as key elements in their assessments (e.g. Frank et al., 2015; Triviño et al., 2015), which can help to visualize and promote the multi-functionality of these systems. Spatial-dependent aspects, such as the scale and the administrative level of analysis, become particularly relevant for planning and management objectives (Hein et al., 2006). In this regard, the municipal domain often offers a good compromise between reasonable spatial resolution and administrative relevance (Rodríguez-Loinaz et al., 2015; Roces-Díaz et al., 2018). Within this management-oriented perspective, the spatial patterns and relationships between ES (trade-offs and synergies) should also be evaluated (Duncker et al., 2012). For instance, negative relationships are frequently reported among materials provision (such as timber) and cultural services (Garcia-Nieto et al., 2013) or biodiversity (Duncker et al., 2012).

In recent years several studies have analysed the ES provided by European Mediterranean landscapes and uncovered their strong relations with social and environmental characteristics (García-Llorente et al., 2015). Some of these studies have focused on the assessment of specific, particularly relevant ES such as water provision (Quintas-Soriano et al., 2014) or erosion

regulation (Guerra et al., 2016), while other works have described and analyzed all the ES provided by specific types of forest ecosystems (e.g., cork oak woodlands (Bugalho et al., 2011)). However, there are still few studies addressing different forest types at the regional scale and including a complete set of ES as a necessary step to address trade-offs and spatial variability in their overall provision (but see Garcia-Nieto et al., 2013).

In this work we define a comprehensive set of bio-physical indicators of forest ES for Catalonia (North-eastern Spain) on the basis of different data sources, and assess them at the municipality level. The specific objectives of this work are: i) to analyze the spatial patterns of these ES and to identify their main hotspot areas; and ii) to assess the spatial relationships of these ES (trade-offs and synergies) and the association between these ES and different socioeconomic, climatic and biodiversity variables that characterize the study area. We hypothesize that the ES analyzed will show clearly differentiated spatial patterns, with a high clustering of provision and regulating services on mountainous municipalities with higher forest cover and lower population density. Other ES (e.g., cultural) will be associated to more populated areas. These disjoint spatial patterns may reflect trade-offs between different ES.

## **2. Material and Methods**

### **2.1. Study area and outline of the experimental approach**

Our study area is Catalonia (North-eastern Spain; Figure 1), an administrative region that covers 32,114 km<sup>2</sup>. It is mainly located in the Mediterranean Biogeographic Region, although a part of its northern area (the Pyrenees Mountains) belongs to the Alpine Region. It is a mountainous area with an altitudinal range from the sea level to more than 3,000 meters on the highest peaks of the Pyrenees. Catalonia had a population of 7,504,008 people in 2015, 43% of them concentrated in the metropolitan area around the capital city (Barcelona, 636 km<sup>2</sup>). It is a highly forested region (43% of its area was covered by forest; LCMC, 2009) where about 33% of the land area was included in the Natura 2000 Network (a system of nature protection areas in the territory of the European Union). It is dominated by tree species of the Pinaceae and Fagaceae families. Forests from coastal and low altitude areas are dominated by *Pinus halepensis* Mill. (Aleppo pine), *Quercus faginea* Lam. (Portuguese oak) and *Quercus ilex* L. (Holm oak). At middle-altitude ranges (from 800 to 1,500 m) the main species are *Pinus sylvestris* L. (Scots pine), *Pinus nigra* J.F.Arnold (Black pine), *Quercus pubescens* Willd. (Downy oak, synonym of *Quercus humilis* Mill.) and also *Fagus sylvatica* L. (European beech) in the wettest zones. Finally, at altitudes higher than 1,500 m the main species are *Pinus uncinata* Raymond ex A.DC. (Mountain pine) and *Abies alba* Mill. (Silver fir). These forests have shown expansion and shrinkage processes over the last millennia in congruence with changes in the

environmental conditions on the Mediterranean Basin and historic land use (Grove and Rackham, 2003). Importantly, recent episodes of forest decline have been detected in the study area, affecting mostly species reaching the southern limit of their distribution in the Iberian peninsula, such as *P. sylvestris* (e.g., Martínez-Vilalta and Piñol, 2002) and *F. sylvatica* (e.g., Peñuelas and Boada, 2003). Approximately 80% of the forests in Catalonia are privately-owned, whereas the remaining 20% are public.




Figure 1. Location of the study area, distribution of the main types of land uses and location of the metropolitan area of Barcelona, together relevant administrative units as counties and municipalities (source: LCMC, 2009).

The spatial unit of our analysis was the municipality (N = 947 municipalities in Catalonia). ES maps were obtained at this municipality level (see below), where values from different sources (including raster format) were aggregated to polygons. As we focused on forest ecosystems and on ES capacity or actual supply (not demand), we restricted our analyses to those municipalities with substantial forest cover. Thus, we selected only those municipalities that contained at least three permanent plots of the Third National Forest Inventory of Spain (NFI; MAGRAMA,

1997-2007), which was considered a minimum sample size to obtain representative estimates and perform statistical comparisons. The Spanish NFI is an intensive program of periodic surveys (every ~10 years) that cover the whole forested area of Spain following a uniform sampling design (Appendix S1). Data of the 3<sup>rd</sup> NFI, conducted in Catalonia in 2000-2001, is used unless otherwise stated. NFI plot density is ~1 plot/km<sup>2</sup> of forest, so that the 3 plot threshold corresponds with an average of (at least) 300 ha of forest per municipality, resulting in a subset of 576 municipalities. Forest cover ranged between 10 % and 95 % in these municipalities.

In addition to using NFI data to delineate the areas (municipalities) of interest to this study, we also used the NFI dataset as a basis for the assessment of most ES. The majority of ES indicators were calculated either directly from NFI data (NFI data, Table 1) or modelled using primary NFI data as input (model-based). These indicators were calculated for individual NFI plots and then aggregated at municipality level. Other ES were estimated based on combining datasets at municipality level, including aerial photo interpretation and GIS analysis (map-based), official statistics, and additional information provided by research centres or other institutions (other statistics) (Table 1). The resulting ES maps were then aggregated to municipal units and resulting values were compared to explore potential trade-offs and synergies between pairs of ES. Finally, we assessed to what extent climatic and socioeconomic conditions explained the spatial variability of forest ES in the study area. These methods are developed in the following sections.

Table 1. Description of the ES indicators, their definitions and the units, transformations, sources and temporal ranges used in this

ES	Indicator	Code	Definition	Units	Supply	Method used	T
<b>Provisioning</b>							
Food provision	Mushrooms production	P1	Expected edible mushroom production for pine, oak and fir forests in one typical year	kg/ha/year	Capacity	Model-based	I
Materials/ Energy provision	Timber and firewood	P2	Wood extractions and firewood harvesting in public and private forests at municipal level per year	t/ha/year	Actual	Official statistics	I
Water provision	Exported water	P3	Water exported yearly by surface runoff or deep drainage into the water table for each forest plot (National Forest Inventory, NFI3) in Catalonia	l/m <sup>2</sup> /year	Capacity	Model-based	I
<b>Regulating</b>							
Climate regulation	Forest Carbon sink	R1	Forest carbon sink on the above and below ground vegetation	t/ha/year	Actual	NFI data	-
Soil fertility regulation	Soil Organic Carbon	R2	Amount of organic carbon in the soil	t/ha	Actual	Model-based	S
Water regulation	Canopies and soil water storage capacity	R3	Sum of canopy water storage capacity and soil water holding capacity for each forest plot (NFI3) in Catalonia	l/m <sup>2</sup> /year	Actual	Model-based	I
Flood protection	Riparian forest cover	R4	Riparian forest cover around watercourses considering a buffer zone of 25m around	%	Actual	Map-based	S
Erosion control	Erosion control	R5	Forest cover of areas with a slope higher than 30%	%	Actual	Map-based	S
<b>Cultural</b>							
Recreational	Rural tourism	C1	Number of beds in rural tourism establishments per municipality	N <sup>o</sup> places/ha	Actual	Official statistics	S
Existence	Natura 2000	C2	Surface of protected areas included in the Natura 2000 Network.	%	Capacity	Map-based	S
Experiential use of organisms	Animal observations	C3	Animal species observations introduced at web portal Ornitho.cat	N <sup>o</sup> obs./ha/year	Actual	Other statistics	I
Physical use of landscape	Wikiloc tracks	C4	Routes recorded in Catalonia and introduced by users at the Wikiloc® app and web portal	N <sup>o</sup> tracks/ha	Actual	Other statistics	I

\* The different data sources referenced in this table are more detailed in the Supplementary Material (Appendix S1)



## 2.2. Ecosystem services assessment

### *Bio-physical indicators and ecosystem services typology*

For this work we defined a specific set of ES indicators adapted for forest ecosystems of the study area and based on widely used ES classifications (i.e. CICES 4.3: Haines-Young and Potschin, 2013). We considered three categories of ES: provisioning, regulating and cultural. For each category we defined different indicators that are related with the supply (actual or capacity) of specific ES. This set of indicators was developed selecting important forest ES in the study area subject to data availability (see below) and, although it is not exhaustive, it is representative of a wide range of ES. All these indicators are presented in Table 1, while further descriptions of data sources and calculation processes are provided in the Supplementary Material (Appendix S1). Although most indicators defined in this assessment represent actual supply of ES, for three of them there was no available data of actual supply and thus their values represent supply capacity (Table 1).

***Provisioning category (three ES).*** We used edible mushrooms production for food provision (P1), as mushroom picking is an important social and economic activity in the study area (Bonet et al., 2010). We estimated edible mushroom production for year 2013 for each municipality in kg/ha/year. For pine forests we combined Third National Forest Inventory (NFI) data and the model developed by de-Miguel et al. (2014). This model accounts for the effects of stand composition, structure and site characteristics in a typical year (Appendix S1). Pine forests are by far the most important forests in terms of cover and, particularly, mushroom production in the study area (Bonet et al., 2010). By contrast, for *Quercus* sp. and *A. alba* forests we used their mean production value per unit of forest area (J.A. Bonet, unpublished) because predictive models based on plot characteristics have not been yet developed. For materials and energy provision (P2) we used the annual timber and firewood removals from both public and privately-owned forests for the period 2006-2014 (in t/ha/year). This information was obtained from official statistics provided by the Catalan Government. Finally, for water provision (P3) we used the amount of water exported from forests, estimated as the sum of surface runoff and deep drainage into the water table for each NFI plot (in l/m<sup>2</sup>/year). Both quantities were estimated using a soil water balance model (De Cáceres et al., 2015). This model calculates daily water balance driven by meteorological data and additional variables characterizing the soil and vegetation structure. Exported water was estimated for year 2010. More detailed information about this model can be found in Appendix S1.

***Regulating category (five ES).*** For climate regulation (R1) we considered the forest Carbon sink capacity in t/ha/year. This value was calculated from the forest Carbon stock change (above- and belowground) using the methodology described in Vayreda et al. (2012) from

consecutive forest surveys (comparing plot-level data from 1989-1990 (2<sup>nd</sup> NFI) and 2000-2001 (3<sup>rd</sup> NFI)). We used the soil organic carbon (SOC, t/ha) to represent the maintenance of soil fertility (R2), as obtained from the map developed by Doblás-Miranda et al. (2013). The capability of forests to regulate water flows (water regulation, R3) was estimated using the sum of canopies' water storage capacity and soil water holding capacity for each NFI3 plot in l/m<sup>2</sup>/year. It was obtained from the same soil water balance model mentioned above (De Cáceres et al., 2015). Flood protection (R4) and erosion control (R5) are strongly dependent on the occurrence of a specific type of ecosystem (riparian forest) or on the detailed distribution of forest cover. For their calculation we used a highly detailed LULC map of Catalonia (LCMC, 2009) with a very high spatial resolution (scale 1:5,000 and minimum mapping unit of 500 m<sup>2</sup>). For flood protection we used the percentage of area alongside water courses covered by riparian forests. To estimate this coverage, we defined a 25 m buffer around watercourses using the LCMC (2009), and calculated the percentage of riparian forests inside this buffer area. Finally, erosion control was assessed by the percentage of slopes steeper than 30% grade that were covered by forests.

***Cultural category (four ES).*** As a proxy of recreational use (C1), we used the number of beds in rural accommodation establishments according to the Official Guide of Touristic Establishments of Catalonia for year 2015 ([www.establimentsturistics.gencat.cat](http://www.establimentsturistics.gencat.cat)), aggregated by municipalities. Existence of landscapes or organisms with conservation interest (C2), was defined as the percentage of municipal surface occupied by protected areas included in the Natura 2000 Network ([www.mediambient.gencat.cat](http://www.mediambient.gencat.cat)). 59.3% of the Natura 2000 Network in the municipalities studied correspond to forests ecosystems. Regarding experiential use of organisms by people (C3) we have calculated the number of animal observations per hectare of municipality using the data stored in the web portal [www.ornitho.cat](http://www.ornitho.cat). This portal stores more than 3,000,000 observations (mostly birds but also mammals, reptiles, amphibians and some groups of invertebrates) from more than 3,500 observers, and includes species based on the perceived interest by the observer (regardless of ecological functionality or any other 'objective' measure of importance). Data were provided by the Catalan Ornithological Institute (ICO) and represent observations uploaded by users of the application all around Catalonia between 2010 and 2015 (including all its municipalities). Finally, in relation to the physical use of the landscape (C4) we calculated the density of routes per municipality introduced in the web portal [www.wikiloc.com](http://www.wikiloc.com), as provided by Wikiloc® (unpublished). Routes include trekking, biking, skiing, running and all types of outdoor routes recorded in Catalonia between 2006 and 2014.

### ***Socioeconomic, climatic and biodiversity predictors of ES supply***

We used a series of socioeconomic, climatic and biodiversity variables to assess if they explained ES variability among municipalities. Socioeconomic variables were obtained from the data provided by the regional administration (Statistical Institute of Catalonia: [www.idescat.cat](http://www.idescat.cat), IDESCAT, 2015), aggregated at the municipality level. We considered six socioeconomic variables: population density (inhabitants/km<sup>2</sup>), unemployment rate (calculated as the number of unemployed people over the total working population), and the percentage of working population occupied in agriculture, industry, construction and tertiary sector (separately). In addition, we included climatic (mean annual temperature (°C) and total annual precipitation (l/m<sup>2</sup>)) and biodiversity information. As biodiversity information we used total forest woody species richness (from NFI3) and forest bird richness (from Estrada et al., 2004) per municipality. Detailed information about these variables is provided in Appendix S1.

### **2.3. Data analysis**

#### ***Data processing***

ES supply needs to be referred to a given land surface to produce useful and comparable indicators. Some ES were expressed in % to land area and no further standardization was required (R4 and R5; Table 1). In all other cases, indicators were expressed in two ways, with the aim of reflecting two types of complementary information: (i) ES supply per unit of municipality area (referred here as land-based indicators); and (ii) supply per unit of forest cover (forest-based indicators). Forest-based indicators are in principle independent of the percentage of the municipalities' area covered by forests, which is likely to be an important driver of the spatial variation of land-based indicators.

Prior to analyze relationships among pairs of ES and with socio-environmental variables, ES values were transformed to make their statistical distribution closer to normality, by applying logarithmic or squared root functions when necessary (Table 1). In addition, proximity-to-target methodology was used to standardize their values to a common 0-1 scale (Rodriguez-Loinaz et al., 2015). In our case, lowest and highest benchmarks were determined from minimum (or maximum) values recorded for each ES and all intermediate values were rescaled linearly between the two extremes. After standardization, they were grouped and averaged for each municipality by category (provisioning, regulating and cultural), which involves giving equal weight to all indicators within a category (i.e., assuming that all indicators are relevant and similarly important).

### *Statistical analysis*

Pearson correlations were used to explore the relationships (trade-offs and synergies; Mouchet et al., 2014) among normalized ES and their categories in space, as well as their relationships with socioeconomic, climatic and biodiversity variables. The spatial aggregation of each ES was explored using Moran's I on a Spatial Autocorrelation Analysis (Moran, 1948; ESRI, 2013b). In addition, each normalized and standardized ES was modelled as a function of socioeconomic, climatic and biodiversity variables using linear mixed-effects models. To avoid multicollinearity issues, correlation and principal component analysis (PCA) were used to select a subgroup of independent variables to be included in the mixed-effects models as explanatory variables. As a result, we finally selected seven variables (population density, unemployment rate, population occupied in tertiary sector, mean annual temperature, total annual rainfall, woody species richness and bird richness) with correlation coefficients between them  $< 0.53$ . PCA confirmed that these variables were relatively orthogonal (Appendix 2, Figure S2.1). County (groups of municipalities, N=41, Figure 1) was incorporated as a random factor to better account for spatial autocorrelation. Preliminary analyses confirmed that including county as a random factor improved model fit (in terms of the Akaike Information Criterion, AIC). All statistical analyses were conducted using the R software environment (v.3.2.0; R Development Core Team, 2014). Finally in order to detect the areas with highest supply of the different categories of ES, a hotspot analysis on ES maps was performed (ESRI, 2013a) using the Getis-Ord  $G_i^*$  clustering method (Getis and Ord, 1992; Schröter and Remme, 2016). A more detailed account of all these analyses is provided in the Supplementary Material (Appendix S2).

## **3. Results**

### **3.1. Spatial distribution of ecosystem services**

The spatial patterns of ES varied from highly clustered to dispersed, and many of them showed a gradient from mountainous areas (in the north) to lowlands (in the south) (Figure 2 for land-based and Figure 3 for forest-based indicators). Land-based indicators of food provision (P1) and water provision (P3) services showed a similar pattern with a concentration of high values in the North of the study area. High supply values for materials provision (P2) were concentrated in the Northeast. On the other hand, climate regulation and soil fertility (R1 and R2) had a rather scattered supply pattern. Other regulating services (R3, R4 and R5) were clustered in the eastern half and the Pyrenees mountains. Most cultural services (C2, C3 and C4) had complex patterns, with high-value supply areas being often close to the coast or the most populated zones. Highest values were observed in Eastern and particularly North-Eastern areas, close to the coast, including the pre-Pyrenees but also southern mountain ranges. The spatial

patterns for forest-based indicators were generally similar to those of land-based indicators. However, some differences could be detected (compare Figure 3 with Figure 2). For instance, when forest-based indicators were used, high values of SOC-soil fertility (R2) and water regulation (R3) were more clustered in the Northwest, on mountainous municipalities.




Figure 2. Maps of all the ES land-based indicators, separated by ES category: provisioning (P), regulating (R) and cultural (C). In each plot municipalities are classified in five levels of supply (20% percentiles, see legends). White colour indicated no supply (non-forested municipalities).

Regarding the spatial aggregation of ES, provisioning and regulating services were more clustered (Moran's I index values between 0.36-0.54 for provisioning and 0.17-0.47 for regulating ES), whereas the spatial patterns of cultural services were more scattered throughout the study area (Moran's I = 0.06-0.25). All these values of Moran's I coefficient correspond to land-based indicators, but similar results were obtained when Moran's I was calculated for forest-based indicators (data not shown). Correlations between of the same ES expressed per unit land (land-based) and per unit forest (forest-based) were positive and highly significant in all cases (p-values < 0.001). These correlations were particularly high for provisioning ES (r = 0.82-0.99) and a little lower for cultural (r = 0.62-0.95) and regulating ES (r = 0.25-0.67).




Figure 3. Maps of all the ES forest-based indicators, separated by ES category: provisioning (P), regulating (R), and cultural (C). Indicators R4 and R5 are omitted because the distinction between land-based and forest-based indicators is not meaningful in their case. In each plot municipalities are classified in five levels of supply (20% percentiles, see legends). White colour indicated no supply (non-forested municipalities).

When the overall distribution of the three categories of ES was compared (land-based indicators), two of them (provisioning and regulating) showed similar spatial patterns, with high supply areas clustered around the Pyrenees (particularly central and eastern areas; Figure 4). Cultural ES were in general less clustered and their highest values occurred close to the Mediterranean coast. The hotspots distribution patterns for land-based indicators confirmed these patterns (Figure 4) and showed that the highest supply areas for provisioning and regulating ES were clustered around the Eastern half of Pyrenees. Cultural ES had a more scattered distribution of hotspots throughout the study area. Hotspots for forest-based indicators are provided in the Supplementary materials (Figure S3.1).




Figure 4. Maps for the three categories of ES using land-based indicators, and location of their hotspot areas. The ES categories are classified in six levels of supply (a no-supply class and five 20% quantiles). Hotspots maps are classified in five categories using the standard deviation of Getis Ord  $G_i^*$  statistic.

### 3.2. Relationships among ecosystem services and with socioeconomic, climatic and biodiversity variables

In general, land-based indicators showed positive pair-wise relationships (Table S3.1) while forest-based ones (Table S3.2) showed in some cases negative relationships, especially between cultural and other indicators. Positive associations were particularly strong among provisioning and regulating services, with highest values ( $r > 0.7$ ) between water storage and mushroom production or water exported.

The comparison between ES categories showed highest correlations between provisioning and regulating ES for both land-based and forest-based indicators (Table 2). The correlation between these two ES categories and cultural ES was positive and significant for land-based indicators, but became non-significant when forest-based indicators were used. Regarding the relationships between categories of ES and climatic variables, temperature was negatively associated with provisioning, regulating and cultural ES (always significantly for land-based indicators). Rainfall was positively related to all ES categories when quantified using forest-based indicators. Biodiversity (woody species and bird richness) showed significant positive relationships with all services (land-based and forest-based, except for forest-based cultural services). Regarding socio-economic variables, the number of significant correlations with ES



categories was lower. The percentage employment in agriculture was by far the variable showing more associations, showing negative relationships with the three categories of ES. Population density showed a negative relationship with regulating services (land-based) and a positive one with cultural ES (forest-based). Finally, the percentage of people employed in the tertiary sector showed positive associations with regulating and cultural services.

Table 2. Correlations among ES categories and socio-environmental variables. Significant values (\*\*\*P<0.001 \*\*0.01 \*P<0.05) are highlighted bold (positive relationships) and italic (negative relationships) characters.

Variables	Land-based type			Forest-based type		
	Prov.	Reg.	Cult.	Prov.	Reg.	Cult.
<b>Provisioning</b>	1.00	-	-	1.00	-	-
<b>Regulating</b>	<b>0.66***</b>	1.00	-	<b>0.50***</b>	1.00	-
<b>Cultural</b>	<b>0.31***</b>	<b>0.35***</b>	1.00	-0.07	0.03	1.00
<b>Population dens.</b>	-0.14	<b>-0.15*</b>	0.01	-0.09	-0.11	<b>0.17**</b>
<b>Unemployment</b>	-0.11	-0.08	0.00	-0.13	<b>-0.16**</b>	0.11
<b>Agriculture</b>	<b>-0.15*</b>	<b>-0.22***</b>	<b>-0.24***</b>	-0.11	<b>-0.18***</b>	<b>-0.24***</b>
<b>Industry</b>	0.13	0.10	-0.11	<b>0.15**</b>	0.01	-0.10
<b>Construction</b>	0.01	0.06	0.07	-0.02	0.07	0.02
<b>Tertiary sector</b>	0.05	0.13	<b>0.28***</b>	0.02	<b>0.15*</b>	<b>0.31***</b>
<b>Temperature</b>	<b>-0.31***</b>	<b>-0.27***</b>	<b>-0.18***</b>	<b>-0.29***</b>	<b>-0.55***</b>	-0.02
<b>Rainfall</b>	<b>0.62***</b>	<b>0.56***</b>	<b>0.45***</b>	<b>0.54***</b>	<b>0.69***</b>	<b>0.18***</b>
<b>Woody sp. richness</b>	<b>0.46***</b>	<b>0.34***</b>	<b>0.26***</b>	<b>0.37***</b>	<b>0.29***</b>	0.10
<b>Bird richness</b>	<b>0.61***</b>	<b>0.47***</b>	<b>0.27***</b>	<b>0.59***</b>	<b>0.61***</b>	0.03

Linear mixed-effects models were used to explore the combined effects of socioeconomic, climatic and biodiversity drivers on the distribution of individual ES (see Table 3 for land-based indicators and Table S3.3 in Appendix S3 for forest-based ones). Explained variance ranged between 9 and 54% (marginal R<sup>2</sup>) and between 13 and 79% (conditional R<sup>2</sup>) for land-based indicators, being generally higher for provisioning ES. The range of explained variance was similar for forest-based indicators. Model results confirmed the dominant role of climatic variables in determining the spatial distribution of ES in the study area. Rainfall showed significant positive effects for 10 out of 12 land-based indicators (7 out of 10 for forest-based ones), while mean temperature was associated with 8 land-based (and 7 forest-based) ES, showing positive and negative relationships depending on the ES (Table 3 and Table S3.3). Woody species richness was the biodiversity variable with highest explanatory power. Its effect on land-based indicators was generally positive (significant in 7 cases), but it was significantly negative on animal observations. The number of negative effects for woody species richness was higher regarding forest-based indicators, including soil fertility, recreational and experiential. Bird richness showed less significant relationships and some of them were negative (i.e., existence-Natura 2000 for both types of indicators, soil fertility and water regulation for



land-based ones, and climate regulation for forest-based ones). Finally, regarding socioeconomic variables, population density had a significant negative effect on 7 land-based indicators, although these relationships did not always remain significant when forest-based indicators were used. Population density was positively associated with two cultural ES (experiential and physical use), regardless of whether land-based or forest-based indices were used. Cultural ES indicators were positively related with the percentage population employed in the tertiary sector, except for recreational.

## **4. Discussion**

### **4.1. Strengths and limitations of this study**

In this work we combined models of ecological processes and different types of databases to produce a set of bio-physical indicators of a wide range of ES. Data sources were as close to underlying ecological processes as possible and included i) field sampling measures, ii) data from statistical models, iii) data from mechanistic models, iv) statistics from official administrative sources, v) information from cartographic sources and vi) data provided by associations for nature conservation and recreation. This set of ES indicators includes primary data and also accounts for structural properties of the ecosystems analyzed, which are often related with ES supply (Gamfeldt et al., 2013).

Most of the indicators used in this work (all except mushroom production, exported water and Natura 2000 network) represent the actual supply (or actual use *sensu* Schröter et al. (2014)) of the ES analyzed, and not their supply capacity (or capacity *sensu* Schröter et al. (2014)) (Table 1). This distinction is important for an accurate assessment of ES (Boerema et al., 2016). It should be noted that the relationship between actual supply and supply capacity differs among ES, reflecting differences in the context of ES delivery and the spatial configurations of supply capacity and demand (Burkhard et al. 2012; Schröter et al., 2014), which was not assessed in this study. In our case the selection of ES (measuring actual vs. supply capacity) primarily reflected limitations in data availability but also inherent differences between ES types (Yahdjian et al., 2015). In particular, the ES categories for which we mix indicators of actual and supply capacity (provisioning and cultural, Table 1) correspond to those for which the overlap between supply capacity and demand is relatively low (Yahdjian et al., 2015). In addition, we assumed that all our indicators of regulating ES represent actual supply, but this could be highly context-dependant (e.g. Andersson et al., 2015; Sutherland et al., 2017).

Table 3. Results of mixed-effects models performed for the 12 land-based indicators and some of the socio-environmental variables. Coefficient estimates plus/minus the standard error followed by their level of significance. \*\*\*P<0.001 \*\*0.01 \*P<0.05. R<sup>2</sup>m: marginal

ES	Intercept	Population Density (x10 <sup>-5</sup> )	Unemployment (x10 <sup>-4</sup> )	Tertiary sector (x10 <sup>-4</sup> )	Mean Temperature (x10 <sup>-4</sup> )	Rainfall (x10 <sup>-4</sup> )
<b>Food provision (P1)</b>	0.55±0.65	-19.5±4.2***	105.8±53.0**	28.6±28.7	-125.5±29.9***	23.7±3.9***
<b>Materials provision (P2)</b>	-11.22±1.86***	-0.1±1.1	-19.3±158.4	-279.6±85.7***	2575.0±852.1***	4.3±1.1***
<b>Water provision (P3)</b>	2.59±0.48***	-14.7 ±3.1***	28.5±39.3	16.4±21.3	-671.7±221.1***	23.7±2.8***
<b>Climate regulation (R1)</b>	-0.96±0.32***	-1.5±2.6	-8.6±39.9	35.7±20.2*	249.5±142.2*	12.3±1.9***
<b>Soil fertility (R2)</b>	5.36±1.32***	-25.9±9.5***	44.2±145.5	82.9±76.2	-825.1±596.4	31.9±8.2***
<b>Water regulation (R3)</b>	-0.83±0.81	-23.9±0.5***	103.9±69.4*	25.5±37.6	193.9±372.1	45.6±4.9***
<b>Flood regulation (R4)</b>	-3.91±1.07***	-10.2±6.9	-133.7±101.5	2.4±54.4	2112.0±484.8***	2.5±6.5
<b>Erosion control (R5)</b>	2.47±1.58	-3.4±11.6	272.5±177.5	-102.3±92.6	533.2±711.9	43.8±9.8***
<b>Recreational (C1)</b>	-0.06±0.04	-0.8±0.3***	-13.0±4.2***	-3.4±2.7	62.2±19.7***	0.7±0.3**
<b>Existence (C2)</b>	0.61±1.96	-14.4±13.5	289.9±203.7	385.0±107.9***	-2194.0±890.2**	63.3±12.2***
<b>Experiential (C3)</b>	-9.49±1.21***	40.5±8.2***	-42.7±122.6	303.7±65.3***	1931.0±55.4***	3.8±7.5
<b>Physical use (C4)</b>	-0.43±0.20**	4.6±1.6***	21.2±25.0	51.9±12.7***	105.8±90.6	2.8±1.3**

Selection of ES indicators is frequently problematic and our study is no exception. Firstly, we did not take into account some potentially important ES (particularly some cultural services such as traditional knowledge, hunting or educational aspects) because there was no information available at the level of analysis and for the type of ecosystems assessed. Secondly, some indicators that are used here to assess a specific ES could in fact be related with more than one ES. For instance, mushroom production is often associated with cultural values while we use it only as an indicator of food provision, and SOC can be also related with climate regulation, although its relationship with forest soil fertility is clearer (Chiti et al., 2012; Rodríguez-Loinaz et al., 2015). Our set of ES indicators represents a heterogeneous ensemble with large differences regarding their sources, calculation methods and spatial patterns. Approaches based on the application of a single, spatially explicit model to estimate forest ES are likely to be more robust and allow more flexibility in accounting for management or developing scenarios, but are normally limited to one or a few ES (e.g. Frank et al., 2015; Triviño et al., 2015; Vauhkonen and Ruotsalainen, 2017). On the other hand, combining data from different sources allows to cover a wider range of ES, potentially resulting in more complete assessments (i.e. Martínez-Harms and Balvanera, 2012; Martínez-Harms et al., 2016). In addition, the ES showed some differences between the temporal periods assessed. This limitation is not easy to avoid when heterogeneous data sources are used, and in our case we prioritized obtaining the best data available for the processes analysed in the study area, even at the cost of small temporal mismatches. Finally, some of the ES estimated here are based on previously tested (and published) ecological models (mushroom production, water exported, etc.), while others are based on simpler approaches using detailed LULC maps (riparian forest, erosion control or Natura 2000). Although we did not assess the accuracy or the uncertainty associated with these last data (Müller and Burkhard, 2012), they are based on best available information that provides an accurate representation of the studied landscape (LCMC, 2009).

Regarding the spatial level of analysis, although the municipality level is often used in ES assessments (e.g., Rodríguez-Loinaz et al., 2015). Municipalities do not necessarily correspond to physical units in terms of environmental characteristics, forest distribution or forest function. Although more detailed, spatially explicit analyses are possible in some cases (e.g. erosion control in Guerra et al., 2016; catchment-level analyses for water-related ES in Stürck et al., 2014), this aggregation at municipality level has several advantages, as it allows: i) focusing the analysis on areas with a significant forest cover; ii) combining plot-level data (from NFI plots by calculation their average value) with other types of spatial information; iii) using administrative information that is not available (or meaningful) at more detailed spatial scales; and iv) an explicit link to the administrative level where most management strategies and land-use policies are decided and applied in general (Kroll et al., 2012; Ariza-Montobbio et al.,

2014), and also in the study area. In addition, recent work in the same study area reports similar spatial patterns of ES at the municipality compared to finer (1x1 km) resolutions (Roces-Díaz et al., 2018).

#### **4.2. Trade-offs and socioeconomic and environmental determinants of ES distribution**

Land-based and forest-based indicators showed broadly similar spatial patterns, but important differences were detected in the analysis of trade-offs and synergies among ES using both approaches, which highlighted their complementary character. Positive and significant relations among ES categories were frequently obtained when using the land-based indicators but not always when using forest-based ones (Table 2). In addition, significant trade-offs between ES appeared only when forest-based indicators were compared (Table S3.2). Thus, referring ES to the total municipal area masked some of the relationships that were detected when they were expressed per unit of forest area. The selection of the type of indicator is likely context-dependant and, thus, deciding which one is more suitable based on first principles appears difficult. However, forest-based indicators better reflect the intrinsic properties of forests and therefore appear more appropriate when the aim of the study is to identify the fundamental trade-offs between different ES.

It is generally accepted that high biodiversity levels are associated to high levels of ES supply (Egoh et al., 2009; Gamfeldt et al., 2013). We used birds and woody species richness as biodiversity descriptors. Although this type of approach may be problematic because richness does not reflect changes in the abundance of species (Van Strien et al., 2012), there is no doubt that species richness is an important determinant of forest ecosystem function (e.g., Vila et al., 2007; Liang et al., 2016). Consistent with previous studies, our results showed positive correlations of provisioning and regulating services with woody species and bird richness (Table 3).

Climatic conditions (mean temperature and annual rainfall) were the main determinants of the spatial variation of the ES analyzed here (Tables 2 and 3 and Table S3.3), including the distribution of hotspots. That is a logical consequence of the relationships between ecological processes and climate, particularly in water-limited regions, but it could be also related with the fact that some of the ecological models used in this work (e.g., Doblas-Miranda et al., 2013; De Cáceres et al., 2015) use climatic conditions as drivers. For example, in the case of the soil water balance model, the exported water (P3) cannot be larger than rainfall

Cultural services often show a high demand from urban populations (Martin-López et al., 2012), which in the study area are closely associated to densely populated municipalities. Considering that all our cultural ES indicators measure ES use (actual supply), it is not surprising that two of them were positively related with population density (Table 3). In addition, animal observations

showed clear positive relations with population density, in agreement with previous studies (Plieninger et al., 2013). The fact that three out of four ES showed positive relations with the percentage of people employed in the tertiary sector (Table 3) likely reflects the economic importance of tourism in the study area (Gary and Cànoves, 2011). Higher supply of cultural ES in forests surrounding urban areas could respond to the demand of these ES from people living in areas where contact with natural ecosystems is often limited (Daniel et al., 2012). Finally, negative relationships between provisioning services from agricultural agroecosystems and other ES are common (Haines-Young et al., 2011; Lee and Lautenbach, 2016), including those provided by forests (Rodriguez et al., 2007).

## **5. Conclusions**

Our results provide a picture of the current supply of several ES by the forests of Catalonia. The integration of information from different sources allowed an assessment of these ES that overcomes some of the limitations and uncertainties derived from approaches based exclusively on land use/cover data. In addition, the calculation of all these ES at the municipal level allowed an analysis that can directly inform management. Although many ES showed highest values in the mountainous and wet areas of the North of the study area, the distribution patterns of some ES and biodiversity variables support a high ecological value of forest-dominated landscapes located close to the Mediterranean coast. The relevance of these landscapes in the Mediterranean Region has been highlighted in previous studies, both from the perspective of biodiversity (e.g., Fattorini et al., 2015) and using ES-based approaches (e.g., Brenner et al., 2010), and may require special attention in conservation strategies and ES management.

Additional research is required to assess possible changes in ES provision as a result of climate change in Mediterranean forested areas. Most of the ES indicators developed in this study allow for an explicit analysis of recent temporal trends (cf. Rodríguez-Loinaz et al., 2015) and can be included in structured forest dynamics models (e.g., de Cáceres et al. 2015). This information, together with the use of state-of-the-art ecological models should improve our capacity to forecast changes in the supply of ES under different climatic and socioeconomic scenarios.

## **Acknowledgments**

We thank to the volunteers from the Catalan Ornithological Institute (ICO) and Wikiloc for providing data for the analyses presented in this study. Funding was obtained from the Catalan Office for Climate Change (OCCC) through project ForESMap, from EU FORESTERRA program (INFORMED project) and from the Spanish government (CGL2013-46808-R, AGL2015-66001-C3-1-R and CGL2014-59742). We also thank the ECOMETAS (CGL2014-53840-REDT) network for support. This study also received funding from the European Union's Horizon 2020 research and innovation programme within the framework of the MultiFUNGtionality Marie Skłodowska-Curie Individual Fellowship (IF-EF) under

grant agreement No655815 and from the Generalitat de Catalunya (Serra-Hunter Fellow). JVRD was supported by the Government of Asturias and the FP7-Marie Curie-COFUND program of the European Commission (Grant ‘Clarín’ ACA17-02). We thank Gabriel Borrás and Gemma Cantos (OCCC) for useful discussion during the elaboration of this work. We are very grateful to all persons who made the two Spanish Forest Inventories possible and, especially, to their main coordinators, Ramon Villaescusa (IFN2) and Jose Antonio Villanueva (IFN3). We also thank four anonymous reviewers for their useful comments, which helped us improve the quality of the manuscript.

## References

- ACDC, Atlas Climático de Cataluña. Departament de Geografia. Unitat de Botànica. Universitat Autònoma de Barcelona. Consulta on-line (05/05/2016). [http://www.opengis.uab.cat/acdc/en\\_index.htm](http://www.opengis.uab.cat/acdc/en_index.htm)
- Alamgir, M., Turton, S.M., Macgregor, C.J., Pert, P.L., 2016. Ecosystem services capacity across heterogeneous forest types: understanding the interactions and suggesting pathways for sustaining multiple ecosystem services. *Sci. Total Environ.* 566, 584-595. doi: 10.1016/j.scitotenv.2016.05.107
- Andersson, E., McPhearson, T., Kremer, P., Gomez-Baggethun, E., Haase, D., Tuvendal, M., Wurster, D., 2014. Scale and context dependence of ecosystem service providing units. *Ecosyst. Serv.* 15, 157-164. doi 10.1016/j.ecoser.2014.08.001
- Andrew, M.E., Wulder, M.A., Nelson, T.A., Coops, N.C., 2015. Spatial data, analysis approaches, and information needs for spatial ecosystem service assessments: a review. *GIScience Remote Sens.* 52, 344–373. doi:10.1080/15481603.2015.1033809
- Ariza-Montobbio, P., Farrell, K.N., Gamboa, G., Ramos-Martin, J., 2014. Integrating energy and land-use planning: socio-metabolic profiles along the rural – urban continuum in Catalonia (Spain). *Environ. Dev. Sustain.* 16, 925–956. doi:10.1007/s10668-014-9533-x
- Boerema, A., Rebelo, A.J., Bodi, M.B., Esler, K.J., Meire, P., 2016. Are ecosystem services adequately quantified? *J. Appl. Ecol.* doi:10.1111/1365-2664.12696
- Bonet, J.A., Palahí, M., Colinas, C., Pukkala, T., Fischer, C.R., Miina, J., Martínez de Aragón, J., 2010. Modelling the production and species richness of wild mushrooms in pine forests of the Central Pyrenees in northeastern Spain. *Can. J. For. Res.* 40, 347–356. doi:10.1139/X09-198
- Brenner, J., Jiménez, J.A., Sardá, R., Garola, A., 2010. An assessment of the non-market value of the ecosystem services provided by the Catalan coastal zone, Spain. *Ocean Coast. Manag.* 53, 27–38. doi:10.1016/j.ocecoaman.2009.10.008
- Brooks, T.M., Mittermeier, R.A., da Fonseca, G.A.B., Gerlach, J., Hoffmann, M., Lamoreux, J.F., Mittermeier, C.G., Pilgrim, J.D., Rodrigues, A.S.L., 2006. Global Biodiversity Conservation Priorities. *Science*, 313, 58–61. doi:10.1126/science.1127609
- Bugalho, M.N., Caldeira, M.C., Pereira, J.S., Aronson, J., Pausas, J.G., 2011. Mediterranean cork oak savannas require human use to sustain biodiversity and ecosystem services. *Front. Ecol. Environ.* 9, 278–286. doi:10.1890/100084
- Burkhard, B., Kroll, F., Nedkov, S., Müller, F., 2012. Mapping ecosystem service supply, demand and budgets. *Ecol. Indic.* 21, 17-29. doi:10.1016/j.ecolind.2011.06.019
- Chiti, T., Díaz-Pinés, E., Rubio, A., 2012. Soil organic carbon stocks of conifers, broadleaf and evergreen broadleaf forests of Spain. *Biol. Fertil. Soils* 48, 817–826. doi:10.1007/s00374-012-0676-3
- Costanza, R., Arge, R., Groot, R. De, Farberk, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., Neill, R.V.O., Paruelo, J., Raskin, R.G., Suttonk, P., 1997. The value of the world ' s ecosystem services and natural capital. *Nature* 387, 253–260.
- Daniel, T.C., Muhar, A., Arnberger, A., Aznar, O., Boyd, J.W., Chan, K.M. a., Costanza, R., Elmqvist, T., Flint, C.G., Gobster, P.H., Gret-Regamey, A., Lave, R., Muhar, S., Penker, M., Ribe, R.G., Schauppenlehner, T., Sikor, T., Soloviy, I., Spierenburg, M., Taczanowska, K., Tam, J., von der Dunk, A., 2012. Contributions of cultural services to the ecosystem services agenda. *Proc. Natl. Acad. Sci.* 109, 8812–8819. doi:10.1073/pnas.1114773109
- De Cáceres, M., Martínez-Vilalta, J., Coll, L., Llorens, P., Casals, P., Poyatos, R., Brotons, L. 2015. Coupling a water balance model with forest inventory data to predict drought stress: the role of forest structural changes vs. climate changes. *Agricultural and Forest Meteorology.* 213: 77-90.
- de-Miguel, S., Bonet, J. A., Pukkala, T., Martínez de Aragón, J. 2014. Impact of forest management intensity on landscape-level mushroom productivity: A regional model-based scenario analysis. *For. Ecol. Manage.* 330, 218–227. <http://doi.org/10.1016/j.foreco.2014.07.014>

- Doblas-Miranda, E., Martínez-Vilalta, J., Lloret, F., Alvarez, A., Avila, A., Bonet, F.J., Brotons, L., Castro, J., Curiel Yuste, J., Diaz, M., Ferrandis, P., García-Hurtado, E., Iriondo, J.M., Keenan, T.F., Latron, J., Llusia, J., Loepfe, L., Mayol, M., More, G., Moya, D., Peñuelas, J., Pons, X., Poyatos, R., Sardans, J., Sus, O., Vallejo, V.R., Vayreda, J., Retana, J., 2015. Reassessing global change research priorities in mediterranean terrestrial ecosystems: How far have we come and where do we go from here? *Glob. Ecol. Biogeogr.* 24, 25–43. doi:10.1111/geb.12224
- Doblas-Miranda, E., Rovira, P., Brotons, L., Martínez-Vilalta, J., Retana, J., Pla, M., Vayreda, J. 2013. Soil carbon stocks and their variability across the forests, shrublands and grasslands of peninsular Spain. *Biogeosciences*, 10(12), 8353–8361. <http://doi.org/10.5194/bg-10-8353-2013>
- Duncker, P.S., Raulund-rasmussen, K., Gundersen, P., Katzensteiner, K., Jong, J. De, Peter, H., 2012. How Forest Management affects Ecosystem Services, including Timber Production and Economic Return : Synergies and Trade-Offs. *Ecol. Soc.* 17 (40), 50.
- Egoh, B., Reyers, B., Rouget, M., Bode, M., Richardson, D., 2009. Spatial congruence between biodiversity and ecosystem services in South Africa. *Biol. Conserv.* 142, 553–562. doi:10.1016/j.biocon.2008.11.009
- Eigenbrod, F., Armsworth, P.R., Anderson, B.J., Heinemeyer, A., Gillings, S., Roy, D.B., Thomas, C.D., Gaston, K.J., 2010. The impact of proxy-based methods on mapping the distribution of ecosystem services. *J. Appl. Ecol.* 47, 377–385. doi:10.1111/j.1365-2664.2010.01777.x
- EME, 2011. Evaluación de los Ecosistemas del Milenio en España, Ecosistemas y Biodiversidad para el Bienestar Humano, Evaluación de los Ecosistemas del Milenio en España, Síntesis de resultados. Fundación Biodiversidad. Ministerio de Medio Ambiente, y Medio Rural y Marino
- ESRI, 2011. ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute.
- ESRI, 2013a. How Hot Spot Analysis (Getis-Ord Gi\*) works, ARCGIS Resources, ArcGIS 10 Help. [online 15 July 2015] URL: [http://resources.arcgis.com/en/help/main/10.1/index.html#/How\\_Hot\\_Spot\\_Analysis\\_Getis\\_Ord\\_Gi\\_works/005p00000011000000/](http://resources.arcgis.com/en/help/main/10.1/index.html#/How_Hot_Spot_Analysis_Getis_Ord_Gi_works/005p00000011000000/)
- ESRI, 2013b. How Spatial Autocorrelation (Global Moran's I) works. [online] URL: <http://resources.arcgis.com/en/help/main/10.1/index.html#/005p0000000t000000>
- Estrada, J., Pedrocchi, V., Brotons, L., Herrando, S., (eds). 2004. *Atles dels Ocells Nidificants de Catalunya 1999-2002*. Institut Català d'Ornitologia (ICO)/Lynx Edicions, Barcelona
- FAO, Food and Agriculture Organization of the United Nations. 2014. *State of the World's Forests: Enhancing the socioeconomic benefits from forests*. Rome. ISSN 1020-5705. 119 pp.
- Fattorini, S., Maltzef, P., Salvati, L., 2015. Use of insect distribution across landscape-soil units to assess conservation priorities in a Mediterranean coastal reserve: the tenebrionid beetles of Castelporziano (Central Italy). *Rend. Lincei* 26, 353–366. doi:10.1007/s12210-015-0391-8
- Frank, S., Fürst, C., Pietzsch, F., 2015. Cross-Sectoral Resource Management: How Forest Management Alternatives Affect the Provision of Biomass and Other Ecosystem Services. *Forests* 6, 533–560. doi:10.3390/f6030533
- Gamfeldt, L., Snäll, T., Bagchi, R., Jonsson, M., Gustafsson, L., Kjellander, P., Ruiz-Jaen, M.C., Fröberg, M., Stendahl, J., Philipson, C.D., Mikusiński, G., Andersson, E., Westerlund, B., Andrén, H., Moberg, F., Moen, J., Bengtsson, J., 2013. Higher levels of multiple ecosystem services are found in forests with more tree species. *Nat. Commun.* 4, 1340. doi:10.1038/ncomms2328
- García-Llorente, M., Iniesta-arandia, I., Willaarts, B.A., Harrison, P.A., Berry, P., Bayo, M. del M., Castro, A.J., Montes, C., Martín-López, B., 2015. Biophysical and sociocultural factors underlying spatial trade-offs of ecosystem services in semiarid watersheds. *Ecol. Soc.* 20, 39. doi:10.5751/ES-07785-200339
- García-Nieto, A.P., García-Llorente, M., Iniesta-Arandia, I., Martín-López, B., 2013. Mapping forest ecosystem services: From providing units to beneficiaries. *Ecosyst. Serv.* 4, 126–138. doi:10.1016/j.ecoser.2013.03.003
- Gary, L., Cànoves, G., 2011. Life cycles, stages and tourism history. The Catalonia (Spain) experience. *Annals of Tourism Research*, 38, 651-671.
- Getis, A., Ord J.K., 1992. The Analysis of Spatial Association. *Geogr Anal* 24:189–206. doi:10.1111/j.1538-4632.1992.tb00261.x
- Grove, T., Rackham, O. 2003. *The Nature of Mediterranean Europe. An Ecological History*. Yale University Press, New Haven
- Guerra, C.A., Maes, J., Geijzendorffer, I., Metzger, M.J., 2016. An assessment of soil erosion prevention by vegetation in Mediterranean Europe: Current trends of ecosystem service provision. *Ecol. Indic.* 60, 213–222. doi:10.1016/j.ecolind.2015.06.043
- Haines-Young, R., Potschin, M., 2013. Towards a common International Classification of Ecosystem Services (CICES): Version 4.3 [online 15 July 2015] URL: [www.cices.eu/](http://www.cices.eu/)

- Haines-Young, R., Potschin, M., Kienast, F., 2011. Indicators of ecosystem service potential at European scales: Mapping marginal changes and trade-offs. *Ecol. Indic.* doi:10.1016/j.ecolind.2011.09.004
- Hampe, A., Petit, R.J., 2005. Conserving biodiversity under climate change: The rear edge matters. *Ecol. Lett.* 8, 461–467. doi:10.1111/j.1461-0248.2005.00739.x
- Hein, L., van Koppen, K., de Groot, R.S., van Ierland, E.C., 2006. Spatial scales, stakeholders and the valuation of ecosystem services. *Ecol. Econ.* 57, 209–228. doi:10.1016/j.ecolecon.2005.04.005
- Hou, Y., Burkhard, B., Müller, F., 2013. Uncertainties in landscape analysis and ecosystem service assessment. *J. Environ. Manage.* doi:10.1016/j.jenvman.2012.12.002
- IDESCAT, Institut d'Estadística de Catalunya. 2015. Información demográfica y económica por municipios. Consulta on-line. <http://www.idescat.cat/es/>
- Kroll, F., Muller, F., Haase, D., Fohrer, N., 2012. Rural-urban gradient analysis of ecosystem services supply and demand dynamics. *Land use policy* 29, 521–535. doi:10.1016/j.landusepol.2011.07.008
- LCMC, Land Cover Map of Catalonia, 2009. Generalitat de Catalunya. CREAL, Universidad Autónoma de Barcelona. <http://www.creaf.uab.es/mcsc/esp/index.htm>
- Lee, H., Lautenbach, S., 2016. A quantitative review of relationships between ecosystem services. *Ecol. Indic.* 66, 340–351. doi:10.1016/j.ecolind.2016.02.004
- Liang, J., Crowther, T.W., Picard, N., Wiser, S., Zhou, M., Alberti, G., Schulze, E.-D., McGuire, D., Bozzato, F., Pretzsch, H., De-Miguel, S., Paquette, A., Hérault, B., Scherer-Lorenzen, M., Barrett, C.B., Glick, H.B., Hengeveld, G.M., Nabuurs, G.-J., Pfautsch, S., Viana, H., Vibrans, A.C., Ammer, C., Schall, P., Verbyla, D., Tchebakova, N., Fischer, M., Watson, J. V., Chen, H.Y.H., Lei, X., Schelhaas, M.-J., Lu, H., Gianelle, D., Parfenova, E.I., Salas, C., Lee, E., Lee, B., Kim, H.S., Bruelheide, H., Coomes, D.A., Piotta, D., Sunderland, T., Schmid, B., Gourlet-Fleury, S., Sonké, B., Tavani, R., Zhu, J., Brandl, S., Vayreda, J., Kitahara, F., Searle, E.B., Neldner, V.J., Ngugi, M.R., Baraloto, C., Frizzera, L., Bałazy, R., Oleksyn, J., Zawila-Niedzwiecki, T., Bouriaud, O., Bussotti, F., Finér, L., Jaroszewicz, B., Jucker, T., Valladares, F., Jagodzinski, A.M., Peri, P.L., Gonmadje, C., Marthy, W., O'Brien, T., Martin, E.H., Marshall, A., Rovero, F., Bitariho, R., Niklaus, P.A., Alvarez-Loayza, P., Chamuya, N., Valencia, R., Mortier, F., Wortel, V., Engone-Obiang, N.L., Ferreira, L. V., Odeke, D.E., Vasquez, R.M., Reich, P.B., 2016. Positive biodiversity–productivity relationship predominant in global forests. *Science* (80-. ). 354, 196–208. doi:10.1126/science.aaf8957
- MAGRAMA, Ministerio de Agricultura, Alimentación y Medio Ambiente. 1997-2007. Segundo y Tercer Inventario Forestal Nacional. Gobierno de España. [online 15 July 2015] URL: [http://www.magrama.gob.es/es/biodiversidad/servicios/banco-datos-naturaleza/informacion-disponible/index\\_inventario\\_forestal.aspx](http://www.magrama.gob.es/es/biodiversidad/servicios/banco-datos-naturaleza/informacion-disponible/index_inventario_forestal.aspx)
- Martínez-Harms, M.J., Balvanera, P., 2012. Methods for mapping ecosystem service supply: A review. *Int. J. Biodivers. Sci. Ecosyst. Serv. Manag.* 8, 17–25. doi:10.1080/21513732.2012.663792
- Martínez-Harms, M.J., Quijas, S., Merenlender, A.M., Balvanera, P., 2016. Enhancing ecosystem services maps combining field and environmental data. *Ecosyst. Serv.* 22, 32–40. doi:10.1016/j.ecoser.2016.09.007
- Martínez-Vilalta, J., Piñol, J., 2002. Drought-induced mortality and hydraulic architecture in pine populations of the NE Iberian Peninsula. *For. Ecol. Manage.* 161, 247–256. doi:10.1016/S0378-1127(01)00495-9
- Martín-López, B., Iniesta-Arandia, I., García-Llorente, M., Palomo, I., Casado-Arzuaga, I., Del Amo, D.G., Gómez-Baggethun, E., Oteros-Rozas, E., Palacios-Agundez, I., Willaarts, B., González, J.A., Santos-Martín, F., Onaindia, M., López-Santiago, C., Montes, C., 2012. Uncovering ecosystem service bundles through social preferences. *PLoS One* 7. doi:10.1371/journal.pone.0038970
- MEA, 2005. *Ecosystems and Human Well-being: Current State and Trends*. Island Press, Washington, DC Millennium Ecosystem Assessment.
- Medail, F., Quezel, P., 1999. Biodiversity Hotspots in the Mediterranean Basin: Setting Global Conservation Priorities. *Conserv. Biol.* 13, 1510–1513. doi:10.1046/j.1523-1739.1999.98467.x
- Miura, S., Amacher, M., Hofer, T., San-Miguel-Ayanz, J., Ernawati, Thackway, R., 2015. Protective functions and ecosystem services of global forests in the past quarter-century. *For. Ecol. Manage.* 352, 35–46. doi:10.1016/j.foreco.2015.03.039
- Moran, P.A.P., 1948. The interpretation of statistical maps. *J. R. Stat. Soc.* 10, 243–251.
- Mouchet, M. a., Lamarque, P., Martín-López, B., Crouzat, E., Gos, P., Byczek, C., Lavorel, S., 2014. An interdisciplinary methodological guide for quantifying associations between ecosystem services. *Glob. Environ. Chang.* 28, 298–308. doi:10.1016/j.gloenvcha.2014.07.012



- Müller, F., Burkhard, B., 2012. The indicator side of ecosystem services. *Ecosyst. Serv.* 1, 26–30. doi:10.1016/j.ecoser.2012.06.001
- Myers, N., Mittermeier, R.A., Mittermeier C.G., da Fonseca, G.A.B., Kent, J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858. doi:10.1038/35002501
- Peñuelas, J., Boada, M., 2003. A global change-induced biome shift in the Montseny mountains (NE Spain). *Glob. Chang. Biol.* 9, 131–140. doi:10.1046/j.1365-2486.2003.00566.x
- Plieninger, T., Dijks, S., Oteros-Rozas, E., Bieling, C., 2013. Assessing, mapping, and quantifying cultural ecosystem services at community level. *Land use policy* 33, 118–129. doi:10.1016/j.landusepol.2012.12.013
- Plummer, M.L., 2009. Assessing benefit transfer for the valuation of ecosystem services. *Frontiers in Ecology and the Environment*, 7, 38–45.
- Quintas-Soriano, C., Castro, A.J., García-Llorente, M., Cabello, J., Castro, H., 2014. From supply to social demand: a landscape-scale analysis of the water regulation service. *Landsc. Ecol.* 29, 1069–1082. doi:10.1007/s10980-014-0032-0
- R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2014. Retrieved April 10, 2014, from [www.r-project.org/](http://www.r-project.org/)
- Roces-Díaz, J.V., Burkhard, B., Kruse, M., Muller, F., Álvarez-Álvarez, P., Diaz-Varela, E. 2017. Use of ecosystem information derived from forest thematic maps for spatial analysis of ecosystem services in northwestern Spain. *Lands. Ecol. Eng.* 13, 45–57. doi:10.1007/s11355-016-0298-2
- Roces-Díaz, J.V., Vayreda, J., Banqué-Casanovas, M., Díaz-Varela, E., Bonet, J.A., Brotons, L., de-Miguel, S., Herrando, S., Martínez-Vilalta, J., 2018. The spatial level of analysis affects the patterns of forest ecosystem services supply and their relationships. *Sci. Total Environ.* 626. doi:10.1016/j.scitotenv.2018.01.150
- Rodríguez, J.P., Beard, T.D., Bennett, E.M., Cumming, G.S., Cork, S.J., Agard, J., Dobson, A.P., Peterson, G.D., 2007. Trade-offs across Space, Time, and Ecosystem Services. *Ecology & Society* 11, 28.
- Rodríguez-Loinaz, G., Alday, J. G., Onaindia, M. 2015. Multiple ecosystem services landscape index: A tool for multifunctional landscapes conservation. *Journal of Environmental Management*, 147, 152–163. <http://doi.org/10.1016/j.jenvman.2014.09.001>
- Ruiz-Benito, P., Gomez-Aparicio, L., Paquette, A., Messier, C., Kattge, J., Zavala, M.A., 2014. Diversity increases carbon storage and tree productivity in Spanish forests. *Glob. Ecol. Biogeogr.* 23, 311–322. doi:10.1111/geb.12126
- Schröter, M., Barton, D.N., Remme, R.P., Hein, L., 2014. Accounting for capacity and flow of ecosystem services: A conceptual model and a case study for Telemark, Norway. *Ecol. Indic.* 36, 539–551. doi:10.1016/j.ecolind.2013.09.018
- Schröter, M., Remme, R.P. 2016. Spatial prioritisation for conserving ecosystem services: comparing hotspots with heuristic optimisation. *Landsc. Ecol.* 31, 431–450. doi: 10.1007/s10980-015-0258-5
- Sutherland, I.J., Gergel, S.E., Bennett, E.M., 2016. Seeing the forest for its multiple ecosystem services: Indicators for cultural services in heterogeneous forests. *Ecol. Indic.* 71, 123–133. doi: 10.1016/j.ecolind.2016.06.037
- Sutherland, I.J., Villamagna, A.M., Dallaire, C.O., Bennett, E.M., Chin, A.T., Yeung, A.C., Lamothe, K.A., Tomscha, S.A., Cormier, R., 2017. Undervalued and under pressure: A plea for greater attention toward regulating ecosystem services. *Ecol. Indic.* (in press). doi: 10.1016/j.ecolind.2017.06.047.
- Stürck, J., Poortinga, A., Verburg, P.H., 2014. Mapping ecosystem services: The supply and demand of flood regulation services in Europe. *Ecol. Indic.* 38, 198–211. doi:10.1016/j.ecolind.2013.11.010
- Thom, D., Seidl, R., 2015. Natural disturbance impacts on ecosystem services and biodiversity in temperate and boreal forests. *Biol. Rev.* doi:10.1111/brv.12193
- Triviño, M., Juutinen, A., Mazziotta, A., Miettinen, K., Podkopaev, D., Reunanen, P., Mönkkönen, M., 2015. Managing a boreal forest landscape for providing timber, storing and sequestering carbon. *Ecosyst. Serv.* 14, 179–189. doi:10.1016/j.ecoser.2015.02.003
- Trumbore, S., Brando, P., Hartmann, H., 2015. Forest health and global change. *Science.* 349, 814–818. doi:10.1126/science.aac6759
- Underwood, E.C., Viers, J.H., Klausmeyer, K.R., Cox, R.L., Shaw, M.R., 2009. Threats and biodiversity in the mediterranean biome. *Divers. Distrib.* 15, 188–197. doi:10.1111/j.1472-4642.2008.00518.x
- Van der Biest, K., Vrebos, D., Staes, J., Boerema, A., Bodi, M.B., Franssen, E., Meire, P., 2015. Evaluation of the accuracy of land-use based ecosystem service assessments for different thematic resolutions. *J. Environ. Manage.* 156, 41–51. doi:10.1016/j.jenvman.2015.03.018

- Van Strien, A.J., Soldaat, L.L., Gregory, R.D., 2012. Desirable mathematical properties of indicators for biodiversity change. *Ecol. Indic.* 14, 202–208. doi:10.1016/j.ecolind.2011.07.007
- Vauhkonen, J., Ruotsalainen, R., 2017. Assessing the provisioning potential of ecosystem services in a Scandinavian boreal forest: Suitability and tradeoff analyses on grid-based wall-to-wall forest inventory data. *For. Ecol. Manage.* 389, 272–284. doi:10.1016/j.foreco.2016.12.005
- Vayreda, J., Martinez-Vilalta, J., Gracia, M., Retana, J., 2012. Recent climate changes interact with stand structure and management to determine changes in tree carbon stocks in Spanish forests. *Glob. Chang. Biol.* 18, 1028–1041. doi:10.1111/j.1365-2486.2011.02606.x
- Vila, M., Basnou, C., Pysek, P., Josefsson, M., Genovesi, P., Gollasch, S., Nentwig, W., Olenin, S., Roques, A., Roy, D., Hulme, P.E., 2010. How well do we understand the impacts of alien species on ecosystem services? A pan-European, cross-taxa assessment. *Front. Ecol. Environ.* 8, 135–144. doi:10.1890/080083
- Vila, M., Vayreda, J., Comas, L., Ibañez, J.J., Mata, T., Obon, B., 2007. Species richness and wood production: A positive association in Mediterranean forests. *Ecol. Lett.* 10, 241–250. doi:10.1111/j.1461-0248.2007.01016.x
- Yahdjian, L., Sala, O.E., Havstad, K.M., 2015. Rangeland ecosystem services: Shifting focus from supply to reconciling supply and demand. *Front. Ecol. Environ.* 13, 44–51. doi:10.1890/140156

## Supplementary material

### *Ecological Indicators*

#### **Assessing the distribution of forest ecosystem services in a highly populated Mediterranean region**

Jose V. Rocés-Díaz, Jordi Vayreda, Mireia Banqué-Casanovas, Martí Cusó, Marc Anton, José A. Bonet, Lluís Brotons, Miquel De Cáceres, Sergi Herrando, Juan Martínez de Aragón, Sergio de-Miguel, Jordi Martínez-Vilalta

#### **Appendix S1. Detailed description of the data sources used in this work**

We have assessed ES using different approaches. Most ES were mapped using previously built models; others ES were calculated using primary data from NFI plots; for both types, ES were calculated at point level and then aggregated at municipality scale. Others ES were estimated based on combing datasets at municipality levels such as aerial photo interpretation and GIS analysis, territorial statistic, information from different databased (including unofficial webpages), etc. The resulting ES maps were then aggregated to municipal units and obtained values were compared to explore their potential trade-offs and synergies between pairs of ES. Finally these values were analysed against climatic and socioeconomic conditions of the studied municipalities.

- The indicator for **climate regulation (R1: Carbon sequestration)** was based in the Spanish National Forest Inventory directly. Other indicators (mushroom production (P1) for food provision ES; water exported (P3) for water provision ES; water storage capacity (R3) for water regulation ES) were based on models that used this data source as an input. The Spanish National Forest Inventory (NFI) is an intensive national database of periodical forest surveys distributed systematically across the forested area of Spain (Villaescusa & Díaz, 1998; Villanueva 2005). The IFN is based on a network of circular plots at a density of 1 plot per 100 ha, which allows forest characterization and includes exhaustive information on the composition of canopy and understory woody species, as well as on forest structure and production. Tree sampling followed a nested design, that is, plot size depends on the diameter at breast height (DBH) of the measured trees to guarantee a representative sampling of the tree size distribution. Thus, all trees with  $DBH \geq 7.5$  cm were measured within 5 m of the centre of the plots, trees with  $DBH \geq 12.5$  cm were also measured between 5 and 10 m around the centre of the plots, whereas trees with  $DBH \geq 22.5$  cm and  $DBH \geq 42.5$  cm were also considered within 10–15 m and 15–25 m around the centre of the plots, respectively. Species identity of all living and standing dead trees was recorded and its height (H) and DBH

were measured. Individual plots are resampled every approximately 10 years. In Catalonia, the 2<sup>nd</sup> NFI was conducted in 1989-1990 and the 3<sup>rd</sup> one (NFI3) in 2000-2001.

*Reference:* MAGRAMA, Ministerio de Agricultura, Alimentación y Medio Ambiente. 1997-2007. Segundo y Tercer Inventario Forestal Nacional. Gobierno de España. [online 15 July 2015] URL: [http://www.magrama.gob.es/es/biodiversidad/servicios/banco-datos-naturaleza/informacion-disponible/index\\_inventario\\_forestal.aspx](http://www.magrama.gob.es/es/biodiversidad/servicios/banco-datos-naturaleza/informacion-disponible/index_inventario_forestal.aspx)

- **Food provision ES. Indicator P1: Mushroom production.** Mushroom production estimates (i.e., food provisioning ES) were based on the mixed-effects models provided by de-Miguel et al. (2014), focusing on edible mushrooms of commercial interest, i.e., widely consumed by people in the study region. These models were developed based on the monitoring, from 1995 to 2012, of weekly mushroom production from permanent sample plots representing most pine forest ecosystems throughout the study region, i.e., pure and mixed stands of *Pinus sylvestris*, *P. nigra*, *P. halepensis* and *P. pinaster*. The models allow for estimating mushroom production for a typical year based on site and stand characteristics. The site conditions driving mushroom production in the models are determined by the elevation above sea level (i.e., a surrogate for typical meteorological conditions) as well as by the interaction between aspect and slope. The models predict increasing mushroom production with increasing elevation and northern aspect. The stand characteristics affecting mushroom yield in the models are both stand composition (i.e., main tree species) and stand structure as described by the stand basal area. *P. sylvestris* stands are expected to provide the highest mushroom production, and the optimal stand basal area maximizing mushroom production is approximately 20 m<sup>2</sup> ha<sup>-1</sup>, although it fluctuates between pine ecosystems.

*Reference:* de-Miguel, S., Bonet, J. A., Pukkala, T., Martínez de Aragón, J. 2014. Impact of forest management intensity on landscape-level mushroom productivity: A regional model-based scenario analysis. *For. Ecol. Manage.* 330, 218–227.

- **Water provision ES and Water regulation ES. Indicators P3: Water exported and R3: Water storage capacity.** Both indicators were provided by **Water balance model**. The purpose of the water balance model presented in De Cáceres et al. (2015) is to predict temporal variations in soil water content and assess drought stress for plants in forest stands. The model has been designed to run in forest inventory plots and it calculates daily water balance for a period given by meteorological input. Soil is represented using two layers and the model keeps track of soil moisture for each layer. Soil water holding capacity includes the effects of soil texture and rock fragment

content. Vegetation is represented as a set of plant cohorts having different height, root distribution, species identity and leaf area index. Every day the model first updates leaf area of deciduous plants according to a simple phenological model. Then, the model recalculates light extinction through the canopy and the water storage capacity of the canopy. For the present study, water storage capacity was calculated assuming full leaf development. After updating the canopy status, the model deals with the water input from rainfall. Before increasing the water content of soil layers, the model subtracts the water lost due to interception and the water lost through surface runoff from rainfall. When refilling a given soil layer, a proportion of water is assumed to directly percolate to the next layer below, and the water percolating from the deepest layer is assumed to be lost via deep drainage. After refilling soil layers, the model determines evaporation from the soil surface and plant transpiration, which depend on potential evapotranspiration, leaf area and soil water potential. More details on the design, calibration and validation of the model can be found in De Cáceres *et al.* (2015). Daily water balance was simulated on all NFI3 plots for year 2010 (details of the simulations can be found in De Cáceres *et al.* 2015). For the present study, exported water (in  $L \cdot m^{-2} \cdot year^{-1}$ ) of each NFI3 plot was defined as the sum (across all days in 2010) of the water predicted to be lost via surface runoff or deep drainage.

*Reference: De Cáceres, M., Martínez-Vilalta, J., Coll, L., Llorens, P., Casals, P., Poyatos, R., Brotons, L. 2015. Coupling a water balance model with forest inventory data to predict drought stress: the role of forest structural changes vs. climate changes. Agricultural and Forest Meteorology. 213: 77-90.*

- **Soil fertility ES. Indicator R2: Organic soil Carbon.** This information is based on a map of Soil Organic Carbon (SOC) of Spanish forests (Doblas-Miranda *et al.*, 2013). This map was based on measurements in more than 900 soil profiles. SOC to a depth of 1m was modelled as a function of vegetation, climatic variables, elevation, and other variables. The statistical model obtained was used to estimate SOC on a grid with a resolution of 200x200 m.

*Reference: Doblas-Miranda, E., Rovira, P., Brotons, L., Martínez-Vilalta, J., Retana, J., Pla, M., Vayreda, J., 2013. Soil carbon stocks and their variability across the forests, shrublands and grasslands of peninsular Spain. Biogeosciences 10, 8353–8361. doi:10.5194/bg-10-8353-2013*

- **Flood protection and Erosion control ES. Indicators R4: Riparian forest cover and R5: Forest cover in areas with high slopes.** The source for these indicators was **Land Cover Map of Catalonia (LCMC, 2009)**. The Land Cover Map of Catalonia (LCMC) is a high resolution thematic cartography of the main land covers of Catalonia. It is a vectorial map generated by photo-interpreting on 1:5000 colour ortho-photomaps from

the year 2005 provided by the Cartographic Institute of Catalonia. The minimum patch area was 500 m<sup>2</sup>, corresponding to the minimum mapping area of the LCMC, and the working scale is 1:1000. This map has a pixel resolution of 0.25 meters and 241 different legend categories. Areas with >30% slope were determined from a Digital Elevation Model at 100 m spatial resolution.

**Reference:** LCMC, *Land Cover Map of Catalonia, 2009*. Generalitat de Catalunya. CREAf, Universidad Autònoma de Barcelona.  
<http://www.creaf.uab.es/mcsc/esp/index.htm>

Supplementary material

**Description and spatial distribution of the socioeconomic, climatic and biodiversity variables**

Table S1.1. Description of the socio-environmental variables and their sources.

<b>Variable</b>	<b>Description</b>	<b>Units</b>	<b>Year</b>
Population density	Number of inhabitants per unit of area in each municipality of the study area	Inhabitants/km <sup>2</sup>	2010
Unemployment	Rate of unemployment of each municipality	%	2010
Agriculture	Percentage of active population of each municipality employed in jobs related to agriculture	%	2010
Industry	Percentage of active population of each municipality employed in jobs related to industry	%	2010
Construction	Percentage of active population of each municipality employed in jobs related to construction	%	2010
Services	Percentage of active population of each municipality employed in jobs related to services	%	2010
Temperature	Mean annual temperature	°C	Mean
Rainfall	Mean annual accumulated rainfall	l/m <sup>2</sup>	Mean
Woody species richness	Total number of woody species per municipality	Nº species	1990
Forest Bird richness	Total number of forest bird species per municipality	Nº species	1990

**Source of climatic data**

Mean annual temperature (°C) and total annual precipitation (l/m<sup>2</sup>) were used to characterize the climate at each municipality. We calculated their average values per municipality using the data from the Digital Climatic Atlas of Catalonia (ACDC, [http://www.opengis.uab.cat/acdc/en\\_index.htm](http://www.opengis.uab.cat/acdc/en_index.htm)).

**Source of biodiversity data**

Finally we defined two biodiversity indicators: the total number of woody species recorded in every municipality and the total number of forest birds in each municipality. Woody species richness was taken from NFI3 plots, and was defined as the total number of woody species observed in all sampled plots by municipality. Richness of forest birds was provided by the Catalan Ornithological Institute (ICO), and includes all breeding forest species (according to species distribution models performed at 1x1 km) from the Atlas of breeding birds in Catalonia 1999-2002 (Estrada *et al.*, 2004). This variable was calculated using the second Catalan Breeding Bird Atlas (CBBA2; Estrada *et al.* 2004). Fieldwork for CBBA2 was conducted between the years 1999 and 2002 and its database represented the most updated data source robustly covering all municipalities in Catalonia when this study was carried out. A total of 3,077 1x1 km squares (corresponding to a stratified selection of ca. 10% of the total number of 1x1 km squares of Catalonia) were surveyed. In order to maximize the overall species detectability two one-hour surveys were done in each 1x1 km square, one in the early and one in the late breeding season, respectively. Presence/absence data for breeding bird species collected in these surveys and a number of environmental predictors were used to develop Species Distribution Models and a cross-validation procedure (70% data for calibration and 30% for validation) was applied for the assessment of model performance. Finally, species' habitat preference was classified in several categories (including forest and some other habitats) according to the information gathered in the same set of 1x1 km squares and the habitats present there.






Figure S1.1. Spatial distribution of the different socioeconomic, climatic and biodiversity variables used in this work.

#### References

- ACDC, Atlas Climático de Cataluña.. Departament de Geografia. Unitat de Botànica. Universitat Autònoma de Barcelona. Consulta on-line. [http://www.opengis.uab.cat/acdc/en\\_index.htm](http://www.opengis.uab.cat/acdc/en_index.htm)
- IDESCAT, Institut d'Estadística de Catalunya. 2015. Informació demogràfica i econòmica per municipis. Consulta on-line. <http://www.idescat.cat/es/>
- Estrada, J.; Pedrocchi, V.; Brotons, L.; Herrando, S. (eds). 2004. Atlas dels Ocells Nidificants de Catalunya 1999-2002. Institut Català d'Ornitologia (ICO)/Lynx Edicions, Barcelona
- MAGRAMA, Ministerio de Agricultura, Alimentación y Medio Ambiente. 1997-2007. Segundo y Tercer Inventario Forestal Nacional. Gobierno de España. [online 15 July 2015] URL: [http://www.magrama.gob.es/es/biodiversidad/servicios/banco-datos-naturaleza/informacion-disponible/index\\_inventario\\_forestal.aspx](http://www.magrama.gob.es/es/biodiversidad/servicios/banco-datos-naturaleza/informacion-disponible/index_inventario_forestal.aspx)

## **Appendix S2. Spatial statistics calculations**

### **Mapping ecosystem services indicators and social, climatic and biodiversity variables**

A database with the value of each indicator was built. As they were calculated at municipality level, they were joined with a spatial shape of the municipalities of the study area. Thus, we produced maps for the different ES indicators, for the types of services and for the social, climatic and environmental variables using the Geographic Information System (GIS) ArcGIS 10.2 (ESRI, 2011). The service type maps were created using the values of the individual ES indicators obtained by averaging the standardized values of the corresponding services. Similar maps were obtained for socioeconomic and climatic variables (Figure S2.1). The values in the map are classified according to the criterion of an equal amount of observations (20% quantiles: 0-20%; 20-40%; 40-60%; 60-80%; 80-100%) in a similar way to most ES geographical assessments at similar scales (Burkhard *et al.*, 2010).

### **Packages used for statistical analyses**

Correlation analyses were conducted using the "corrplot" package (Wei and Simko, 2016). PCA analysis and graphics were conducted using the "FactoMineR" package (Le *et al.*, 2008), and linear mixed models were fitted using "lme4" (Bates *et al.*, 2015) and "lmerTest" (Kuznetsova *et al.*, 2015) packages. Additionally, we followed Nakagawa and Schielzeth (2013) to calculate the proportion of variation accounted by fixed effects alone (marginal  $R^2$ ,  $R^2_m$ ) and the proportion of variation accounted by fixed and random effects together (conditional  $R^2$ ,  $R^2_c$ ) for each model and also Akaike Information Criterion (AIC), as implemented in the "MuMIn" package (Barton, 2014).

## Principal Component Analysis (PCA)




Figure S2.1. Representation on two main axes of PCA analysis.

## Statistical methods for Hotspots calculation

In order to identify areas of high and low FES supply (respectively hotspots and coldspots) in each FES map, the Getis-Ord  $G_i^*$  statistic (Getis and Ord, 1992; Ord and Getis, 1995; ESRI, 2013) was calculated (Eq.1).

$$G_i^* = \frac{\sum_{j=1}^n w_{i,j} x_j - \left[ \frac{\sum_{j=1}^n x_j}{n} \right] \sum_{j=1}^n w_{i,j}}{S \sqrt{\frac{n \sum_{j=1}^n w_{i,j}^2 - (\sum_{j=1}^n w_{i,j})^2}{n-1}}} \quad \text{Eq.1}$$

Where  $n$  is the number of features;  $w_{i,j}$  is the distance between the features  $i$  and  $j$ ;  $x_j$  is the value of potential supply of the ES, and  $S$  is calculated as follows (Eq.2):

$$S = \sqrt{\frac{\sum_{j=1}^n x_j^2}{n} - \left[ \frac{\sum_{j=1}^n x_j}{n} \right]^2} \quad \text{Eq.2}$$

For a given dataset, the Getis-Ord  $G_i^*$  statistic identifies those clusters of spatial features with values (of potential FES supply) higher (or lower) of those expected to be found by random chance (ESRI 2013a). The output of the  $G_i^*$  is a Z-score for each feature that represents the statistical significance of clustering for a specified distance. Higher Z-scores indicate higher intensity of feature value (FES supply) clustering and hotspots of FES supply. Negative Z-scores indicate clusters with low FES supply values (coldspots). The analyses were performed using ArcGIS 10.1 spatial statistic tools.

### Statistical methods for Moran's I calculation

The spatial autocorrelation between the indicators were was by using Moran's I coefficient (Eq.3; Moran 1948; ESRI 2013a).

$$I = \frac{n}{\sum_{i=1}^n \sum_{j=1}^n w_{ij}} \frac{\sum_{i=1}^n \sum_{j=1}^n w_{ij} z_i z_j}{\sum_{i=1}^n z_i^2} \quad \text{Eq.3.}$$

Where  $n$  is the number of features (points);  $w_{ij}$  is the distance between the features  $i$  and  $j$ ;  $z_i$  is the deviation of the attribute (here the value of each indicator) from feature  $i$  from the mean value.

The coefficient indicates the variation in values of one variable, based on the distance between the elements (the centroid of each municipality). As a result, Moran's I enables classification of the spatial patterns of the points on the basis of the degree of clustering or dispersion that the values of potential supply of the ES show. This index can vary from -1 (highly negative spatial autocorrelation) to +1 (highly positive autocorrelation).

### References

- Bartoń K. 2014. MuMIn: Multi-model inference. R package version 1.10.0. Retrieved May 14, 2014, from <http://cran.r-project.org/package=MuMIn>.
- Bates, D., Mächler, M., Bolker, B. M., Walker, S. C. 2015. Fitting Linear Mixed-Effects Models Using lme. *Journal of Statistical Software* 4, 67(1). <http://doi.org/10.18637/jss.v067.i01>
- Burkhard, B., Kroll, F., Müller, F. 2010. Landscapes' Capacities to Provide Ecosystem Services – a Concept for Land-Cover Based Assessments. *Landsc. Online* 1–22. doi: 10.3097/LO.200915
- ESRI, 2011. ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute.
- ESRI, 2013a. How Hot Spot Analysis (Getis-Ord  $G_i^*$ ) works, ARCGIS Resources, ArcGIS 10 Help. [online 15 July 2015] URL: [http://resources.arcgis.com/en/help/main/10.1/index.html#/How\\_Hot\\_Spot\\_Analysis\\_Getis\\_Ord\\_Gi\\_works/005p00000011000000/](http://resources.arcgis.com/en/help/main/10.1/index.html#/How_Hot_Spot_Analysis_Getis_Ord_Gi_works/005p00000011000000/)
- ESRI, 2013b How Spatial Autocorrelation (Global Moran's I) works. [online] URL: <http://resources.arcgis.com/en/help/main/10.1/index.html#/005p0000000t000000>
- Getis, A., Ord, J.K., 1992. The Analysis of Spatial Association. *Geogr Anal* 24:189–206. doi: 10.1111/j.1538-4632.1992.tb00261.x
- Kuznetsova, A., Bruun Brockhoff, P., Haubo Bojesen Christensen, R. 2015. lmerTest: Tests for random and fixed effects for linear mixed effect models (lmer objects of lme4 package). R package version 2.0-29.
- Lê, S., Josse, J., Husson, F. 2008. FactoMineR: An R Package for Multivariate Analysis. *Journal of Statistical Software*. 25(1): 1-18.
- Moran, P.A.P., 1948. The interpretation of statistical maps. *J. R. Stat. Soc.* 10, 243–251.

- Nakagawa, S., Schielzeth, H., 2013. A general and simple method for obtaining  $R^2$  from generalized linear mixed-effects models, *Methods in Ecology and Evolution* 2, 133–142. <http://doi.org/10.1111/j.2041-210x.2012.00261.x>
- Ord, J.K., Getis, A., 1995. Local Spatial Autocorrelation Statistics: Distributional Issues and an Application. *Geogr Anal* 27:286–306. doi: 10.1111/j.1538-4632.1995.tb00912.x
- Wei, T., Simko, V., 2016. Package Corrplot. R package version 0.77. April 2016

## Appendix S3. Supplementary results

### Correlations among ES

#### *Land-based type indicators*

Table S3.1. Correlations among land-based ES indicators. Significant values (\*\*\*) $P < 0.001$  (\*\*) $0.01 < P < 0.05$  (\*) $P < 0.05$  are highlighted (positive relationships) and italic (negative relationships) characters.

ES indicators	P1	P2	P3	R1	R2	R3	R4	R5	C1	C2	C3	C4
<b>Food provision (P1)</b>	1.00											
<b>Materials provision (P2)</b>	<b>0.39***</b>	1.00										
<b>Water provision (P3)</b>	<b>0.69***</b>	<b>0.41***</b>	1.00									
<b>Climate regulation (R1)</b>	<b>0.25***</b>	<b>0.18***</b>	<b>0.22***</b>	1.00								
<b>Soil fertility (R2)</b>	<b>0.31***</b>	<b>0.24***</b>	<b>0.35***</b>	<b>0.20***</b>	1.00							
<b>Water regulation (R3)</b>	<b>0.78***</b>	<b>0.47***</b>	<b>0.72***</b>	<b>0.36***</b>	<b>0.34***</b>	1.00						
<b>Flood regulation (R4)</b>	<b>0.20***</b>	<b>0.26***</b>	<b>0.29***</b>	<b>0.16***</b>	0.12	<b>0.25***</b>	1.00					
<b>Erosion control (R5)</b>	<b>0.63***</b>	<b>0.32***</b>	<b>0.41***</b>	<b>0.25***</b>	<b>0.21***</b>	<b>0.62***</b>	0.12	1.00				
<b>Recreational (C1)</b>	<b>0.17***</b>	<b>0.17***</b>	<b>0.20***</b>	0.11	0.08	<b>0.18**</b>	<b>0.29***</b>	0.06	1.00			
<b>Existence (C2)</b>	<b>0.41***</b>	0.06	<b>0.31***</b>	0.14*	0.13*	<b>0.45***</b>	-0.01	<b>0.36***</b>	-0.01	1.00		
<b>Experiential (C3)</b>	-0.07	0.03	-0.03	0.02	-0.01	-0.01	0.06	-0.01	-0.01	0.06	1.00	
<b>Physical use (C4)</b>	0.04	0.02	0.04	0.07	0.02	0.11	-0.09	0.08	0.08	0.08	0.08	1.00

Supplementary material

*Forest-based type indicators*

Table S3.2. Correlations among forest-based ES indicators. Significant values (\*\*P<0.01 \*P<0.05) are highlighted (bold relationships) and italic (negative relationships) characters.

ES indicators	P1	P2	P3	R1	R2	R3	R4	R5
<b>Food provision (P1)</b>	1.00							
<b>Materials provision (P2)</b>	<b>0.20***</b>	1.00						
<b>Water provision (P3)</b>	<b>0.33***</b>	<b>0.18***</b>	1.00					
<b>Climate regulation (R1)</b>	<b>0.26***</b>	<b>0.17***</b>	0.08	1.00				
<b>Soil fertility (R2)</b>	<b>0.41***</b>	-0.01	<b>0.55***</b>	<b>0.24***</b>	1.00			
<b>Water regulation (R3)</b>	<b>0.28***</b>	<b>0.19***</b>	<b>0.18***</b>	<b>0.41***</b>	<b>0.26***</b>	1.00		
<b>Flood regulation (R4)</b>	0.14	<b>0.26***</b>	<b>0.22***</b>	0.06	0.07	<b>0.18**</b>	1.00	
<b>Erosion control (R5)</b>	<b>0.47***</b>	<b>0.26***</b>	0.04	<b>0.30***</b>	<b>0.16**</b>	<b>0.25***</b>	0.12	1.00
<b>Recreational (C1)</b>	-0.01	0.06	0.11	0.03	0.06	-0.01	<b>0.23***</b>	-0.13
<b>Existence (C2)</b>	<b>0.17**</b>	<i>-0.14*</i>	0.04	0.17	0.08	<b>0.18***</b>	-0.13	<b>0.22***</b>
<b>Experiential (C3)</b>	<i>-0.21***</i>	-0.08	-0.07	<i>-0.09*</i>	<i>-0.16**</i>	-0.03	0.00	<i>-0.20***</i>
<b>Physical use (C4)</b>	-0.06	-0.04	-0.03	0.02	-0.05	0.04	-0.04	-0.02

**Distribution of the forest-based ES categories**




Figure S3.1. Distribution of forest-based ES categories on the study area and their hotspots.



Supplementary material

**Mixed models for the forest-based type ES**

Table S3.3. Results of Mixed models performed for the 10 forest-based indicators and some of the social, climatic and biodiversity indicators. Values for each variable are equal in land-based and forest-based thus their values are in Table 3. Columns for each variable show the coefficient estimates plus standard error and level of significance. \*\*\*P<0.001 \*\*0.01 \*P<0.05 R<sup>2</sup>m: marginal R-squared, R<sup>2</sup>c: conditional R-squared.

ES	Intercept	Population Density (x10 <sup>-5</sup> )	Unemployment (x10 <sup>-4</sup> )	Tertiary sector (x10 <sup>-4</sup> )	Mean Temperature (x10 <sup>-4</sup> )	Rainfall (x10 <sup>-4</sup> )	Woody sp. richness (x10 <sup>-4</sup> )	Biodiversity (x10 <sup>-4</sup> )
<b>Food provision (P1)</b>	3.11±0.50***	-8.5±3.4**	54.7±39.8	17.7±21.6	-1385.0±213.1***	5.7±2.9*	83.4±15.9***	-
<b>Materials provision (P2)</b>	-12.57±2.06***	3.9±13.1	-69.6±176.4	-299.4±95.3***	3148.0±944.8***	36.8±12.4***	395.6±70.4***	4
<b>Water provision (P3)</b>	5.10±0.42***	-2.0±2.7	-25.1±35.9	13.9±19.4	-747.0±194.8***	4.8±2.5*	-8.9±14.3	1
<b>Climate regulation (R1)</b>	-1.34±0.66**	2.3±4.5	-7.4±67.1	65.5±35.6*	347.2±297.6	30.4±4.1***	2.9±26.9	-
<b>Soil fertility (R2)</b>	12.88±0.29***	0.2±1.9	28.2±27.9	13.2±14.9	-3247.0±136.3***	5.1±1.8***	-31.3±1.11***	-2
<b>Water regulation (R3)</b>	4.38±0.24***	-0.3±1.5	9.1±20.9	11.5±11.3	-14.8±108.7	7.9±1.4***	12.9±8.4	-3
<b>Recreational (C1)</b>	-0.02±0.07	-1.1±0.5**	-22.9±6.8***	-5.8±3.6	99.5±32.9***	-0.1±0.4	-10.1±2.7***	2
<b>Existence (C2)</b>	3.24±2.83	7.3±1.9	331.3±276.3	487.7±147.4***	-3072.0±1288.0**	59.1±17.4***	620.2±110.6***	-
<b>Experiential (C3)</b>	-6.78±1.29***	51.3±8.8***	-85.2±132.6	294.5±70.3***	1774.0±588.1***	-12.9±8.0	-251.6±53.1***	5
<b>Physical use (C4)</b>	-8.29±1.61***	41.3±12.6***	208.8±193.1	443.1±98.9***	753.8±722.1	13.5±10.0	42.1±77.9	-