

# *A 4D feature tracking algorithm: a multidimensional view of cyclone systems*

Article

Accepted Version

Lakkis, G., Canziani, P., Yuchechen, A., Rocamora, L., Caferri, A., Hodges, K. and O'Neill, A. (2019) A 4D feature tracking algorithm: a multidimensional view of cyclone systems. *Quarterly Journal of the Royal Meteorological Society*, 145 (719). pp. 395-417. ISSN 1477-870X doi: <https://doi.org/10.1002/qj.3436> Available at <http://centaur.reading.ac.uk/80947/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/qj.3436>

Publisher: Royal Meteorological Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

## A 4D Feature Tracking Algorithm: a multidimensional view of cyclone systems

Journal:	<i>QJRMS</i>
Manuscript ID	QJ-18-0133.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	26-Sep-2018
Complete List of Authors:	Lakkis, Susan; Pontificia Universidad Catolica Argentina, Facultad de Ingeniería y Ciencias Agrarias; Universidad Tecnologica Nacional Facultad Regional Buenos Aires, Unidad de Desarrollo e Investigación para las Ingenierías Canziani, Pablo; CONICET Yuchechen, Adrian; Universidad Tecnológica Nacional, Unidad de Investigación y Desarrollo de las Ingenierías Rocamora, Leandro; Universidad Tecnologica Nacional Facultad Regional Buenos Aires, Unidad de Investigación y Desarrollo de las Ingenierías Caferrí, Agustín; Universidad Tecnologica Nacional Facultad Regional Buenos Aires, Unidad de Investigación y Desarrollo de las Ingenierías Hodges, Kevin; The University of Reading, Dept. of Meterology O'Neill, Alan; The University of Reading, Dept. of Meterology
Keywords:	Dynamic/Processes < 1. Tools and methods, Statistical methods < 1. Tools and methods, Climate < 2. Scale, Global < 2. Scale, Synoptic < 2. Scale, Dynamics < 3. Physical phenomenon, Mid-latitude < 5. Geographic/climatic zone, Polar < 5. Geographic/climatic zone
Country Keywords:	Argentina, United Kingdom Of Great Britain And Northern Ireland

## A 4D Feature Tracking Algorithm: a multidimensional view of cyclone systems

S. Gabriela Lakkis<sup>b,a</sup>, Pablo Canziani<sup>a,c</sup>, Adrián Yuchechen<sup>b,a,c</sup>, Leandro Rocamora<sup>a</sup>, Agustin Caferri<sup>a</sup>, Kevin Hodges<sup>D</sup>, Alan O'Neill<sup>d</sup>

- a. Unidad de Investigación y Desarrollo de las Ingenierías, Facultad Regional Buenos Aires, Universidad Tecnológica Nacional, Buenos Aires, Argentina
- b. Facultad de Ingeniería y Ciencias Agrarias, Pontificia Universidad Católica Argentina, Buenos Aires, Argentina
- c. Consejo Nacional de Investigaciones Científicas y Técnicas, CONICET, Buenos Aires, Argentina
- d. Department of Meteorology, University of Reading, Reading, U.K.

Contact:

Dr. Gabriela Lakkis

UIDI

Secretaría de Ciencia, Tecnología e Innovación Productiva

FRBA-UTN

Medrano 951

C1179AAQ Capital Federal

República Argentina

Email: [gabylakkisetul@gmail.com](mailto:gabylakkisetul@gmail.com)

Review



## Abstract

An objective 4D algorithm developed to track extratropical relative vorticity anomaly 3D structure over time is introduced and validated. The STACKER algorithm, structured with the TRACKER single level tracking algorithm as source of the single-level raw tracks, objectively combines tracks from various levels to determine the 3D structure of the cyclone (or anticyclone) events throughout their life cycle. Stacker works progressively, beginning with two initial levels and then adding additional levels to the stack in a bottom-up and/or top-down approach. This allows an iterative stacking approach, adding one level at a time, resulting in an optimized 4D determination of relative vorticity anomaly events.

A two-stage validation process is carried out with the ERA-Interim dataset for the 2015 austral winter. First the overall tracking capability during an austral winter, taking into account a set of climate indicators and their impacts on Southern Hemisphere circulation, was compared to previous climatologies, in order to verify the density and distribution of the cyclone events detected by STACKER. Results show the cyclone density distribution is in very good agreement with previous climatologies, after taking into account potential differences due to climate variability and different tracking methodologies. The second stage focuses on three different long-lived events over the Southern Hemisphere, during the winter of 2015 spanning seven different pressure levels. Both GOES satellite imagery, infrared and water vapour channels, and ERA-Interim cloud cover products are used in order to validate the tracks obtained as well as the algorithm's capability and reliability. The observed 3D cyclone structures and their time evolution are consistent with current understanding of cyclone system development. Thus, the two-stage validation confirms that the algorithm is suitable to track multilevel events, and can follow and analyse their 3-D life cycle and develop full 3D climatologies and climate variability studies.

Keywords: feature tracking, cyclones, relative vorticity, multilevel structures, dynamic programming, optimal algorithm

## 1. Introduction.

Extratropical cyclones are synoptic scale systems, typically defined as the region surrounding a local minimum in sea level pressure, with strong associated circulation around this minimum. They are fundamental meteorological features, significantly involved in a number of weather phenomena and play a key role in the distribution of moisture, energy and momentum at extratropical latitudes of both hemispheres. They are also linked to many severe weather events at mid and high latitudes affecting natural environments and human infrastructure and property (Neu et al, 2013). While tropical cyclones, often referred to as hurricanes or typhoons depending on their location, have strong seasonality and characteristic tracks in the tropics, extratropical cyclones have different life cycles and displacements, are much more frequent and can vary both in shape and structure spanning a diverse range of sizes. Ayrault and Joly (2000) identified 12 cyclone families over the extratropical North Atlantic in a study of approximately 6000 extratropical cyclone events. Furthermore, extratropical cyclones can develop in diverse synoptic situations at mid to high latitudes depending on their relationship with the evolution of the jet stream, polar ice coverage and mid-latitude sea surface temperature (SST) gradients. Some cyclones are confined to lower tropospheric levels and others may develop to extend through the troposphere (e.g. shallow and deep cyclones, [Lim and Simmonds, 2007](#)). Chen et al. (2014) found that the cyclone and anticyclone activity over China are linked to upper tropospheric jets over East Asia.

The detection of cyclone tracks is a challenge that dates back to the 19th century: its origin can be found in Van Bebber (1891). Petterssen (1956) studied the occurrence and behaviour of moving low pressure centres in the northern hemisphere. Subsequent analyses were carried by several authors in the same region, focusing on climatologies of cyclone development (Homar, 2003) and their relationship with atmospheric dynamics (Hoskins and Valdes, 1990). To efficiently study and to quantify cyclone systems, a number of algorithms have been proposed in recent years, especially considering the renewed interest in the context of climate change. The Intergovernmental Panel on Climate Change has noted the relevance of understanding the dynamic climatology of cyclones and its relation to climate change in several assessments (IPCC, 2007, 2014). [Grieger et al. \(2018\)](#) also highlight the relevance of such studies over the southern hemisphere (SH) given the crucial role extra cyclones play in the poleward transport of heat and moisture to polar regions.

1 One of the most frequent ways to assess the cyclonic activity involves statistical criteria, which in general are  
2 focused on the occurrence and position of the systems, but not generally on the tracking or their physical  
3 properties (e.g. Jansa et al., 2001; Finnis et al., 2007). A more specific direct approach to identify and  
4 analyses these systems is to use objective feature tracking algorithms that detect features belonging to  
5 cyclones in gridded reanalysis products. A number of cyclone tracking algorithms have been developed  
6 using different features to detect and track cyclones and anticyclones (Neu et al, 2013 and references therein,  
7 **Grieger et al. (2018)** and references therein, Hodges, 1995, 1999, 2008). A widely used tracking method, the  
8 neighbour point tracking (NPT), consists of a two steps approach in which first, the cyclonic system centres,  
9 defined as a local maximum or minimum value of a climatic variable, are detected at each timestep and then  
10 the track based on the nearest neighbour concept (Kelemen et al., 2014) is obtained through the temporal  
11 connection between cyclones centres (Murray and Simmonds, 1991; Satake et al., 2013; Flaounas, 2014).  
12 Most frequently used climate variables for tracking purposes include mean sea level pressure (MSLP),  
13 MSLP gradients, geopotential height, potential and relative vorticity, singly or in combination (hybrid  
14 approach), e.g., **Trigo et al., 1999; Picornell et al., 2001; Jansa et al., 2001; Hoskins and Hodges, 2002;**  
15 **Hanson et al., 2004; Bartholy et al., 2008; Hewson and Tittley, 2010.** Such algorithms imply the use of a  
16 decision tree in order to apply necessary constraints regarding the detection and evolution of the feature  
17 under study. Thus, a robust and efficient tracking algorithm has to be able to identify feature values, usually  
18 below or above a defined threshold, their path over time, and also the link between the detected centres as  
19 well as their life-cycle. But in addition, the algorithm also has to consider that cyclones can split or merge  
20 with other cyclones.

21 **Grieger et al. (2018) carried out an assessment on the degree of consensus among 14 state-of-the-art cyclone**  
22 **identification and tracking methods, following Neu et al. (2013). Both studies found that the selected feature**  
23 **tracking algorithms yielded similar overall climatic structures, but notable differences remain, e.g., in**  
24 **intensity of events and event counts. Tracking results tend to group into vorticity-based and**  
25 **MSLP/geopotential based clusters. Different cyclone tracking schemes yielded different cyclones trends,**  
26 **sometime differing even in sign.**

27 Currently available algorithms mostly track atmospheric events on a single geopotential/pressure level, a few  
28 of them considering vertically averaged levels, multivariable ensemble or models (**Serra et al., 2010;**  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 **Kelemen et al., 2014; Hodges et al., 2017).** Because cyclones can have diverse characteristics and evolution  
2  
3 at various levels, single-level or average tracking may impact climate analysis. For example, over the SH the  
4  
5 Andes mountain range and the African Plateau in the SH (Inatsu and Hoskins, 2004, Hoskins and Hodges  
6  
7 2005), different choices of tracking level can give different results for storm track and cyclone evolution.  
8  
9 Thus determining cyclone track structure is an important issue for climate analysis. Lim and Simmonds  
10  
11 (2007) compared cyclone tracks at six levels, between MSLP and 500hPa to assess the vertical climatology  
12  
13 of SH cyclones for the austral winter months (JJA), for the period 1979-2001. Detected cyclone positions at  
14  
15 the various levels were combined by considering the distance between cyclone centres at a given time, as  
16  
17 given by the pressure anomaly (depth) at MSLP and geopotential height anomaly centres, from one level to  
18  
19 the next above, using a reference radius of 444km. Starting at sea level, cyclone centres detected at two  
20  
21 adjacent levels, within this reference radius, were considered as belonging to the same cyclone event. At  
22  
23 each time-step this vertical coupling was carried out up to the level that complied with this criterion. Cyclone  
24  
25 tracks were then defined in terms of multiple level events, by considering their variations in height, at the  
26  
27 times of overlap *only*. Results of this bottom-to-top approach composition were further separated into  
28  
29 shallow (up to 700hPa) and well-organized cyclones (at least up to 500hPa) in a SH cyclone climatology  
30  
31 study. Extending and improving the track coupling process to several more pressure levels would provide  
32  
33 new insights on cyclone types and cyclone life cycles as well as help assess linkages with other weather and  
34  
35 climate processes. In addition, a multilevel tracking developed in this way should take into consideration the  
36  
37 different track lengths at various levels, including track points existing at different levels, prior to or after  
38  
39 period of vertical overlapping. In other words, the description of the multilevel system should not be  
40  
41 restricted only to the period of strict overlapping at different heights, **in order to fully describe the system's**  
42  
43 **evolution.**  
44

45  
46 The method developed by Lim and Simmonds (2007) provides the starting point for the implementation of a  
47  
48 more comprehensive approach to the processing and combination of cyclone tracks at multiple levels for  
49  
50 detailed studies. First, it is necessary to extend the detection process to a larger number of pressure levels  
51  
52 throughout the troposphere. Secondly, given the different kinds of cyclone and anti-cyclone processes and  
53  
54 life cycles, it is necessary to apply both a bottom-to-top approach as well as a top-to-bottom one. Thirdly it is  
55  
56 necessary to extend the climatology to span recent years as well as all seasons. This implies the processing of  
57  
58 vast amounts of data for the detection of tracks at each level and the subsequent combination of numerous  
59  
60

1 tracks from levels, both bottom-to-top and top-to-bottom, to determine the vertical structure and space-time  
2 evolution of events. This requires the implementation of algorithms that use an optimized, automated and  
3 efficient approach to assess and define the vertical structure of cyclones and anti-cyclones. **From now on**  
4 **reference will exclusively be made to cyclones, though the method may also be used for anticyclones.**  
5  
6  
7  
8  
9

10 The present study is based on the cyclone tracking code developed by Hodges (1995, 1999) which uses  
11 relative vorticity rather than pressure or geopotential anomalies computed from MSLP and geopotential as in  
12 Lim and Simmonds (2007) **or Keable et al., (2002), the latter being based on geostrophic vorticity computed**  
13 **from the geopotential.** Following a dynamic programming approach, an automated algorithm, called  
14 STACKER, was developed to determine which cyclone events detected at the different levels are to be  
15 combined into single events, preserving all the track features detected at each level. The STACKER  
16 algorithm is built using a smoothness cost function approach. European Centre for Medium-Range Weather  
17 Forecasts (ECMWF) Reanalysis ERA-Interim products for the SH were used. Not only are there fewer  
18 studies for the SH, but the oceanic nature of the hemisphere and major orographic barriers - such as the  
19 Andes mountain range, the well-established regions of cyclogenesis on the eastern seaboard of southern  
20 South America and cyclolysis on its western seaboard, on either side of the Andes mountain range (Inatsu  
21 and Hoskins, 2004, Hoskins & Hodges 2005), provide interesting tests for the algorithm and its validation.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

34 The STACKER algorithm is introduced in section 2. Multilevel cyclone tracking for the 2015 winter (JJA)  
35 was carried out with STACKER. Section 3 presents the validation of the STACKER algorithm using the  
36 2015 JJA mean state and cyclone density. Furthermore, three selected multilevel events developing near  
37 southern South America are introduced and compared with GOES imagery, ERA-I cloud cover and winds  
38 for further algorithm validation. Conclusions are presented in Section 4.  
39  
40  
41  
42  
43  
44  
45  
46

## 47 **2. Methodology.**

### 48 **2.1 Data**

49 In this section the 4D STACKER algorithm, developed to track significant extratropical anomaly  
50 structures by using relative vorticity anomalies and multilevel track coupling, is presented. The reanalysis  
51 data product use to help develop and test the algorithm is the ERAI 6-hourly relative vorticity (RV), with a  
52 horizontal resolution of  $1.5^{\circ} \times 1.5^{\circ}$  at various levels between the surface and 100hPa (lower stratosphere). In  
53  
54  
55  
56  
57  
58  
59  
60

1 terms of choice of ERAI for the testing of STACKER, it should be noted that Hodges et al. (2011) compared  
2  
3  
4 4 different reanalyses (JR25, ERA Interim, NASA MERRA reanalysis and the NCEP CFSR) for composite  
5  
6 cyclone diagnostics and showed that the detected cyclones compared well between the reanalyses in terms of  
7  
8 numbers, location and spatial distribution, with differences being primarily found in the cyclone intensities.,  
9  
10 Froude (2010) working with a set of reanalysis resolutions, concluded that the relative performance of the  
11  
12 different Ensemble Prediction Systems (EPS) in predicting cyclones seem to agree with the results obtained  
13  
14 using more conventional forecast verification measures (Park et al. 2008) and results comparing the skill of  
15  
16 the ensemble mean with that of the control forecasts show that for cyclone position there is very little  
17  
18 difference between all of the EPSs, independent of resolution.

19 All tests with STACKER were carried out over the southern hemisphere during the winter season (JJA) of  
20  
21 2015.  
22  
23  
24

## 25 **2.2. Tracking Algorithm.**

26  
27  
28 The base algorithm selected to track the cyclones at different levels in the present study is the TRACK  
29  
30 algorithm (Hodges, 1995) the details of which can be found in several papers (Hodges 1994, 1995, 1999,  
31  
32 2008). This algorithm has been used in numerous studies to track cyclones at a single pressure level not only  
33  
34 near sea-level but also in the upper troposphere (e.g., Hoskins and Hodges, 2002, Hoskins and Hodges, 2005,  
35  
36 Mailier et al, 2006, Froude et al, 2007, Serra et al., 2010, Catto et al., 2010, Dacre et al, 2012, Zappa et al.,  
37  
38 2013, Zappa et al, 2014; Pinheiro et al, 2017) and its implementation is well documented. Greiger et al.  
39  
40 (2018) also conclude that vorticity-based tracking algorithms can identify cyclone structures at an early stage  
41  
42 in their lifecycle, hence its use in the current implementation of the STACKER algorithm.  
43

44  
45 A brief review of TRACK is included here in order to understand the subsequent steps. TRACK was  
46  
47 designed to track feature points in a gridded dataset over consecutive time steps on a single level. The first  
48  
49 step of the algorithm spatially filters the data to remove the planetary scales and to reduce the noise in the  
50  
51 RV field to allow more reliable tracking of the features (Hoskins and Hodges, 2002). The filter band is  
52  
53 chosen for the appropriate scales of the feature of interest. This is done by transforming the field to spectral  
54  
55 space using spherical harmonics, applying a triangular truncation filter and transforming back to real space.  
56  
57 For synoptic scale cyclones the filter band chosen is T5-42 which has been used in the studies referenced  
58  
59  
60

1 above. A tapering filter is also applied to spectral coefficients to reduce Gibbs oscillations (Hoskins and  
2 Sardeshmukh, 1984). The efficacy of this approach is highlighted in Froude (2010) (their Figure 1) which  
3 compares the T42 RV filtered field and the feature points identified with the much higher resolution T799  
4 (25km) MSLP field. When using RV for synoptic scale cyclones the same filtering is performed irrespective  
5 of the original higher data resolution.  
6  
7  
8  
9  
10

11 The next step of the algorithm identifies local minimum or maximum values of RV as feature points  
12 representing cyclonic and anti-cyclonic events in the SH. These feature points are then formed into tracks by  
13 first initialising a set of tracks based on a nearest neighbour method and then using the minimization of a  
14 Smoothness Cost Function (SMF) subject to adaptive constraints (Hodges, 1999) suitable for extra-tropical  
15 cyclones, to produce an optimal smoothest set of tracks. This whole procedure can be summarized in three  
16 main steps: Data pre-processing, Feature Point Detection and Tracking.  
17  
18  
19  
20  
21  
22  
23

24 Previous studies have compared the automated tracking using TRACK with a more subjective manual  
25 tracking approach, though this can be sensitive to different observers. Colle et al (2013) found a success rate  
26 ~85% using RV over a North American-Western Atlantic region that improves if short lived systems, ~1day,  
27 are excluded. The success rate was found to increase to ~90-95% using MSLP. The study by Gramscianinov,  
28 Hodges and Camargo ( “The Properties and Genesis Environments of South Atlantic Cyclones”, submitted,  
29 2018), using RV, found a success rate over a region of the South Atlantic of ~89%. These success rates  
30 compare favourably with other schemes.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

### 41 2.3. STACKER Algorithm.

42 The STACKER algorithm is designed so as to associate all tracks obtained from a given feature-tracking  
43 algorithm that represent the same synoptic event at different vertical levels, here pressure levels are used. In  
44 order to expand the analysis to include more pressure levels and hence to describe the four-dimensional  
45 evolution of cyclones a binary linear programming model is developed, in which each possible combination  
46 of simultaneous detections on adjacent pressure levels would become an optimization variable to represent  
47 the most coherent relationship between the given set of detections. This algorithm named “*STACKER*  
48 *algorithm*” is a progressive model optimization algorithm. It considers tracks from TRACK outputs at two  
49 adjacent pressure levels and links them according to an objective function criterion. Subsequently, the  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1 resulting stacked tracks are linked with the next adjacent pressure level tracks, until the stacking is complete.  
2  
3 The name STACKER comes from the idea of stacking tracks from upper levels over tracks from lower  
4 levels. The algorithm works in either directions, i.e., bottom-to-top and top-to-bottom, i.e., including the  
5 possibility of stacking lower level tracks under higher level tracks, such as upper level troughs or cut off  
6 lows  
7  
8  
9  
10

11 STACKER only works with track detections from two levels at a time and iterates over the levels adding a  
12 single level to the stack each time. Figure 1 presents the STACKER algorithm flowchart. The stacking  
13 process is highlighted in blue, while the single level tracking input is highlighted in green. The flowchart  
14 shows the decision-making process and the optimization process, given all the possible outcomes of the  
15 decision process. The optimization process should create a link for every possible pair of combinations  
16 between two consecutive levels, i.e., track-to-track linking. However, this approach assumes that each lower  
17 track corresponds to only one upper track, but this may not be entirely true. Indeed, it is possible that an  
18 event changes during the evolution of its life cycle as a consequence of changes in atmospheric circulation,  
19 land-sea transitions, or it may be impacted by significant orography, resulting either in lysis or, occasionally,  
20 modification of part of the vertical structure while maintaining its nature. To avoid this limitation a point-to-  
21 point linking is used. This method considers a link for every possible combination of a lower (upper) point  
22 with an upper (lower) point, while constraining the problem such that a link between two points can be made  
23 only if, all of the other links can be made simultaneously. Furthermore, points must not be associated with  
24 more than one active link. Thus the optimization process returns a track-to-track association born from a  
25 point-to-point correspondence subject to chosen constraints (Fig. 1). Specific details of the STACKER  
26 algorithm can be found in the Appendix.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 Using this approach, the algorithm maintains the coherence of linking two potentially associated tracks  
45 assuming they belong to the same object and thereby gains the flexibility of linking more than one track on  
46 each pressure level. Nonetheless, the combinatorial problem becomes far more complex and the amount of  
47 optimization variables, problem constraints and processing resources needed increases considerably. In order  
48 to avoid highly improbable associations between very dissimilar detections, a “linkage” threshold value, i.e.,  
49 a strong constraint, will discern which associations will be relevant for the study and thereby decrease the  
50 problem complexity. Also, some filters can be applied, like maximum radius distance between points,  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1 minimum time duration, minimum distance travelled, region of origin, etc. Fig. 2 shows how a maximum  
2 radius threshold is considered at each point along a track to constrain the coupling of points from the upper  
3 track (blue disk) in a manner similar to that used Lim and Simmonds (2007). The plot also highlights how  
4 this helps discriminate the coupling process for tracks crossing each other in the lower level.  
5  
6  
7  
8  
9

10 Due to the algorithm's stacking nature, which operates in both directions, bottom-to-top and top-to-bottom,  
11 from now on and for the sake of simplicity the section that has already been stacked or is the result of a  
12 previous stage of the algorithm will usually be referred to as "lower", independently of the direction of the  
13 stacking process. Similarly, what it is commonly referred as "upper" is the new section that is to be added or  
14 stacked onto an already existing stack. Figure 3 explains the iterative stacking process through a simple  
15 example considering track segments detected at three adjacent levels. "Upper" tracks are usually one-level  
16 tracks, while "lower" tracks are considered to contain tracks from more than one pressure level, resulting  
17 from the stacking process, i.e., stacks, except during the initial stack iteration where one "lower" and one  
18 "upper" track are combined. While the same process can be used in either stacking direction, bottom-to-top  
19 or top-to-bottom oriented, the final solution, while similar may not exactly be identical for a given multi-  
20 levelled event stacked in different directions. Minor differences may result from potential fragmentation of  
21 tracks (Figure 3). The differences between both directions could be attributed to the physical nature of the  
22 cyclonic system. Thus the best direction would be associated with what physical processes are driving  
23 cyclone development or maintenance. For instance, at upper levels in sub-tropical latitudes, cyclogenesis  
24 could be due to baroclinic instability, while at lower levels at polar latitudes, sea-ice to open water thermal  
25 contrast could contribute to the forcing at lower levels (Hakim, 2003, Joly et al., 2003 and references  
26 therein). Bottom-to-top and top-to-bottom tracks which overlap both in space and time are merged to obtain  
27 an integrated view of the event. This merger feature has been included here as an automatic process given the  
28 use of STACKER for climatology studies (see further details in Appendix). Studies focusing on case study  
29 dynamics may consider bottom-to-top and top-to-bottom, presenting results independently if desired.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 Before proceeding further, it is important to note an issue linked to the relative vorticity threshold value used  
51 in track determination. While this threshold is defined for TRACK, it is also relevant to STACKER, using a  
52 vorticity-based feature-tracking, because its choice may impact the stacking process. This happens because  
53 tracks obtained at various levels are combined in the stacking process. The relative vorticity threshold as  
54  
55  
56  
57  
58  
59  
60

1 defined in TRACK is fixed at  $-1.0 \times 10^{-5} \text{ s}^{-1}$ , without height or latitude dependence. This is set at a  
2 comparatively low value to capture as much of the cyclone lifecycle as possible for any season. A more  
3 restrictive threshold will reduce the number of systems identified but will also shorten the lifecycles of those  
4 that are identified, more so for the weaker summer systems. Furthermore, it is important to note that  
5 TRACK, as well as other vorticity-based tracking algorithms (Greiger et al., 2018), spectrally filters the  
6 vorticity to remove the small-scale noise (see Methodology section). This will reduce the tendency to  
7 identify a large number of small scale systems at all levels. RV exhibits a somewhat larger mean value at  
8 500hPa, the non-divergence level. Inspection of monthly mean RV fields shows that the range of RV values  
9 does not change significantly with height, nor with season. RV threshold levels for the lowermost and  
10 uppermost levels are similar. Furthermore, variations are primarily spatial. RV standard deviation fields  
11 show that the amplitude of RV perturbations increase with height. Hence a fixed threshold at a low value  
12 does not significantly impact the detection of cyclonic systems.

13 Furthermore, a fixed RV threshold value such as the one chosen here does not affect cyclone event counts in  
14 the lower levels near the surface nor in levels near the tropopause. When stacks are constructed by  
15 STACKER bottom-up or top-down and the stacks contain tracks near the upper or lower levels this does not  
16 pose a problem because those initial levels, with similar threshold values, will constrain the number of tracks  
17 from the middle troposphere which are stacked. Concern about this value could arise for stacks which begin  
18 at or around 500hPa, i.e., mid tropospheric systems, where this threshold may enhance the number of tracks  
19 detected there, since in principle relative vorticity maximizes there. If this is so, and if the threshold value  
20 becomes more restrictive, the number of systems identified would be reduced, and, as already noted, result in  
21 shorter lifecycles for those identified. This criterion has precedents in for example, Pinheiro et al., (2017).  
22 Even so, and in order to test the threshold sensitivity, the TRACK algorithm was run at 500 hPa, with a set of  
23 more restrictive RV thresholds values. Results indicated that the difference may be limited to less than 2%  
24 fewer detected events, which is negligible. Even more so if the discrepancies between the different tracking  
25 algorithms compared in Neu et al. (2013) and Greiger et al. (2018) are taken into consideration. Note that a  
26 similar issue could arise with other tracking algorithms when used with STACKER, depending on the  
27 criteria used to define the anomalies to be tracked at different levels, even with pressure or geopotential-  
28 based tracking.

1 The detailed definitions and concepts of essential aspects, as well as the mathematical components of the  
2 STACKER Algorithm, are available in the Appendix.  
3  
4

5  
6 The STACKER algorithm only uses detected cyclone tracks as inputs, i.e., cyclone position at a given time  
7 and height, as the above paragraphs show. The issues regarding RV pertain to the use of the RV-based  
8 TRACK algorithm. STACKER, however, can be used in principle with tracks obtained by any reliable  
9 cyclone tracking algorithm. As was the case here, given the RV threshold dependence with height, care will  
10 be required to determine the adequate height-dependent threshold for the variable used in the selected  
11 tracking algorithm, be it pressure, geopotential, temperature or a combination thereof.  
12  
13  
14  
15  
16  
17  
18  
19  
20

### 21 **3. Validation Studies**

22  
23 The fields used for testing the combined tracking and stacking algorithms came from the ERA-Interim  
24 reanalysis data set. As explained above each event was first independently tracked with TRACK on each  
25 pressure level during the 2015 JJA (austral winter) period and then stacked with the STACKER Algorithm  
26 (SA). A total of 753 multilevel, stacked extratropical cyclone events over the SH (south of 30°S), lasting at  
27 least **two** days, were detected during this period, with vertical structures spanning two levels or more, i.e.,  
28 multilevel events, during part of or for all their tracked life cycle. The number of cyclone events detected that  
29 corresponded at 2 levels is 347, at 3 levels, 167, at 4 levels, 99, at 5 levels or more, 140. This does not mean  
30 that each event spanned all the levels by which it was categorized throughout its life cycle, rather that the  
31 category assigned represents the largest number of levels spanned at some stage in its evolution.  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 The validation is carried out in two steps. First, the overall seasonal tracking capability during the 2015  
42 austral winter is compared to previous climatologies, in order to verify that the detected stacked cyclone  
43 events were geographically distributed, within reasonable limits of expected variability, in agreement with  
44 these climatologies. During 2015, a strong El Niño event took place with Multivariate ENSO index (MEI)  
45 values between 2 and 2.5 during the sampled period (<https://www.esrl.noaa.gov/psd/enso/mei/table.html>).  
46 The seasonal Southern Annular Mode (SAM) index reached a high positive value of 3.19 (<http://www.nerc-bas.ac.uk/public/icd/gjma/newsam.1957.2007.seas.txt>), which may also imply changes to cyclone  
47 trajectories. The Indian Ocean Dipole index at this time also indicates a positive phase with comparatively  
48 large values ([https://www.esrl.noaa.gov/psd/gcos\\_wgsp/Timeseries/Data/dmi.long.data](https://www.esrl.noaa.gov/psd/gcos_wgsp/Timeseries/Data/dmi.long.data)), while the  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 Interdecadal Pacific Oscillation has weak negative values  
2  
3 (<https://www.esrl.noaa.gov/psd/data/timeseries/IPOTPI/>). These variability indicators, known to impact SH  
4 weather systems, e.g., Pezza and Ambrizzi, (2003), Pezza et al. (2007), Pezza et al, (2008), Pezza et al,  
5 (2012), Reboita et al. (2009, 2015), are considered here qualitatively since there may be non-linear  
6 relationships with cyclone development and lifecycle, as these studies point out. Their impacts on cyclone  
7 occurrence and development can only be evaluated through a fully developed climatology analysis.  
8  
9

10  
11  
12  
13  
14 The second validation step is focused on three different long lived events over the SH during the winter  
15 (JJA) of 2015 spanning seven different pressure levels (925, 850, 700, 600, 500, 400 and 250 hPa). Two  
16 originate at subtropical/midlatitudes and one begins near Antarctica. These three events selected for SA  
17 validation were located at some point in their evolution in the vicinity of the Andes mountain range, which is  
18 a major orographic feature in the SH circulation (Inatsu and Hoskins, 2004). Following Gozzo et al. (2014)  
19 who included GOES satellite Infrared (IR) imagery in their case study, both GOES IR, Water Vapour -WV-  
20 channels and ERA-Interim cloud cover products are used here in order to validate the algorithm's capability  
21 and reliability.  
22  
23  
24  
25  
26  
27  
28  
29

### 30 31 32 33 **3.1 Mean JJA 2015 State compared with Climatology.**

34  
35 The results for the 2015 austral winter are presented in Figs. 4 through 8 and primarily but not exclusively  
36 refer to the SH cyclone climatology of Hoskins and Hodges (2005), which was based on TRACK and with  
37 the climatology of Lim and Simmonds (2007), which was based on the University of Melbourne pressure-  
38 based feature tracking system (Murray and Simmonds, 1991a,b). Fig.4 (and Animation 1 - supplementary  
39 material) shows the wind speed at 300hPa for JJA 2015 to highlight the mean position and variability of the  
40 jets during the period. The well-known SH jet structure can be seen with the spiralling of the subtropical jet  
41 into the polar jet, beginning over the eastern Pacific, over Chile and Argentina and into the South Atlantic,  
42 merging with the polar jet near the Greenwich Meridian. Overall features agree with the Peña-Ortiz et al.  
43 (2013) jetstream climatology using data from the NCEP/NCAR Reanalysis. The main differences are in the  
44 position of the transition of the subtropical to polar jet, which in 2015 occurs further south over southern  
45 South America and in a stronger polar jet close to Antarctica. These differences can be expected from the El  
46 Niño state and positive SAM during this particular austral winter.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 Fig. 5 shows the cyclone event distribution during JJA 2015 for all multilevel cyclones detected. The largest  
2 cyclone density is located off the coast of Rio Negro and Chubut provinces, Argentina, over the South  
3 Atlantic, somewhat poleward of the axis of the poleward spiralling jet, in a region where it accelerates. Over  
4 the South Atlantic, close to the Greenwich meridian, two additional small high cyclone density areas can be  
5 seen, one equatorward and the other poleward of the jet axis. The other prominent cyclone density region is  
6 found close to Antarctica almost under the axis of the polar jet extending from the eastern South Atlantic  
7 across the southern Indian Ocean. Another small cyclone density region is located over the central Pacific  
8 under the subtropical jet, where it reaches its highest speed.

9  
10  
11  
12  
13  
14  
15  
16  
17  
18 Fig. 6 specifically shows the cyclone density for “shallow” cyclones (spanning 2 and 3 levels) at low, mid  
19 and upper troposphere levels, respectively, “intermediate” events (Fig. 7) spanning lower or upper  
20 troposphere, respectively and vertically “well-organized” cyclones (Fig. 8) spanning most if not all the levels  
21 considered in this study, building on Lim and Simmonds (2007) definitions and discussion. Even though  
22 only a single season is considered, which as noted is particularly impacted by a strong ENSO event, its mean  
23 state is comparable with the above-mentioned climatology studies, spanning at least 45 and 22 years  
24 respectively. Overall results found here agree with the broad features of cyclone occurrence over the SH  
25 winter, after taking into account the known consequences of the previously described state of the climate  
26 system.

27  
28  
29  
30  
31  
32  
33  
34  
35  
36 Total shallow cyclone (events detected at 2 to 3 levels, Figure 6) densities show that, while present  
37 throughout the extratropical SH, the more frequent occurrence of such systems is on the Atlantic coast of  
38 southern South America. Densities also increase near the subtropical jet over the South Pacific and near the  
39 entrance of the subtropical jet over the South Atlantic. When considering the occurrence of shallow events in  
40 different height ranges, near-surface (925, 825, and 700 hPa surfaces, Fig. 6b) densities maximize off the  
41 coast of Argentine Patagonia in the South Atlantic near Golfo Nuevo and Golfo San Matías between 40° and  
42 45°S, just poleward of the spiralling jet. This is a prominent cyclogenetic region in the SH (Hoskins and  
43 Hodges, 2005; Simmonds and Keay, 2000). This region appears to extend into the central South Atlantic.  
44 Another region of prominent cyclone density would have been expected over the Rio de la Plata Basin in  
45 subtropical South Eastern South America. However, present results agree with Pezza and Ambrizzi (2003)  
46 and Reboita et al (2015) who noted (also using the University of Melbourne tracking code applied to MSLP  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 alone) that surface cyclone occurrence is strengthened over the Patagonian Atlantic coast during El Niño  
2 events. The lack of a cyclogenetic region near the Rio de la Plata basin (e.g., Hoskins and Hodges, 2005)  
3 and such a strong feature along the coast of Patagonia could be linked to the widening of the Hadley Cell  
4 during El Niño events and the associated southward displacement of the subtropical jet, as seen in Fig. 4.  
5 Another prominent region is found near Antarctica due south of Australia in the vicinity of Terre Adélie and  
6 George V Land, where the polar jet has weakened significantly (Fig. 4). Some shallow cyclone density  
7 secondary areas are found in the Indian Ocean extra-tropics, seemingly along the equatorward edge of the  
8 polar jet, which is strong in that longitude region. Note the central South Pacific region with reduced cyclone  
9 density, also in agreement with previous studies.

10 Mid tropospheric (700, 600 and 500 hPa surfaces, Fig. 6c) shallow systems primarily occur off the Atlantic  
11 coast of Patagonia, extending into the Central South Atlantic, but with a narrower latitude extent than the  
12 surface shallow cyclone density. This region extends under the subtropical jet, where jet wind speed begins  
13 to increase prior to its spiralling transition to the polar jet, over the South Atlantic (Fig. 4). There is a  
14 secondary area of cyclone activity in the subtropical central South Pacific, in the vicinity of subtropical jet  
15 where the jet begins to slow down as it extends into southern South America.

16 Upper level (500, 400 and 250hPa, Fig. 6d) shallow systems maximize to the west of South Africa between  
17 20° and 30°S. These occur in the vicinity of the subtropical jet entry region west of Africa. A secondary  
18 maximum is also observed in the central South Pacific subtropics where maximum subtropical jet speeds are  
19 reached. As in the lower levels there is reduced activity in the central South Pacific. Upper level shallow  
20 cyclone density (Fig. 6d) off the coast of Patagonia is reduced with respect to the mid and lower stratosphere.  
21 The first two density areas are in agreement with cut-off low (COL) occurrence in Pinheiro et al (2017).  
22 However, as the authors note, the cyclone centres detected by TRACK in the upper troposphere may include  
23 pressure troughs as well as closed circulation systems at upper tropospheric pressure levels, and hence  
24 further criteria need to be applied to confirm the occurrence of COLs. Once more, reduced shallow cyclone  
25 occurrence is observed in the central South Pacific at extratropical latitudes.

26 Intermediate cyclones are less frequent than shallow events (Fig. 7a). They appear to occur more frequently  
27 over the Indian Ocean and over northern Chile, Argentina, Bolivia and Paraguay. While intermediate  
28 cyclones can be detected anywhere between subtropical and high latitudes, these systems (events detected at  
29  
30

1  
2 4 to 5 levels) in the low to mid troposphere (925, 850, 700, 600 and 500 hPa surfaces, Fig. 7b) have larger  
3 densities only in the southern Indian Ocean between high and extratropical latitudes, primarily on the  
4 equatorward edge of the polar jet. Mid to upper level (700, 600, 500, 400 and 250 hPa surfaces, Fig. 7c)  
5 intermediate events occur in the same areas. As was the case with shallow events, there is a reduced density  
6 region in the extratropical central South Pacific.  
7  
8  
9  
10

11 Well-organized cyclone events considered here are those that span most heights, i.e., 6 or 7 levels (Fig. 8)  
12 and are mostly found between 45° and 60°S, spanning an extended longitude range from the central South  
13 Atlantic into the southern Indian Ocean, as far as Australia. This result agrees with Lim and Simmonds  
14 (2007). These well-organized cyclone events have their largest density in the vicinity of the jet entry region  
15 and strongest section of the polar jet. A secondary area occurs where the polar jet weakens considerably.  
16 While such well-organized cyclone events may arise throughout the subtropical and extratropical SH, they  
17 appear to be essentially higher extratropical latitude features with very limited occurrence in the subtropics,  
18 at least during an El Niño year. Inspection of sea-ice satellite maps for the 2015 winter (not shown) appear to  
19 suggest that both the polar jet and well-organized cyclones are located near the edge of the sea-ice/open  
20 ocean transition.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31  
32 Summarising, the detected cyclone systems cluster as expected in the vicinity of the subtropical and polar  
33 jets as well as near the spiralling transition of the jet over the South Atlantic. Most of the higher density  
34 clusters occur in regions where the upper level jets accelerate or decelerate. Well organized cyclones  
35 spanning the whole troposphere appear to occur also under the strongest winds in the polar jet. The SA  
36 results show that the cyclone systems spanning different height ranges tend to occur in different regions of  
37 the SH. The reasons for the varying spatial distribution of cyclones occurring at different height ranges, in  
38 particular for shallow and intermediate cyclones, is beyond the scope of the current study, requiring a further  
39 climatology study using a larger number of years.  
40  
41  
42  
43  
44  
45  
46  
47

### 48 3.2 Validation Cases

49  
50  
51 Well-organized, long-lived events are considered here, given that they represent the full analysis of SA  
52 capability. The events analysed are merged bottom-to-top and top-to-bottom stacks. Even though the well-  
53 organized events span most or all of the 7 pressure levels, SA results are displayed here at five levels only  
54 (925, 850, 700, 500, and 400 hPa) only for the sake of clarity. In each case, a first set of images shows the  
55  
56  
57  
58  
59  
60



1 stacking on a 3D display. The top image presents a map view of the stacking and the second image shows an  
2 oblique view of the same display, so that tracks at different heights can be viewed without overlapping. A  
3 second set of plots and images present both ERA-Interim cloud cover at 500hPa and NOAA's GOES  
4 Satellite imagery from the IR and WV imagery channels, respectively, as a 4 timestep sequence of evolution  
5 from the events. Each set also includes a close-up of the geopotential field at 300hPa and 1000-500hPa  
6 thickness field, centred in the position of the RV anomaly at each timestep. Finally, animation loops, for each  
7 of the 3 events, are provided as supplementary material to facilitate the visualization of cyclonic system  
8 evolution as captured by SA in ERA-I cloud cover and 300hPa wind intensity together with GOES IR and  
9 WV images.

### 3.2.1. Event 1 July 29<sup>th</sup> – August 10<sup>th</sup> 2015.

10 The system for this validation case developed between July 29<sup>th</sup> and August 10<sup>th</sup>, 2015, i.e., the RV anomaly  
11 lifecycle lasted 12 days. It is based on 236 independent cyclone event detections by TRACK, combined by  
12 SA (Fig 9, 10). The system begins its development close to New Zealand's eastern coast (36.01°S,  
13 178.85°E), travels east through the Pacific Ocean and twelve days later rapidly decays over the Andes  
14 mountain range (32.62°S, 285.40°E). During the early stages of this event a train of 4 other cyclone events  
15 develop to the south and flow across the South Pacific and South Atlantic (Animation 2 - supplementary  
16 material). This event initially appears to develop in the upper levels (400 and 500hPa) and a few days later  
17 appears in the lower levels, moving east along the poleward edge of the subtropical jet (Figs. 9 and 10 a). At  
18 this point the RV anomaly stretches under the poleward edge of the subtropical jet between a confluent ridge  
19 to the north and a diffluent trough, aligned in a southeast-northwest axis, to the south. These features are  
20 visible both in the 1000-500hPa thickness and 300hPa geopotential, though somewhat less intense in the  
21 latter. All these features suggest significant baroclinicity. Two days later this circulation structure continues  
22 as the system moves east. Note the existence of the semi-permanent anticyclone near tropical latitudes off the  
23 coast of Peru in this and subsequent timesteps. By August 7<sup>th</sup> the trough sharpens as it becomes narrower,  
24 particularly in the 100-500hPa thickness. (Fig. 10c). Inspection of Fig. 10c shows both in clouds (ERA-I and  
25 GOES), winds and RV the presence of a small cold front along the axis of the trough. Fig. 10d, shows a  
26 sharp trough aligned with the coast of Chile as the cyclone reaches maturity, as it moves close to the highest



1 part of the Andes mountain range, with peaks above 6000m. The system rapidly veers south when it comes  
2 almost on top of the Andes, and it decays, following the jet. **This southward motion is** probably induced by a  
3 high pressure system over Central South America, visible in the WV image **geopotential and thickness** and  
4 animation. It follows the coast for about a couple of days until it vanishes. The **RV anomaly showing a well-**  
5 **developed cyclone** is very organized through its vertical structure and the cyclone is clearly visible in the  
6 infrared and water vapour channel imagery.  
7  
8  
9  
10  
11  
12

13  
14 This event lacks a track at 250 hPa. It is assumed that the event did not develop at that altitude and evolved  
15 below that level. It may also be that either TRACK did not detect a track at that altitude or because the SA  
16 did not find during this period a track to link with the rest of the tracks in the stack. It is nevertheless tracked  
17 through six pressure levels, making it a distinct well organized structure for a case study.  
18  
19  
20  
21  
22  
23  
24

### 25 **3.2.2. Event June 27<sup>th</sup> -July 11<sup>th</sup>, 2015.**

26  
27 This validation case is tracked through all seven pressure levels and appears on all five levels displayed in  
28 **Figs. 11 and 12**. This event is not always as distinct in GOES imagery as is the previous one, **because of**  
29 **more dispersed cloud features**. Hence it has a better signature in WV imagery than in the IR imagery, but it is  
30 still traceable and appears to respond differently to the presence of the Andes Mountain Range compared  
31 with the previous case. **Initially the 300hPa winds exhibit weak jet features (Fig. 12a)**. It is first detected at  
32 925hPa on 27<sup>th</sup> of June over the Pacific Ocean (32.95°S – 225.17°E), 6hrs later at 850hPa and next day at all  
33 other levels (Fig. 11, 12). **During the early stages it is associated with a weak diffluent trough followed by a**  
34 **somewhat stronger ridge in 300hPa geopotential height and 100-500hPa thickness fields (Fig 12a), but with**  
35 **no confluent ridge as in the previous event**. It moves eastward and slightly equatorward, reaching southern  
36 South America in the vicinity of Puerto Montt (~ 41°S, Chile, **Fig. 12b**) and the diffluent trough is somewhat  
37 stronger. During this stage the system evolves along the poleward edge of a weak polar jet (Animation 3 -  
38 supplementary material). At this **somewhat higher** latitude the average height of the Andes is **limited to**  
39 **2500-3000m** and thus the system is able to continue its development over and beyond the Andes. It crosses  
40 Argentina diagonally towards the Rio de la Plata (34°S), reaching the South Atlantic near the border between  
41 Uruguay and Brazil. During this stage the system is tracked at 850hPa and levels above. **The GOES images**  
42 **and ERA-I cloud cover plots show a weak mature cyclone signature over the South Pacific, approximately**  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 24 hours before reaching the Andes. The cloud signature associated to the RV anomaly is weak after it  
2 crosses the continent, most probably because of low levels in atmospheric water content as it travels over the  
3 semiarid Patagonia steppes particularly during winter. The RV anomaly couples over Buenos Aires with the  
4 poleward edge of the subtropical jet after becoming separated from the edge of a polar jet branch over  
5 Argentina. At this time the trough is sharper in the 1000-500hPa thickness map than at 300hPa geopotential  
6 (Fig. 12 c). Also, a cloud signature begins to develop probably due to enhanced atmospheric water content as  
7 the anomaly travelled over the humid Pampas Region and the warm waters of the Rio de La Plata estuary.  
8 The cyclone cloud signature then reappears over the South Atlantic at a distance from the coast during the  
9 latter part of the event (Fig. 12 d). During this period the trough changes from a southeast-northwest slanted  
10 feature to a south to north feature suggesting cyclone maturity. The system evolves on the poleward side of  
11 the subtropical jet, under the influence of enhanced baroclinicity and probably atmospheric water vapour.  
12 The system vanishes over the mid-eastern Atlantic Ocean (35.73°E, 20.98°S). It is important to highlight the  
13 difference between the results of a stack analysis versus TRACK applied to a single level in the lower  
14 troposphere. In the latter case this particular event would have been detected as two independent events. In  
15 the first case, the STACKER algorithm is able to link them as an evolving single system which undergoes a  
16 first development over the South Pacific, almost has no signature in the drier atmosphere over Patagonia and  
17 then starts developing once more over the subtropical South Atlantic.

### 3.2.3. Event July 10<sup>th</sup> - July 24<sup>th</sup>, 2015

18 This validation event is first detected close to the coast of eastern Antarctica. It is first tracked over the  
19 Antarctic coastline (18.69°E, 69.06°S) due south of South Africa on July 10<sup>th</sup>, and is present at the beginning  
20 at all levels with a few hours difference (Fig 13, 14, Animation 4 - supplementary material). Of the three  
21 validation cases, it is the event which shows the most change in the levels of the tracks as the system evolves  
22 as well as the longest overall track since it virtually circumnavigates all the extratropical Southern  
23 Hemisphere. The lowermost 925hPa track lasts a couple of days and vanishes by the time the system travels  
24 south of New Zealand (Fig. 13b). By the time it reaches the mid South Pacific, it is limited to 500hPa and  
25 above (including 250hPa (not shown)). A probable cause for the initiation of this event could be found in the  
26 thermal difference between sea-ice and open waters near the initial tracking point, which is close to the edge

1 of the sea-ice in the mean July 2015 satellite sea-ice retrievals. The system is tracked at and above 500hPa as  
2 the system veers away from Antarctica, north of the Bellingshausen Sea, following the polar jet on its  
3 poleward side, as the latter moves north. As the RV anomaly nears southern South America (July 18<sup>th</sup>, Fig.  
4 14a) a distinct cyclone cloud pattern can be seen both in IR and WV images. At this stage, the cyclone is  
5 surrounded by the polar jet. A diffluent trough can be observed both in the 1000-500hPa thickness and in  
6 300hPa geopotential height. Also note the ridge to the northwest of this trough near 35°S. By the 19th July  
7 (Fig. 14b) the system path reaches the continent and begins to cross the Andes north of 40°S, where the  
8 Andes rapidly vary in height from 3000 and 5000m towards the Equator. At this point the RV anomaly is no  
9 longer linked to the polar jet and reaches the subtropical jet. There appears to be a small front behind the  
10 system visible in the 300hPa winds and a much stronger one preceding the system extending over Argentina  
11 and into the South Atlantic. Both have comparatively weak cloud structure, even more so the latter one. The  
12 cyclone system itself is only tenuously visible in the WV image. The tracks cross Argentina north of the Rio  
13 de la Plata near 32°S and cross Uruguay and Brazil into the South Atlantic, veering to the Southeast (July  
14 21<sup>st</sup>, Fig. 14c). The cyclone structure begins to reappear as the system reaches the Atlantic Ocean, following  
15 the front. The system reaches 45-50°S by July 22<sup>nd</sup> (fig. 14d) and continues its life cycle at that latitude (Fig.  
16 13). The subtropical jet resumes its more zonal behaviour near the tropic of Capricorn. A high pressure  
17 system develops to the northeast of the cyclone, with a confluent ridge. The cyclonic cloud pattern develops  
18 as the system moves over the South Atlantic, reaching maturity. The event finally decays fourteen days after  
19 being first tracked, July 24<sup>th</sup>, very close to the original detection region (3.1°W - 52.0°S).

#### 3.2.4 Additional comments about STACKER based on validation cases

20 A number of issues regarding SA come out from the above event analysis. Each validation case discussed  
21 above has a different, distinct evolution which SA was able to determine and show the differences. In the  
22 first place, note that the SA track points determined using RV are well co-located with respect to the  
23 evolving troughs in all 3 events. While this is linked to the specific tracking algorithm used, it shows that the  
24 tracks at various levels are well-correlated with relevant circulation features. Furthermore, the RV anomaly  
25 features found at the different levels, for a given event, are consistent with these observed regional/local  
26 circulation features. Fig. 15 shows 3 such RV anomaly stacks at a given time, one for each of the 3 events

1 discussed above. The figure shows that, be it a snapshot of a 2-layer state of an originally multi-level event,  
2 or a multilevel event active at all levels, the RV anomalies in each case have a high degree of congruence  
3 with corresponding circulation features present at each level of an event as well as with each other at the  
4 different levels, highlighting how RV anomaly features vary with height within a cyclonic system.  
5 Furthermore the comparison with both infrared and water vapour imagery which convey information  
6 cloudiness and circulation on different height ranges validates the observed vertical structure found by SA  
7 for the cyclone events.  
8  
9

10 Inspection of the coordinates (latitude and longitude) at the various stacked levels for each event shows that  
11 as the initial RV anomalies evolve within the stack, the initial vertical slant of the system (the RV anomalies  
12 at upper levels lag behind those at lower levels) the diminishes as cyclonic features develop and mature. The  
13 differences in latitude and longitude between upper and lower level tracks become very small, in agreement  
14 with the baroclinic instability theory for cyclogenesis. Furthermore, the inspection of the troughs observed in  
15 1000-500hPa thickness and 300hPa geopotential shows that as the RV anomalies evolve into distinct  
16 cyclonic features, their inclination with respect to a meridian decrease until the trough finally becomes  
17 parallel to it, when the systems reach full development.  
18  
19

20 Consequently, the SA is correctly combining tracks at different levels corresponding to the same cyclone  
21 event and provides accurate 3D views of the evolving systems.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

#### 32 **4. Conclusion.**

33 A 4D-STACKER algorithm has been developed, following dynamic programming criteria, to track  
34 extratropical RV anomaly 3D structures over time with an objective systematic approach. The algorithm is  
35 based on the TRACK algorithm which provides the raw tracks at single levels, which must then be combined  
36 objectively to determine the 3D structure and its evolution. Because of the extremely large number of  
37 possible single level track combinations, an optimal solution is reached according to a defined criterion  
38 necessary to reduce options and better define the track “stacks” representing a given cyclone event.  
39  
40

41 The SA is progressive, beginning with two initial levels and then adding additional levels to the stack. This  
42 allows an iterative stacking approach, adding one level at a time, which furthermore allows for a bottom-up  
43 or top-down approach. Furthermore bottom-up and top-down results can be objectively merged to obtain a  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 better description of the detected events. This leads to an improved 4D view of the RV anomaly events, and  
2 hence of the structure of cyclones necessary for the advancement of atmospheric system dynamics and  
3 dynamic meteorology.  
4

5 SA is validated through its application to a winter season (JJA 2015) over the extratropical southern  
6 hemisphere, and in particular southern South America, where the Andes mountain range provides an  
7 orographic barrier that strongly impacts on the SH atmospheric circulation (Inatsu and Hoskins, 2004). SA  
8 validation is specifically carried out using cyclone tracks and features. Their behaviour is impacted by the  
9 Andes in different ways depending on latitude, giving valuable information on the capability of SA to  
10 capture such features.  
11

12 After taking into account specific issues pertaining to currently known impacts of major variability processes  
13 such as ENSO on the SH extratropical cyclone development, SA results are in very good agreement with  
14 current cyclone climatologies mostly based on single level studies and limited to the lower troposphere. The  
15 behaviour of the cyclones and their tracks during the sampled period verifies the known dynamic behaviour  
16 of cyclones and their link with upper atmospheric jets. Cyclone densities for all events agree well with the  
17 climatologies, and the specific analyses for the shallow, intermediate and well-developed cyclones show that  
18 the different vertical organizations have each preferred zones of occurrence.  
19

20 The three case studies, based on long term well-developed systems highlight the impact of the Andes  
21 depending on where the system reaches the mountain range. The Altas Cumbres region of the Andes (6000m  
22 and above) appears to disrupt the system. Hoskins and Hodges (2005) and Kreable et al. (2002) have  
23 identified this as a 500hPa cyclolysis region. The lower Patagonian Andes have a reduced impact limiting the  
24 development of the well-developed cyclone structure but once the systems cross the mountain range and  
25 enter the South Atlantic the development of the cyclone structures starts once more. Such development  
26 appears to take place near well-known cyclogenetic regions. These validation studies highlight the  
27 importance looking at the 3D temporal evolution of individual cyclones and other vorticity anomaly events  
28 in order to adequately understand their behaviour.  
29

30 Consequently, the STACKER Algorithm, using TRACK as the feature-tracking algorithm, demonstrates the  
31 applicability of dynamic programming for both 4D dynamic meteorology and dynamic climatology studies,  
32 as evidenced here by the two-stage validation. Understanding how complex 3D structures such as cyclones  
33 evolve in dynamically, spatially varying flows, and the classification of their main multilevel aspects and  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 behaviours remains a significant challenge. Furthermore, understanding how climate variability and climate  
2 change impact on cyclone and severe weather events and their distribution also requires a 4D perspective.  
3 SA is therefore a valuable tool for studies of vortex dynamics. A 4D-Stacker-based SH dynamic climatology  
4 is currently under way.  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

## 19 **5. Acknowledgments**

20  
21  
22  
23 The authors would like to thank CONICET, Universidad Tecnológica Nacional, Facultad Regional Buenos  
24 Aires, and Pontificia Universidad Católica Argentina. We would also like to thank UTN PID 4500, and  
25 Ministerio de Defensa PIDDEF 2014 No. 26 grants, and ECMWF for the ERAI reanalysis products as well  
26 to The NOAA Comprehensive Large Array-data Stewardship System for the GOES images. Finally we wish  
27 to thank the useful comments and suggestions made by the reviewers.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Appendix

The following paragraphs include the main definitions and concepts in the design and implementation of the STACKER algorithm. These can be references to Figures 1, 2 and 3 in the main body of text.

### 1. Points.

Points are zero-dimensional objects in a four-dimensional space. These are defined by a single value of longitude, latitude, height and time. These objects are generated at each selected height as detections of local maxima/minima of the relative vorticity (or other anomalies in T, Z or P) by TRACK during the preliminary stage of the analysis. Detections are assumed to be meteorological events, e.g., cyclones, anti-cyclones or other significant perturbations in the atmosphere, at a given instant or time frame. Detections in TRACK are not directly related to each other until the tracking optimization process is completed. Points are associated into Tracks by the tracking algorithm, e.g., TRACK.

### 2. Tracks.

Tracks are the main object obtained by the tracking algorithm. They consist of a series of points, from consecutive time-steps associated according to the criteria that weather features cannot change their position, direction, velocity or intensity abruptly. It is safe to assume that a given Track represents at least a fraction of a feature's translational behaviour during its life cycle. A set of points defining a Track will always be consecutive.

### 3. Stacks.

Stacks are the main output of the STACKER Algorithm. Stacks consist of a series of associated Tracks on several levels. These are four-dimensional objects, in a combination of discrete values within the longitude, latitude, height and time ranges chosen for the study (cf. Fig. 3). Stacks are always composed by Tracks stacked from consecutive levels (levels may correspond to pressure, model, PV or theta levels), i.e., the stacked detected event must be continuous in the vertical axis. It is possible to find several non-overlapping Tracks associated on the same level, because of the Point-to-Point linking approach, e.g., Stack 4 at the 850 hPa level in Fig.3. Tracks on a given Stack are directly related only to Tracks on the first-lower and first-upper level and are indirectly related to the rest of the Tracks in the Stack.

### 4 Similitude Function and Threshold Value.

1 The most important feature of the method is the design of the objective cost function and the selection of  
2 threshold values. The criterion to determine the likelihood of two Tracks being associated is defined in the  
3 Similitude Function (SF). It assigns a value to each optimization variable, or Link, and penalize, as a cost,  
4 dissimilar behaviour between detections, e.g., distance between Points, significant deviations in velocity or  
5 directions between Tracks, differences in size or intensity of the events, etc. This “Similitude Cost” must  
6 have a Threshold which states that any Link between two Points that have a high SF value is not relevant to  
7 the problem. If the two Points in a Link belong to two Tracks such that, at some point, two of its Points are  
8 possible links according to the SF criteria, then the Link should be considered relevant for the study (Fig. 3).  
9 The threshold will act both as a filter to avoid the build-up of a massive numerical problem to solve, and  
10 provide a “default” SF value for the solver to choose if there are no better options to associate a Point with  
11 another one.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

## 24 **5. Links.**

25  
26 Links represent the hypothesis that two single and simultaneous detections from two consecutive pressure  
27 levels, correspond to the same meteorological event. These are the association between any lower Point and  
28 any simultaneous upper Point. Each link is assigned a cost value according to the Similitude Cost Function  
29 (SCF), depending on the classification of the Link. There are as many optimization variables as there are  
30 Links created in the analysis. The solving process will decide which Links are more likely to couple two  
31 detections of the same meteorological event at adjacent levels. Since Links are rather complex compared to  
32 the other objects in the algorithm, they are classified according to their behaviour as True, False and Ghost  
33 Links.  
34  
35  
36  
37  
38  
39  
40  
41  
42

## 43 **6. Similitude Cost Function.**

44  
45 The Similitude Cost Function (SCF) sets the priority value on Links to be activated later in the solving  
46 process and is derived from the SF. Its value is assigned to a Link and later used as a coefficient in the  
47 Objective Function. Since the objective function must be minimized, lower cost values will be prioritized  
48 over others. Hence the importance of its criterion and Threshold value. Its desired behaviour is to prioritise  
49 True Links over the rest, penalize False Links, which exceed the present Threshold and present Ghost Links  
50 as “default” option to the solver. More specifically:  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1 **Threshold value.** In this study, Points are limited to longitude and latitude, in order to calculate distance. For  
2 more complex studies the core set of parameters can include other variables, e.g., altitude, wind speed,  
3 pressure, temperature, humidity, etc. The Threshold value is arbitrary and is specifically chosen for each  
4 study according to its aims, precision required and processing resources available. Here, following Lim and  
5 Simmonds (2007) the SF is equal to the “distance” between a lower and an upper Point, so the Threshold  
6 value can be understood as the radius of a circular surface centred at the projection of the Lower Point’s  
7 position on the upper level. “Distance” is defined as the arc length between the projections of the Points on a  
8 sphere with Earth’s mean radius.  
9

10  
11  
12  
13  
14  
15  
16  
17  
18 **True Links.** These are the essence of what Links represent. A True Link is generated between two  
19 simultaneous Points when its SF value does not exceed the Threshold value. It means that the relation  
20 between these two Points is relevant for the algorithm according to the current study criteria. Its Cost value is  
21 assigned directly from the Similitude Function and it will be smaller than the Threshold value.  
22  
23

24  
25  
26 **False Links.** False Links complement True Links. False Links are Links between two simultaneous Points  
27 where their SF value exceeds the Threshold value. The algorithm generates them because each Point belongs  
28 to a Stack that, at some time-step, is related by a True Link with the other Point’s Stack. Since the Stacker  
29 Algorithm is not allowed to associate single simultaneous Points without all of the other simultaneous Points  
30 between two Stacks, the algorithm False links complement what could not be True Linked. Its Cost value is  
31 equally assigned by the SF and it will be larger than the Threshold value, thus penalizing associations with a  
32 large number of these Links and prioritizing associations with larger numbers of True Links.  
33  
34  
35  
36  
37  
38  
39

40  
41 **Ghost Links.** Ghost Links relate two simultaneous Points, but one of them does not represent a local  
42 maximum/minimum detection from the Tracking Algorithm. These objects are artificially generated in order  
43 to give the linear programming solver a default alternative to choose when there are no better options to  
44 associate a Point with. Ghost Points are essentially a projection of a lower Point in the upper level in a given  
45 stage of the algorithm and do not belong to any Stack. The Stacker Algorithm specifically assigns the  
46 Threshold value to its Cost value in this case.  
47  
48  
49  
50  
51

## 52 **7. Binary Linear Programming.**

53  
54  
55 The STACKER Algorithm is a progressive optimization method. The whole problem is modelled so that it  
56 can be solved through Dynamic Programming by disassembling it into smaller problems. To achieve its final  
57  
58  
59  
60

1 solution, it iterates through a series of local stages which only includes the association of Points from two  
2 consecutive pressure levels. Each of these local problems are proposed as Linear Programming problems to  
3 guarantee the existence of an optimal solution. These problems can be solved by a specific case of Linear  
4 Integer Programming, called Binary Programming, in which the optimization variables can only take two  
5 possible values, either 1 or 0, i.e., activated or deactivated. This way the solver is proposed with several  
6 options to activate or deactivate (Links) according to the criteria of minimizing a Similitude Cost Function,  
7 respecting the restrictions that constraints provide.  
8  
9  
10  
11  
12  
13  
14

## 15 **10. Constraints.**

16 Constraints are, by far, the most complex and delicate features of the algorithm because of their  
17 progressively increasing size. These constraints condition the solver's behaviour in order to achieve the  
18 desired results, each addressing a specific need:  
19  
20  
21  
22  
23

24 **Must Be Together Constraint (MBT).** This constraint is responsible for assuring that Stacks will not be  
25 disassembled into smaller groups of Points by the solver. Since the aim of the algorithm is a Stack-to-Stack  
26 association using Point-to-Point links, if a single Link between an upper and lower Stack has been activated  
27 by the solver, all Links between those two same Stacks must be activated too.  
28  
29  
30  
31  
32

33 **Must Not Be Together Constraint (MNBT).** This set of constraints prevents the solver from associating  
34 Stacks that overlap with each other at some point (Fig. 2), i.e., two different Stacks that contain at least a  
35 single Point on the same surface and time-step, must be different events, and thus, must not be combined into  
36 the same Stack.  
37  
38  
39  
40

41 **Do Not Repeat Lower Tracks (DRLT).** This set of constraints forces the solver to activate at least one Link  
42 for every lower Point and makes it impossible for the solver to associate the same lower Point to more than  
43 one upper Point. In this case, for every set of Links containing the same lower Point, only one must be  
44 activated at the same time. The activation of a True or False Link represents the association between two  
45 Stacks. The activation of a Ghost Link indicates that the solver took the "default option" and any other Links  
46 available for its lower Point were not optimal for the solution of the local problem.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 **Do Not Repeat Higher Tracks.** Like the previous set of constraints, DRHT makes it impossible for the  
2 solver to associate the same upper Point to more than one lower Point. The only difference lies in the fact  
3 that it will not force the solver to activate at least one Link for every upper Point.  
4  
5  
6

### 7 **11. Binary Solution Parsing.**

8  
9  
10 Once the Linear Programming process is complete, the solver returns all activated and deactivated Links.  
11 Deactivated Links are irrelevant to the parsing, as well as active Ghost Links and have no practical meaning.  
12 Active True and False Links represent the most coherent associations between lower and upper Stacks  
13 according to the SF criteria. Parsing must take all of the associated Stacks' Points and build a new set of  
14 Stacks on as many levels as the previous two sets combined. Point-to-Point linkage allows the association of  
15 any number of Stacks together:  
16  
17  
18  
19  
20  
21

- 22 • A Stack may have no coherent associations, and hence no changes are required.
- 23 • Any number of lower Stacks can be associated with any number of higher Stacks, building a new  
24 Stack with as many levels as the previous Stacks combined had and all of their Points.  
25  
26  
27  
28

29  
30 New Stacks will become lower (upper) Stacks on the next iteration of the problem or will become the final  
31 solution if there is no further stacking possible. If the process is repeated in the opposite vertical direction  
32 (Top-to-Bottom or Bottom-to-Top) the Stacker Algorithm will provide a similar but somewhat different  
33 solution. Merging Top-to-Bottom and Bottom-to-top stacks combines all detected features into a single stack  
34 for subsequent analysis.  
35  
36  
37  
38  
39

### 40 **12. Stack Concatenation.**

41  
42 In this study, the Linear Programming package from the R programming language was used to implement  
43 the algorithm. Due to computational limitations, the size of the problem becomes unsolvable for large  
44 amounts of variables (over 10000) and a Stack concatenator process is proposed. Concatenation consists of  
45 identifying the same meteorological event through several multi-level Stacks obtained in short samples and  
46 combining all of the sections into a single, longer Stack. To this end, we used an overlap of five days  
47 between datasets and assumed that Stacks that shared at least one detection (Point) during the overlapping  
48 period must be the same event and thus must be combined. This is possible since in principle the analysis of  
49 fields by tracking algorithms does not suffer from time sampling edge effects. The concatenation process is  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 an approximation and may produce some overlapping and occasional discrepancies in the combined Stacks.  
2  
3 Since this impacts less than 0.005% of the whole Stack set, it does not compromise the coherence of the  
4  
5 algorithm's results.  
6  
7  
8  
9  
10  
11  
12

## 13 6. References.

14  
15  
16 Ayrault F. et A. Joly. 2000. L'origine des dépressions météorologiques sur l'Atlantique: nouvelle perspective  
17 climatologique Comptes-Rendus de l'Académie des Sciences de Paris, Sciences de la Terre et des Planètes,  
18  
19 **330**, 173--178.  
20

21  
22  
23 Bartholy, J., Pongrácz, R., and Pattantyús-Ábrahám, M.. 2008. Analyzing the genesis, intensity, and  
24  
25 tracks of western Mediterranean cyclones. *Theor. Appl. Climatol.*, **96**, 133–144.  
26

27  
28 Catto, J.L., L.C. Shaffrey, and K.I. Hodges. 2010. Can Climate Models Capture the Structure of  
29  
30 Extratropical Cyclones?, *J. Climate*, **23**, 1621–1635, doi:10.1175/2009JCLI3318.1.  
31

32  
33 Chen, L., B. Tan, N. G. Kvamsto and O. M. Johannessen. 2014. Wintertime cyclone/anticyclone activity  
34  
35 over China and its relation to upper tropospheric jets. *Tellus*, **66A**, 21889, doi: 10.3402/tellusa.v66.21889.  
36

37  
38 Dacre, H.F., M.K. Hawcroft, M.A. Stringer, and K.I. Hodges. 2012. An Extratropical Cyclone Atlas: A Tool  
39  
40 for Illustrating Cyclone Structure and Evolution Characteristics. *Bull. Amer. Meteor. Soc.*, **93**, 1497–1502  
41

42  
43 Finnis, J., Holland, M.M., Serreze, M C., and Cassano, J J. 2007. Response of Northern Hemisphere  
44  
45 extratropical cyclone activity and associated precipitation to climate change, as represented by the  
46  
47 Community Climate System Model. *J. Geophys. Res.* **112**, G04S42.  
48

49  
50 Flaounas, E., Kotroni, V., Lagouvardos, K., and Flaounas, I.: CycloTRACK. 2014. Tracking winter  
51  
52 extratropical cyclones based on relative vorticity: sensitivity to data filtering and other relevant parameters,  
53  
54 *Geosci. Model Dev.*, **7**, 1841-1853, doi:10.5194/gmd-7-1841-2014.  
55

56  
57 Froude, L.S. 2010. TIGGE: Comparison of the Prediction of Northern Hemisphere Extratropical Cyclones by  
58  
59 Different Ensemble Prediction Systems. *Wea. Forecasting*, **25**, 819–836, doi:10.1175/2010WAF2222326.1  
60

1 Froude, L.S., L. Bengtsson, and K.I. Hodges. 2007. The Prediction of Extratropical Storm Tracks by the  
2 ECMWF and NCEP Ensemble Prediction Systems, *Mon. Wea. Rev.*, **135**, 2545–2567.  
3  
4

5  
6  
7  
8 Gozzo, L.F., R.P. da Rocha, M.S. Reboita, and S. Sugahara. 2014. Subtropical Cyclones over the  
9 Southwestern South Atlantic: Climatological Aspects and Case Study. *J. Climate*, **27**, 8543–8562,  
10 doi:10.1175/JCLI-D-14-00149.1.  
11  
12  
13

14 Grieger, J., Leckebusch, Gregor C., Raible, Christoph C., Rudeva, I and Simmonds, I. 2018. Subantarctic  
15 cyclones identified by 14 tracking methods, and their role for moisture transports into the continent, *Tellus*  
16 *A: Dynamic Meteorology and Oceanography*, **70**, doi: 10.1080/16000870.2018.1454808  
17  
18  
19

20  
21 Hakim, G.J. 2003. Cyclogenesis, in *Encyclopedia of Atmospheric Sciences*, 589-594, Holton, J.R., Curry,  
22 J.A., Pyle, J.A., eds., Academic Press.  
23  
24

25  
26 Hanson, C.E., Palutikof, J.P., and Davies, T.D. 2004. Objective cyclone climatologies of the North  
27 Atlantic: A comparison between the ECMWF and NCEP Reanalyses. *Climate Dynamics* **22**, 757–769.  
28  
29

30  
31 Hewson, T. D. and Titley, H. A. 2010. Objective identification, typing and tracking of the complete life-  
32 cycles of cyclonic features at high spatial resolution. *Meteor. Appl.* **17**, 355–381. doi:10.1002/met.204.  
33  
34

35  
36 Hodges, K. I. 1994. A general-method for tracking analysis and its application to meteorological data.  
37 *Monthly Weather Review*, **122**, 2573-2586. doi:10.1175/1520-0493(1994)122<2573:AGMFTA>2.0.CO;2  
38

39  
40 Hodges, K. I. 1995. Feature tracking on the unit-sphere. *Monthly Weather Review*, **123**, 3458-3465,  
41 doi: 10.1175/1520-0493(1995)123<3458:FTOTUS>2.0.CO;2.  
42

43  
44 Hodges, K. I. 1999. Adaptive constraints for feature tracking. *Monthly Weather Review*, **127**, 1362-1373,  
45 doi: 10.1175/1520-0493(1999)127<1362:ACFFT>2.0.CO;2  
46  
47

48  
49 Hodges, K.I. 2008. Confidence intervals and significance tests for spherical data derived from feature  
50 tracking. *Monthly Weather Review*, **136**, 1758-1777. doi:10.1175/2007MWR2299.1  
51

52  
53 Hodges KI, Hoskins BJ, Boyle J, Thorncroft C. 2003. A comparison of recent reanalysis datasets using  
54 objective feature tracking: Storm tracks and tropical easterly waves, **131**, 2012, *Mon. Wea. Rev.*, 132, 1325-  
55 1327, doi: 10.1175/1520-0493(2003)131<2012:ACORRD>2.0.CO;2.  
56  
57  
58  
59  
60

1 Homar, V., R., Romero, D. J. Stensrud, C. Ramis, and S. Alonso. 2003. Numerical diagnosis of a small,  
2 quasi-tropical cyclone over the western Mediterranean: Dynamical vs. boundary factors, *Q. J. R. Met. Soc.*,  
3 **129**, 1469–1490.

4  
5  
6  
7  
8 [Hoskins, B. J. and P. D. Sardeshmukh. 1984. Spectral smoothing on the sphere. \*Mon. Wea. Rev.\*, \*\*112\*\*, 2524–  
9 2529.](#)

10 Hoskins, B. J., & Valdes, P. J. 1990. On the Existence of Storm Tracks. *J. Atmos. Sci.*, **47**, 1854 - 1864.

11  
12  
13 Hoskins, B. J. and Hodges, K. I. 2002. New perspectives on the Northern Hemisphere winter storm tracks, *J.*  
14 *Atmos. Sci.*, **59**, 1041–1061, 2002.

15  
16  
17  
18 Hoskins BJ, Hodges K. I. 2005. A new perspective on Southern Hemisphere storm tracks, *J. Clim.*, **18**, 4108–  
19 4129.

20  
21  
22 Inatsu M. and Hoskins B.J. 2004. The zonal asymmetry of the Southern Hemisphere winter storm track, *J.*  
23 *Clim.*, **17**, 4882-4892.

24  
25  
26  
27 IPCC, 2007: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the  
28 Fourth Assessment Report of the Intergovernmental Panel on Climate Change [Solomon, S., D. Qin, M.  
29 Manning, Z. Chen, M. Marquis, K.B. Averyt, M.Tignor and H.L. Miller (eds.)]. Cambridge University Press,  
30 Cambridge, United Kingdom and New York, NY, USA.

31  
32  
33  
34  
35 IPCC, 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the  
36 Fifth Assessment Report of the Intergovernmental Panel on Climate Change, R.K. Pachauri and L.A. Meyer  
37 (eds.), IPCC, Geneva, Switzerland, 151 pp.

38  
39  
40  
41 Jansa, A., Genoves, A., Picornell, M.A., Campins, J., Riosalido, R., and Carretero, O. 2001. Western  
42 Mediterranean cyclones and heavy rain. Part 2: Statistical approach. *Meteorol. Appl.* **8**, 43–56.

43  
44  
45  
46 [Joly, A., Ayrault, F. and Malardel, S. 2003. Cyclones, Extratropical, in \*Encyclopedia of Atmospheric\*  
47 \*Sciences\*, 594-615, Holton, J.R., Curry, J.A., Pyle, J.A., eds., Academic Press.](#)

48  
49  
50  
51 Keable, M. , Simmonds, I. and Keay, K. 2002. Distribution and temporal variability of 500 hPa cyclone  
52 characteristics in the Southern Hemisphere. *Int. J. Climatol.*, **22**, 131-150. doi:10.1002/joc.728.

53  
54  
55 Kelemen FD, Bartholy J, Pongracz R. 2015. Multivariable cyclone analysis in the Mediterranean region. *Q.*  
56 *J. Hung Meteorol Serv* **119**, 159–184

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Lim, E.-P., and I. Simmonds. 2007. Southern Hemisphere winter extra-tropical cyclone characteristics and vertical organization observed with the ERA-40 reanalysis data in 1979–2001, *J. Clim.*, **20**, 2675–2690, doi:10.1175/JCLI4135.1.

Mailier, P.J., D.B. Stephenson, C.A. Ferro, and K.I. Hodges. 2006. Serial Clustering of Extratropical Cyclones. *Mon. Wea. Rev.*, **134**, 2224–2240.

Murray, R. J. and Simmonds, I. 1991a. A numerical scheme for tracking cyclone centres from digital data. Part I: Development and operation of the scheme. *Aust. Meteor. Mag.* **39**, 155–166.

Murray, R. J. and Simmonds, I. 1991b. A numerical scheme for tracking cyclone centres from digital data. Part II: Application to January and July GCM simulations. *Aust. Meteor. Mag.* **39**, 167–180.

Neu, U., M.G. Akperov, N. Bellenbaum, R. Benestad, R. Blender, R. Caballero, A. Coccozza, H.F. Dacre, Y. Feng, K. Fraedrich, J. Grieger, S. Gulev, J. Hanley, T. Hewson, M. Inatsu, K. Keay, S.F. Kew, I. Kindem, G.C. Leckebusch, M.L. Liberato, P. Lionello, I.I. Mokhov, J.G. Pinto, C.C. Raible, M. Reale, I. Rudeva, M. Schuster, I. Simmonds, M. Sinclair, M. Sprenger, N.D. Tilinina, I.F. Trigo, S. Ulbrich, U. Ulbrich, X.L. Wang, and H. Wernli. 2013. IMILAST: A Community Effort to Intercompare Extratropical Cyclone Detection and Tracking Algorithms. *Bull. Amer. Meteor. Soc.*, **94**, 529–547, <https://doi.org/10.1175/BAMS-D-11-00154.1>

Park, S. K., D. -L. Zhang, and H. -H. Kim. 2008. Impact of dropwindsonde data on the track forecasts of a tropical cyclone: An observing-systems simulation experiment study. *Asia-Pacific J. Atmos. Sci.*, **44**, 85-92.

Petterssen, S. 1956. *Weather Analysis and Forecasting*. 2nd Edition, Vol. I, McGraw-Hill Book Co., Inc., New York.

Pezza, A. B. & Ambrizzi, T. 2003. Variability of Southern Hemisphere cyclone and anticyclone behavior: Further analysis. *J. Clim.* **16**, 1075–1083.

Pezza, A.B., and Simmonds, I. and Renwick, J.A. 2007. Southern hemisphere cyclones and anticyclones: recent trends and links with decadal variability in the Pacific Ocean, *Int. J. Climatol.*, **27**, 1403-1419, doi: 10.1002/joc.1477.

Pezza, A. B., Durrant, T., Simmonds, I. and Smith, I. 2008. Southern Hemisphere synoptic behavior in extreme phases of SAM, ENSO, sea ice extent, and southern Australia rainfall. *J. Clim.* **21**, 5566–5584.

1 Pezza, A. B., Rashid, H. and Simmonds, I. 2012. Climate links and recent extremes in Antarctic sea ice,  
2 high-latitude cyclones, Southern Annular Mode and ENSO. *Clim. Dyn.* **38**, 57–73. doi:10.1007/s00382-011-  
3 1044-y.  
4  
5  
6

7  
8 Picornell, M.A., Jansá, A., Genovés, A., and Campins, J. 2001. Automated database of mesocyclones from  
9 the HIRLAM(INM)-0.5° analyses in the western Mediterranean. *Int. J. Climatol.* **21**, 335–354.  
10  
11

12 Pinheiro, H.R., Hodges, K.I., Gan, M.A. et al., 2017, A new perspective of the climatological features of  
13 upper-level cut-off lows in the Southern Hemisphere, *Clim. Dyn.* **48**, 541-559. doi:10.1007/s00382-016-  
14 3093-8.  
15  
16  
17

18 Reboita, M.S., Ambrizzi, T., da Rocha, R.P. 2009. Relationship between the southern annular mode and  
19 southern hemisphere atmospheric systems, *Revista Brasileira de Meteorologia*, **24**, 48-55  
20  
21  
22

23 Reboita, M.S., da Rocha, R.P., Ambrizzi, T., and Gouveia, C. 2015. Trend and teleconnection patterns in the  
24 climatology of extratropical cyclones over the Southern Hemisphere, *Clim. Dyn.*, **45**, 1929-1944,  
25 doi:10.1007/s00382-014-2447-3.  
26  
27  
28

29 Satake, K., Inatsu M., Mori M., Hasegawa A. 2013. Tropical Cyclone Tracking Using a Neighbor Enclosed  
30 Area Tracking Algorithm, *Mon. Wea. Rev.*, **141**, 3539-3553, doi:10.1175/MWR-D-12-00092.1  
31  
32  
33

34 Serra, Y.L., Kiladis, G.N., and Hodges, K.I. 2010. Tracking and Mean Structure of Easterly Waves over the  
35 Intra-Americas Sea, *J. Clim.*, **23**, 4823-4840.  
36  
37  
38

39 Trigo, I.F., Davies, T.D., and Bigg, G.R. 1999. Objective Climatology of Cyclones in the Mediterranean  
40 Region. *J. Clim.*, **12**, 1685–1696.  
41  
42

43 van Bebber WJ. 1891. Die Zugstrassen der barometrischen Minima nach den Bahnenkarten der Deutschen  
44 Seewarte für den Zeitraum von 1870 – 1890. *Meteorolog. Zeitschrift*, **8**, 361-366.  
45  
46

47 Zappa, G., L.C. Shaffrey, K.I. Hodges, P.G. Sansom, and D.B. Stephenson. 2013. A Multimodel Assessment  
48 of Future Projections of North Atlantic and European Extratropical Cyclones in the CMIP5 Climate Models.  
49 *J. Clim.*, **26**, 5846–5862.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1 Zappa, G., G. Masato, L. Shaffrey, T. Woollings, and K. Hodges. 2014. Linking Northern Hemisphere  
2 blocking and storm track biases in the CMIP5 climate models, *Geophys. Res. Lett.*, **41**, 135–139, doi:  
3  
4 10.1002/2013GL058480.  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

1 **Figure Captions**

2  
3  
4  
5 **Figure 1**

6  
7 Stacker algorithm flow chart, highlighting the main steps in the track sorting and stacking process  
8 optimization approach.  
9

10  
11  
12  
13 **Figure 2**

14  
15 Schematic of the linkage criteria for track stacking. The figure highlights the maximum radius threshold used  
16 in determining which tracks are linked.  
17  
18

19  
20  
21 **Figure 3**

22  
23 Process diagram showing successive steps in the stacking process (one way) with a simple example which  
24 includes fragmented tracks at one level.  
25  
26

27  
28  
29 **Figure 4**

30  
31 JJA 2015 mean wind velocity at 300hPa.  
32  
33

34  
35 **Figure 5.**

36  
37 JJA 2015 SH Multilevel extratropical cyclone density field (cyclones detected at two or more levels).  
38  
39

40  
41 **Figure 6**

42  
43 JJA 2015 Shallow cyclone event densities over the SH: a. all two/three-level events, b. near-surface events,  
44 c. middle troposphere events and d. upper troposphere events.  
45  
46

47  
48  
49 **Figure 7**

50  
51 JJA 2015 Intermediate cyclone event densities over the SH: a. all intermediate events, b. lower to middle  
52 troposphere events and c. middle to upper troposphere events.  
53  
54

55  
56  
57 **Figure 8**

1 JJA 2015 Well-developed cyclone event densities over the SH, spanning most of the troposphere.  
2  
3  
4

5 **Figure 9**

6 Validation Event 1 July 29<sup>th</sup> – August 10<sup>th</sup> 2015. a. Cyclone trajectory map. b. oblique-view perspective of  
7 cyclone trajectory showing the vertical stack as the system evolves.  
8  
9  
10

11  
12  
13 **Figure 10**

14 Validation Event 1 July 29<sup>th</sup> – August 10<sup>th</sup> 2015. Snapshots of the evolution of the event as seen in i. GOES  
15 IR channel image, ii. GOES WV channel image, iii. ERA-I 500hPa cloud cover, 500hPa RV anomaly and  
16 Stacker trajectory iv. ERA-I 300hPa winds, with velocities over 40ms-1 highlighted in shades of blue,  
17 500hPa RV anomaly and Stacker trajectory., v. 300hPa geopotential height and vi. 1000-500hPa thickness  
18 map, both centred at the times shown. Track position shown with a large cross in GOES imagery. Anomaly  
19 structure at 500hPa is shown together with track in iii. And iv. a. 4<sup>th</sup> August, 2015, at 12:00 UTC, b. 6<sup>th</sup>  
20 August, 00:00 UTC, c. 7<sup>th</sup> August 2015, 12:00 UTC and d. 9<sup>th</sup> August, 2015, at 00:00 UTC.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 **Figure 11**

32 Validation Event 2 June 27<sup>th</sup> -July 11<sup>th</sup>, 2015. Same as Fig. 9 but for validation event 2.  
33  
34  
35  
36

37 **Figure 12**

38 Validation Event 2 June 27<sup>th</sup> -July 11<sup>th</sup>, 2015. Same as Fig. 11 but for validation event 2. a. June 30<sup>th</sup>, 2015 at  
39 18:00 UTC, b. July 2<sup>nd</sup>, 2015 at 06:00 UTC, c. July 3<sup>rd</sup>, 2015, at 18:00 UTC and d. July 5<sup>th</sup>, 2015, at 06:00  
40 UTC.  
41  
42  
43  
44  
45  
46

47 **Figure 13**

48 Validation Event 3 July 10<sup>th</sup> - July 24<sup>th</sup>, 2015. Same as Fig. 9 but for validation event 3.  
49  
50  
51  
52

53 **Figure 14.**  
54  
55  
56  
57  
58  
59  
60

1 Validation Event 3 July 10<sup>th</sup> - July 24<sup>th</sup>, 2015. Same as Fig. 11 but for validation event 3. a. July 18<sup>th</sup>, 2015, at  
2  
3 06:00 UTC, b. July 19<sup>th</sup>, 2015, at 18:00 UTC, c. July 21<sup>st</sup>, 2015 at 06:00 UTC, and d. July 22<sup>nd</sup>, 2015, at  
4  
5 18:00 UTC.  
6  
7  
8

9  
10 **Figure 15**

11 Snapshots of the vertical structure of the associated RV anomalies at different levels at a given time: a.  
12 validation event 1, 8<sup>th</sup> August, 2015, 18:00 UTC b. validation event 2, 5<sup>th</sup> July, 2015, 06:00 UTC and c.  
13 validation event 3, 22<sup>nd</sup> July, 2015, 12:00 UTC.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


For Peer Review

## A 4D Feature Tracking Algorithm: a multidimensional view of cyclone systems

S. Gabriela Lakkis<sup>b,a</sup>, Pablo Canziani<sup>a,c</sup>, Adrián Yucheche<sup>a,c</sup>, Leandro Rocamora<sup>a</sup>,  
Agustin Caferri<sup>a</sup>, Kevin Hodges<sup>d</sup>, Alan O'Neill<sup>d</sup>

- a. Unidad de Investigación y Desarrollo de las Ingenierías, Facultad Regional Buenos Aires, Universidad Tecnológica Nacional, Buenos Aires, Argentina
- b. Facultad de Ingeniería y Ciencias Agrarias, Pontifica Universidad Católica Argentina, Buenos Aires, Argentina
- c. Consejo Nacional de Investigaciones Científicas y Técnicas, CONICET, Buenos Aires, Argentina
- d. Department of Meteorology, University of Reading, Reading, U.K.


Contact: Dr. Gabriela Lakkis, mail: [gabylakkisetul@gmail.com](mailto:gabylakkisetul@gmail.com)



An objective 4D algorithm developed to track extratropical relative vorticity anomaly 3D structure over time is introduced and validated. The mean distribution of SH cyclone systems detected by Stacker during 2015 winter is shown in the upper image. Stacker provides the detected anomaly tracks corresponding to a given RV anomaly event as well as the anomaly itself. The lower image shows a cyclone event highlighting the tracks at levels where the event was detected as well as the RV anomalies, which show the anomaly 3D structure.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


For Peer Review



Stacker algorithm flow chart, highlighting the main steps in the track sorting and stacking process optimization approach.

210x297mm (96 x 96 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Schematic of the linkage criteria for track stacking. The figure highlights the maximum radius threshold used in determining which tracks are linked.

177x86mm (96 x 96 DPI)

Peer Review






Process diagram showing successive steps in the stacking process (one way) with a simple example which includes fragmented tracks at one level.

190x190mm (96 x 96 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



JJA 2015 mean wind velocity at 300hPa.

231x231mm (96 x 96 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



JJA 2015 SH Multilevel extratropical cyclone density field (cyclones detected at two or more levels).


231x231mm (96 x 96 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



JJA 2015 Shallow cyclone event densities over the SH: a. all two/three-level events,  
231x231mm (96 x 96 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




JA 2015 Shallow cyclone event densities over the SH:b. near-surface events

231x231mm (96 x 96 DPI)




1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



JA 2015 Shallow cyclone event densities over the SH:c. middle troposphere events  
231x231mm (96 x 96 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



JA 2015 Shallow cyclone event densities over the SH:d. upper troposphere events.

231x231mm (96 x 96 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



JJA 2015 Intermediate cyclone event densities over the SH: a. all intermediate events,  
231x231mm (96 x 96 DPI)




1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



JJA 2015 Intermediate cyclone event densities over the SH: b. lower to middle troposphere events

231x231mm (96 x 96 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



JJA 2015 Intermediate cyclone event densities over the SH: c. middle to upper troposphere events.

231x231mm (96 x 96 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



JJA 2015 Well-developed cyclone event densities over the SH, spanning most of the troposphere.

231x231mm (96 x 96 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Validation Event 1 July 29th – August 10th 2015. a. Cyclone trajectory map.

371x172mm (70 x 70 DPI)

Peer Review




Validation Event 1 July 29th – August 10th 2015. b. oblique-view perspective of cyclone trajectory showing the vertical stack as the system evolves.

371x165mm (70 x 70 DPI)

Peer Review


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Validation Event 1 July 29th – August 10th 2015. Snapshots of the evolution of the event as seen in i. GOES IR channel image, ii. GOES WV channel image, iii. ERA-I 500hPa cloud cover, 500hPa RV anomaly and Stacker trajectory iv. ERA-I 300hPa winds, with velocities over 40ms<sup>-1</sup> highlighted in shades of blue, 500hPa RV anomaly and Stacker trajectory., v. 300hPa geopotential height and vi. 1000-500hPa thickness map, both centred at the times shown. Track position shown with a large cross in GOES imagery. Anomaly structure at 500hPa is shown together with track in iii. And iv. a. 4th August, 2015, at 12:00 UTC,


240x222mm (108 x 108 DPI)





Validation Event 1 July 29th – August 10th 2015. Snapshots of the evolution of the event as seen in i. GOES IR channel image, ii. GOES WV channel image, iii. ERA-I 500hPa cloud cover, 500hPa RV anomaly and Stacker trajectory iv. ERA-I 300hPa winds, with velocities over 40ms<sup>-1</sup> highlighted in shades of blue, 500hPa RV anomaly and Stacker trajectory., v. 300hPa geopotential height and vi. 1000-500hPa thickness map, both centred at the times shown. Track position shown with a large cross in GOES imagery. Anomaly structure at 500hPa is shown together with track in iii. And iv. b. 6th August, 00:00 UTC,


240x222mm (108 x 108 DPI)



Validation Event 1 July 29th – August 10th 2015. Snapshots of the evolution of the event as seen in i. GOES IR channel image, ii. GOES WV channel image, iii. ERA-I 500hPa cloud cover, 500hPa RV anomaly and Stacker trajectory iv. ERA-I 300hPa winds, with velocities over 40ms<sup>-1</sup> highlighted in shades of blue, 500hPa RV anomaly and Stacker trajectory., v. 300hPa geopotential height and vi. 1000-500hPa thickness map, both centred at the times shown. Track position shown with a large cross in GOES imagery. Anomaly structure at 500hPa is shown together with track in iii. And iv. c. 7th August 2015, 12:00 UTC and


240x222mm (108 x 108 DPI)





Validation Event 1 July 29th – August 10th 2015. Snapshots of the evolution of the event as seen in i. GOES IR channel image, ii. GOES WV channel image, iii. ERA-I 500hPa cloud cover, 500hPa RV anomaly and Stacker trajectory iv. ERA-I 300hPa winds, with velocities over 40ms<sup>-1</sup> highlighted in shades of blue, 500hPa RV anomaly and Stacker trajectory., v. 300hPa geopotential height and vi. 1000-500hPa thickness map, both centred at the times shown. Track position shown with a large cross in GOES imagery. Anomaly structure at 500hPa is shown together with track in iii. And iv. d. 9th August, 2015, at 00:00 UTC.


240x222mm (108 x 108 DPI)



Validation Event 1 July 29th – August 10th 2015. Snapshots of the evolution of the event as seen in i. GOES IR channel image, ii. GOES WV channel image, iii. ERA-I 500hPa cloud cover, 500hPa RV anomaly and Stacker trajectory iv. ERA-I 300hPa winds, with velocities over 40ms<sup>-1</sup> highlighted in shades of blue, 500hPa RV anomaly and Stacker trajectory., v. 300hPa geopotential height and vi. 1000-500hPa thickness map, both centred at the times shown. Track position shown with a large cross in GOES imagery. Anomaly structure at 500hPa is shown together with track in iii. And iv. a. 4th August, 2015, at 12:00 UTC

371x174mm (70 x 70 DPI)


Review



Validation Event 1 July 29th – August 10th 2015. Snapshots of the evolution of the event as seen in i. GOES IR channel image, ii. GOES WV channel image, iii. ERA-I 500hPa cloud cover, 500hPa RV anomaly and Stacker trajectory iv. ERA-I 300hPa winds, with velocities over 40ms<sup>-1</sup> highlighted in shades of blue, 500hPa RV anomaly and Stacker trajectory., v. 300hPa geopotential height and vi. 1000-500hPa thickness map, both centred at the times shown. Track position shown with a large cross in GOES imagery. Anomaly structure at 500hPa is shown together with track in iii. And iv. a. 4th August, 2015, at 12:00 UTC, b. 6th August, 00:00 UTC,

371x171mm (70 x 70 DPI)


Review



Validation Event 2 June 27th -July 11th, 2015. Same as Fig. 11 but for validation event 2. a. June 30th, 2015 at 18:00 UTC

240x222mm (108 x 108 DPI)






Validation Event 2 June 27th -July 11th, 2015. Same as Fig. 11 but for validation event 2. b. July 2nd, 2015 at 06:00 UTC


240x222mm (108 x 108 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Validation Event 2 June 27th -July 11th, 2015. Same as Fig. 11 but for validation event 2. c. July 3rd, 2015, at 18:00 UTC.


240x222mm (108 x 108 DPI)



Validation Event 2 June 27th -July 11th, 2015. Same as Fig. 11 but for validation event 2. d. July 5th, 2015, at 06:00 UTC.

240x222mm (108 x 108 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Validation Event 3 July 10th- July 24th, 2015. Same as Fig. 9 but for validation event 3.

371x172mm (70 x 70 DPI)

Peer Review






Validation Event 3 July 10th- July 24th, 2015. Same as Fig. 9 but for validation event 3.

371x168mm (70 x 70 DPI)


Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Validation Event 3 July 10th- July 24th, 2015. Same as Fig. 11 but for validation event 3. a. July 18th, 2015, at 06:00 UTC,


240x222mm (108 x 108 DPI)



Validation Event 3 July 10th- July 24th, 2015. Same as Fig. 11 but for validation event 3. b. July 19th, 2015, at 18:00 UTC,

240x222mm (108 x 108 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Validation Event 3 July 10th- July 24th, 2015. Same as Fig. 11 but for validation event 3. , c. July 21st, 2015 at 06:00 UTC,

240x222mm (108 x 108 DPI)




Wed Jul 22 18:00:00 UTC 2015



Validation Event 3 July 10th- July 24th, 2015. Same as Fig. 11 but for validation event 3. d. July 22nd, 2015, at 18:00 UTC.

240x222mm (108 x 108 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Snapshots of the vertical structure of the associated RV anomalies at different levels at a given time: a. validation event 1, 8th August, 2015, 18:00 UTC

371x166mm (70 x 70 DPI)

Peer Review




Snapshots of the vertical structure of the associated RV anomalies at different levels at a given time: b. validation event 2, 5th July, 2015, 06:00 UTC

371x163mm (70 x 70 DPI)

Peer Review



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Snapshots of the vertical structure of the associated RV anomalies at different levels at a given time: c. validation event 3, 22nd July, 2015, 12:00 UTC.

371x168mm (70 x 70 DPI)

Peer Review