

Introduction

Over the last decade, researchers have suggested that endurance performance can be further enhanced by consuming a carbohydrate and protein (CHO-PRO) supplement when compared to a carbohydrate only supplement (CHO) (Ivy, Res, Sprague & Widzer, 2003; Saunders, Kane & Todd, 2004). However, the practical implications of these studies are held back by a number of methodological issues (Van Essen & Gibala, 2006).

Aims

The purpose of this study was to determine whether the inclusion of protein into a carbohydrate drink would improve endurance performance, when compared to a carbohydrate drink and a nonenergetic sweetened placebo (PL) in a cycle to exhaustion at 70% $\text{VO}_{2\text{max}}$.

Method

Seven male university students (age: 21 ± 1 years, height: 174.1 ± 3.4 cm, mass: 71.9 ± 3.5 kg, $\text{VO}_{2\text{max}}$: 44 ± 10 $\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) performed a cycle to exhaustion on three occasions separated by 7 days. In a double-blinded, placebo controlled, repeated measures crossover study, participants ingested CHO-PRO, CHO, PL at a rate of 500 ml, 50 min before and 500 ml during exercise. Both CHO and CHO-PRO beverages were matched for total calories (148 kcal). Carbohydrate content was 34.7 g & 35.3 g for the CHO-PRO and CHO respectively. The CHO-PRO beverages contained an addition of 3.9 g of protein. Data analysed included distance travelled, time to exhaustion, blood glucose and blood lactate concentration. Statistical analyses include a one way repeated measures ANOVA with a pairwise comparison.




Figure 5. Blood glucose levels before and during exercise. Group means are presented at each time point. * $P < 0.05$ between supplemented drinks and PL.




Figure 2. Mean (\pm SD) time to exhaustion at 70% $\text{VO}_{2\text{max}}$ for participants that consumed a PL, CHO and CHO-PRO drink before and during exercise. * $P = 0.04$ between PRO and PL; ** $P = 0.05$ between CHO and PL.

Results

As predicted for time to exhaustion, distance travelled and glucose levels, CHO and CHO-PRO significantly increased results when compared to the PL condition ($P < 0.05$). However no significant difference was observed between CHO and CHO-PRO in time to exhaustion (45.23 ± 25.32 min vs 48.92 ± 28.42 min; $P = 0.39$) (Fig. 2), distance travelled (24.09 ± 13.43 km vs 24.88 ± 13.45 km; $P = 0.44$), average blood glucose (5.687 ± 0.28 $\text{mmol}\cdot\text{l}^{-1}$ vs 5.690 ± 0.37 $\text{mmol}\cdot\text{l}^{-1}$; $P = 1.000$) and glucose over time ($P = 0.95$) (Fig. 1). Variables such as lactate, heart rate and RPM were not significantly different ($P > 0.05$) between all three trials.

Discussion

Ingesting a CHO and CHO-PRO beverage 50 minutes before and during exercise will improve endurance performance when compared to a PL condition ($P < 0.05$), however a CHO-PRO beverage will not significantly further enhance performance when compared to a CHO only supplement ($P > 0.05$). Further research needs to be studied in this area, as there are clear conflicting conclusions amongst the literature.

Acknowledgements

The authors thank David Sims, Matthew Bradley, Craig Boyd, Garry Pheasey and Douglas Herbet for their excellent technical assistance. We also thank Orbana for providing the supplemented CHO-PRO powder. This work was supported by Manchester Metropolitan University.

References

- Ivy, J. L., Res, P. T., Sprague, R. C., & Widzer, M. O. (2003). Effect of a carbohydrate-protein supplement on endurance performance during exercise of varying intensity. *International Journal of Sport Nutrition and Exercise Metabolism*, 13(3), 382-395.
- Saunders, M. J., Kane, M. D., & Kent Todd, M. (2004). Effects of a carbohydrate-protein beverage on cycling endurance and muscle damage. *Medicine and Science in Sports and Exercise*, 36(7), 1233-1238.
- Van Essen, M., & Gibala, M. J. (2006). Failure of protein to improve time trial performance when added to a sports drink. *Medicine and Science in Sports and Exercise*, 38(8), 1476-1483.