

Abstract:

Background: *Managing chronic wounds is associated with a burden to patients, caregivers, health services and society. The cost of treating these conditions is increasing and there is a lack of clarity regarding the role of dressings in improving outcomes. With chronic wounds becoming more prevalent, and the cost of wound management increasing, strategies using dressings to minimise the healthcare burden are essential.*

Methods: *A systematic review of the literature was carried out on the role of dressings in DFU, and VLU management strategies, their effectiveness, associated resource use/cost, and quality of life burden for patients. From this evidence base statements were written, regarding chronicity in wounds, burden of illness, healing time, and the role of matrix metalloproteinases (MMPs), early interventions and dressings. A modified Delphi methodology involving two iterations of email questionnaires followed by a face to face meeting was used to validate the statements, in order to arrive at a consensus for each. Clinical experts were selected; representing nurses, surgeons, podiatrists, academics, and policy experts.*

Results: *In the first round, 38/47 statements reached or exceeded the consensus threshold of 80% and none were rejected. According to the protocol, any statement not confirmed or rejected had to be modified using the comments from participants and resubmitted. In the second round, 5/9 remaining statements were confirmed and none rejected, leaving 4 to discuss at the meeting. All final statements were confirmed with at least 80% consensus.*

Conclusions: *This modified Delphi panel sought to gain clarity from clinical experts surrounding the use of dressings in the management of chronic wounds. A full consensus statement was developed to help clinicians and policy makers improve the management of patients with these conditions.*

Background

Diabetic Foot Ulcers (DFUs) and Venous Leg Ulcers (VLUs) are two of the most common lower limb wounds.¹ A growing global epidemic of chronic wounds not only leaves patients in pain and with a reduced quality of life, but also causes a significant financial burden to healthcare providers worldwide.^{2,3} In 2016, independent research funded by the National Health Service's (NHS) National Institute of Health Research (NIHR) stated that the prevalence of long lasting ulcers below the knee that take longer than six weeks to heal is seen in 15 out of every 10,000 people,⁴ which is an increase of threefold on a previous estimate. The impact of these wounds, including DFU and VLU is likely to continue to rise, with an aging population and increasing incidence of diabetes⁵ accelerating the growth.

The burden of these wounds is felt not only by patients, but also by carers, families, employers, and by the healthcare system. Should a DFU remain unhealed and eventually

require amputation, this is devastating for the patient and their subsequent decreased level of independence, will place a strain on the family or carers. The financial burden to the healthcare system is substantial; an estimate of the cost of chronic wounds to the NHS is between £2.3-3.1 billion (for the year 2005/6).⁶ Diabetes UK estimated that in 2014-15 around £1 billion (or approximately £1 in every £140 the NHS spends) is spent on foot ulcers or amputations each year.⁷ Prescribing costs are also rising, in 2004 £122 million was spent on wound dressings, and 8 years later in 2012, the prescribing costs for wound dressings had risen by 51% to £184 million.⁸ The rise in spending on chronic wounds can be partially attributed to the increasing numbers of people presenting with DFU and VLU. The increased demand has led to a need to create an efficient treatment pathway that will both improve the welfare of substantial numbers of patients and also reduce overall NHS spending. The Scottish Intercollegiate Guidelines Network (SIGN) guideline on the management of Venous Leg Ulcer states that "*Simple non-adherent dressings are recommended in the management of venous leg ulcers*".⁹ However, this guidance was issued in 2010, and the more recent National Institute for Health and Care Excellence (NICE) guidelines on the treatment of DFU, updated in January 2016, asks the research question: *What is the clinical effectiveness of different dressing types in treating diabetic foot problems?*¹⁰

To help improve outcomes, it is important to understand the expected healing process of a wound and being able to identify exactly when a wound deviates from this could reduce costs. Recently, there has been an increase in the understanding of wound physiology and how the micro-environment of a wound is important to achieving wound healing. It has been found that a key family of enzymes, MMPs, have a fundamental role in wound healing. As a result of this understanding, there are new treatment options that work to change the wound environment to promote and stimulate healing.

The uncertainty regarding the use of dressings in wound management and their place in the treatment pathway is clear; many recent Cochrane reviews have not been able to recommend a single type of dressing.¹¹ To address this, it was proposed to convene a panel of clinical experts to produce a consensus statement using a modified Delphi methodology. This is an anonymous, iterative process, where a group of multidisciplinary experts aimed to reach agreement in areas where there is a lack of explicit and clear guidance for clinical practice.¹² In addition to the role of MMPs, this study aimed to provide understanding on a range of topics including; the definition of chronicity in wounds, the burden of illness, clinical outcomes of reducing healing time and the impact of early interventions on clinical and economic outcomes.

Methods

A systematic literature review (SLR) was the first part of this study. A search strategy consistent with the PICO framework, focusing on Population, Intervention, Comparison and Outcomes, was formulated for the area of wound management. A full list of search terms is available in Table 1.

Table 1: Search terms used

Search terms	Item
(Wound* AND Chronic) OR (Ulcer AND (Pressure OR Diabetic Foot OR Venous Leg))	Population
Management OR Treatment OR Care	Intervention
Dressing*	Intervention
Resource AND (Use OR Utilisation) OR Cost	Outcome
Quality of Life OR Patient Outcomes OR Burden OR Impact	Outcome
Effectiveness OR Efficacy	Outcome

Online databases were then searched using these search terms for publications looking at clinical, economic and quality of life outcomes in patients with chronic wounds such as DFU, VLU and PU.

Table 2: List of databases used

Search tool	Count
Science Direct	2479
NICE Evidence search	805
Medline (PubMed)	78
CRD (University of York)	47
Cochrane	8
Total exported to EndNote:	3417

Of the 3417 articles retrieved, 827 were included at initial screening. The rest were duplicate of deemed irrelevant at this early stage. After this, a pre-defined set of inclusion and exclusion criteria were applied to the search results. These criteria are shown in Table 3. These criteria were applied by 2 reviewers from the Manchester Met project team.

A broad range of study types was included to allow a large evidence base for the statements that were to be generated.

Eventually, 145 full texts were used in order to inform the development of the statements. A schematic of the literature search is shown in Figure 1.

Table 3: Inclusion and Exclusion Criteria

Inclusion criteria	
Population	Diabetic Foot Ulcer, Venous Leg Ulcer, or a study of mixed wounds that included the aforementioned.
Interventions	Dressings
Outcomes	Wound healing, Wound Area Reduction, healing rate, Quality of Life Outcomes, Economic outcomes
Study design	<ul style="list-style-type: none"> o Randomised Controlled Trials o Patient Reported Outcomes o Observational studies o Epidemiology Studies o Modelling o Case Studies o Economic studies o Database Studies o Systematic/ Literature Reviews o Treatment pathway/guidelines
Language restrictions	English Language
Search dates	After 1987
Exclusion criteria	
Population	Paediatrics (<18), Acute wounds (including Burns, Trauma, Surgery)
Interventions	Surgical Novel non-surgical (including electrical stimulation, hyperbaric treatment, electrical stimulation) Infection control measures (including silver, iodine or honey) Debridement (including, surgical, maggot) Bioengineered skin substitutes Offloading
Outcomes	Not meeting inclusion criteria
Study design	In vitro studies, review or discussion articles
Language restrictions	Non-English language (if the abstract was available in English and enough data was available, this was included in the data extraction, otherwise these articles were excluded).
Search dates	Before 1987




Figure 1: PRISMA Search strategy flow diagram

In order to develop the statements that would be taken forward to the Delphi panel for review, a thematic analysis of the papers was undertaken. 145 texts were reviewed by the project lead, and 304 direct quotations from 131 of the were extracted in four set categories; epidemiology, clinical effectiveness, quality of life, and economics and cost. These quotations were reviewed and agreed as representative by the Manchester Met project team.

A lack of clarity presented itself in many ways, including; an inconclusive systematic literature review, a dressing being deemed as not having enough robust evidence, or as opposing results being published on the same subject.

An assessment of the quotations highlighted many sub categories, which were aggregated under the themes shown in Table 4, and used to develop the 47 statements to put forward to the Delphi panel for voting and further refinement.

Table 4: Themes

Theme	Number of statements
Definition of chronicity	3
Burden of illness	10
Reduce healing time	4
The role of matrix metalloproteinases	13

Early interventions lead to better outcomes	7
The use of dressings and treatments	10

In order to validate the statements, and assess the evidence using a group of experts and their combined wealth of clinical and academic expertise, a modified Delphi methodology was carried out. The Delphi method was developed by the RAND Corporation in the 1950's, and aims to arrive at an expert consensus using an iterative process. The method consists of a group of experts anonymously replying to a questionnaire; then receiving the group feedback, after which this process repeats itself.

The modified process that was used for this study included two rounds of anonymous email voting followed by a face-to-face meeting. The meeting was face to face with all participants and was a very structured round-table meeting with strict agenda. The threshold for consensus was set at 80%, and participants had the option of voting yes or no against the statements, thereby confirming or rejecting the statements respectively. Using previous Delphi methodology studies as a guide, 80% consensus was a relatively high threshold.

The participants were sent an excel sheet workbook that consisted of 6 sheets:

1. Cover sheet: For participants to record their name, affiliation and job title.
2. Introduction: An overview of the workbook and the process.
3. Instructions: An overview of the tasks needed to be completed by the participant.
4. Voting sheet: For the participants to record their responses.
5. References: Full listing of quotations, with bibliographic information and classification of evidence using a modified SIGN system.
6. Search methodology: An overview of the search strategy and results of the SLR.

The voting sheet allowed participants to click on hyperlinks to review the evidence base for each statement; each study was also given a level of evidence classification using a modified version of the SIGN Evidence classification shown in table 5. Participants were invited to review the evidence base for the statements, and were given the full bibliographic information and evidence classification for each source used.

Table 5: Modified SIGN Evidence Classification:

Level	Description
1	Guidelines, Meta-analyses, systematic reviews of RCTs, or RCTs

2	Economic Evaluations, Systematic reviews of case control or cohort studies. Case control or cohort studies
3	Non-analytic studies, e.g. case reports, case series, in vivo or in vitro studies
4	Expert opinion

The panel of 12 members were identified and approached for their experience treating and managing wounds such as DFU, VLU and PU. A range of different specialities were included on the panel; this is to reflect the multidisciplinary care pathway for patients with these chronic wounds. The final clinical experts included in the panel are listed in Table 6.

Any statement that fell in between 80% 'yes' and 80% 'no' was amended by the Manchester Met project team using the comments made by the participants and resubmitted to them in the following round.

This modified methodology was chosen for its iterative and impartial rigour that allowed each participant a fair chance to voice their opinions in the anonymous voting rounds, a step that is important in empowering panel members to voice their opinions amongst the multidisciplinary group.

This study was reviewed and approved by Manchester Metropolitan University Faculty Academic Ethics Committee, with number 1486. Panel members gave informed consent to participate both verbally and in writing.

Results

Twelve panel members were approached, however one panel member dropped out of the process and another was unable to complete the workbook in time for their comments to be included, yet joined the discussion and endorsed the consensus. A final ten participants completed the first workbook and the results of the first anonymous round of voting were as follows: 38 statements confirmed, 9 statements did not reach the 80% consensus threshold and 0 statements were rejected. 18 statements were agreed by 100% of the panel.

The same 10 participants completed the second workbook which consisted of the 9 statements that had been amended and resubmitted, 5 statements were agreed, and 4 did not reach the 80% consensus threshold, 0 were rejected.

At the meeting, the remaining 4 statements were amended and presented to the panel, where they gained consensus. Due to the large number of statements confirmed before the

final round; it was considered prudent to revisit comments on statements which had been confirmed with a level of 80-99%, in order to increase the level of agreement and ensure semantic clarity.

After the meeting, the statements were collected, ordered, and presented in the below consensus statement for dissemination. The statements themselves are identified with bold text, and underlined words or phrases are defined in Table 5 at the end of the statement.

Consensus statement

There is a need for consensus when the literature or guidance does not provide clarity. This lack of clarity can be identified by: contradictory information in the literature, a lack of robust evidence or systematic reviews that prove inconclusive. Recent reports and guidelines on wound management are not specific and do not make recommendations on treatment options. The Cochrane Review "Protease-modulating matrix treatments for healing venous leg ulcers" identifies the need for further research into these dressings.¹³

Contents of this consensus statement:

1. The role of matrix metalloproteinases (MMPs)
2. Quality of life for patients with DFU, VLU and PU
3. Time to healing and NHS burden
4. Early intervention and economic impact
5. Conclusions
6. Definitions
7. Panel Members

1. The role of MMPs in chronic wounds:

Wounds are deemed chronic when they do not follow a normal healing pattern and can be perpetuated by having an underlying aetiology.¹⁴⁻¹⁹ A normal healing pattern contains four phases of healing categorised according to the activity of their cellular components: haemostasis phase, inflammatory phase, proliferative phase, and maturation (or remodelling) phase. Wounds with underlying aetiologies include Diabetic Foot Ulcers (DFUs), Venous Leg Ulcers, (VLUs) and Pressure Ulcers (PUs).

Matrix metalloproteinases (MMPs) are a part of healthy healing, expressed at the inflammatory phase of early wound healing.²⁰⁻²⁴ MMPs are enzymes that are responsible for degradation of the extracellular matrix and also play a pivotal role in regulation of cell proliferation, migration, differentiation, and death. **When a wound moves to the proliferative phase of healing, the level of MMPs fall.**²⁵ If the wound does not advance to the proliferative phase of healing in an expected time period, it can be considered chronic. **These chronic wounds have been shown to have up to 30 times the level of MMPs than an acute wound.**²⁶⁻³¹

Wounds such as DFU, VLU and PU are shown to have raised levels of MMPs from first presentation to a wound care

specialist.³²⁻³⁶ With raised levels of MMPs, the wound is stuck in the inflammation phase, leading to the destruction of new tissues,³⁷⁻⁴⁰ thus preventing progression to the next stage of healing.⁴¹⁻⁴³

Persistently elevated levels of MMP are predictive of non-healing⁴⁴⁻⁴⁸ and specifically, of the 24 known MMPs, MMP-9 has been shown to be detrimental to healing, killing growth factors.^{49 - 53} Interventions that modulate the wound environment may enhance healing^{54 - 60} because evidence suggests removing excess MMPs from wounds improves healing.⁶¹⁻⁶⁵ A specific MMP-9 inhibitor is potentially more effective in stimulating healing⁶⁶⁻⁶⁸ than standard care alone. In addition to modulating the wound environment, the ideal dressing should be cost-effective, acceptable to the patient and also be effective on older and larger wounds.⁶⁹⁻⁸⁴

The lipido-colloid nano-oligosaccharide factor (TLC-NOSF) technology inhibits MMPs and accelerates healing,⁸⁵⁻⁸⁸ it has been shown as superior to basic foam dressings in reducing healing time^{89, 90} and as superior to oxidized regenerated cellulose and collagen, especially in non-responsive, older wounds.⁹¹ Further to this, TLC- NOSF has been shown to reduce levels of MMP-9 in vitro.^{92, 93}

2. Quality of life for patients with DFU, VLU and PU

Wounds such as DFU, VLU and PU are associated with increased morbidity and mortality.⁹⁴⁻⁹⁹ In addition to this increased risk of death and high likelihood of comorbidities, patients with these conditions suffer significantly reduced health related quality of life across dimensions such as pain, physical limitation, social isolation, and anxiety/depression.¹⁰⁰⁻¹⁰⁴ The psychological impact of these wounds can be severe, with patients reporting a loss of self, poor self-image, feelings of being a burden and hopelessness for the future.¹⁰⁵⁻¹⁰⁹ These wounds can take a long time to heal and have a high likelihood of recurrence, which again detracts from quality of life.¹¹⁰⁻¹¹⁶ Clinician focus tends to be on the treatment of the wound, which fails to account for the large psychological and social burden experienced by some patients.¹¹⁷⁻¹²⁰

The pain caused by chronic wounds impacts quality of life.¹²¹⁻¹²⁴ Dressing changes can be a cause of pain: products and techniques to minimise this are recommended.¹²⁵⁻¹²⁹ Dressing changes and local management of the wound site is considered easy in most cases with the TLC-NOSF dressing,^{130, 131} which has also been shown to significantly reduce pain/discomfort and anxiety/depression for a patient.^{132, 133}

In addition to the health related quality of life burden, the patient also faces financial costs such as time away from work, early retirement, medications, dressings, and transport costs.¹³⁴⁻¹³⁷ Chronic wounds are a burden to both the patient and to the carer¹³⁸ and this cost is often excluded or underestimated in cost-effectiveness models.¹³⁹⁻¹⁴⁴

3. Time to healing and NHS burden

As well as a quality of life burden to patients, DFU, VLU and PU are a significant workload burden for healthcare providers.¹⁴⁵⁻¹⁵⁵ Home visits are a key driver of the cost to treat chronic wounds.^{156 - 160} Advanced dressings require fewer changes and therefore fewer visits are more likely to reduce costs, especially when the dressing also reduces healing time.¹⁶¹⁻¹⁶⁵ Protease inhibitors have been shown to be a cost-effective option.^{166 - 169} Management plans associated with shorter treatment periods and fewer adverse events are more cost-effective.¹⁷⁰⁻¹⁷⁵ Ulcers can be slow to heal, with wound size and duration affecting healing.¹⁷⁶⁻¹⁸⁰ The initial wound area reduction at 4 weeks is predictive of healing by 24 weeks.¹⁸¹⁻¹⁸³

4. Early intervention and economic impact

Early diagnosis and treatment of a DFU, VLU or PU can improve quality of life for a patient.^{184, 185} This early investment in treatment provides a reduction in long-term costs; prolonged futile treatment is more costly.¹⁸⁶⁻¹⁸⁹ There is a need for a long-term view from decision makers, for example, the purchase price of a dressing is not indicative of cost-effectiveness.¹⁹⁰

Some ulcers are more expensive to manage, these include: chronic wounds, recurrent wounds, and older wounds.¹⁹¹⁻¹⁹⁵ Older wounds are harder and more expensive to heal so early intervention will reduce the healing time and cost.¹⁹⁶⁻²⁰⁰ VLU is more prevalent in older populations who may benefit from less invasive treatment options.²⁰¹⁻²⁰⁴ An adjunctive therapy such as a dressing that modulates the microenvironment can promote faster healing in complicated wounds.^{205 - 211} An adjunctive therapy to standard wound care should be considered in cases where you anticipate wound healing may be compromised.²¹²⁻²¹⁷

5. Conclusions

This consensus process seeks to provide clarity for the management of chronic wounds. We have agreed that:

- Chronic wounds including DFU, VLU and PU significantly impair a patient's health and quality of life and this needs to be taken into consideration in patient care with the aim of reducing healing time.
- Inhibiting MMPs plays an important role in wound healing and raised levels of these enzymes have been shown to be present in DFU, VLU and PU.
- Early interventions are a more cost-effective option, both in terms of health and quality of life improvement for a patient and in financial savings to the healthcare system

6. Definitions

Table 5: Consensus statement definitions

Term (in order of appearance)	Definition
Normal healing pattern	A normal healing pattern contains four phases of healing categorised

	according to the activity of their cellular components. The phases are haemostasis phase, inflammatory phase, proliferative phase, and maturation (or remodelling) phase. Normal healing will move through these phases naturally at a predictable rate.
Aetiology	The cause or origin of a disease or disorder as determined by medical diagnosis. (The American Heritage® Medical Dictionary Copyright © 2007, 2004 by Houghton Mifflin Company)
Matrix metalloproteinases (MMPs)	By regulating the integrity and composition of the extracellular matrix, these enzymes play a pivotal role in the control of signals elicited by matrix molecules that regulate cell proliferation, differentiation, and death. (Farlex Partner Medical Dictionary © Farlex 2012)
Acute wound	An acute wound is an injury to the skin that occurs suddenly rather than over time. It heals at a predictable and expected rate according to the normal wound healing process: (http://www.woundcarecenters.org/article/wound-types/acute-wounds)
Basic foam dressings	A foam dressing with no active agents.
Morbidity	A diseased condition or state.
Mortality	Likelihood of death, or death rate.
Significantly	Having reached statistical significance.
Carer	An unpaid carer; a relative, friend or neighbour.
Healthcare Providers	Any individual, institution, or agency that provides health services.
Advanced dressings	Dressings that regulate wound healing by simple physicochemical means, typically by controlling moisture levels. (NICE Evidence summary [ESMPB2] March 2016)

Adjunctive therapy	Another treatment used together with the primary treatment. Its purpose is to assist the primary treatment. (PubMed Health Glossary: Source: NIH - National Cancer Institute)
Standard wound care	Standard care used to promote wound healing, which can be achieved through off-loading in DFU, compression in VLU and/ or repositioning in PU

7. Panel Members:

The panel was made up of a multidisciplinary group and was supported by a group of technical experts to advise on the methodology. The panel members with voting rights are

Name	Title	Place of work
Professor Nancy Devlin	Panel Chair Person	Office of Health Economics
April Betts	Project Manager	Manchester Metropolitan University
Professor Isaac Odeyemi	Visiting Professor of Health Technology Assessment and Health Policy	Manchester Metropolitan University
Professor Francis Fatoye	Professor of Health Economics and Outcomes	Manchester Metropolitan University
Dr Gillian Yeowell	MSc Advanced Physiotherapy programme leader	Manchester Metropolitan University
Richard Shorney	Meeting Facilitator	Real Healthcare Solutions

listed in Table 7, and the technical experts in Table 6.

Table 6: Technical experts

Table 7: Clinical experts

Name	Title	Place of work
Dr Leanne Atkin	Vascular Nurse Specialist	Mid Yorks NHS
Dr Caroline Dowsett	Nurse Consultant Tissue Viability	East London NHS Foundation Trust, London
Sarah Gardner	Clinical Lead, Tissue Viability	Oxford Health NHS Foundation Trust
Dr Julie Green	Senior Lecturer in Nursing, Director of Postgraduate Programmes	Keele University, School of Nursing and Midwifery
Dr Chris Manu	Consultant Diabetologist and Clinical Researcher in Diabetic Foot	Kings College Hospital, London
Tracey McKenzie	Head of Tissue Viability Services	Torbay and Southern Devon NHS Foundation Trust
Helena Meally	Hospital Podiatrist	Leeds Teaching Hospitals NHS Trust
Louise Mitchell	Clinical Lead Podiatrist	Birmingham Community HealthCare
Julie Mullings	Lead Tissue Viability Nurse	University Hospital South Manchester
David Russell	Consultant Vascular Surgeon and Honorary Clinical Associate Professor	Leeds Teaching Hospitals NHS Trust
Andrew Sharpe	Advanced Podiatrist and Lecturer Practitioner	West Lancashire Community Service, Virgin Care Ltd and University of Huddersfield

Discussion

A lack of clear evidence supporting a single treatment strategy, or mandated clinical guidance can have a detrimental impact on emerging technologies. Reviews relying solely on the literature, and not clinical opinion, often present uncertainty, such as the review carried out by Cochrane in 2016, *Protease-modulating matrix treatments for healing venous leg ulcers*. This systematic literature review declared the evidence not conclusive and the certainty judged as low. This however, does not mean that the available evidence is of no use to clinicians, as demonstrated by the consensus panel. New studies, published and ongoing, have been designed to assess the efficacy and tolerability of specific protease modulating matrix treatments.

New evidence available has shown promising results. A manuscript exploring the real world usage of a protease modulating dressing has now been published (Munter 2017). The quality of life endpoints associated with this dressing have been explored in Meaume 2017, and ClinicalTrials.gov shows that a trial titled “Assessment of the Efficacy and Safety of a New Wound Dressing in the Local Treatment of Diabetic Foot Ulcers” is due to report. The Cochrane Collaboration has also just registered a protocol looking into the efficacy of using protease levels to predict healing outcomes in VLU patients.

The objective of this project was to provide clear guidance for clinical practice on a range of topics where there is a lack of clarity in the literature. The rigorous process that was followed has generated a consensus statement, agreed by a multidisciplinary panel.

The certainty of evidence for wound care dressings is low, as evidenced by a series of inconclusive Cochrane reviews (hydrocolloid, alginate, hydrogel, foam, protease-modulating matrix treatments) that found low levels of evidence and high risk of bias.^{218, 219, 220, 221, 222}. Cochrane risk of bias tools judge using blinding criteria that are difficult to meet in any wound trial; due to practical issues with packaging, nurse involvement etc. In light of this; this consensus panel allowed the participants to judge the validity of the evidence in the context of their own clinical expertise.

The consensus statement agrees that chronic wounds have a significant impact on a patient, regarding both their health and quality of life. In order to mitigate this for the patient and the healthcare provider, early intervention is key to successful treatment. The role of MMPs in wound healing is important, and in wounds with raised MMP levels, such as DFU, VLU and PU, a MMP inhibitor can expedite wound healing.

The modified Delphi process has many benefits, such as the anonymity enjoyed by the participants in the first two rounds. This helped to ensure a wide range of expert opinions were collected with the return of the workbooks. The face to face meeting after this was to allow the panel to come together as a group and review the study output. It is possible that the face-to-face element may weaken the strength of the methodology, however an individual’s earlier comments remained anonymous and the Chairperson present ensured that the review of comments was without derision. The process is also iterative, and the systematic review of the literature carried out prior to the Delphi process ensured it is supported by evidence, repeatable and transparent.

The strength of the process became apparent after the first round, with 80% of the statements reaching the consensus threshold. This could be attributed to a number of factors,

including the body of evidence presented in the workbook, the anonymity provided preventing individuals unduly influencing others, or perhaps the lack of clarity in the literature is not reflected in clinical practice. At the end of the process, all of the original statements had been confirmed, with modifications. This can once again be attributed to the fact that the methodology allowed the participants' comments to inform amendment of the statements when resubmitted. This flexibility in the approach allowed for more participation and elicited more expertise from the panel members. However, a limitation of the methodology relates to the binary yes/no structure of the questions. This process could be further improved by amending the voting to a scale which would allow for more ranking of the statements.

The Delphi process differs from a traditional expert panel or advisory board, the participants of an advisory board are likely to meet once, for a few hours and have a semi-structured discussion, often based on some pre-work. The modified Delphi methodology used in this study allowed direct access to the evidence base for the statements, and

participants were granted anonymity when sharing their opinions. Due to the high consensus levels, it is possible that the statements generated from the SLR were more based on fact and evidence than they were opinion, however the Delphi method proves to be a good tool for validating the output of a literature review with a multidisciplinary panel. Perhaps in future studies, the Delphi methodology can be used in more subjective areas such as guidelines and treatment pathways.

It is hoped that the dissemination of the consensus statement will lead to an improvement in patient care, and a reduction in costs for the healthcare system when tackling the issue of ulcers of varying aetiologies. The increasing prevalence of these wounds, especially DFU, calls for more research into wound management, the mode of action of MMP inhibitors, and how to maximise efficiencies in the healthcare system whilst maintaining a gold standard of care for patients.

¹ Guest J., Ayoub N., McIlwraith T., et al. Health economic burden that wounds impose on the National Health Service in the UK. *BMJ Open*. 2015; 5: 12. Article: E009283.

² Posnett J., Gottrup F., Lundgren H., Saal G.. The resource impact of wounds on healthcare providers in Europe. *Journal of Wound Care*. 2009; 18:4, 154-161.

³ Fife C., Carter, M. Walker, D., Thomson, B. Wound Care Outcomes and Associated Cost Among Patients Treated in US Outpatient Wound Centers: Data From the US Wound Registry. *Wounds*. 2012; 24:1, 10-17.

⁴ Cullum N, Buckley H, Dumville J, et al. Wounds research for patient benefit: a 5-year programme of research. *Programme Grants for Applied Research*. 2016; 4:13, DOI 10.3310/pgfar04130.

⁵ Diabetes UK. State of the Nation 2016.

<https://www.diabetes.org.uk/professionals/position-statements-reports/statistics/state-of-the-nation-2016-time-to-take-control-of-diabetes>. (Accessed August 2017)

⁶ Posnett, J., Franks, P. The burden of chronic wounds in the UK. *Nursing Times*. 2008; 104: 3, 44-45.

⁷ Kerr M. <https://www.improvingfootcareforpeoplewithdiabetesand savingmoney: an economic study in England. 2017. https://diabetes-resources-production.s3-eu-west-1.amazonaws.com/diabetes-storage/migration/pdf/Improving%2520footcare%2520economic%2520study%2520%28January%25202017%29.pdf> (Accessed August 2017)

⁸ NIHR Signal. Long lasting ulcers below the knee are more common than previously thought. 2016. <https://discover.dc.nihr.ac.uk/portal/article/4000656/long-lasting-ulcers-below-the-knee-are-more-common-than-previously-thought> (Accessed August 2017)

⁹ Scottish Intercollegiate Guidelines Network (SIGN). 120: Management of chronic venous leg ulcers: A national clinical guideline. 2010.

<http://www.sign.ac.uk/assets/sign120.pdf> (Accessed May 2017)

¹⁰ National Institute for Health and Care Excellence (NICE). NG19: Diabetic foot problems: prevention and management. 2015. [nice.org.uk/guidance/ng19](https://www.nice.org.uk/guidance/ng19) (Accessed August 2017)

¹¹ Dumville, J., Deshpande, S., O'Meara, S. and Speak, K. Foam dressings for healing diabetic foot ulcers. *Cochrane Database of Systematic Reviews*. 2013; 6: 6, Article: CD009111.

¹² Dumville, J., O'Meara, S., Deshpande, S. and Speak, K. Alginate dressings for healing diabetic foot ulcers. *Cochrane Database of Systematic Reviews*. 2013; 25:6, Article: CD009110.

¹³ Westby, M., Norman, G., Dumville, J., Stubbs, N. and Cullum, N. Protease-modulating matrix treatments for healing venous leg ulcers. *Cochrane Database of Systematic Reviews*. 2016; 12. Article: CD011918.

¹⁴ Helmer-Hirschberg, O. Analysis of the Future: The Delphi Method. RAND Corporation, 1967. <https://www.rand.org/pubs/papers/P3558.html>. (Accessed May 2017).

¹⁵ Westby, M., Norman, G., Dumville, J., Stubbs, N. and Cullum, N. Protease-modulating matrix treatments for healing venous leg ulcers. *Cochrane Database of Systematic Reviews*. 2016; 12. Article: CD011918.

Statement 1

¹⁶ Levine, S., Myerson, M. Diabetic foot ulceration. *The Foot*. 1995; 5: 4, 157-164

¹⁷ Flegg, J., Kasza, J., Darby, I., Weller, C. Healing of venous ulcers using compression therapy: Predictions of a mathematical model. *Journal of Theoretical Biology*. 2015; 379, 1-9.

¹⁸ Schuren, J., Becker, A., Sibbald, R. A liquid film-forming acrylate for peri-wound protection: a systematic review and meta-analysis (3M Cavilon no-sting barrier film). *International Wound Journal*. 2005; 2:3, 230-8.

¹⁹ Kelechi, T., Johnson, J., Yates, S. Chronic venous disease and venous leg ulcers: An evidence-based update. *Journal of Vascular Nursing*. 2015; 33:2, 36-46.

²⁰ Frykberg, R., Banks, J. Challenges in the Treatment of Chronic Wounds, *Advances in Wound Care*. 2015; 4:9, 560-582

²¹ Demidova-Rice, T., Hamblin, M., Herman, I. Acute and Impaired Wound Healing: Pathophysiology and Current Methods for Drug Delivery, Part 1: Normal and Chronic Wounds: Biology, Causes, and Approaches to Care, *Advances in Skin and Wound Care*. 2012; 25:7, 304-314.

Statement 18

²² Crovetti, G., Martinelli, G., Issi, M., et al. Platelet gel for healing cutaneous chronic wounds. *Transfusion and Apheresis Science*. 2004; 30:2, 145-151.

²³ Olczyk, P., Mencner, L., Komosinska-Vasves, K. The Role of the Extracellular Matrix Components in Cutaneous Wound Healing, *BioMed Research International*. 2014, Article: 747584.

²⁴ Broughton G., Janis, J., Attinger, C. Wound Healing: An Overview. *Plastic and Reconstructive Surgery*. 2006; 117:7, 1-32.

Statement 19

²⁵ Moffatta, C., Stantonon, J., Murray, S., et al. A randomised trial to compare the performance of Oxymyze® and Iodozyme® with standard care in the treatment of patients with venous and mixed venous/arterial ulceration. *Wound Medicine*. 2014, 6, 1-10.

²⁶ Heublein, H., Bader, A., Giri, S. Preclinical and clinical evidence for stem cell therapies as treatment for diabetic wounds. *Drug Discovery Today*. 2015; 20:6, 645-780.

Statement 12

²⁷ Trengove N., Stacey M., MacAuley S et al. Analysis of the acute and chronic wound environments: the role of proteases and their inhibitors. *Wound Repair and Regeneration*. 1999; 7:6, 442-452.

Statement 2

²⁸ Westby, M., Norman, G., Dumville, J., Stubbs, N. and Cullum, N. Protease-modulating matrix treatments for healing venous leg ulcers. *Cochrane Database of Systematic Reviews*. 2016; 12. Article: CD011918.

²⁹ Eming S., Krieg T., Davidson J. Inflammation in wound repair: molecular and cellular mechanisms. *Journal of Investigative Dermatology*. 2007; 127:3, 514-525.

³⁰ Trengove N., Stacey M., MacAuley S et al. Analysis of the acute and chronic wound environments: the role of proteases and their inhibitors. *Wound Repair and Regeneration*. 1999; 7:6, 442-452.

³¹ Zelen, C., Gould, L., Serena, T., et al. A prospective, randomised, controlled, multi-centre comparative effectiveness study of healing using dehydrated human amnion/chorion membrane allograft, bioengineered skin substitute or standard of care for treatment of chronic lower extremity diabetic ulcers. *International Wound Journal*. 2015; 12, 724-732.

³² Ahmad, J. The diabetic foot. *Diabetes & Metabolic Syndrome*. 2016; 10: 1, 48-60.

³³ Health Quality Ontario. Management of chronic pressure ulcers: an evidence-based analysis. *Ontario Health Technology Assessment Series*. 2009; 9: 3, 1-203.

Statement 3

³⁴ Heublein, H., Bader, A., Giri, S. Preclinical and clinical evidence for stem cell therapies as treatment for diabetic wounds. *Drug Discovery Today*. 2015; 20:6, 645-780.

³³ Lazaro J., Izzo, V., Meaume, S., et al. Elevated levels of matrix metalloproteinases and chronic wound healing: an updated review of clinical evidence *Journal of wound care*. 2016; 25: 5, 277-287.

³⁴ Menghini, R., Uccioli, L., Vainieri, E., et al. Expression of tissue inhibitor of metalloproteinase 3 is reduced in ischemic but not neuropathic ulcers from patients with type 2 diabetes mellitus. *Acta Diabetologica*. 2013; 50: 6, 907-910.

³⁵ Rayment, E., Upton, Z., Shooter, G. Increased matrix metalloproteinase-9 (MMP-9) activity observed in chronic wound fluid is related to the clinical severity of the ulcer. *British Journal of Dermatology*. 2008; 158: 5, 951-61.

³⁶ Beidler, S, Douillet, C, Berndt, D., et al. Multiplexed analysis of matrix metalloproteinases in leg ulcer tissue of patients with chronic venous insufficiency before and after compression therapy. *Wound Repair and Regeneration*. 2008; 16: 5, 642-8.

Statement 22

³⁷ Ravari H., Hamidi-Almadari, D., Salimifar, M., et al. Treatment of non-healing wounds with autologous bone marrow cells, platelets, fibrin glue and collagen matrix. *Cytotherapy*. 2011; 13:6, 701-11.

³⁸ Wysocki A., Staiano-Coico L., Grinnell F. Wound fluid from chronic leg ulcers contains elevated levels of metalloproteinases MMP-2 and MMP-9. *Journal of Investigative Dermatology*, 1993; 101:1, 64-8.

³⁹ Schmutz J., Meaume S., Fays S. et al. Evaluation of the nano-oligosaccharide factor lipido-colloid matrix in the local management of venous leg ulcers: results of a randomised, controlled trial. *International Wound Journal*. 2008; 5: 172-182.

⁴⁰ Snyder, R. Treatment of nonhealing ulcers with allografts. *Clinics in Dermatology*. 2005; 24:4, 388-95.

Statement 21

⁴¹ PrescQIPP. B115. Wound care: Protease-modulating matrix dressings. 2016. <https://www.prescqipp.info/component/jdownloads/send/235-wound-care-protecting-modulating-matrix-dressings/2385-b115-wound-care-protecting-modulating-matrix-dressings-briefing> (Accessed May 2017)

⁴² Ravari H., Hamidi-Almadari, D., Salimifar, M., et al. Treatment of non-healing wounds with autologous bone marrow cells, platelets, fibrin glue and collagen matrix. *Cytotherapy*. 2011; 13:6, 701-11.

⁴³ Heublein, H., Bader, A., Giri, S. Preclinical and clinical evidence for stem cell therapies as treatment for diabetic wounds. *Drug Discovery Today*. 2015; 20:6, 645-780.

Statement 23

⁴⁴ Westby, M., Norman, G., Dumville, J., Stubbs, N. and Cullum, N. Protease-modulating matrix treatments for healing venous leg ulcers. *Cochrane Database of Systematic Reviews*. 2016; 12. Article: CD011918.

⁴⁵ Wysocki A., Staiano-Coico L., Grinnell F. Wound fluid from chronic leg ulcers contains elevated levels of metalloproteinases MMP-2 and MMP-9. *Journal of Investigative Dermatology*, 1993; 101:1, 64-8.

⁴⁶ National Institute for Health and Care Excellence. MIB82. Woundchek Protease Status for assessing elevated protease status in chronic wounds. 2016. <https://www.nice.org.uk/advice/mib83>. (Accessed May 2017)

⁴⁷ Snyder, R. Treatment of nonhealing ulcers with allografts. *Clinics in Dermatology*. 2005; 24:4, 388-95.

⁴⁸ Schuren, J., Becker, A., Sibbald, R. A liquid film-forming acrylate for peri-wound protection: a systematic review and meta-analysis (3M Cavilon no-sting barrier film). *International Wound Journal*. 2005; 2:3, 230-8.

Statement 27

⁴⁹ Trengove N., Stacey M., MacAuley S et al. Analysis of the acute and chronic wound environments: the role of proteases and their inhibitors. *Wound Repair and Regeneration*. 1999; 7:6, 442-452.

⁵⁰ Moffatta, C., Stantonb, J., Murray, S., et al. A randomised trial to compare the performance of Oxzyme® and Iodozyme® with standard care in the treatment of patients with venous and mixed venous/arterial ulceration. *Wound Medicine*. 2014, 6, 1-10.

⁵¹ Campbell, C., Parish, L. The decubitus ulcer: Facts and controversies. *Clinics in Dermatology*. 2010; 28: 5, 527-32.

⁵² Heublein, H., Bader, A., Giri, S. Preclinical and clinical evidence for stem cell therapies as treatment for diabetic wounds. *Drug Discovery Today*. 2015; 20:6, 645-780.

⁵³ Kelechi, T., Johnson, J., Yates, S. Chronic venous disease and venous leg ulcers: An evidence-based update. *Journal of Vascular Nursing*. 2015; 33: 2, 36-46.

Statement 24

⁵⁴ Meaume, S., Truchetet F., Cambazard F., et al. A randomized, controlled, double-blind prospective trial with a Lipido-Colloid Technology-Nano-OligoSaccharide Factor wound dressing in the local management of venous leg ulcers. *Wound Repair and Regeneration*. 2012; 20: 4, 500-11.

⁵⁵ Humbert, P., Favier, B., Veran, Y., et al. On behalf of the CLEANSITE study group. Protease-modulating polyacrylate-based hydrogel stimulates wound bed preparation in venous leg ulcers – a randomized controlled trial. *Journal of the European Academy of Dermatology and Venereology*. 2014; 28: 12, 1742-50.

⁵⁶ Trengove N., Stacey M., MacAuley S et al. Analysis of the acute and chronic wound environments: the role of proteases and their inhibitors. *Wound Repair and Regeneration*. 1999; 7:6, 442-452.

⁵⁷ Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640.

⁵⁸ Heublein, H., Bader, A., Giri, S. Preclinical and clinical evidence for stem cell therapies as treatment for diabetic wounds. *Drug Discovery Today*. 2015; 20:6, 645-780.

⁵⁹ Moffatta, C., Stantonb, J., Murray, S., et al. A randomised trial to compare the performance of Oxzyme® and Iodozyme® with standard care in the treatment of

patients with venous and mixed venous/arterial ulceration. *Wound Medicine*. 2014, 6, 1-10.

⁶⁰ Schuren, J., Becker, A., Sibbald, R. A liquid film-forming acrylate for peri-wound protection: a systematic review and meta-analysis (3M Cavilon no-sting barrier film). *International Wound Journal*. 2005; 2:3, 230-8.

Statement 25

⁶¹ Grier J., Hunter C., Oboh L. Top tips QIPP messages for prescribing dressings. East & South East England Specialist Pharmacy Services. 2013. <https://www.sps.nhs.uk/wp-content/uploads/2015/09/Top20Tips20QIPP20messages20for20prescribing20dressing.s.pdf>. (Accessed May 2017)

⁶² PrescQIPP. B115. Wound care: Protease-modulating matrix dressings. 2016. <https://www.prescqipp.info/component/jdownloads/send/235-wound-care-protecting-modulating-matrix-dressings/2385-b115-wound-care-protecting-modulating-matrix-dressings-briefing> (Accessed May 2017)

⁶³ Meaume, S., Truchetet F., Cambazard F., et al. A randomized, controlled, double-blind prospective trial with a Lipido-Colloid Technology-Nano-OligoSaccharide Factor wound dressing in the local management of venous leg ulcers. *Wound Repair and Regeneration*. 2012; 20: 4, 500-11.

⁶⁴ Humbert, P., Favier, B., Veran, Y., et al. On behalf of the CLEANSITE study group. Protease-modulating polyacrylate-based hydrogel stimulates wound bed preparation in venous leg ulcers – a randomized controlled trial. *Journal of the European Academy of Dermatology and Venereology*. 2014; 28: 12, 1742-50.

⁶⁵ Westby, M., Norman, G., Dumville, J., Stubbs, N. and Cullum, N. Protease-modulating matrix treatments for healing venous leg ulcers. *Cochrane Database of Systematic Reviews*. 2016; 12. Article: CD011918.

Statement 28

⁶⁶ Rayment, E., Upton, Z., Shooter, G. Increased matrix metalloproteinase-9 (MMP-9) activity observed in chronic wound fluid is related to the clinical severity of the ulcer. *British Journal of Dermatology*. 2008; 158: 5, 951-61.

⁶⁷ Campbell, C., Parish, L. The decubitus ulcer: Facts and controversies. *Clinics in Dermatology*. 2010; 28: 5, 527-32.

⁶⁸ Heublein, H., Bader, A., Giri, S. Preclinical and clinical evidence for stem cell therapies as treatment for diabetic wounds. *Drug Discovery Today*. 2015; 20:6, 645-780.

Statement 41

⁶⁹ CADTH Rapid Response Reports. Dressing Materials for the Treatment of Pressure Ulcers in Patients in Long-Term Care Facilities: A Review of the Comparative Clinical Effectiveness and Guidelines. Canadian Agency for Drugs and Technologies in Health. 2013.

https://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0064797/pdf/PubMedHealth_PMH0064797.pdf. (Accessed May 2017)

⁷⁰ National Institute for Health and Care Excellence (NICE). CG179. Pressure ulcers: prevention and management. 2014. <https://www.nice.org.uk/guidance/cg179> (Accessed May 2017)

⁷¹ National Institute for Health and Care Excellence (NICE). NG19: Diabetic foot problems: prevention and management. 2015. [nice.org.uk/guidance/ng19](https://www.nice.org.uk/guidance/ng19) (Accessed August 2017)

⁷² O'Donnell Jr, T., Balk, E. The need for an Intersociety Consensus Guideline for venous ulcer. *Journal of Vascular Surgery*. 2011; 54: 6, 83-90.

⁷³ National Health and Medical Research Council (NHMRC). Australian and New Zealand clinical practice guideline for prevention and management of venous leg ulcers. 2011.

https://www.nhmrc.gov.au/files/nhmrc/publications/attachments/ext003_venous_leg_ulcers_aust_nz_0.pdf (Accessed May 2017)

⁷⁴ Widener, J. Venous leg ulcers: Summary of new clinical practice guidelines published August 2014 in the *Journal of Vascular Surgery*. *Journal of Vascular Nursing*. 2015; 33: 2, 60-7.

⁷⁵ Ouahes, N., Phillips, T. Leg ulcers. *Current Problems in Dermatology*. 1995; 7: 4, 113-42.

⁷⁶ Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640.

⁷⁷ Kelechi, T., Johnson, J., Yates, S. Chronic venous disease and venous leg ulcers: An evidence-based update. *Journal of Vascular Nursing*. 2015; 33: 2, 36-46.

⁷⁸ Kim, P., Steinberg, J. Wound Care: Biofilm and Its Impact on the Latest Treatment Modalities for Ulcerations of the Diabetic Foot. *Seminars in Vascular Surgery*. 2012; 25: 2, 70-4.

⁷⁹ Reagan, M., Teasell, R., Wolfe, D., et al. A systematic review of therapeutic interventions for pressure ulcers after spinal cord injury. *Archives of Physical Medicine and Rehabilitation*. 2009; 90: 2, 213-31.

⁸⁰ Wu L., Norman G., Dumville J. Dressings for treating foot ulcers in people with diabetes: an overview of systematic reviews. *Cochrane Database of Systematic Reviews*. 2015; 14: 7, Article: CD010471.

⁸¹ Health Quality Ontario. Management of chronic pressure ulcers: an evidence-based analysis. *Ontario Health Technology Assessment Series*. 2009; 9: 3, 1-203.

⁸² O'Donnell Jr, T., Lau, J. A systematic review of randomized controlled trials of wound dressings for chronic venous ulcer. *Journal of Vascular Surgery*. 2006; 44: 5, 1118-25.

⁸³ PrescQIPP. B115. Wound care: Protease-modulating matrix dressings. 2016. <https://www.prescqipp.info/component/jdownloads/send/235-wound-care-protecting-modulating-matrix-dressings/2385-b115-wound-care-protecting-modulating-matrix-dressings-briefing> (Accessed May 2017)

⁸⁴ Phillips T., Stanton, B., Provan, A., Lew, R. A study of the impact of leg ulcers on quality of life: Financial, social, and psychological implications. *Journal of the American Academy of Dermatology*. 1994; 31: 1, 49-53.

Statement 26

⁸⁵ Augustin, M., Herberger K., Kroeger K et al. Cost-effectiveness of treating vascular leg ulcers with UrgoStart and UrgoCell Contact. *International Wound Journal*. 2016; 13: 1, 82-7.

⁸⁶ Schmutz J., Meaume S., Fays S. et al. Evaluation of the nano-oligosaccharide factor lipido-colloid matrix in the local management of venous leg ulcers: results of a randomised, controlled trial. *International Wound Journal*. 2008; 5: 172–182.

⁸⁸ Meaume, S., Truchetet F., Cambazard F., et al. A randomized, controlled, double-blind prospective trial with a Lipido-Colloid Technology-Nano-OligoSaccharide Factor wound dressing in the local management of venous leg ulcers. *Wound Repair and Regeneration*. 2012; 20: 4, 500-11.

Statement 30

⁸⁹ Meaume, S., Truchetet F., Cambazard F., et al. A randomized, controlled, double-blind prospective trial with a Lipido-Colloid Technology-Nano-OligoSaccharide Factor wound dressing in the local management of venous leg ulcers. *Wound Repair and Regeneration*. 2012; 20: 4, 500-11.

⁹⁰ Augustin, M., Herberger K., Kroeger K et al. Cost-effectiveness of treating vascular leg ulcers with UrgoStart and UrgoCell Contact. *International Wound Journal*. 2016; 13: 1, 82-7.

Statement 42

⁹¹ Schmutz J., Meaume S., Fays S. et al. Evaluation of the nano-oligosaccharide factor lipido-colloid matrix in the local management of venous leg ulcers: results of a randomised, controlled trial. *International Wound Journal*. 2008; 5: 172–182.

Statement 29

⁹² Bernard, F., Juchaux, F., Laurensou, C., et al. In vitro effect of a new metalloproteinase inhibitor. Poster presented at EWMA, Lisbon, May 2008.

⁹³ Coulomb, B., Couty, L., Fournier, B. et al. Evaluation of the matrix impregnated with NOSF (Nano Oligo Saccharide Factor) in an in vitro dermal reconstruction model. *JPC* 2008; 63: 8, 54-57.

Statement 31

⁹⁴ Zelen, C., Gould, L., Serena, T., et al. A prospective, randomised, controlled, multi-centre comparative effectiveness study of healing using dehydrated human amnion/chorion membrane allograft, bioengineered skin substitute or standard of care for treatment of chronic lower extremity diabetic ulcers. *International Wound Journal*. 2015; 12, 724-732.

⁹⁵ Diabetes UK. Fast track for a foot attack- reducing amputations. 2013. <https://diabetes-resources-production.s3-eu-west-1.amazonaws.com/diabetes-storage/migration/pdf/putting-feet-first-foot-attack-report022013.pdf> (Accessed May 2017)

⁹⁶ Health Quality Ontario. Management of chronic pressure ulcers: an evidence-based analysis. *Ontario Health Technology Assessment Series*. 2009; 9: 3, 1-203.

⁹⁷ Mousa, A., Broce, M., Yacoub, M., et al. Validation of venous duplex ultrasound imaging in determining iliac vein stenosis after standard treatment of active chronic venous ulcers. *Journal of Vascular Surgery: Venous and Lymphatic Disorders*. 2016; 4: 3, 307-12.

⁹⁸ Nelson, E., O'Meara, S., Craig, D., et al. A series of systematic reviews to inform a decision analysis for sampling and treating infected diabetic foot ulcers. *Health Technology Assessment*. 2006; 10: 12, 1-221.

⁹⁹ Frykberg, R. Diabetic foot ulcers: Current concepts. *The Journal of Foot and Ankle Surgery*. 1998; 37: 5, 440-6.

Statement 6

¹⁰⁰ Purwins, S., Herberger, K., Debus, E. et al. Cost-of-illness of chronic leg ulcers in Germany. *International Wound Journal*. 2010; 7: 2, 97-102.

¹⁰¹ Gorecki, C., Nixon, J., Madill, A., et al. What influences the impact of pressure ulcers on health-related quality of life? A qualitative patient-focused exploration of contributory factors. *Journal of Tissue Viability*. 2012; 21: 1, 3-12.

¹⁰² Green J., Jester R., McKinley R., Pooler A. The impact of chronic venous leg ulcers: a systematic review. *Journal of Wound Care*. 2014; 23: 12, 601-12.

¹⁰³ Krasner, D. Painful Venous Leg Ulcers: Themes and stories about living with the Pain and Suffering. *Ostomy Wound Management*. 1998; 25:3, 158-68.

¹⁰⁴ Hogg F., Peach, G., Price, P., Thompson M., Hinchliffe R. Measures of health-related quality of life in diabetes-related foot disease: a systematic review. *Diabetologia*. 2012; 55: 3, 552-65.

Statement 7

¹⁰⁵ Gorecki, C., Nixon, J., Madill, A., et al. What influences the impact of pressure ulcers on health-related quality of life? A qualitative patient-focused exploration of contributory factors. *Journal of Tissue Viability*. 2012; 21: 1, 3-12.

¹⁰⁶ Kinmond, K., McGee, P., Gough, S., Ashford, R. 'Loss of self': a psychosocial study of the quality of life of adults with diabetic foot ulceration. *Journal of Tissue Viability*. 2003; 13: 1, 6-16.

¹⁰⁷ Phillips T., Stanton, B., Provan, A., Lew, R. A study of the impact of leg ulcers on quality of life: Financial, social, and psychologic implications. *Journal of the American Academy of Dermatology*. 1994; 31: 1, 49-53.

¹⁰⁸ Green J., Jester R., McKinley R., Pooler A. The impact of chronic venous leg ulcers: a systematic review. *Journal of Wound Care*. 2014; 23: 12, 601-12.

¹⁰⁹ Herber, O., Schnepf, W., Rieger, M. A systematic review on the impact of leg ulceration on patients' quality of life. *Health and Quality of Life Outcomes*. 2007; 25:5, 44

Statement 13

¹¹⁰ National Institute for Health and Care Excellence (NICE). Leg Ulcer- Venous. 2016. <https://cks.nice.org.uk/leg-ulcer-venous#!topicsummary> (Accessed May 2017)

¹¹¹ Etufugh, C. Phillips, T. Venous ulcers. *Clinics in Dermatology*. 2007; 21: 1, 121-30.

¹¹² Jones K. Why do chronic venous leg ulcers not heal? *Journal of Nursing Care*. 2009; 24: 2, 116-24.

¹¹³ Wu B., Lu J., Yang, M., Xu, T. Sulodexide for treating venous leg ulcers. *Cochrane Database of Systematic Reviews*. 2016; 2: 6, Article: CD010694

¹¹⁴ Hogg F., Peach, G., Price, P., Thompson M., Hinchliffe R. Measures of health-related quality of life in diabetes-related foot disease: a systematic review. *Diabetologia*. 2012; 55: 3, 552-65.

¹¹⁵ Korn, P., Patel, S., Heller, J., et al. Why insurers should reimburse for compression stockings in patients with chronic venous stasis. *Journal of Vascular Surgery*. 2002; 35: 5, 950-7.

¹¹⁶ Green J., Jester R., McKinley R., Pooler A. The impact of chronic venous leg ulcers: a systematic review. *Journal of Wound Care*. 2014; 23: 12, 601-12.

Statement 8

¹¹⁷ National Institute for Health and Care Excellence (NICE). Leg Ulcer- Venous. 2016. <https://cks.nice.org.uk/leg-ulcer-venous#!topicsummary> (Accessed May 2017)

¹¹⁸ NIHR Signal. Long lasting ulcers below the knee are more common than previously thought. 2016. <https://discover.dc.nihr.ac.uk/portal/article/4000656/long-lasting-ulcers-below-the-knee-are-more-common-than-previously-thought> (Accessed August 2017)

¹¹⁹ Dealey, C. Case study methodology in tissue viability. Part 2: a study to determine the levels of knowledge of nurses providing care for patients with leg ulcers in an acute hospital setting. *Journal of Tissue Viability*. 2001; 11: 1, 28-34.

¹²⁰ Green J., Jester R., McKinley R., Pooler A. The impact of chronic venous leg ulcers: a systematic review. *Journal of Wound Care*. 2014; 23: 12, 601-12.

Statement 4

¹²¹ Domingues, E., Cavalcanti, M., Costa, P., et al. Pain prevalence, socio-demographic and clinical features in patients with chronic ulcers. *Journal of Tissue Viability*. 2016; 25: 3, 180-4.

¹²² Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640.

¹²³ Mousa, A., Broce, M., Yacoub, M., et al. Validation of venous duplex ultrasound imaging in determining iliac vein stenosis after standard treatment of active chronic venous ulcers. *Journal of Vascular Surgery: Venous and Lymphatic Disorders*. 2016; 4: 3, 307-12.

¹²⁴ Herber, O., Schnepf, W., Rieger, M. A systematic review on the impact of leg ulceration on patients' quality of life. *Health and Quality of Life Outcomes*. 2007; 25:5, 44

Statement 46

¹²⁵ Tabolli, S., Tinelli, G., Guarnera, G., et al. Measuring the Health Status of Patients with Vascular Leg Ulcers and the Burden for their Caregivers. *European Journal of Vascular and Endovascular Surgery*. 2007; 34: 5, 613-8.

¹²⁶ Ciliberti, M., De Lara, F., Serra, G., et al. Effective management of pressure ulcers using Hydrofibre technology with silver ions. *Wound Medicine*. 2014; 5, 20-4.

¹²⁷ Krasner, D. Painful Venous Leg Ulcers: Themes and stories about living with the Pain and Suffering. *Ostomy Wound Management*. 1998; 25:3, 158-68.

¹²⁸ Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640.

¹²⁹ Phillips T., Stanton, B., Provan, A., Lew, R. A study of the impact of leg ulcers on quality of life: Financial, social, and psychologic implications. *Journal of the American Academy of Dermatology*. 1994; 31: 1, 49-53.

Statement 47

¹³⁰ Stevens J., Chaloner D. Urgosorb dressing: management of acute and chronic wounds. *British Journal of Nursing*. 2005; 14: 15, Supplement.

¹³¹ Meaume, S. Domp Martin, A., Lok, C., et al. Quality of life in patients with leg ulcers: results from CHALLENGE, a double-blind randomised controlled trial. *Journal of Wound Care*. 2017; 26: 7, 1-10.

Statement 9

¹³² Meaume, S., Truchetet F., Cambazard F., et al. A randomized, controlled, double-blind prospective trial with a Lipido-Colloid Technology-Nano-OligoSaccharide Factor wound dressing in the local management of venous leg ulcers. *Wound Repair and Regeneration*. 2012; 20: 4, 500-11.

¹³³ Schmutz J., Meaume S., Fays S. et al. Evaluation of the nano-oligosaccharide factor lipido-colloid matrix in the local management of venous leg ulcers: results of a randomised, controlled trial. *International Wound Journal*. 2008; 5: 172–182.

Statement 12

¹³⁴ Ouahes, N. Phillips, T. Leg ulcers. *Current Problems in Dermatology*. 1995; 7: 4, 114-42

¹³⁵ Kelechi, T., Johnson, J., Yates, S. Chronic venous disease and venous leg ulcers: An evidence-based update. *Journal of Vascular Nursing*. 2015; 33: 2, 36-46.

¹³⁶ Reichardt, L. Venous ulceration: Compression as the mainstay of therapy. *Journal of Wound, Ostomy and Continence Nursing*. 1999; 26: 1, 39-47.

¹³⁷ Etufugh, C. Phillips, T. Venous ulcers. *Clinics in Dermatology*. 2007; 21: 1, 121-30.

Statement 10

¹³⁸ Tabolli, S., Tinelli, G., Guarnera, G., et al. Measuring the Health Status of Patients with Vascular Leg Ulcers and the Burden for their Caregivers. *European Journal of Vascular and Endovascular Surgery*. 2007; 34: 5, 613-8.

Statement 11

¹³⁹ Ouahes, N. Phillips, T. Leg ulcers. *Current Problems in Dermatology*. 1995; 7: 4, 113-42

¹⁴⁰ Tabolli, S., Tinelli, G., Guarnera, G., et al. Measuring the Health Status of Patients with Vascular Leg Ulcers and the Burden for their Caregivers. *European Journal of Vascular and Endovascular Surgery*. 2007; 34: 5, 613-8

¹⁴¹ Phillips T., Stanton, B., Provan, A., Lew, R. A study of the impact of leg ulcers on quality of life: Financial, social, and psychologic implications. *Journal of the American Academy of Dermatology*. 1994; 31: 1, 49-53

¹⁴² Souliotis, K., Kalemikerakis, I., Saridi M., Papageorgiou, M., Kalokerinou, A. A cost and clinical effectiveness analysis among moist wound healing dressings versus traditional methods in home care patients with pressure ulcers. *Wound Repair and Regeneration*. 2016; 24:3, 596-601.

- 143 O'Brien J., Grace, P., Perry, I., et al. Randomized clinical trial and economic analysis of four-layer compression bandaging for venous ulcers. *British Journal of Surgery*. 2003; 90: 7, 794-8.
- 144 Jeffcoate, W., Price, P., Phillips, C., et al. Randomised controlled trial of the use of three dressing preparations in the management of chronic ulceration of the foot in diabetes. *Health Technology Assessment*. 2009; 13: 54, 1-86.
- Statement 5**
- 145 Ohura T., Sanada H., Mino Y. Clinical study using activity-based costing to assess cost-effectiveness of a wound management system utilizing modern dressings in comparison with traditional wound care. *Nihon Ronen Igakkai Zasshi*. 2004; 41: 1, 82-91.
- 146 Ghatnekar, O., Persson, U., Willis, M., Odegaard, K. Cost effectiveness of becaplermin in the treatment of diabetic foot ulcers in four European countries. *Pharmacoeconomics*. 2001; 19: 7, 767-78.
- 147 Guillén-Solà, M., Soler Mieras, A., Tomàs-Vidal A., and GAUPP-Expert Panel. A multi-center, randomized, clinical trial comparing adhesive polyurethane foam dressing and adhesive hydrocolloid dressing in patients with grade II pressure ulcers in primary care and nursing homes. *BMC Family Practice*. 2013; 14:196.
- 148 Hopkins, A., Worboys, F. Establishing community wound prevalence within an inner London borough: Exploring the complexities. *Journal of Tissue Viability*. 2014; 23: 4, 121-8.
- 149 NHS Right Care. NHS Right Care Scenario. Long term condition scenario: The variation between sub-optimal and optimal pathways. Betty's Story: Leg ulcer wound care. 2017. <https://www.england.nhs.uk/rightcare/wp-content/uploads/sites/40/2017/01/nhs-rightcare-bettys-story-narrative-full.pdf> (Accessed May 2017)
- 150 National Institute for Health and Care Excellence (NICE). Leg Ulcer- Venous. 2016. <https://cks.nice.org.uk/leg-ulcer-venous/#topicsummary> (Accessed May 2017)
- 151 Moffatta, C., Stantonb, J., Murray, S., et al. A randomised trial to compare the performance of Oxzyme® and Iodozyme® with standard care in the treatment of patients with venous and mixed venous/arterial ulceration. *Wound Medicine*. 2014, 6, 1-10.
- 152 Vikatmaa P., Juutilainenb, V., Kuukasjärvic P., Malmivaarac, A. Negative Pressure Wound Therapy: a Systematic Review on Effectiveness and Safety. *European Journal of Vascular and Endovascular Surgery*. 2008; 36: 4, 438-48.
- 153 Martínez-Sánchez, G., Al-Dalaina, S., Menéndezb S., et al. Therapeutic efficacy of ozone in patients with diabetic foot. *European Journal of Pharmacology*. 2005; 523: 1-3, 151-61.
- 154 Ahmad, J. The diabetic foot. *Diabetes & Metabolic Syndrome*. 2016; 10: 1, 48-60.
- 155 Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640.
- Statement 45**
- 156 Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640
- 157 Rudolph, D Standards of care for venous leg ulcers: Compression therapy and moist wound healing. *Journal of Vascular Nursing*. 2001; 19: 1, 20-27.
- 158 Foglia E., Restelli U., Napoletano A., et al. Pressure ulcers management: an economic evaluation. *Journal of Preventative Medicine and Hygiene*. 2012; 53-1, 30-6.
- 159 Guest J., Taylor R., Vowden K., Vowden P. Relative cost-effectiveness of a skin protectant in managing venous leg ulcers in the UK. *Journal of Wound Care*. 2012; 21: 8, 389-94.
- 160 Short, K., Bull, R. Leg ulcers and lymphoedema. *Medicine*. 2009; 37:6, 269-72.
- Statement 45**
- 161 Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640.
- 162 Rudolph, D Standards of care for venous leg ulcers: Compression therapy and moist wound healing. *Journal of Vascular Nursing*. 2001; 19: 1, 20-27.
- 163 Foglia E., Restelli U., Napoletano A., et al. Pressure ulcers management: an economic evaluation. *Journal of Preventative Medicine and Hygiene*. 2012; 53-1, 30-6.
- 164 Guest J., Taylor R., Vowden K., Vowden P. Relative cost-effectiveness of a skin protectant in managing venous leg ulcers in the UK. *Journal of Wound Care*. 2012; 21: 8, 389-94
- 165 Short, K., Bull, R. Leg ulcers and lymphoedema. *Medicine*. 2009; 37:6, 269-72
- 166 Augustin, M., Herberger K., Kroeger K et al. Cost-effectiveness of treating vascular leg ulcers with UrgoStart and UrgoCell Contact. *International Wound Journal*. 2016; 13: 1, 82-7.
- 167 Augustin, M., Herberger K., Kroeger K et al. Cost-effectiveness of treating vascular leg ulcers with UrgoStart and UrgoCell Contact. *International Wound Journal*. 2016; 13: 1, 82-7.
- 168 Moffatta, C., Stantonb, J., Murray, S., et al. A randomised trial to compare the performance of Oxzyme® and Iodozyme® with standard care in the treatment of patients with venous and mixed venous/arterial ulceration. *Wound Medicine*. 2014, 6, 1-10.
- 169 Nisi, G., Brandi, C., Grimaldi, L., et al. Use of a protease-modulating matrix in the treatment of pressure sores. *Chirurgia Italia*. 2005; 57: 4, 465-8.
- Statement 36**
- 170 Augustin, M., Herberger K., Kroeger K et al. Cost-effectiveness of treating vascular leg ulcers with UrgoStart and UrgoCell Contact. *International Wound Journal*. 2016; 13: 1, 82-7
- 171 Ohura T., Sanada H., Mino Y. Clinical study using activity-based costing to assess cost-effectiveness of a wound management system utilizing modern dressings in comparison with traditional wound care. *Nihon Ronen Igakkai Zasshi*. 2004; 41: 1, 82-91.
- 172 Franks, P., Bosanquet, N. Cost-effectiveness: seeking value for money in lower extremity wound management. *International Journal of Lower Extremity Wounds*. 2004; 3: 2, 87-95.
- 173 Zelen, C., Gould, L., Serena, T., et al. A prospective, randomised, controlled, multi-centre comparative effectiveness study of healing using dehydrated human amnion/chorion membrane allograft, bioengineered skin substitute or standard of care for treatment of chronic lower extremity diabetic ulcers. *International Wound Journal*. 2015; 12, 724-732.
- 174 Souliotis, K., Kalemikerakis, I., Saridi M., Papageorgiou, M., Kalokerinou, A. A cost and clinical effectiveness analysis among moist wound healing dressings versus traditional methods in home care patients with pressure ulcers. *Wound Repair and Regeneration*. 2016; 24:3, 596-601.
- 175 Augustin, M., Herberger K., Kroeger K et al. Cost-effectiveness of treating vascular leg ulcers with UrgoStart and UrgoCell Contact. *International Wound Journal*. 2016; 13: 1, 82-7
- Statement 14**
- 176 O'Meara, S., Cullum, N., Nelson, E. Compression for venous leg ulcers Cochrane Database of Systematic Reviews. 2009; 21: 1, Article: CD000265.
- 177 Margolis, D., Allen-Taylor, L., Hoffstad Ma, O., Berlin J. The accuracy of venous leg ulcer prognostic models in a wound care system. *Wound Repair and Regeneration*. 2004; 12: 2, 163-28.
- 178 O'Brien, J., Grace, P., Perry, I., et al. Randomized clinical trial and economic analysis of four-layer compression bandaging for venous ulcers. *British Journal of Surgery*. 2004; 90: 7, 794-8.
- 179 Watson J., Kang'ombe A., Soares M., et al. VenUS III: a randomised controlled trial of therapeutic ultrasound in the management of venous leg ulcers. *Health Technology Assessment* 2011; 15: 13, 1-192.
- 180 Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640.
- Statement 16**
- 181 PrescQIPP. B115. Wound care: Protease-modulating matrix dressings. 2016. <https://www.prescqipp.info/component/jdownloads/send/235-wound-care-protease-modulating-matrix-dressings-2385-b115-wound-care-protease-modulating-matrix-dressings-briefing> (Accessed May 2017)
- 182 Meaume, S., Truchetet F., Cambazard F., et al. A randomized, controlled, double-blind prospective trial with a Lipido-Colloid Technology-Nano-OligoSaccharide Factor wound dressing in the local management of venous leg ulcers. *Wound Repair and Regeneration*. 2012; 20: 4, 500-11.
- 183 Gelfand, J., Hoffstad, O., Margolis, D. Surrogate Endpoints for the Treatment of Venous Leg Ulcers. *Journal of Investigative Dermatology*. 2002; 119: 6, 1420-5.
- Statement 33**
- 184 Meissner, M., Eklof, B., Smith, P., et al. Secondary chronic venous disorders. *Journal of Vascular Surgery*. 2007; 46: 6, S63-8.
- 185 Kelechi, T., Johnson, J., Yates, S. Chronic venous disease and venous leg ulcers: An evidence-based update. *Journal of Vascular Nursing*. 2015; 33: 2, 36-46.
- Statement 37**
- 186 Raju, S., Lurie, F., O'Donnell Jr, T. Compression use in the era of endovenous interventions and wound care centers. *Journal of Vascular Surgery: Venous and Lymphatic Disorder*. 2016; 4: 3, 346-54.
- 187 Augustin, M., Herberger K., Kroeger K et al. Cost-effectiveness of treating vascular leg ulcers with UrgoStart and UrgoCell Contact. *International Wound Journal*. 2016; 13: 1, 82-7
- 188 Augustin, M., Vanscheidt, W. Comment: Chronic venous leg ulcers: the future of cell-based therapies. *The Lancet*. 2010; 380: 9848, 953-5
- 189 Kelechi, T., Mueller, M., Hankin, C., et al. A randomized, investigator-blinded, controlled pilot study to evaluate the safety and efficacy of a poly-N-acetyl glucosamine-derived membrane material in patients with venous leg ulcers. *Journal of the American Academy of Dermatology*. 2012; 66: 6, 209-215.
- Statement 43**
- 190 Schmutz J., Meaume S., Fays S. et al. Evaluation of the nano-oligosaccharide factor lipido-colloid matrix in the local management of venous leg ulcers: results of a randomised, controlled trial. *International Wound Journal*. 2008; 5: 172-182.
- Statement 15**
- 191 Guidelines and Audit Implementation Network (GAIN). An investigation into lower leg ulceration in Northern Ireland. 2013. <https://rqia.org.uk/RQIA/files/dc/dc8d08a4-8560-4fbc-9784-6204607e985b.pdf> (Accessed May 2017)
- 192 Currie C, Morgan C, Peters J. The epidemiology and cost of inpatient care for peripheral vascular disease, infection, neuropathy, and ulceration in Diabetes. *Diabetes Care*. 1998; 21: 42-8.
- 193 Dealey C., Posnett J., Walker A. The cost of pressure ulcers in the United Kingdom. *Journal of Wound Care*. 2012; 21: 6, 261-6
- 194 Smith, G., Ingram, A. Clinical and cost effectiveness evaluation of low friction and shear garments. *Journal of Wound Care*. 2010; 19: 12, 535-42.
- 195 Demarré, L., Van Lancker, A., Van Hecke, A., et al. The cost of prevention and treatment of pressure ulcers: A systematic review. *International Journal of Nursing Studies*. 2015; 52: 11, 1754-74.
- Statement 35**
- 196 Raju, S., Lurie, F., O'Donnell Jr, T. Compression use in the era of endovenous interventions and wound care centers. *Journal of Vascular Surgery: Venous and Lymphatic Disorder*. 2016; 4: 3, 346-54.
- 197 Hopkins, A., Worboys, F. Establishing community wound prevalence within an inner London borough: Exploring the complexities. *Journal of Tissue Viability*. 2014; 23: 4, 121-8.
- 198 Augustin, M., Herberger K., Kroeger K et al. Cost-effectiveness of treating vascular leg ulcers with UrgoStart and UrgoCell Contact. *International Wound Journal*. 2016; 13: 1, 82-7

¹⁹⁹ Augustin, M., Vanscheidt, W. Comment: Chronic venous leg ulcers: the future of cell-based therapies. *The Lancet*. 2010; 380: 9848, 953-5

²⁰⁰ Smith, G., Ingram, A. Clinical and cost effectiveness evaluation of low friction and shear garments. *Journal of Wound Care*. 2010; 19: 12, 535-42.

Statement 32

²⁰¹ National Health and Medical Research Council (NHMRC). Australian and New Zealand clinical practice guideline for prevention and management of venous leg ulcers. 2011.

https://www.nhmrc.gov.au/files/nhmrc/publications/attachments/ext003_venous_leg_ulcers_aust_nz_0.pdf (Accessed May 2017)

²⁰² Pang, K., Bate, G., Darvall, K et al. Healing and Recurrence Rates Following Ultrasound-guided Foam Sclerotherapy of Superficial Venous Reflux in Patients with Chronic Venous Ulceration. *European Journal of Vascular and Endovascular Surgery*. 2010; 40: 6, 790-5.

²⁰³ Ouahes, N. Phillips, T. Leg ulcers. *Current Problems in Dermatology*. 1995; 7: 4, 113-42

²⁰⁴ Rudolph, D. Pathophysiology and management of venous ulcers. *Journal of Wound Ostomy and Continence Nursing*. 1998; 25: 5, 248-55.

Statement 39

²⁰⁵ Carter, M., Waycaster, C., Schaum, K., Gilligan, A. Cost-Effectiveness of Three Adjunct Cellular/Tissue-Derived Products Used in the Management of Chronic Venous Leg Ulcers. *Value in Health*. 2014; 17: 8, 801-13.

²⁰⁶ Meaume, S., Truchetet F., Cambazard F., et al. A randomized, controlled, double-blind prospective trial with a Lipido-Colloid Technology-Nano-OligoSaccharide Factor wound dressing in the local management of venous leg ulcers. *Wound Repair and Regeneration*. 2012; 20: 4, 500-11.

²⁰⁷ Meissner, M., Eklof, B., Smith, P., et al. Secondary chronic venous disorders. *Journal of Vascular Surgery*. 2007; 46: 6, S63-8.

²⁰⁸ Reichardt, L. Venous ulceration: Compression as the mainstay of therapy. *Journal of Wound, Ostomy and Continence Nursing*. 1999; 26: 1, 39-47.

²⁰⁹ Ghatnekar, G., Grek, C., Armstrong, D., et al. The Effect of a Connexin43-Based Peptide on the Healing of Chronic Venous Leg Ulcers: A Multicenter, Randomized Trial. *Journal of Investigative Dermatology*. 2015; 135: 1, 289-98.

²¹⁰ Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640.

²¹¹ Snyder, R. Treatment of nonhealing ulcers with allografts. *Clinics in Dermatology*. 2005; 24:4, 388-95.

Statement 34

²¹² Meaume, S., Truchetet F., Cambazard F., et al. A randomized, controlled, double-blind prospective trial with a Lipido-Colloid Technology-Nano-OligoSaccharide Factor wound dressing in the local management of venous leg ulcers. *Wound Repair and Regeneration*. 2012; 20: 4, 500-11.

²¹³ Meissner, M., Eklof, B., Smith, P., et al. Secondary chronic venous disorders. *Journal of Vascular Surgery*. 2007; 46: 6, S63-8.

²¹⁴ Reichardt, L. Venous ulceration: Compression as the mainstay of therapy. *Journal of Wound, Ostomy and Continence Nursing*. 1999; 26: 1, 39-47.

²¹⁵ Ghatnekar, G., Grek, C., Armstrong, D., et al. The Effect of a Connexin43-Based Peptide on the Healing of Chronic Venous Leg Ulcers: A Multicenter, Randomized Trial. *Journal of Investigative Dermatology*. 2015; 135: 1, 289-98.

²¹⁶ Alavi, A., Sibbald, R., Phillips, T., et al. What's new: Management of venous leg ulcers: Treating venous leg ulcers. *Journal of the American Academy of Dermatology*. 2016; 74: 4, 627-640.

²¹⁷ Snyder, R. Treatment of nonhealing ulcers with allografts. *Clinics in Dermatology*. 2005; 24:4, 388-95.

²¹⁸ Dumville, J., O'Meara, S., Deshpande, S. and Speak, K. Hydrogel dressings for healing diabetic foot ulcers. *Cochrane Database of Systematic Reviews*. 2013; 12: 7, CD009101

²¹⁹ Dumville, J., O'Meara, S., Deshpande, S. and Speak, K. Alginate dressings for healing diabetic foot ulcers. *Cochrane Database of Systematic Reviews*. 2013; 25:6, Article: CD009110.

²²⁰ Dumville, J., O'Meara, S., Deshpande, S. and Speak, K. Hydrocolloid dressings for healing diabetic foot ulcers. *Cochrane Database of Systematic Reviews*. 2013; 6: 8, CD009099

²²¹ Dumville, J., Deshpande, S., O'Meara, S. and Speak, K. Foam dressings for healing diabetic foot ulcers. *Cochrane Database of Systematic Reviews*. 2013; 6: 6, Article: CD009111.

²²² Westby, M., Norman, G., Dumville, J., Stubbs, N. and Cullum, N. Protease-modulating matrix treatments for healing venous leg ulcers. *Cochrane Database of Systematic Reviews*. 2016; 12. Article: CD011918.