



**Manchester  
Metropolitan  
University**

---

Šimunic, Boštjan and Pišot, Rado and Rittweger, Jörn and Degens, Hans (2018) *Age-related Slowing of Contractile Properties Differs between Power-, Endurance- and non-athletes; a Tensiomyographic Assessment*. *Journals of Gerontology, Series A*, 73 (12). pp. 1602-1608. ISSN 1079-5006

---

**Downloaded from:** <http://e-space.mmu.ac.uk/620427/>

**Version:** Accepted Version

**Publisher:** Oxford University Press on behalf of The Gerontological Society of America

**DOI:** <https://doi.org/10.1093/gerona/gly069>

Please cite the published version

<https://e-space.mmu.ac.uk>

## **Title page**

**TITLE:** Age-related slowing of contractile properties differs between power-, endurance- and non-athletes; a tensiomyographic assessment

## **Authors:**

Assoc. Prof. Boštjan Šimunič, PhD <sup>1</sup> (email: [bostjan.simunic@zrs-kp.si](mailto:bostjan.simunic@zrs-kp.si))

Prof. dr. Rado Pišot, PhD <sup>1</sup>

Prof. Jörn Rittweger, PhD, MD <sup>2,3</sup>

Prof. Hans Degens, PhD <sup>4,5</sup>

## **Affiliations:**

<sup>1</sup> Science and Research Centre Koper, Institute for Kinesiology Research, Koper, Garibaldijeva 1, SI- 6000 Koper, Slovenia;

<sup>2</sup> Institute of Aerospace Medicine, German Aerospace Center, Linder Hoehe, D-51147 Cologne, Germany;

<sup>3</sup> Department of Pediatrics and Adolescent Medicine, University of Cologne, Cologne, Germany;

<sup>4</sup> School of Healthcare Science, Manchester Metropolitan University, Manchester, UK;

<sup>5</sup> Institute of Sport Science and Innovations, Lithuanian Sports University, Kaunas, Lithuania

## **Corresponding author:**

Boštjan Šimunič

Science and Research Centre Koper

Garibaldijeva 1

6000 Koper

Email: [bostjan.simunic@zrs-kp.si](mailto:bostjan.simunic@zrs-kp.si)

© The Author(s) 2018. Published by Oxford University Press on behalf of The Gerontological Society of America. All rights reserved. For permissions, please e-mail: [journals.permissions@oup.com](mailto:journals.permissions@oup.com).

## Abstract

Although master athletes maintain high levels of physical activity, they also suffer from an age-related decline in skeletal muscle function. There are indications of disproportional age- and physical inactivity-induced muscle wasting between muscles. Tensiomyography is a non-invasive tool that has been used to study the effects of a variety of sports on the contraction time (Tc) in different skeletal muscles. The aim of this cross-sectional study was to assess age-related changes in the Tc of the vastus lateralis, gastrocnemius medialis and biceps femoris muscles with Tensiomyography in older non-athletes (age =  $62.1 \pm 12.7$  years;  $N_{\text{MALES}} = 133$ ;  $N_{\text{FEMALES}} = 246$ ), and power (age =  $56.9 \pm 13.5$  years;  $N_{\text{MALES}} = 100$ ;  $N_{\text{FEMALES}} = 78$ ) and endurance master athletes (age =  $56.5 \pm 14.5$  years;  $N_{\text{MALES}} = 76$ ;  $N_{\text{FEMALES}} = 73$ ). We found an age-related slowing in all muscles, irrespective of discipline, where endurance master athletes had the longest and power master athletes had the shortest Tc. The longer Tc in endurance master athletes than in non-athletes suggests that regular endurance sport activity aggravates slowing of skeletal muscles during ageing.

**Key words:** Skeletal muscle, TMG, Contraction time, Master athletes, MHC

## Introduction

In non-athletes muscle mass shows a progressive decline of as much as 1 to 1.5% per year after about the age of 50<sup>1</sup>. The age-related loss of muscle mass is a consequence of loss and atrophy of muscle fibres<sup>2-4</sup>. Since many of the age-related changes in skeletal muscle are similar to those induced by disuse<sup>5</sup> it is likely that the decrease in physical activity in old age<sup>6,7</sup> is a major contributor to the muscle wasting during ageing. Master athletes, however, maintain high levels of physical activity<sup>8-10</sup> and suffer from fewer morbidities<sup>11</sup>, thereby providing a unique human research model to disentangle the effects of disuse and co-morbidities from ageing *per se*.

Master athletes are athletes older than 35 years who train for and participate in athletic competitions. Their training regimes range from 6 to over 20 hours per week<sup>12</sup> and they may well work at the limits of physiology<sup>10</sup>. Even though their athletic performance exceeds that in age-matched non-athletes, also master athletes suffer from an age-related decline in muscle force and power<sup>2,13,14</sup>, ventilatory function<sup>15</sup> and maximal oxygen consumption<sup>16</sup>.

In line with these decreases in physiological parameters, jump, sprint and endurance performance decline with age<sup>13,17</sup>, something seen even at the individual level<sup>18</sup>, and after the age of 70 it even appears to be an exponential decline<sup>19</sup>. This age-related decline in performance is not only caused by a loss of muscle mass and force generating capacity, but also by slowing of movement as a consequence of slower muscle contractile properties and increased tendon compliance<sup>13</sup>.

Part of the slowing of skeletal muscle is attributable to a preferential atrophy of fast type II fibres<sup>3,4,20,21</sup>, which produce at least 5-times as much power as slow type I fibres<sup>22</sup>. Single-

fibre studies have also provided evidence of an age-related decrease in shortening velocity independent of fibre type switching in fibres expressing types I and IIa myosin heavy chain (MHC) isoforms<sup>23,24</sup>. Thus, the loss of muscle mass, changes in fibre type composition and slowing of muscle fibres may all contribute to the decline in anaerobic performance.

Tensiomyography (TMG) is a non-invasive tool to assess the contractile properties (e.g. contraction time – Tc) of a muscle and can be used to estimate the percentage of type I myosin heavy chain at least in the vastus lateralis (VL) muscle<sup>25</sup>. We have recently shown in a large cohort of adolescents that regular sport exercise decreased Tc in the non-postural biceps femoris (BF), but not in the postural VL<sup>26</sup>. In addition, we observed that 8 weeks of plyometric training in young athletes results in an 8-26% shorter Tc in five lower limb skeletal muscles that explained ~30% of the improvement in explosive power<sup>27</sup>. This suggests that Tc is indeed related to muscle contractile properties and secondly that regular exercise may modulate the effects of ageing on skeletal muscle properties.

Previously we found in older people that a lower performance in the timed-up-and-go and 6-minute walk tests was to a significant extent related to a lower maximal shortening velocity of the muscle<sup>28</sup>. A longer Tc in older people may likewise have similar implications for functions of daily life. Yet, despite numerous studies reporting TMG results in children, adults and athletes, there is only one study<sup>29</sup> that investigated the impact of ageing on the TMG-derived velocity of contraction. They reported that the oldest group had the lowest velocity of contraction, but it is unclear whether this was due to a lower Tc and/or higher Dm. Therefore, the aim of this study was to assess age-related changes in VL, BF and gastrocnemius medialis (GM) muscle contractile properties with TMG in male and female older non-athletes, and master endurance and power athletes in a cross-sectional study.

## Methods

### Participants

Altogether 706 participants (311 men; aged 35 to 90 years) were recruited. Master athletes were recruited during: (i) European Veteran Athletics Championships in Ljubljana 2008 (EVACS; N=253); (ii) European Master Games in Lignano 2011 (EMG; N=56); and (iii) Slovenian Championships in Master Athletics Ljubljana 2015 (VAD; N=18). From the 176 male master athletes 40 were sprinters ( $\leq 400$  m), 38 were throwers, 22 were jumpers, 57 runners ( $\geq 800$  m) and 19 competed in walking events. From the 151 female master athletes 29% were sprinters ( $\leq 400$  m), 29% were throwers, 20% were jumpers, 51 runners ( $\geq 800$  m) and 22% competed in walking events. All master athletes had competed for more than 3 years in master championships. Data of non-athletes were obtained from mass measurements during two international projects led by the Institute for Kinesiology Research at the Science and Research Centre Koper: (i) Physical Activity and Nutrition for Great Ageing 2013-15 (PANGEA; N=259); and (ii) Active and Quality Ageing in Home Environment 2015-16 (A-Qu-A; N=120). Non-athletes were not members of any sport clubs and did not compete in sport championships. Participants were excluded if they suffered from major skeletal, muscular or nervous disorders. Basic anthropometric data and age, sex and sport group distribution are presented in Table 1. The study was reviewed and approved by the Republic of Slovenia National Medical Ethics Committee (EVACS, PANGEA, and AQUA), Italian Ethics Committee at regional level (EMG), and Institutional Ethics Committee of the University of Primorska (VAD). Data collection conformed to the standards set by the Declaration of Helsinki (2002) and its amendments. All participants were fully informed of any risks and benefits associated with the study, and written informed consent was obtained from each participant prior to data collection.

<< Table 1 around here >>

## **Research design**

The study was cross-sectional and participants were divided by sex, sport and age. We defined three sport groups: non-athletes (NA), power athletes (PowA: sprints  $\leq$  400m, jumps and throws) and endurance athletes (EndA: runs  $\geq$  800m; walking). We defined three age groups: 35-49 years, 50-64 years and  $\geq$ 65 years, where the oldest participant was 90 years old.

## **Anthropometric measurements**

The participants were asked to abstain from intense physical activity for 24 hrs (NA) or competition (PowA and EndA) before the assessment. After determination of body height and body mass (Seca Instruments Ltd., Hamburg, Germany) TMG was performed.

## **TMG**

TMG was used to assess skeletal muscle contractile properties in the VL, BF and GM muscles. All measurements were performed during electrically-evoked isometric contractions. For VL TMG, participants were in a supine position with the knee angle at  $30^{\circ}$  flexion (where  $0^{\circ}$  represents a fully extended knee joint). For the BF they were in a prone position with the knee at  $5^{\circ}$  flexion and for the GM they were in a prone position with the ankle in neutral position as previously reported<sup>25,30</sup>. Foam pads were used to support the joints.

A single 1-ms maximal monophasic electrical impulse was used to elicit a twitch that caused the muscle belly to oscillate. These oscillations were recorded using a sensitive digital displacement sensor (TMG-BMC, Ljubljana, Slovenia) that was placed on the surface of the skin over the mid belly of the muscle of interest. If needed, the measuring point and electrode

---

positions were adjusted to obtain maximal Dm of the muscle belly. Initially, the stimulation amplitude was set just above the threshold and then gradually increased until the Dm of the radial twitch displacement increased no further. From two maximal twitch responses, Tc was calculated and the average used for further analysis. Tc was defined (Figure 1) as the time for the amplitude to increase from 10% to 90% of Dm <sup>25</sup>.

<< Figure 1 around here >>

### **Muscle biopsies**

In a subsample of master athletes (N=17; 4 EndA + 13 PowA) we have taken VL muscle biopsies after TMG assessment and determined the myosin heavy chain (MHC) composition as described previously <sup>25</sup>. Briefly, VL muscle biopsies were obtained with a conchotome after skin sterilization and local anaesthesia (2% lidocaine) and sterile conditions. The sample (~100 mg) was taken at 40% from the distance of the knee joint (0%) and the spina iliaca anterior superior (100%). The samples were placed on cork perpendicular to the long axis of the fibres, frozen in liquid nitrogen with vigorous shaking and stored at -80 °C until analysis. For MHC determination, 10- $\mu$ m cross-sections were cut in a cryostat, collected in a centrifuge tube with Laemli sample buffer (3 sections in 100  $\mu$ L) and boiled for 2 minutes to denature the proteins. Twelve- $\mu$ L samples were loaded on SDS poly-acrylamide gels run at 15 °C, 120 V for 27 hours. The stacking gel contained 4% acrylamide and the separating gel had 7% acrylamide with both containing 30% glycerol. After the run, the gels were stained with the Silver Stain Plus kit following the instructions of the manufacturer (BIORAD LABORATORIES, United Kingdom). Bands (Figure 2) were identified based on the migration distance, and relative quantities determined with Quantity One software (BIORAD LABORATORIES, United Kingdom).



<< Figure 2 around here >>

## Statistics

SPSS (IBM, USA) software was used for all statistical analyses. All data in text and tables are presented as mean  $\pm$  standard deviation, while in figures standard errors were used. Visual inspection and the Shapiro-Wilk test indicated that all data were normally distributed. Sphericity (homogeneity of covariance) was verified by the Mauchly's test. When the assumption of sphericity was not met, the significance of the F-ratios was adjusted according to the Greenhouse-Geisser procedure. Main effects were studied with a General Linear Model repeated-measures ANOVA with muscle (VL, GM, and BF) as within factor, and three between factors: sport group (three levels: NA, PowA, EndA), sex (man or woman) and age (three levels: 35-49 years, 50-64 years and  $\geq 65$  years). Where significant effects were found for sport or age effects or 2-way interactions (three-way interactions were excluded from the analysis), post-hoc analysis with Bonferroni corrections was used to locate the differences between sport and age groups. We used the Pearson correlation coefficient to assess associations between age and Tc in each muscle and sex. We also performed stepwise multiple regression analysis of Tc with age, sex, muscle and sport as predictors, with entry criteria at  $p \leq .05$  and removal criteria at  $p \geq .1$ . In a subsample of 17 master athletes who also gave a VL muscle biopsy we performed Pearson correlation analysis to assess to what extent VL Tc is affected by the MHC-I content of the muscle, age and sport. Statistical significance was accepted at p-values  $< .05$ . Additionally, the effect size for dependent variables was assessed as partial eta-squared ( $\eta^2$ ).

## Results

Table 1 shows the anthropometric data. We found that PowA were the tallest ( $P=.005$ ), and that they and NA had a higher body mass than EndA ( $P<.001$ ). NA had a higher body mass index than PowA and EndA ( $P<.001$ ). Women were smaller ( $P<.001$ ), lighter ( $P<.001$ ) and had a lower body mass index ( $P=.001$ ) than men.

Tc was positively correlated with age, irrespective of muscle and sport group (Figure 3). This correlation was weakest in VL (even not significant for female NA), and strongest in BF for PowA and EndA, where EndA had the strongest correlation in GM.

<< Figure 3 around here >>

A multiple regression analysis (Table 2) confirmed an overall correlation of  $R=.690$  ( $p<.001$ ) with 47.6% of the variance in Tc explained by Age, BF, VL, and athletic discipline as predictors, but not sex. The largest contributors (part correlations) to the explained variance in Tc are BF and Age. The multiple regression prediction model is as follows:

$$Tc = 11.754 \times BF + .193 \times Age - 3.292 \times An - 3.185 \times VL + 3.142 \times Ae + 18.098$$

where Tc is a muscle contraction time (in ms), BF is the biceps femoris muscle (0 or 1), VL is the vastus lateralis muscle (0 or 1), Age is an age (in years), An is Power athletes (0 or 1), and Ae is Endurance athletes (0 or 1).

<< Table 2 around here >>

In a subsample of master athletes ( $N_{POW}=13 + N_{END}=4$ ) we assessed the MHC composition of the VL (Table 3) and found a significant bi-variate correlation between Tc and MHC-I ( $r=.864$ ;  $P<.001$ ), while there was no significant correlation between age and Tc ( $P=.309$ ) or MHC-I ( $P=0.719$ ). A correlation was found also between VL Tc and MHC-IIa ( $r=-.760$ ;  $P<.001$ ) but not between VL Tc and MHC-IIx ( $r=-.326$ ;  $P=.201$ ).

<< Table 3 around here >>

After dividing the participants into three age groups we found significant effects for muscle ( $P<.001$ ;  $\eta^2=.395$ ), sport ( $P<.001$ ;  $\eta^2=.133$ ), age ( $P<.001$ ;  $\eta^2=.153$ ), and sport\*age ( $P<.001$ ;  $\eta^2=.036$ ), muscle\*sex ( $P=.048$ ;  $\eta^2=.005$ ), muscle\*sport ( $P<.001$ ;  $\eta^2=.033$ ), muscle\*age ( $P<.001$ ;  $\eta^2=.043$ ) interactions. As there were interactions between muscle with sport, age and sex, we performed subsequent post-hoc ANOVAs first for the non-athletes only and then for each muscle separately (Figure 4).

<< Figure 4 around here >>

Looking at the NA only, we found the lowest Tc in VL ( $26.1 \pm 4.2$  ms), the longest in BF ( $43.1 \pm 11.3$  ms) and that of the GM ( $30.0 \pm 7.7$  ms) in between the two.

In the VL, there were effects of age ( $P<.001$ ;  $\eta^2=.039$ ), sex ( $P<.001$ ;  $\eta^2=.017$ ), sport ( $P<.001$ ;  $\eta^2=.065$ ), and sport\*age ( $P=.003$ ;  $\eta^2=.023$ ) and sex\*age ( $P=.072$ ;  $\eta^2=.008$ ) interactions. The interactions indicated that the effects of age differed between sports and sexes. The sex\*age interaction was reflected by an age-related increase in Tc in men only ( $P=0.002$ ). The sport\*age interaction was reflected by the shorter Tc in PowA than NA at all ages in men only

( $P=.004$ ). In both male PowA and NA Tc increased from 35-49 to 50-64 years by 8.7% ( $P=.004$ ) and 10.5% ( $P<.001$ ), respectively. The Tc was higher in 65-90-year-old EndA than 50-64-year-old EndA ( $P=.015$ ) and EndA had a longer Tc than NA ( $P=.008$ ).

In the GM, there were effects of age ( $P<.001$ ;  $\eta^2=.066$ ) and sport ( $P<.001$ ;  $\eta^2=.039$ ), and a sport\*age interaction ( $P=.009$ ;  $\eta^2=.022$ ). The latter indicated that the effect of age differed between the different sports. Although in all groups Tc was longer at 65-90 years in comparison to 35-49 years (10.2%;  $P=.017$ ), PowA had shorter Tc than NA at 35-49 years (-9.7%;  $P=.014$ ), and EndA had longer Tc at 65-90 years than NA (22.6%;  $P<.001$ ).

In the BF there were effects of age ( $P<.001$ ;  $\eta^2=.129$ ) and sport ( $P<.001$ ;  $\eta^2=.099$ ), and sport\*age ( $P=.086$ ;  $\eta^2=.012$ ) and sex\*sport ( $P=.087$ ;  $\eta^2=.007$ ) interactions. The interactions indicated that the effects of sport differed between sports, and between men and women. Tc was consistently increased with increasing age in PowA, but PowA always had a shorter Tc than NA in both men and women. However, the relative difference of Tc between NA and PowA was decreasing with increasing age (in men: from 20.5% to 12.3%; in women from 27.4% to 11.3%). In EndA Tc was higher than in NA at 35-49 years (only in men: 22.5%;  $P=.001$ ), then stabilised at 50-64 years and increased again at 65-90 years (in men: 33.2%;  $P<.001$ , in women: 26.5%;  $P<.001$ ), when EndA had higher Tc than NA (only in men: 20.5%;  $P<.001$ ). We found that body mass and body mass index explained at best 1% of the variance (data not shown).

## Discussion

In the present cross-sectional study, we used TMG to non-invasively assess the contractile properties of leg skeletal muscles in master endurance and power athletes, and age-matched

non-athletes. The main observation of the present study is the age-related slowing of all muscles. Endurance athletes had slower muscles than age-matched non-athletes, while power athletes had the fastest contractile properties, irrespective of sex. For all muscles, an age\*sport interaction was found, indicating that the effects of age differ between sports. Given that endurance athletes had slower contractile properties than non-athletes, this suggests that regular long-distance running accelerates the age-related slowing of skeletal muscle.

### *TMG*

TMG was developed to non-invasively measure skeletal muscle contractile properties<sup>31</sup>. We have previously shown in a population of 20-83-year olds that Tc predicted 77% of the variance of the proportion of type I MHC in the VL<sup>25</sup>, which was confirmed by our results obtained in a sub group of 17 master athletes (75%). Although this regression model has not been validated in other muscles, there are no obvious reasons to believe that it would not apply to other muscles, albeit with different regression coefficients. Here we show that Tc was lowest in the VL ( $26.1 \pm 4.2$  ms), then in GM ( $30.0 \pm 7.7$  ms) and longest in BF ( $43.1 \pm 11.3$  ms), corresponding with the reported differences in type I proportions between these muscles (50%, 54-63% and 67%, respectively)<sup>32,33</sup>.

### *Sex and sport related differences in contractile properties*

We did not observe significant differences between men and women in the contractile properties of the VL, BF and GM. This corresponds with the similar fibre type distribution in men and women<sup>21,34,35</sup>.

Power athletes had a lower Tc than endurance and non-athletes, particularly in the non-postural BF, and to a lesser extent in the postural VL and GM. The shorter Tc in power

athletes is most likely attributable to a higher shortening velocity in both type I (75%) and type IIa (45%) fibres, as reported after 12 weeks of resistance exercise in older men <sup>36</sup>. Interestingly, the same authors <sup>37</sup> demonstrated that the same programme in women induced a decrease in the shortening velocity of type I fibres, with no change in that of type IIa fibres. We did not observe a significant sex\*sport interaction for VL, but such an interaction for the BF was reflected by a longer Tc in male, rather than female, endurance than non-athletes, while female endurance and non-athletes had a similar Tc. We have no explanation for these differences, but the effects of resistance exercise on single fibre contractile properties are not unequivocal, with at least one study showing no change in single fibre contractile properties in young men <sup>38</sup>.

#### *Age-related slowing*

In the present study, we observed an age-related slowing of the muscle contractile properties, confirming the general observation of an age-related slowing of muscle in both non-athletes and master athletes <sup>13,39</sup>. This may be attributable to a preferential loss of type II fibres <sup>40</sup>. That, however, is not an unequivocal observation as other studies did not see such a change <sup>4,21,41</sup> and one longitudinal study even reported an increased proportion of type II fibres over a 12-year period <sup>24</sup>. Even if type II fibres are preserved, their contribution to the contractile properties of the muscle may decrease if they undergo a larger age-related atrophy than type I fibres. Indeed, some studies reported a protection of type I fibres as compared with type II fibres <sup>4,21,40,42</sup>, but again, others have reported that type I and II fibres atrophy similarly with age <sup>41,43</sup>. The discrepancy between studies could be attributed to several factors: (i) pooling of muscle fibre phenotypes; (ii) different muscles investigated; (iii) small study samples; and (iv) by necessity when using biopsies, a relatively small amount of muscle tissue being analysed.

Age-related slowing may not only be due to fibre type shifts, but also slowing of fibres. In fact, it has been reported in both rodents<sup>44</sup> and humans<sup>43,45</sup> that particularly type I and IIa fibres are exhibiting an age-related slowing independent of shifts in myosin heavy and light chain isoform composition. Such an age-related slowing maybe caused by glycation of the myosin molecule<sup>46</sup> that may have a mitochondrial origin<sup>47</sup>. Given that endurance athletes have a larger proportion of type I and IIa fibres than power and non-athletes<sup>14</sup>, we expected the slowing to be more pronounced in the endurance than power and non-athletes, something we did indeed observe. While this may to some extent, as in ageing, be related to glycation of myosin in these fibres, such an effect will be attenuated, but probably not entirely abolished, in endurance athletes, who have a higher insulin sensitivity than non-athletes (Goodpaster, 2001). The role of myosin glycation is admittedly speculative, but is an interesting avenue of investigation. If this pathway is indeed involved then consumption of anti-glycating agents<sup>48</sup> may alleviate some of the age-related slowing, particularly in endurance athletes.

### *Conclusion and Perspective*

TMG revealed that the age-related slowing of muscle contractile properties occurs particularly in endurance athletes. Here we suggest that this may be related to their high proportion of type I and IIa fibres that have been reported to exhibit an age-related slowing independent of shifts in myosin heavy and light chain composition.

### **Funding**

This work was supported by the Cross-border Cooperation Program Slovenia–Italy 2007–2013 with European Regional Development Fund (PANGEA – Physical Activity and Nutrition for Quality Ageing, grant number 042-2/2009-18/052012); by the Slovenian Research Agency (L5-5550 – Development of Non-invasive Marker for Muscle Atrophy,

grant number 1000-15-1988); by the Norwegian Financial Mechanism and the Republic of Slovenia Government Office for Development and European Cohesion (A-Qu-A – Active and Quality Ageing in Home Environment, grant number 4300-472/2014); and by Slovenian Sport Foundation (The Effect of Regular Sport Exercise on Muscle Composition in Master Athletes, grant number RR-14-419).

### **Acknowledgements**

We acknowledge the excellent assistance of the entire staff of the European Veteran Athletics Championships (Ljubljana, Slovenia). We would like to thank all project partners and other researchers involved in data collection, students, participants and everyone who helped.

### **Literature**

1. Janssen I, Heymsfield SB, Wang Z, Ross R. Skeletal muscle mass and distribution in 468 men and women aged 18–88 yr. *J Appl Physiol*. 2000;89(1):81-88.  
doi:10.1093/oxfordjournals.aje.a009520
  2. Degens H. Determinants of skeletal muscle hypertrophy and the attenuated hypertrophic response at old age. *J Sports Med Doping Stud*. 2012;S1:003:2-9.  
doi:10.4172/2161-0673.S1-003
  3. Sato T, Akatsuka H, Kito K, Tokoro Y, Tauchi H, Kato K. Age changes in size and number of muscle fibers in human minor pectoral muscle. *Mech Ageing Dev*. 1984;28(1):99-109.
  4. Lexell J, Taylor CC, Sjöström M. What is the cause of the ageing atrophy?. Total number, size and proportion of different fiber types studied in whole vastus lateralis muscle from 15- to 83-year-old men. *J Neurol Sci*. 1988;84(2-3):275-294.  
doi:10.1016/0022-510X(88)90132-3
-



5. Degens H, Alway SE. Control of Muscle Size During Disuse, Disease, and Aging. *Int J Sports Med.* 2006;27(2):94-99. doi:10.1055/s-2005-837571
6. Wullems JA, Verschueren SMP, Degens H, Morse CI, Onambélé GL. A review of the assessment and prevalence of sedentarism in older adults, its physiology/health impact and non-exercise mobility counter-measures. *Biogerontology.* 2016;17(3):547-565. doi:10.1007/s10522-016-9640-1
7. McPhee JS, French DP, Jackson D, Nazroo J, Pendleton N, Degens H. Physical activity in older age: perspectives for healthy ageing and frailty. *Biogerontology.* 2016;17(3):567-580. doi:10.1007/s10522-016-9641-0
8. Suominen H, Korhonen M. Sport Performance in Master Athletes: Age-Associated Changes and Underlying Neuromuscular Factors. In: *Neuromuscular Aspects of Sport Performance.* Oxford, UK: Wiley-Blackwell; 2010:270-282. doi:10.1002/9781444324822.ch15
9. Hawkins S, Wiswell R, Marcell T. Exercise and the master athlete--a model of successful aging? *J Gerontol A Biol Sci Med Sci.* 2003;58(11):1009-1011. doi:10.1093/gerona/58.11.M1009
10. Rittweger J, Kwiet A, Felsenberg D. Physical performance in aging elite athletes - Challenging the limits of physiology. *J Musculoskelet Neuronal Interact.* 2004;4(2):159-160.
11. Kettunen JA, Kujala UM, Kaprio J, Sarna S. Health of Master Track and Field Athletes. *Clin J Sport Med.* 2006;16(2):142-148. doi:10.1097/00042752-200603000-00010
12. Kasch FW, Boyer JL, Van Camp SP, Verity LS, Wallace JP. Effect of exercise on cardiovascular ageing. *Age Ageing.* 1993;22(1):5-10. <http://www.ncbi.nlm.nih.gov/pubmed/8438667>.

13. Arampatzis A, Degens H, Baltzopoulos V, Rittweger J. Why do older sprinters reach the finish line later? *Exerc Sport Sci Rev.* 2011;39(1):18-22.  
doi:10.1097/JES.0b013e318201efe0
14. Korhonen MT. Aging, muscle fiber type, and contractile function in sprint-trained athletes. *J Appl Physiol.* 2006;101(3):906-917. doi:10.1152/jappphysiol.00299.2006
15. Degens H, Maden-Wilkinson TM, Ireland A, et al. Relationship between ventilatory function and age in master athletes and a sedentary reference population. *Age (Omaha).* 2013;35(3):1007-1015. doi:10.1007/s11357-012-9409-7
16. Tanaka H, Seals DR. Endurance exercise performance in Masters athletes: age-associated changes and underlying physiological mechanisms. *J Physiol.* 2008;586(1):55-63. doi:10.1113/jphysiol.2007.141879
17. Rittweger J, di Prampero PE, Maffulli N, Narici M V. Sprint and endurance power and ageing: an analysis of master athletic world records. *Proc R Soc B Biol Sci.* 2009;276(1657):683-689. doi:10.1098/rspb.2008.1319
18. Cheng S, Degens H, Evans M, et al. What makes a 97-year-old man cycle 5,000 km a year? *Gerontology.* 2016;62(5):508-512. doi:10.1159/000443390
19. Lazarus NR, Harridge SDR. Declining performance of master athletes: silhouettes of the trajectory of healthy human ageing? *J Physiol.* 2017;595(9):2941-2948.  
doi:10.1113/JP272443
20. Klitgaard H, Mantoni M, Schiaffino S, et al. Function, morphology and protein expression of ageing skeletal muscle: A cross-sectional study of elderly men with different training backgrounds. *Acta Physiol Scand.* 1990;140(1):41-54.  
doi:10.1111/j.1748-1716.1990.tb08974.x
21. Barnouin Y, McPhee JS, Butler-Browne G, et al. Coupling between skeletal muscle fiber size and capillarization is maintained during healthy aging. *J Cachexia*

- Sarcopenia Muscle*. 2017;8(4):647-659. doi:10.1002/jcsm.12194
22. Gilliver SF, Jones DA, Rittweger J, Degens H. Variation in the determinants of power of chemically skinned type I rat soleus muscle fibres. *J Comp Physiol A Neuroethol Sensory, Neural, Behav Physiol*. 2011;197(4):311-319. doi:10.1007/s00359-010-0613-6
  23. Larsson L, Li X, Yu F, Degens H. Age-related changes in contractile properties and expression of myosin isoforms in single skeletal muscle cells. *Muscle Nerve Suppl*. 1997;5:S74-8. <http://www.ncbi.nlm.nih.gov/pubmed/9331390>.
  24. Frontera WR, Suh D, Krivickas LS, Hughes VA, Goldstein R, Roubenoff R. Skeletal muscle fiber quality in older men and women. *Am J Physiol - Cell Physiol*. 2000;279(3):C611-C618.
  25. Šimunic B, Degens H, Rittweger J. Noninvasive Estimation of Myosin Heavy Chain Composition in Human Skeletal Muscle. *Med Sci Sport Exerc Sport Exerc*. 2011;d(February):27-30. doi:10.1249/MSS.0b013e31821522d0
  26. Šimunič B, Degens H, Završnik J, Koren K, Volmut T, Pišot R. Tensiomyographic Assessment of Muscle Contractile Properties in 9- to 14-Year Old Children. *Int J Sports Med*. 2017;38(9):659-665. doi:10.1055/s-0043-110679
  27. Zubac D, Šimunič B. Skeletal Muscle Contraction Time and Tone Decrease After 8 Weeks of Plyometric Training. *J strength Cond Res*. 2017;31(6):1610-1619. doi:10.1519/JSC.0000000000001626
  28. Maden-Wilkinson TM, McPhee JS, Jones DA, Degens H. Age-related loss of muscle mass, strength, and power and their association with mobility in recreationally-active older adults in the United Kingdom. *J Aging Phys Act*. 2015;23(3):352-360. doi:10.1123/japa.2013-0219
  29. Rodríguez-Ruiz D, García-Manso JM, Rodríguez-Matoso D, Sarmiento S, Da Silva-

- Grigoletto M, Pisot R. Effects of age and physical activity on response speed in knee flexor and extensor muscles. *Eur Rev Aging Phys Act.* 2013;10(2):127-132.  
doi:10.1007/s11556-013-0127-7
30. Šimunič B. Between-day reliability of a method for non-invasive estimation of muscle composition. *J Electromyogr Kinesiol.* 2012;22(4):527-530.  
doi:10.1016/j.jelekin.2012.04.003
31. Valenčič V. Direct measurement of the skeletal muscle tonus. *Adv Extern Control Hub Extrem.* 1990:102-108.
32. Johnson MA, Polgar J, Weightman D, Appleton D. Data on the distribution of fibre types in thirty-six human muscles. An autopsy study. *J Neurol Sci.* 1973;18(1):111-129. doi:10.1016/0022-510X(73)90023-3
33. Dahmane R, Djordjevič S, Šimunič B, Valenčič V. Spatial fiber type distribution in normal human muscle: Histochemical and tensiomyographical evaluation. *J Biomech.* 2005;38(12):2451-2459. doi:10.1016/j.jbiomech.2004.10.020
34. Simoneau J, Bouchard C. Human variation in skeletal muscle fiber-type proportion and enzyme activities. *Am J Physiol* .... 1989;257:567-572.  
doi:10.1152/AJPENDO.1989.257.4.E567
35. Staron RS, Hagerman FC, Hikida RS, et al. Fiber Type Composition of the Vastus Lateralis Muscle of Young Men and Women. *J Histochem Cytochem.* 2000;48(5):623-629. doi:10.1177/002215540004800506
36. Trappe S, Williamson D, Godard M, Porter D, Rowden G, Costill D. Effect of resistance training on single muscle fiber contractile function in older men. *J Appl Physiol.* 2000;89:143-152.
37. Trappe S, Godard M, Gallagher P, Carroll C, Rowden G, Porter D. Resistance training improves single muscle fiber contractile function in older women. *Am J Physiol Cell*
-

- Physiol.* 2001;281(2):C398-406. doi:10.1111/j.1475-097X.1981.tb00876.x
38. Erskine RM, Jones DA, Maffulli N, Williams AG, Stewart CE, Degens H. What causes in vivo muscle specific tension to increase following resistance training? *Exp Physiol.* 2011;96(2):145-155. doi:10.1113/expphysiol.2010.053975
  39. Degens H. Age-related skeletal muscle dysfunction: Causes and mechanisms. In: *Journal of Musculoskeletal Neuronal Interactions*. Vol 7. ; 2007:246-252.
  40. Larsson L, Sjödén B, Karlsson J. Histochemical and biochemical changes in human skeletal muscle with age in sedentary males, age 22–65 years. *Acta Physiol Scand.* 1978;103(1):31-39. doi:10.1111/j.1748-1716.1978.tb06187.x
  41. Gouspillou G, Sgarioto N, Kapchinsky S, et al. Increased sensitivity to mitochondrial permeability transition and myonuclear translocation of endonuclease G in atrophied muscle of physically active older humans. *FASEB J.* 2014;28(4):1621-1633. doi:10.1096/fj.13-242750
  42. Nilwik R, Snijders T, Leenders M, et al. The decline in skeletal muscle mass with aging is mainly attributed to a reduction in type II muscle fiber size. *Exp Gerontol.* 2013;48(5):492-498. doi:10.1016/j.exger.2013.02.012
  43. D'Antona G, Pellegrino MA, Adami R, et al. The effect of ageing and immobilization on structure and function of human skeletal muscle fibres. *J Physiol.* 2003;552(Pt 2):499-511. doi:10.1113/jphysiol.2003.046276
  44. Degens H, Yu F, Li X, Larsson L. Effects of age and gender on shortening velocity and myosin isoforms in single rat muscle fibres. *Acta Physiol Scand.* 1998;163(1):33-40. doi:10.1046/j.1365-201x.1998.00329.x
  45. Larsson L, Li X, Frontera WR. Effects of aging on shortening velocity and myosin isoform composition in single human skeletal muscle cells. *Am J Physiol.* 1997;272(2 Pt 1):C638--49. doi:10.1152/ajpcell.1997.272.2.C638
-

46. Ramamurthy B. Changes in myosin structure and function in response to glycation. *FASEB J.* 2001;15(13):2415-2422. doi:10.1096/fj.01-0183com
47. Nishikawa T, Edelstein D, Du XL, et al. Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. *Nature.* 2000;404(6779):787-790. doi:10.1038/35008121
48. Yamagishi S, Nakamura K, Inoue H. Therapeutic potentials of unicellular green alga *Chlorella* in advanced glycation end product (AGE)-related disorders. *Med Hypotheses.* 2005;65(5):953-955. doi:10.1016/j.mehy.2005.05.006

Accepted Manuscript

## Tables

**Table 1:** Anthropometric data of participants, grouped by age, sex and sport.

	Non-athletes		Power athletes		Endurance athletes	
	Men	Women	Men	Women	Men	Women
N	133	246	100	78	76	73
N <sub>35-44</sub>	31	28	32	26	20	27
N <sub>45-64</sub>	45	68	33	28	25	22
N <sub>65-90</sub>	57	150	35	24	31	24
Age/years	57.8±14.0	63.4±11.4	57.5±14.1	55.5±12.2	56.8±14.7	56.2±14.6
Age <sub>35-49</sub> /years	42.0±7.4	41.3±5.2	40.8±4.5	41.4±5.1	40.0±4.1	41.4±3.7
Age <sub>50-64</sub> /years	57.2±4.6	60.3±4.8	58.8±5.3	56.4±4.5	56.5±5.3	56.7±4.9
Age <sub>65-90</sub> /years	72.4±4.7	71.8±5.1	71.6±8.0	70.6±2.5	71.8±5.1	73.9±4.8
Body height/m	1.78±.07	1.62±.07	1.78±.08	1.69±.07	1.78±.07	1.66±.07
Body height <sub>35-49</sub> /m	1.81±.07	1.68±.08	1.81±.07	1.73±.06	1.81±.07	1.70±.09
Body height <sub>50-64</sub> /m	1.79±.07	1.63±.06	1.78±.05	1.67±.08	1.75±.08	1.66±.05
Body height <sub>65-90</sub> /m	1.74±.06	1.61±.06	1.73±.08	1.67±.06	1.73±.06	1.59±.04
Body mass/kg	85±12	70±12	79±13	64±6	72±12	60±9
Body mass <sub>35-49</sub> /kg	86±12	66±10	83±16	64±6	75±9	63±8
Body mass <sub>50-64</sub> /kg	87±13	70±13	79±13	63±8	72±16	59±10
Body mass <sub>65-90</sub> /kg	83±11	70±11	72±12	65±11	64±6	55±8
BMI/kg/m <sup>2</sup>	26.6±3.4	26.2±4.4	24.9±2.9	22.4±3.0	22.9±3.7	21.7±2.8
BMI <sub>35-49</sub> /kg/m <sup>2</sup>	26.1±3.4	23.6±3.3	25.5±3.6	21.3±2.5	23.1±2.9	21.9±2.9
BMI <sub>50-64</sub> /kg/m <sup>2</sup>	26.3±3.3	25.9±4.6	24.9±2.6	22.6±3.1	23.4±4.2	21.5±3.0
BMI <sub>65-90</sub> /kg/m <sup>2</sup>	27.2±3.1	27.0±4.3	24.1±2.1	23.1±3.2	21.6±1.7	21.9±2.5

*BMI: body mass index; indexes <sub>35-49</sub>, <sub>50-64</sub>, <sub>65-90</sub> represent the age range in an age group. For statistical analysis see Results.*

**Table 2:** Multiple linear regression analysis of age, sex (removed by regression), sport (Aerobic and Anaerobic), and muscle (biceps femoris and vastus lateralis) as predictors of tensiomyographic contraction time.

Predictors	Unstandardized coefficients	Standardized coefficients	P	Part r	VIF
Constant	18.098 ms	.907	<.001		
Biceps femoris	11.754 ms	.421	<.001	.453	1.371
Age	.193 ms/years	.013	<.001	.236	1.050
Anaerobic	-3.292 ms	.426	<.001	-.125	1.088
Vastus lateralis	-3.185 ms	.419	<.001	-.123	1.372
Aerobic	3.142 ms	.509	<.001	.100	1.069

*Part r: Part correlation of each predictor to contraction time when controlling for effects of other predictors; VIF: variance inflation factor.*

Accepted Manuscript



**Table 3:** Myosin heavy chain (MHC) composition of the vastus lateralis muscle for a subsample of master athletes (master athlete's data from Šimunič et al., 2011).

	Total	Power athletes	Endurance athletes
	N=17	N=13	N=4
MHC-I / %	35.6±18.4	31.7±16.3	48.1±21.5
MHC-II / %	64.5±18.4	68.3±16.3	52.0±21.5
MHC-IIa / %	44.7±14.4	45.2±12.4	43.3±22.0
MHC-IIx / %	19.7±15.1	23.1±15.0	8.7±10.3

Accepted Manuscript

## Figure captions:

**Figure 1:** Typical tensiomyographic responses of the biceps femoris of a non-athlete, power athlete and endurance athlete (left) and of a non-athlete in different age groups (right). Tc: contraction time defined as the time from 10% to 90% of the maximal displacement amplitude (Dm).

**Figure 2:** Representative SDS-PAGE gel to determine the myosin heavy chain composition of the vastus lateralis muscle from master athletes.



*Legend: Lane 1-3, 5, 7-11 contain all three myosin heavy chain isoforms (from top to bottom IIX, IIA, I) and lane 4 and 6 only contain type I and IIA myosin heavy chain isoforms.*

**Figure 3:** Pearson correlation (r) between age and contraction time (Tc) for men (left) and women (right) in each muscle and sport group.

**Figure 4:** Contraction time (Tc) differences in three skeletal muscles between athletes and non-athletes in different age groups for men (left column graphs) and women (right column graphs).

*Legend: a, b, c...significantly different from non-athletes at  $P < .05$ ;  $P < .01$ , and  $P < .001$ , respectively; i, j, k... significantly different from previous age group at  $P < .05$ ;  $P < .01$ , and  $P < .001$ , respectively; G. ... Gastrocnemius.*

Figure 1.



cript


AC

*Figure 2.*




Accepted Manuscript

Figure 3.



cript

Figure 4.



cript

Accepted