



**Manchester
Metropolitan
University**

Dean, AP and Hartley, A and McIntosh, OA and Smith, A and Feord, HK and Holmberg, NH and King, T and Yardley, E and White, KN and Pittman, JK (2018) *Metabolic adaptation of a Chlamydomonas acidophila strain isolated from acid mine drainage ponds with low eukaryotic diversity*. Science of the Total Environment, 647. pp. 75-87. ISSN 0048-9697

Downloaded from: <http://e-space.mmu.ac.uk/621624/>

Publisher: Elsevier

DOI: <https://doi.org/10.1016/j.scitotenv.2018.07.445>

Usage rights: Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Please cite the published version

<https://e-space.mmu.ac.uk>

1 **Metabolic adaptation of a *Chlamydomonas acidophila* strain isolated from acid**
2 **mine drainage ponds with low eukaryotic diversity**

3

4 Andrew P. Dean^a, Antoni Hartley^d, Owen A. McIntosh^d, Alyssa Smith^{b,1}, Helen Feord^{b,2},
5 Nicolas H. Holmberg^b, Thomas King^b, Ellen Yardley^c, Keith N. White^{b,d}, and Jon K.
6 Pittman^{b,d,*}

7

8 ^aSchool of Science and the Environment, Manchester Metropolitan University, Oxford Road,
9 Manchester, M1 5GD, UK; ^bFaculty of Life Sciences, University of Manchester, Michael
10 Smith Building, Oxford Road, Manchester M13 9PT, UK; ^cDepartment of Geography,
11 University of Sheffield, Sheffield S10 2TN, UK; ^dSchool of Earth and Environmental
12 Sciences, Faculty of Science and Engineering, University of Manchester, Michael Smith
13 Building, Oxford Road, Manchester M13 9PT, UK

14

15 *Corresponding author.

16 Email address: jon.pittman@manchester.ac.uk (J.K. Pittman)

17

18 ¹Present address: Department of Chemistry, University of Cambridge, Cambridge, UK

19 ²Present address: School of Biological Sciences, University of Edinburgh, Edinburgh, UK

20

21 **Abstract**

22 The diversity and biological characteristics of eukaryotic communities within acid mine
23 drainage (AMD) sites is less well studied than for prokaryotic communities. Furthermore, for
24 many eukaryotic extremophiles the potential mechanisms of adaptation are unclear. This
25 study describes an evaluation of eight highly acidic (pH 1.6 – 3.1) and one moderately acidic
26 (pH 5.6) metal-rich acid mine drainage ponds at a disused copper mine. The severity of
27 AMD pollution on eukaryote biodiversity was examined, and while the most species-rich site
28 was less acidic, biodiversity did not only correlate with pH but also with the concentration of
29 dissolved and particulate metals. Acid-tolerant microalgae were present in all ponds,
30 including the species *Chlamydomonas acidophila*, abundance of which was high in one very
31 metal-rich and highly acidic (pH 1.6) pond, which had a particularly high PO₄-P
32 concentration. The *C. acidophila* strain named PM01 had a broad-range pH tolerance and
33 tolerance to high concentrations of Cd, Cu and Zn, with bioaccumulation of these metals
34 within the cell. Comparison of metal tolerance between the isolated strain and other *C.*
35 *acidophila* strains previously isolated from different acidic environments found that the new
36 strain exhibited much higher Cu tolerance, suggesting adaptation by *C. acidophila* PM01 to
37 excess Cu. An analysis of the metabolic profile of the strains in response to increasing
38 concentrations of Cu suggests that this tolerance by PM01 is in part due to metabolic
39 adaptation and changes in protein content and secondary structure.

40

41 **Keywords:** *Chlamydomonas acidophila*, acid tolerance, metal tolerance, acid mine
42 drainage, bioremediation, copper, zinc, cadmium

43

44 **1. Introduction**

45 Many freshwater bodies worldwide are highly acidic either due to natural causes or
46 anthropogenic activities such as mining (Schultze, 2013; Smucker et al., 2014). Acid mine
47 drainage (AMD) due principally to pyrite oxidation, is the cause of significant acidity in lakes
48 and ponds situated in areas impacted by mining, and in rivers receiving mine water

49 discharge (Johnson, 2003; Nordstrom, 2000). Because decreasing pH causes increased
50 solubility of metals, AMD results in high concentrations of dissolved Fe, S and various trace
51 metals such as Cu, Cd and Zn in the contaminated waters. Concentrations of nutrients,
52 especially inorganic phosphate (PO₄-P), are also frequently very low (Nixdorf et al., 1998).
53 The combination of toxic metals and nutrient limitation limits biodiversity and can cause
54 significant ecosystem damage (Deneke, 2000; Smucker et al., 2014). Evaluation of the
55 biological impacts of AMD allows quantification of pollution damage, allows understanding of
56 fundamental processes of adaptation and can identify AMD-tolerant species that have
57 biotechnological applications, such as bioremediation (Ñancuqueo and Johnson, 2011; Yun
58 et al., 2014).

59 While prokaryotes in AMD environments has been extensively studied and reviewed
60 (Johnson and Hallberg, 2003; Mendez-Garcia et al., 2015), there is still limited knowledge
61 regarding the presence and roles of eukaryotes in these aquatic environments (Aguilera et
62 al., 2006; Baker et al., 2004; Nixdorf et al., 1998). Photosynthetic microorganisms are found
63 in many AMD ecosystems; however, the biodiversity of phytoplankton in such waters is
64 severely limited and dominated by just a few acid-tolerant genera, such as *Chlamydomonas*,
65 *Dunaliella*, *Euglena* and *Ochromonas* (Aguilera et al., 2006; Hargreaves et al., 1975;
66 Ñancuqueo and Johnson, 2012; Nixdorf et al., 1998; Pedrozo et al., 2001). Despite being
67 able to tolerate the highly acidic and metal-rich conditions, productivity of these extremophile
68 microalgae is often limited by low inorganic carbon and nutrient availability in acidic waters
69 (Beamud et al., 2007; Spijkerman et al., 2007b). A fairly broad diversity of heterotrophic fungi
70 and protists has also been observed in acidic waters (Baker et al., 2004; Das et al., 2009),
71 while the diversity and abundance of zooplankton is typically very low as most species are
72 unable to tolerate these environments (Deneke, 2000).

73 The high concentrations of dissolved metals in AMD can cause toxicity to
74 microorganisms through a wide variety of mechanisms, some of which are shared between
75 metals and across different organisms, such as competition with essential metals, direct
76 interactions with proteins and other molecules within the cell, and induction of oxidative

77 stress (Sharma and Dietz, 2009). Metals such as Cu are particularly efficient at inducing the
78 formation of reactive oxygen species (ROS) in contrast to non-redox active metals such as
79 Zn and Cd (Valko et al., 2005). In most photosynthetic organisms, excess Cu has many
80 detrimental effects with the photosynthetic apparatus, including direct inhibition of
81 photosynthetic activity and degradation of chloroplast structures (Bernal et al., 2006; Küpper
82 et al., 2003). Furthermore, non-extremophile microalgae exposed to high Cu conditions
83 exhibit high concentrations of ROS and subsequent ROS-induced damage including lipid
84 membrane peroxidation (Jamers et al., 2013; Jiang et al., 2016; Sabatini et al., 2009).

85 The adaptive mechanisms by which eukaryotic microorganisms including
86 extremophile microalgae can survive in acid and metal rich conditions are still poorly
87 researched but potential insights into these mechanisms are increasing. For example,
88 proteomic approaches have indicated the importance of metal and acidity tolerance proteins,
89 such as molecular chaperones of the Heat Shock Protein family (Cid et al., 2010; Gerloff-
90 Elias et al., 2006). Likewise, genome sequencing and transcriptomics studies are beginning
91 to identify the array of genes that might explain extremophile functional characteristics, some
92 of which may have been obtained by horizontal gene transfer from bacteria. Genome
93 sequences of the acidophiles *Chlamydomonas eustigma* (Hirooka et al., 2017) and *Galdieria*
94 *sulphuraria* (Schönknecht et al., 2013) have recently been determined. Furthermore,
95 transcriptomic approaches are beginning to provide insight into the molecular mechanisms
96 of *Chlamydomonas acidophila* tolerance in response to Cd and Cu exposure (Olsson et al.,
97 2015; Puente-Sánchez et al., 2018), and *Dunaliella acidophila* in response to Cd (Olsson et
98 al., 2017; Puente-Sánchez et al., 2016), although further experimental analyses of these
99 transcriptomic datasets are needed.

100 AMD tolerant biota might have potential for bioremediation, with biological-based
101 processes potentially more cost effective and sustainable than chemical based methods
102 such as anoxic limestone drains and chemical addition (Geller and Schultze, 2013; Hedin et
103 al., 2010; Johnson and Hallberg, 2005). Bioremediation methods can include utilisation of
104 bacterial SO₄ reduction and neutralisation (Neculita et al., 2007) or aerobic wetlands that can

105 oxidise and precipitate dissolved metals (Dean et al., 2013). However, eukaryotic algae that
106 can tolerate AMD conditions may be an alternative bioremediation agent (Abinandan et al.,
107 2018; Das et al., 2009). Novel extremophile algal strains that show high acid and metal
108 tolerance, and metal bioaccumulation traits are therefore needed for such applications. In
109 addition, extremophile algae may have other biotechnological applications, such as a source
110 of novel high-value chemicals including nutritional vitamins and anti-oxidants, food additives,
111 and biofuels (Varshney et al., 2015).

112 The aim of this study was to identify eukaryotes, especially extremophile microalgae,
113 in a series of standing waters affected by AMD with the intention to characterise a strain of
114 microalgae for evidence of AMD adaptation. Following a screen of eukaryotic biota within
115 nine Cu-rich AMD ponds, an extremophile chlorophyte microalgal strain identified as *C.*
116 *acidophila* was examined in detail due to its abundance and ubiquity across the site and its
117 high tolerance to acidity and dissolved metal concentrations, especially to Cu.

118

119 **2. Materials and Methods**

120

121 *2.1. Study site*

122 The site for this study is Parys Mountain, a disused Cu mine, in Anglesey North Wales, UK.
123 The site has been mined for Cu from the Bronze Age, until mining activities ceased in the
124 early 1900s (Dean et al., 2013). The area consists of large amounts of exposed spoil, with
125 large pits and depressions that have filled with rainwater, and now retain large amounts of
126 metal-rich and acidic water (Fig. 1). In addition, precipitation ponds and lagoons were
127 constructed at the base of Parys Mountain, which were built in order to extract metals from
128 the water as part of the mining process, and also contain large volumes of AMD polluted
129 water (Younger and Potter, 2012). All the ponds are situated at close proximity within a
130 similar geology with the rocks naturally rich in Cu, Pb and Zn. It is the only known example of
131 Kuroko type volcanogenic massive sulfides in the UK, though the geology has been

132 disturbed by many millennia of underground and surface mining activities (Younger and
133 Potter, 2012).

134 Of the various mining ponds and lagoons at the Parys Mountain site, nine ponds
135 were examined (Fig. 1). Ponds 1 – 4 are located at an elevated position on the spoil outcrop,
136 with one of these (Pond 4) on the side of a steep incline of the now drained large opencast
137 (Fig. 1). The remaining five larger ponds (Ponds 5 – 9) are the precipitation ponds and
138 lagoons at the base of the Parys Mountain outcrop, each adjacent to agricultural land. Ponds
139 1 – 4 and 9 are shallow and less than 1 m depth and subject to rapid variation in depth due
140 to seasonal evaporation and rainfall. Ponds 5 – 8 are deeper and are typically 2 m in depth.
141 All ponds showed little variation in depth across each pond. Ponds 1, 2 and 4 had no
142 vegetation in or surrounding them, whereas the other ponds had surrounding vegetation and
143 marginal wetland plants.

144

145 *2.2. Field site sampling*

146 Sampling at the nine AMD ponds was carried out in 2013 to 2015, including a spring
147 (February and March), summer (June) and autumn (October) sampling regime in 2015.
148 Water chemistry samples were taken in triplicate at each pond on each sampling occasion
149 and were taken at approximately 15 cm depth 1 – 2 m from the edge of the pond. Water pH,
150 conductivity, temperature and dissolved oxygen were measured using a YSI 556 probe
151 (Xylem Analytics). For analysis of dissolved water chemistry metals 250 mL of pond water
152 was filtered through a 0.45 µm cellulose acetate filter, as described previously (August et al.,
153 2002; Boulton et al., 1994), and a 50 mL volume was retained for the analysis of dissolved
154 nutrients (PO₃-P, NO₃-N, NH₄-N). This dissolved fraction will also include some colloidal
155 metals (Florence et al., 2016). A further 50 mL volume was acidified to 1% (v/v) nitric acid
156 final concentration for the analysis of dissolved metals (Al, As, Cd, Cu, Fe, Mn, Pb, S, Zn).
157 The filter paper was retained, and used for the determination of particulate metals.

158 For the analysis of algae 250 mL of unfiltered water was collected and aliquots were
159 preserved with Lugol's iodine for enumeration of algae cells and further unfiltered water

160 sample was taken in a sterile container for the isolation and identification of algae. Algal
161 samples were taken from all ponds, including samples taken from where algal biofilms were
162 observed (Ponds 5 and 6). For chlorophyll-a measurement, 250 mL of pond water was
163 filtered onto a GF/C filter paper.

164 Sediment samples were also taken to a depth of approximately 2 cm depth for
165 determination of acid-extractable metals. For invertebrate sampling, sediment samples were
166 taken from the littoral, approximately 1 m from the pond edge and 20 – 50 cm depth
167 depending on the pond and sieved to remove organisms. This was followed by a 3 min
168 sweep using a hand-held net with a mesh size of 1 mm, and a 3 min examination under
169 large stones. Ethanol (70%) was added to preserve the invertebrate biota for identification to
170 family level using standard keys (Greenhaigh and Ovenden, 2007; Quigley, 1977).

171

172 *2.3 Nutrient, chlorophyll and metal analysis*

173 Dissolved $\text{PO}_3\text{-P}$, $\text{NO}_3\text{-N}$, and $\text{NH}_4\text{-N}$ were measured from the 0.45 μm filtered water
174 samples using a Skalar Sans Plus autoanalyser. Chlorophyll-a concentrations were
175 determined by absorbance spectroscopy following extraction in 96% (v/v) ethanol, as
176 described previously (Dean et al., 2010). Dissolved metals were determined by inductively
177 coupled plasma atomic emission spectroscopy (ICP-AES) using a Perkin-Elmer Optima
178 5300, exactly as described previously (Dean et al., 2013). Acid extractable sediment metals
179 and suspended particulate metals were determined by acid digestion of 0.1 g of 250 μm
180 sieved dried sediments and digestion of the pond water filter papers, respectively.

181 Sediments and filter papers were digested in 5 mL ultrapure-grade nitric acid at 70°C for 4 h,
182 diluted to 2% (v/v) nitric acid, and the metals measured by ICP-AES. Certified Reference
183 Standard TM25.5 was used for all ICP-AES analyses. All samples were calibrated using a
184 matrix-matched serial dilution of Specpure multi-element plasma standard solution 4 (Alfa
185 Aesar) set by linear regression. Only results with a relative standard deviation < 20% were
186 considered.

187

188 *2.4. Microalgae cultivation and analysis*

189 Microalgae was visually identified to genus level and enumerated from the Lugol's iodine
190 preserved water samples by light microscopy using a Sedgewick-Rafter cell counting slide
191 and a morphological taxonomy key (John et al., 2002). Isolation of individual algae strains
192 was carried out by incubating serial dilutions of water samples on modified acid medium
193 (MAM) agar plates at 22 °C with a 16-h light:8-h dark light regime and a photon flux of
194 approximately 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$. MAM is a defined inorganic medium developed previously
195 (Olaveson and Stokes, 1989), and used here as described by the Canadian Phycological
196 Culture Centre, University of Waterloo, Canada containing 0.5 g L⁻¹ (NH₄)₂SO₄, 0.01 g L⁻¹
197 CaCl₂·2H₂O, 0.5 g L⁻¹ MgSO₄·7H₂O, 0.3 g L⁻¹ KH₂PO₄, 0.03 g L⁻¹ NaCl, 0.01 g L⁻¹
198 Na₂EDTA·2H₂O, 4.98 mg L⁻¹ FeSO₄·7H₂O, 2.86 mg L⁻¹ H₃BO₃, 1.81 mg L⁻¹ MnCl₂·4H₂O, 0.22
199 mg L⁻¹ ZnSO₄·7H₂O, 0.39 mg L⁻¹ NaMoO₄·2H₂O, 79 $\mu\text{g L}^{-1}$ CuSO₄·5H₂O, 49.4 $\mu\text{g L}^{-1}$
200 Co(NO₃)₂·6H₂O, 1 $\mu\text{g L}^{-1}$ vitamin B12, 1 $\mu\text{g L}^{-1}$ biotin, and 0.2 mg L⁻¹ thiamine-HCl, adjusted
201 to pH 3.0. Individual colonies were then extracted and grown in liquid MAM.

202 For identification of isolates using 18S rRNA gene amplification, DNA was extracted
203 using an UltraClean Tissue & Cells DNA isolation kit (Mo Bio) and 18S rRNA gene amplicon
204 sequences amplified using universal primers EukF and EukR (DeLong, 1992). DNA
205 amplicons were purified (Qiagen PCR Purification kit) before sequencing (to give ~1 kb
206 sequence reads) by GATC-Biotech, with subsequent sequence analysis performed by
207 BLAST, using the NCBI GenBank database (Table S1). The species identification of putative
208 *C. acidophila* strain was further confirmed by partial length 18S rDNA gene amplicon
209 sequencing, using PCR primers (18SFOR 5'-WAC CTG GTT GAT CCT GCC AGT-3', and
210 18SREV 5'-GAT CCT TCY GCA GGT TCA CCT AC-3') and PCR conditions as described
211 (Huss et al., 1999), and sequenced as described above but with sequence reads of ~1.7 kb
212 size. Phylogenetic analysis was then performed essentially as described previously
213 (Osundeko et al., 2013). 18S rRNA nucleotide sequences of selected unicellular Chlorophyta
214 microalgae of the *Chlamydomonas moewusii* clade were obtained from GenBank and
215 sequences were aligned using ClustalW. The phylogenetic tree was generated using the

216 maximum likelihood method using RAxML-GUI and the GTR-GAMMA model (Stamatakis,
217 2006). Confidence in the tree was assessed using the thorough bootstrap method by
218 performing 10 runs of 100 replications.

219 An isolate of *C. acidophila* from Pond 1 (named PM01) was cultivated to quantify its
220 tolerance to high metal concentrations and pH ranges alongside two strains of *C. acidophila*
221 previously isolated from an acidic pond near Fratikovy Lazne, Czech Republic (CCAP
222 11/136) (Fott and McCarthy, 1964), and an acidic mining Lake 111 in eastern Germany
223 (CCAP 11/137) (Gerloff-Elias et al., 2005). All cultures were maintained in MAM pH 3.0.
224 Cultures were serially inoculated in fresh media and were grown in batch culture conditions
225 on an orbital shaker at 120 rpm at 22 °C with a 16-h light:8-h dark light regime and a photon
226 flux of approximately 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$. For metal exposure treatments, various
227 concentrations of metals as chloride salts were added to liquid MAM pH 3.0 or solid MAM pH
228 3.0 agar plates. For pH range treatments, MAM was adjusted and buffered to the desired pH
229 (from pH 1.0 to 7.0) as described previously (Gerloff-Elias et al., 2006). Essentially the
230 medium was buffered through the presence of Fe in the medium for pH 1.0 to 3.0, with 10
231 mM citric acid for pH 4.0 to 5.0, and with 10 mM HEPES for pH 6.0 to 7.0. The pH could be
232 maintained to within ~0.5 pH unit during the growth period. Starting cell densities were
233 normalised by optical density measurement at 680 nm ($\text{OD}_{680\text{nm}}$). Algae growth was
234 determined by cell number, and in some instances by $\text{OD}_{680\text{nm}}$, total chlorophyll or growth
235 rate measurement, exactly as described previously (Osundeko et al., 2013). Algae samples
236 were prepared for metal content measurement by ICP-AES, as described previously
237 (Webster et al., 2011). Cells were EDTA washed to remove externally bound metals, as
238 validated elsewhere for Cd, Cu, Pb and Zn (Hassler et al., 2004). Cells were centrifuged for
239 10 min at 3000 g, followed by resuspension of the cell pellet in 10 mL of 1 mM EDTA for 5
240 min, then re-centrifuged and washed with a 15 mL volume of Milli-Q water. After further
241 centrifugation, cell pellets were oven-dried at 60 °C for 24 h and then digested in 0.5 mL of
242 ultrapure concentrated nitric acid at 70 °C for 3 h. Samples were diluted in Milli-Q water to
243 2% (v/v) concentration of acid and analysed by ICP-AES as described above. Cells of *C.*

244 *acidophila* were measured (width and length) using an eyepiece graticule, and cell volume
245 was calculated according the prolate spheroid formula (Hillebrand et al., 1999). The cell
246 volume measurements were used to calculate the internal (cellular) concentration of metal
247 (determined from EDTA-washed cells) on a volume basis. This value was then divided by
248 the external (MAM or pond water) concentration of metal in order to calculate concentration
249 factor (K_{conc}) values.

250

251 *2.5. Fourier transform-infrared (FT-IR) spectroscopy*

252 Each algal strain was grown in liquid MAM (pH 3.0), with Cu concentrations of 0, 6.5, 13 and
253 130 mg L⁻¹ for 14 days, at which point cultures had an OD_{750nm} of between 0.3 and 0.5.
254 Cultures were normalised to an OD_{750nm} of 0.3 and 10 mL was centrifuged at 1800 *g* for 5
255 min. The supernatant was removed and the cells washed in 1.5 mL of 0.9% (v/v) NaCl, then
256 centrifuged and washed again before final resuspension of the cell pellet in 1 mL of 0.9%
257 (v/v) NaCl before 20 µL of this suspension was deposited onto a 96-well silicon microplate.
258 The samples were then oven dried at 40°C for 1 h, and an additional 20 µL sample was
259 added and the samples once again dried at 40°C for 1 h. The plate was placed in a HTS-XT
260 high-throughput microplate extension and FT-IR spectra collected using an Equinox 55 FT-
261 IR spectrometer (Bruker Corporation), equipped with a deuterated triglycerine sulphate
262 detector. Spectra were collected over the wavenumber range 4000-600 cm⁻¹. Each sample
263 was analysed as nine technical replicates. Spectra were pre-processed using extended
264 multiplicative signal correction (Martens and Stark, 1991) prior to multivariate analysis. Band
265 assignments were determined as described previously (Driver et al., 2015).

266

267 *2.6. Modelling and statistical analysis*

268 Metal speciation modelling was performed using Visual Minteq version 3.0 (Gustafsson,
269 2010). All data was statistically analyzed by one-way ANOVA using Tukey post-hoc test
270 performed using Prism v.6.04 (GraphPad). Principal component analysis (PCA) of
271 environmental data was performed using PRIMER v.6 (Primer-E) and plotted using XLSTAT

272 (Addinsoft), while PCA of FT-IR spectra was performed using MATLAB version R2016a. All
273 environmental data (except for pH values) were natural log transformed for PCA and linear
274 regression analysis.

275

276 **3. Results**

277

278 *3.1. Chemical characteristics of extremely acidic metal-rich mining ponds*

279 The nine ponds were situated in close proximity (Fig. 1) but differed in water chemistry (Fig.
280 2). In all ponds, concentrations of suspended particulate metals were present at much lower
281 concentrations compared to dissolved metals (Fig. S1). None of the metal or nutrient
282 concentrations in the ponds differed significantly across the spring, summer and autumn
283 samples. Ponds 1, 2 and 4 were situated on top of the Parys Mountain site in depressions
284 of the spoil waste. These ponds were characterised by very low pH (< 2.0) and high
285 conductivity values (Fig. 2A). Pond 1 in particular was highly acidic (mean pH of 1.6) and
286 had a very high conductivity (mean 10.03 mS), which is predominantly controlled by the high
287 dissolved S concentration, as well as high concentrations of dissolved Fe, Al, Cd, Cu and Zn
288 (Fig. 2B, C). In addition, Pond 1 also had a very high PO₄-P concentration (Fig. 2D) and the
289 highest dissolved As concentration (Fig. S1). Ponds 2 and 4 also had high dissolved metal
290 and S concentrations but differed from Pond 1 due to lower conductivity (~3 mS) and much
291 lower PO₄-P. Pond 3 was also situated on the top of the mine site, however, this pond stood
292 out as the least polluted, with a weakly acidic pH (mean pH 5.3), and low conductivity and
293 dissolved metals and S (Fig. 2A; Fig. S1). The precipitation ponds and lagoons (Ponds 5 –
294 9) were also highly acidic but had lower conductivity due to much lower dissolved metal
295 concentrations, with the exception of dissolved Mn (Fig. S1). In most ponds, ammonium
296 (NH₄-N) was the dominant form of N rather than nitrate (NO₃-N), as is expected in acidic
297 waters where nitrification rates are low (Baffico et al., 2004). Pond 6 had unusually high
298 NO₃-N (Fig. 2D) possibly because the pond is directly adjacent to fertilised farmland.
299 Chlorophyll-a was detectable in all ponds, indicating the presence of photosynthetic

300 microorganisms, but Pond 1 also differed from the other ponds with respect to having
301 significantly higher concentration of chlorophyll-a, which increased substantially during the
302 year as the water temperature and insolation increased (Fig. 2E).

303 The differences in water chemistry between the ponds situated within the mine spoil
304 waste (Ponds 1, 2 and 4), the lagoons and precipitation ponds (Ponds 5 – 9), and the less
305 polluted Pond 3, are reflected in the PCA plot (Fig. 3). In particular, Pond 1 and Pond 3 were
306 distinguished from each other and from the other ponds through difference in pH.
307 Furthermore, the high sediment Al and Mn concentrations explained the differentiation of
308 Pond 3, likely a consequence of the high pH of this water resulting in the precipitation of
309 dissolved Al. Pond 1 was situated within the PC space particularly on the basis of high
310 chlorophyll-a, PO₄-P and dissolved As concentration. It is therefore apparent that many of
311 these ponds are highly toxic environments, particularly Pond 1 with its very high acidity and
312 high concentration of toxic dissolved trace metals including As, Al, Cd, and Cu.

313

314 3.2. *Taxa diversity*

315 Overall taxa diversity, as determined by number of different taxonomic families, was lowest
316 in the highly acidic, high conductivity Pond 1 (with two taxa) and also low in the Ponds 2 and
317 4, which are also situated among open mine spoil. The highest number of taxa were found in
318 the low conductivity, mildly acidic (pH 5.3) Pond 3, with 15 taxa, while the
319 lagoons/precipitation ponds had intermediate taxa diversity (7-10 families). Linear regression
320 analyses of the relationships between taxa diversity and water chemistry (Fig. S2) showed a
321 significant positive correlation between increasing pH values and increasing taxa number (R^2
322 = 0.59; $p = 0.03$) but also a very strong negative correlation between decreasing conductivity
323 and increasing taxa number ($R^2 = 0.85$; $p = 0.001$), indicative of the dissolved metal
324 concentration. In particular, there was significant correlation between dissolved As, Cd, Cu,
325 Fe and S and taxa number, as well as between particulate Fe and S with taxa number, but
326 no significant correlation on the basis of sediment metal concentration (Fig. S2).

327 Invertebrates were observed in all ponds, though only Chironomidae was identified in
328 all, although there were only one or two individuals recorded from Pond 2 and 4 (Fig. 5A). In
329 Ponds 1 and 4, this was the only invertebrate taxa present, indicating that the conditions in
330 these ponds are not suited for a diverse invertebrate community, but are able to support a
331 few species that have adapted to the extreme AMD conditions. Other invertebrate taxa that
332 were abundant are Corixidae, present in five of the ponds, and Sialidae, present in four of
333 the ponds, but both absent in the three most acidic ponds (Fig. 4). The observed biota in
334 Pond 3 included 14 invertebrate families, including many that are pollution-sensitive. Ponds
335 5 – 9 contained between 5 and 7 invertebrate taxa. All ponds were devoid of macrophytes
336 within the open water.

337 A number of distinct eukaryotic microorganisms were isolated from the open water
338 including microalgal, fungal and protozoan species (Table S1). The highest eukaryotic
339 diversity was in Ponds 2, 5 and 6 (Fig. 4). In Ponds 5 and 6 and their adjacent channels
340 much of the microorganisms were observed to be associated in biofilms. These included a
341 strain which showed 99% identity based on the 18S rRNA sequence to *Euglena mutabilis*
342 (Table S1), a well-known acid-tolerant species that has been previously observed in AMD
343 environments, including coal mine waste sites (Brake et al., 2001), and including at the adit
344 draining from the Parys Mountain mine (Ñancuqueo and Johnson, 2012). In addition, a non-
345 motile chlorophyte (likely to be *Koliella corcontica*; 96% identity) that has been previously
346 observed in mildly acidic lakes (Vrba et al., 2003), and a diatom (likely to be *Eunotia*
347 *naegelii*; 98% identity) was found associated with biofilm adjacent to Pond 6. However,
348 diatoms were not found in open water in any of the Parys Mountain ponds. Ponds 1, 3 and 7
349 had the lowest number of eukaryotic microorganisms, with just one chlorophyte algae taxa
350 identified in each pond (Fig. 4). There was substantial heterogeneity between the ponds,
351 with all microbial taxa present in just one pond sample with the exception of the *Euglena* sp.
352 present in two ponds, and a *Chlamydomonas* sp. identified in all ponds (100% abundance).
353 Sequencing of the ~1 kb 18S rRNA gene amplicon from the *Chlamydomonas* sp. showed
354 highest sequence identity (99%) to a strain annotated as *C. acidophila* (CCAP 11/134)

355 (Table S1). Quantification of cell density of this strain found highest density in Pond 1 (Fig.
356 5A), which correlated significantly with the highest pond PO₄-P concentration ($R^2 = 0.60$; $p =$
357 0.02), but cell density was also lowest in the most alkaline pond water where there was the
358 lowest dissolved metal concentrations, likely explaining the positive correlation between cell
359 density and conductivity or dissolved metal concentrations, such as Cu (Fig. 5B).

360

361 3.3. Phylogeny of a *C. acidophila* strain

362 *C. acidophila* was found across the Parys Mountain site in all nine ponds and was highly
363 abundant in the most acidic and metal rich Pond 1. Therefore this strain was studied further
364 in more detail. The isolate of *C. acidophila* from Pond 1 (named PM01) was examined by
365 longer (1.7 kb) 18S rRNA gene read sequence and phylogenetic analysis (Fig. 6). PM01 18S
366 rRNA sequence was identical apart from one nucleotide within this 1.7 kb region to
367 sequences from three strains annotated as *C. acidophila* (Gerloff-Elias et al., 2005): CCAP
368 11/134 (an isolate from Argentina), CCAP 11/136 (an isolate from Czech Republic) and
369 CCAP 11/137 (an isolate from Germany). PM01 was also identical within this 18S region to
370 an unidentified strain (Rt1n1) originally isolated from the acidic Rio Tinto river in Spain
371 (Amaral Zettler et al., 2002). The CCAP 11/136 strain (Fott and McCarthy, 1964), previously
372 regarded as an authentic strain of *C. acidophila* (Gerloff-Elias et al., 2005), is also deposited
373 in a different culture collection as strain UTCC 354 (Pollio et al., 2005), but this has distinct
374 18S rDNA sequence (Fig. 6), indicating that CCAP 11/136 and UTCC 354 are not in fact
375 identical. It was previously argued that the UTCC 354 strain assigned as *C. acidophila* may
376 be more appropriately assigned as *Chlamydomonas pitschmannii*, another highly
377 acidotolerant species (Pollio et al., 2005). Other strains recorded as *C. acidophila* including
378 UTCC 121 (an isolate from Canada) (Twiss, 1990) and OU 030/a (an isolate from Japan)
379 (Nishikawa and Tominaga, 2001), were also distinct from PM01 and CCAP 11/136 but
380 grouped more closely to the UTCC 354 strain and *C. pitschmannii* (Fig. 6). Although strain
381 OU 030/a has also been previously regarded as an authentic strain of *C. acidophila*
382 (Nishikawa and Tominaga, 2001; Pollio et al., 2005), here we refer to the CCAP 11/136 and

383 11/137 strains as *C. acidophila* species, in line with previous analysis (Gerloff-Elias et al.,
384 2005), and thus the PM01 strain is further referred to as a strain of this species.

385

386 3.4. AMD tolerance by *C. acidophila* PM01

387 To examine the relationship between pH and *C. acidophila* further, strain PM01 from Pond 1
388 was grown in MAM artificial pond water at a range of pH values from pH 1.0 to 7.0. PM01
389 exhibited a very broad pH tolerance range, with optimal growth at pH 3.0 – 5.0 (mean growth
390 rate ranging between 0.158 – 0.175 d⁻¹; no significant difference between treatments; p >
391 0.05), and with the highest cell density after 25 d obtained in pH 3.0 conditions. Strong
392 growth was still observed in pH 7.0 (mean growth rate 0.153 d⁻¹) and pH 2.0 (mean growth
393 rate 0.157 d⁻¹) conditions, but with a significant reduction (p < 0.05) in growth after 25 d in pH
394 2.0 and pH 7.0 by 25% and 37%, respectively compared to pH 3.0 MAM. and could grow at
395 pH 1.0 after an extended lag phase of 13 d (growth rate 0.018 d⁻¹), but with a significant
396 reduction in growth by 73% after 25 days compared to pH 3.0 MAM. The pH characteristics
397 of the ponds may explain in part the microalgae biodiversity and cell density profiles, and as
398 described above, there was a significant negative correlation ($R^2 = 0.57$; p = 0.02) between
399 pH and *C. acidophila* cell density (Fig. 5B).

400 Samples of PM01 taken directly from Pond 1 showed high concentration of absorbed
401 and internalised metals, as determined by measurement of EDTA-washed cells, to remove
402 externally cell wall-bound metals. There was a high concentration of Cu and Zn accumulated
403 almost entirely within the cell (no significant difference between EDTA washed versus
404 unwashed cells) (Table 1). Relative to mean Pond 1 dissolved Cu concentration of 58.6 mg
405 L⁻¹, the internal cellular Cu concentration in pond 1 cells was 84.3 fg cell⁻¹ (equivalent to
406 185.5 mg L⁻¹). The mean concentration of dissolved Zn in Pond 1 was 37.9 mg L⁻¹, while the
407 internal cellular Zn concentration was 115.0 fg cell⁻¹ (253.0 mg L⁻¹). This gives concentration
408 factor (K_{conc}) values of 3.2 for Cu and 6.7 for Zn. Substantial bioconcentration was also
409 observed for Pb and Cd. Relative to a mean Pb concentration in Pond 1 of 0.8 mg L⁻¹, the
410 accumulation and bioconcentration of Pb by the strain was particularly high, with a cellular

411 concentration of $146.3 \text{ fg cell}^{-1}$ (321.8 mg L^{-1}) giving a K_{conc} value of 421.8. Cd was also
412 accumulated almost entirely within the cell to a concentration of 2.8 fg cell^{-1} (6.2 mg L^{-1}), and
413 relative to pond water concentration of 0.3 mg L^{-1} gives a K_{conc} value of 24.5. In contrast,
414 only 56% of accumulated Fe was taken up into the cell (Fig. S3).

415 Metal tolerance and accumulation by *C. acidophila* PM01 was further examined in an
416 artificial growth medium to assess the tolerance range of three of the trace metals found
417 within the Parys Mountain ponds. PM01 was grown in increasing concentrations of Cd, Cu
418 and Zn in MAM at pH 3.0. For cells grown under controlled conditions, the maximum cell
419 density achieved and the total chlorophyll concentration per cell (as a measure of cell
420 physiological status) was higher than that observed for the cells analysed *in situ* (in Pond 1)
421 (Table 1). The PM01 strain displayed tolerance to high concentrations of these three metals.
422 To allow comparison with Pond 1 water concentrations, the free ionic Cd^{2+} , Cu^{2+} and Zn^{2+}
423 concentrations were calculated using the Visual Minteq speciation model. PM01 could
424 tolerate up to $2.6 \text{ mg L}^{-1} \text{ Cd}^{2+}$, which was 16-times higher than present in Pond 1 water, with
425 no significant inhibition of growth rate, maximum cell density or chlorophyll content when
426 compared to no added Cd^{2+} (Table 1). However, at a concentration of $6.0 \text{ mg L}^{-1} \text{ Cd}^{2+}$ all
427 three parameters (growth rate, cell density, chlorophyll content) were significantly reduced (p
428 < 0.05). PM01 was highly tolerant to Cu, and none of the concentrations up to 78.3 mg L^{-1}
429 Cu^{2+} treatment significantly inhibited growth or chlorophyll-a concentration, the apparent
430 reduction in cell density was not significant. Substantial Zn tolerance was also observed,
431 with no significant inhibition to growth rate at $855.4 \text{ mg L}^{-1} \text{ Zn}^{2+}$, which was nearly 30-times
432 higher than the Zn^{2+} concentration in Pond 1, but at $1760.8 \text{ mg L}^{-1} \text{ Zn}^{2+}$ there was a
433 significant ($p < 0.05$) reduction in maximum cell density, although growth rate and chlorophyll
434 concentration was not significantly inhibited. The strain was still growing in 3002.8 mg L^{-1}
435 Zn^{2+} despite cell density being inhibited by 94%, with also a significant ($p < 0.05$) reduction in
436 growth rate and chlorophyll concentration (Table 1).

437 Accumulation of Cd, Cu and Zn was also quantified in the metal-treated PM01 strain
438 after 25 d growth in MAM pH 3.0 (Table 1). There was a concentration-dependent increase

439 in cellular accumulation of Cu and Cd with no significant difference between the values with
440 or without EDTA washing, indicating that almost all of the Cu and Cd was taken up within the
441 cell. In contrast, there were significant ($p < 0.05$) differences in Zn concentration following
442 EDTA washing, indicating that a smaller proportion of Zn was internalised (Table 1). As the
443 Zn concentration in the medium increased, the relative concentration within the cell
444 decreased, suggesting that Zn transport into the cell saturated at higher concentrations.
445 Overall, the characteristics of metal accumulation in artificial media were broadly similar to
446 those observed in the pond.

447

448 3.5. Metabolic adaptation to Cu tolerance by *C. acidophila* PM01

449 It was unknown whether the metal tolerance properties of *C. acidophila* differ between
450 strains isolated from AMD sites with differing water chemistry. To begin to assess this, the
451 tolerance of PM01 to a range of metals (Al, Cd, Cu, Fe, Mn and Zn) was compared to two
452 other strains of *C. acidophila* (CCAP 11/136 and CCAP 11/137) that had previously been
453 isolated from different field sites. Strains CCAP 11/136 and CCAP 11/137 were validated by
454 18S rDNA sequencing and phylogenetic analysis as *C. acidophila* species (Fig. 6). The
455 CCAP 11/136 strain was originally isolated from a highly acidic (~pH 1.0) humic acid-rich
456 peat water environment but the metal characteristics of the site were not described (Fott and
457 McCarthy, 1964). In contrast, strain CCAP 11/137 originated from German mining Lake 111
458 at pH 2.6, very low total P ($8 \mu\text{g L}^{-1}$) and fairly high levels of Zn (0.75 mg L^{-1}) (Spijkerman et
459 al., 2007a).

460 For Al, Cd and Mn there was no growth difference between the three strains.
461 However, Fe treatment substantially inhibited growth of CCAP 11/136, while Zn treatment
462 slightly inhibited growth of PM01 and CCAP 11/137 (Fig. S4). However, there was a very
463 clear-cut difference in growth between the three strains following Cu exposure (Fig. 7). On
464 solid media containing 130 mg L^{-1} Cu, the CCAP 11/137 strain was unable to grow, and the
465 CCAP 11/136 strain grew very weakly in contrast to strong growth by the PM01 strain (Fig.
466 7A). In liquid media with addition of 0.0 mg L^{-1} , 6.5 mg L^{-1} and 3 mg L^{-1} growth rate and cell

467 density after 14 days was identical between all three strains. However, with 130 mg L⁻¹ Cu
468 addition, cell growth was unchanged for PM01 but both CCAP 11/136 and 11/137 strains
469 were barely able to grow (Fig. 7B).

470 To examine whether there were any macromolecular changes within the strains in
471 response to the increasing copper treatment, and to examine whether the different strains
472 could be distinguished on the basis of their metabolic 'fingerprint', the FT-IR spectroscopy
473 technique was used. FT-IR spectra were collected for replicates of each strain cultivated in the
474 absence of added Cu or with the addition of 6.5 and 13 mg L⁻¹ Cu (Fig. S5). In addition,
475 PM01 was tested at the 130 mg L⁻¹ concentration that inhibited growth of 11/136 and 11/137,
476 thereby preventing FT-IR spectroscopy analysis of these strains at the higher Cu
477 concentration. PCA of all FT-IR spectra showed that the CCAP 11/137 strain samples
478 cluster separately from CCAP 11/136 and PM01 in all treatments, with the close clustering of
479 spectra indicating that Cu addition did not significantly alter the metabolic fingerprint of
480 CCAP 11/137 (Fig. 8A). Likewise Cu addition did not significantly alter the metabolic
481 fingerprint of the CCAP 11/136 strain, with all samples clustering with the PM01 control
482 samples. However, there was a clear difference in the FT-IR spectra-derived metabolic
483 profile of PM01 samples following addition of increasing Cu concentrations, with the sample
484 position within the PCA plot changing on the basis of both PC1 and PC2. At the highest
485 (130 mg L⁻¹) Cu concentration all replicate PM01 samples are distinct from the control
486 samples (Fig. 8A). The PC loading plots (Fig. 8B) indicate that along PC1 the spectral
487 changes are based predominantly on increased abundance of amide peaks at 1655 and
488 1545 cm⁻¹, as well as a decrease in one of the carbohydrate peaks at approximately 1036
489 cm⁻¹.

490

491 **4. Discussion**

492 This study has demonstrated that while AMD substantially impacts biodiversity in aquatic
493 environments, there is still substantial taxa abundance in these extreme locations. The
494 scarcity of invertebrates across the ponds clearly indicates the severity of pollution at this

495 abandoned mine site. With the exception of the near-neutral pH Pond 3, invertebrate
496 diversity was very poor in all ponds. Taxa diversity overall, including invertebrate diversity in
497 particular, was strongly correlated with pH and with dissolved metal concentration, which
498 was in line with expectations and previous studies (Courtney and Clements, 2002; Malmqvist
499 and Hoffsten, 1999). Nevertheless, some invertebrate species can adapt to extreme AMD
500 conditions (De Bisthoven et al., 2005; Deneke, 2000), and can be used as indicators of
501 mine-waste pollution (Gray and Delaney, 2008). Here Chironomidae were observed in all
502 ponds, including the very acidic Pond 1 and 4, and highly abundant in Pond 9, probably
503 because this pond is fairly shallow and has fine sediment in contrast to the other ponds.
504 Certain Chironomidae species are able to survive in highly acidic waters, such as the acid-
505 tolerant *Chironomus acidophilus* that has been found nearby in the highly acidic (pH 2.4) and
506 metal-rich Afon Goch river draining Parys Mountain (Michailova et al., 2009).

507 While Chironomidae were present in all ponds so there was also the consistent
508 presence of chlorophyte microalgae, and in particular a strain confirmed as *C. acidophila*.
509 The widespread occurrence of *C. acidophila* at Parys Mountain reflects the extremely acidic
510 and metal-rich pond water and is consistent with other acidic and metal-rich sites (Fott and
511 McCarthy, 1964; Gerloff-Elias et al., 2005; Hargreaves et al., 1975; Twiss, 1990). The widely
512 differing *C. acidophila* abundance between ponds may reflect nutrient availability. AMD
513 environments typically have very low productivity, due in part to low nutrients such as PO₄-P
514 (Spijkerman et al., 2007a; Spijkerman et al., 2007b), and in this study nutrients, and in
515 particular PO₄-P, were at low concentration in all but Pond 1. The low cell density of this
516 species in Ponds 2 – 9, where PO₄-P concentrations were low suggests growth limitation
517 due to low PO₄-P availability. In contrast, Pond 1 had very high PO₄-P and a high
518 concentration of *C. acidophila* cells. Other studies have shown high PO₄-P levels associated
519 with acidic lakes in areas with abundant PO₄-P-containing FeS minerals (Spijkerman, 2008);
520 however, as Pond 2 is within 5 m of Pond 1 and has the same mineralogy but significantly
521 lower PO₄-P levels, the high concentration of PO₄-P in Pond 1 is unlikely to be due to
522 minerology of the surrounding area. It may be that the high PO₄-P concentration is due to a

523 eutrophication event, as there is evidence of construction waste deposition that is likely to
524 explain the PO₄-P entry into the pond.

525 The high microalgae productivity of Pond 1 is due almost exclusively to *C. acidophila*
526 abundance. Acidophilic microalgae present in the other ponds, such as *K. corcontica*, *E.*
527 *naegelii* and *E. mutabilis*, were absent from Pond 1. *E. naegelii* and other acid-tolerant
528 diatoms are increasingly used as indicator species for acid polluted environments (Zalack et
529 al., 2010) and have been previously found in surface sediment from one of the rivers
530 draining from the Parys Mountain site (Dean et al., 2013), as well as in acid pit lakes with
531 equivalent water chemistry (Geller, 2013). However, their apparent low abundance in these
532 ponds might partly be due to the sampling regime used here from the near-surface open
533 water rather than from the bottom of the ponds. *E. mutabilis* was observed in two of the nine
534 ponds (Ponds 5 and 6), which are both ~pH 3.0, a pH range that has been shown to be
535 preferential for this organism (Brake et al., 2001). Furthermore, other acidophilic microalgae
536 such as *Ochromonas* sp. and *Dunaliella* sp., that are widely abundant in many acidic sites
537 (Aguilera et al., 2006; Nixdorf et al., 1998), were not identified in Pond 1, or in any other
538 Parys Mountain ponds. That *C. acidophila* was the only microalgal species identified in Pond
539 1 is likely to be due to the extreme conditions that restricts microalgal diversity. One of these
540 factors appears to be water pH, and the *C. acidophila* strain studied here was particularly
541 acid tolerant. In waters with pH > 3.0, it has been previously observed that algal biodiversity
542 is generally higher (Smucker et al., 2014), yet the least acidic pond studied here, Pond 3 (pH
543 5.3), still had very low algal biodiversity and very low *C. acidophila* abundance, suggesting
544 that other factors are also important, such as nutrient availability or water chemistry.

545 Although the dissolved Fe, Zn, Cu and Al concentrations in Pond 1 did not exceed
546 the very high concentrations seen in some AMD lakes and rivers such as those of the
547 Iberian Pyrite Belt (Sánchez España et al., 2008), they did exceed those seen in mine pit
548 lakes in Germany, Poland, Australia and USA (Geller, 2013). The metal concentrations of
549 the shallow Pond 1 can thus be considered high. The strain of *C. acidophila* isolated from
550 Pond 1 is therefore not just extremely acid-tolerant, but can also tolerate high metal

551 concentrations. The PM01 strain showed substantial tolerance to Cd, Cu, and Zn. A
552 comparison of PM01 to other genera of acid tolerant algae shows the Zn tolerance of PM01
553 was higher than that reported for *Chlorella protothecoides* var. *acidicola* isolated from AMD
554 sites in a Spanish mine and *E. mutabilis* isolated from the adit flowing from Parys Mountain.
555 However, both of these strains showed a higher Cu tolerance than PM01 (Ñancuqueo and
556 Johnson, 2012).

557 Other studies looking at metal tolerance in *C. acidophila* are fairly scarce, though a
558 study looking at a putative strain of *C. acidophila* OU 030/a isolated from a volcanic acid lake
559 also showed high tolerance to Cd, Cu and Zn in metal-rich minimal media (at pH 4.0) when
560 compared to other algal species (Nishikawa and Tominaga, 2001). Furthermore, a strain of
561 *C. acidophila* RT46 isolated from Río Tinto in Spain, showed unaffected photosynthetic
562 activity in response to 0.5 mM Cu (~32 mg L⁻¹ Cu) exposure (Olsson et al., 2015). A putative
563 *C. acidophila* strain (UTCC 121) isolated from Cu contaminated soil was previously shown to
564 tolerate up to 100 mg L⁻¹ Cu. However, in contrast, a laboratory strain of *C. acidophila*
565 (CCAP 11/96) was Cu sensitive, as was the non-acidophilic freshwater alga
566 *Chlamydomonas reinhardtii* (Twiss, 1990), indicating that the *Chlamydomonas* genera is not
567 intrinsically Cu tolerant. In this study we demonstrate that the *C. acidophila* strain isolated in
568 this study (PM01) has higher Cu tolerance than strains (CCAP 11/136 and 11/137) from
569 AMD field sites with less Cu pollution. This shows that the PM01 strain has adapted to the
570 high dissolved Cu concentrations of Pond 1, rather than this species having innate Cu
571 tolerance properties.

572 FT-IR spectroscopy analysis demonstrates that the tolerance of PM01 to copper is
573 partly due to its ability to modulate its metabolism in response to increasing Cu exposure, as
574 indicated by a dose-dependent change in spectra characteristics, which does not occur with
575 the other *C. acidophila* strains. Examination of the FT-IR spectra indicates that this metabolic
576 adaptation is predominantly due to protein increase and potential modification of protein
577 secondary structure, as shown by significant increase in amide I peak height associated with
578 C=O stretching, and in amide II peak height associated with N-H bending and C-N stretching

579 (Giordano et al., 2001). A previous study using FT-IR spectroscopy to examine sensitivity
580 and subsequent acclimation of microalgae to wastewater treatment, also found that the
581 acclimation process was coincident with a relative increase in amide I and amide II peak
582 height (Osundeko et al., 2014). Moreover, this same study showed that particularly sensitive
583 strains including *Chlamydomonas debaryana* and *Desmodesmus intermedius* exhibited
584 accumulation of carbon storage products including glycerolipids and starch, while acclimated
585 strains that could tolerate the wastewater conditions did not show this response. Likewise,
586 other metabolic indicators of stress such as increased carbohydrate and lipid peaks within
587 the FT-IR spectra were not observed in response to high Cu treatment in PM01.

588 Because Cu exerts toxicity in part through inhibition of key cellular processes
589 including photosynthesis, either directly due to Cu binding or indirectly via accumulation of
590 ROS (Jamers et al., 2013; Küpper et al., 2003; Sabatini et al., 2009), it would be expected
591 that adaptive mechanisms would counteract these processes in some ways. Proteomic
592 responses to Cu stress linked to Cu tolerance have previously been observed in other
593 organisms. Cu exposure experiments in plants and fungi have observed increases in soluble
594 protein that has been linked to induction of anti-oxidant enzymes (Cavalcanti Luna et al.,
595 2015; Gao et al., 2008; Rout et al., 2013) while induction of Cu-binding proteins has been
596 demonstrated in a Cu-tolerant variety of rice (Chen et al., 2015). Mechanisms of Cu
597 tolerance in algae are not well understood, and may involve differential Cu uptake and
598 internalisation in some cases (Levy et al., 2008). Although the molecular mechanisms of
599 stress tolerance by *C. acidophila* are also poorly understood but there have been some
600 recent insights. Heat shock proteins, which are a family of evolutionarily conserved stress
601 tolerance molecular chaperones, have been previously found to increase in abundance in *C.*
602 *acidophila* CCAP 11/137 in response to very low pH and metal-rich lake water treatment,
603 partly in response to high Fe concentration (Gerloff-Elias et al., 2006; Spijkerman et al.,
604 2007a). To date, no proteomic or enzymatic analysis has been performed in *C. acidophila* in
605 response to Cu stress, but a transcriptomics approach observed differences in mRNA
606 transcript profiles following Cu treatment in *C. acidophila* RT46. A range of gene transcripts

607 were up-regulated in response to Cu treatment including those involved in photosynthesis,
608 signaling and stress-response (Olsson et al., 2015). Future experiments will aim to examine
609 the proteomic adaptive response of *C. acidophila* PM01 in more detail in order to enhance
610 our fundamental understanding of metal tolerance in microalgae, also to appreciate the
611 effects that environmental pollution has on adaptive evolution and the ecological
612 consequences of such adaptation.

613 A potential application of highly metal tolerant microalgae such as *C. acidophila*
614 PM01 is the potential use of such organisms for metal bioremediation. Indeed PM01 was
615 confirmed to bioconcentrate metals through a combination of cell wall binding and
616 internalisation. Algae strains, such as those isolated in this study, may be used for *in situ*
617 lake bioremediation in surface water mesocosms by controlled eutrophication and harvesting
618 (Dessouki et al., 2005), or for *ex situ* bioremediation, such as immobilised algae in
619 bioreactors (Mehta and Gaur, 2005). The accumulated metals may then be harvested and
620 processed to allow metal recovery (Minoda et al., 2015; Raikova et al., 2016). Alternatively,
621 highly acid and metal tolerant microalgae such as the PM01 strain, may have an important
622 role in sustaining SO₄-reducing bacteria by providing organic carbon and thus increasing the
623 efficiency of AMD remediation microbial bioreactors (Diez-Ercilla et al., 2014; Ñancucheo
624 and Johnson, 2012; Totsche et al., 2006). For example, it was demonstrated that microalgal
625 addition to mine tailing mesocosms containing pyrite-oxidizing bacteria caused higher
626 production of alkalinity, higher concentrations of ferrous Fe, and increased immobilization of
627 Cu and Zn (Ñancucheo and Johnson, 2011).

628 The main aim of this study was to identify microorganisms from the open, standing
629 waters of the AMD ponds. Some of the ponds were surrounded by vegetation including
630 wetland plant species, which will harbor associated microorganisms (Aguinaga et al., 2018).
631 Although not the scope of this study, future research can examine the role of the plants on
632 microbial communities within isolated AMD pond environments, as well an examination of
633 biota along spatial transects of the ponds including within the sediment. Moreover, future
634 studies will be needed to examine how other microorganisms found in these environments

635 have adapted to AMD stresses, and whether there are common mechanisms between
636 different extremophile species.

637

638 **5. Conclusions**

639 AMD is a major source of freshwater pollution worldwide for which restoration is very
640 important. However, as found in this study, while extremely acidic and metal rich AMD
641 substantially impacts biota, there is still substantial biodiversity, with tolerance derived
642 through natural adaptation. In particular, a strain of *C. acidophila* is abundant in all ponds at
643 this Cu mine site, especially in waters with high acidity and coupled with high PO₄-P
644 concentration. Although Cu toxicity is a significant challenge to most photosynthetic
645 organisms, this strain of *C. acidophila* has specifically adapted to the high Cu status of the
646 ponds in contrast to other strains of the same species isolated from field sites elsewhere.
647 Moreover, the *C. acidophila* strain displays evidence of Cu-dependent metabolic plasticity.
648 The marked metal tolerance and metal accumulation characteristics of *C. acidophila* PM01
649 indicates that organisms from these environments have biotechnological potential, such as
650 bioremediation.

651

652 **Acknowledgments**

653 We thank Oscar Aguinaga and Rachel Webster for assistance with sampling work, Lorna
654 Slater and Shunzheng Wang for performing preliminary laboratory analysis of microalgae,
655 Claude-Eric Souquieres for assistance with sequence data analysis, and Paul Lythgoe for
656 ICP-AES analysis. This work was supported in part by BBSRC DTP PhD studentship
657 funding to O.A.M. (grant number BB/M011208/1).

658

659 **References**

660 Abinandan S, Subashchandrabose SR, Venkateswarlu K, Megharaj M. Microalgae–bacteria
661 biofilms: a sustainable synergistic approach in remediation of acid mine drainage.
662 Applied Microbiology and Biotechnology 2018; 102: 1131-1144.

663 Aguilera A, Manrubia SC, Gómez F, Rodríguez N, Amils R. Eukaryotic community
664 distribution and its relationship to water physicochemical parameters in an extreme
665 acidic environment, Río Tinto (Southwestern Spain). Applied and Environmental
666 Microbiology 2006; 72: 5325-5330.

667 Aguinaga OE, McMahon A, White KN, Dean AP, Pittman JK. Microbial community shifts in
668 response to acid mine drainage pollution within a natural wetland ecosystem.
669 Frontiers in Microbiology 2018; 9: 1445.

670 Amaral Zettler LA, Gomez F, Zettler E, Keenan BG, Amils R, Sogin ML. Microbiology:
671 eukaryotic diversity in Spain's River of Fire. Nature 2002; 417: 137-137.

672 August EE, McKnight DM, Hrncir DC, Garhart KS. Seasonal variability of metals transport
673 through a wetland impacted by mine drainage in the Rocky Mountains.
674 Environmental Science & Technology 2002; 36: 3779-3786.

675 Baffico GD, Diaz MM, Wenzel MT, Koschorreck M, Schimmele M, Neu TR, et al. Community
676 structure and photosynthetic activity of epilithon from a highly acidic (pH \leq 2) mountain
677 stream in Patagonia, Argentina. Extremophiles 2004; 8: 463-473.

678 Baker BJ, Lutz MA, Dawson SC, Bond PL, Banfield JF. Metabolically active eukaryotic
679 communities in extremely acidic mine drainage. Applied and Environmental
680 Microbiology 2004; 70: 6264-6271.

681 Beamud SG, Diaz MM, Pedrozo FL. Summer phytoplankton composition and nitrogen
682 limitation of the deep, naturally-acidic (pH \sim 2.2) Lake Caviahue, Patagonia,
683 Argentina. Limnologica 2007; 37: 37-48.

684 Bernal M, Ramiro MV, Cases R, Picorel R, Yruela I. Excess copper effect on growth,
685 chloroplast ultrastructure, oxygen-evolution activity and chlorophyll fluorescence in
686 *Glycine max* cell suspensions. Physiologia Plantarum 2006; 127: 312-325.

687 Boulton S, Collins DN, White KN, Curtis CD. Metal transport in a stream polluted by acid mine
688 drainage - the Afon Goch, Anglesey, UK. Environmental Pollution 1994; 84: 279-284.

689 Brake SS, Dannelly HK, Connors KA. Controls on the nature and distribution of an alga in
690 coal mine-waste environments and its potential impact on water quality.
691 Environmental Geology 2001; 40: 458-469.

692 Cavalcanti Luna MA, Vieira ER, Okada K, Campos-Takaki GM, do Nascimento AE. Copper-
693 induced adaptation, oxidative stress and its tolerance in *Aspergillus niger* UCP1261.
694 Electronic Journal of Biotechnology 2015; 18: 418-427.

695 Chen C, Song Y, Zhuang K, Li L, Xia Y, Shen Z. Proteomic analysis of copper-binding
696 proteins in excess copper-stressed roots of two rice (*Oryza sativa* L.) varieties with
697 different Cu tolerances. PLoS ONE 2015; 10: e0125367.

698 Cid C, Garcia-Descalzo L, Casado-Lafuente V, Amils R, Aguilera A. Proteomic analysis of
699 the response of an acidophilic strain of *Chlamydomonas* sp. (Chlorophyta) to natural
700 metal-rich water. Proteomics 2010; 10: 2026-2036.

701 Courtney LA, Clements WH. Assessing the influence of water and substratum quality on
702 benthic macroinvertebrate communities in a metal-polluted stream: an experimental
703 approach. Freshwater Biology 2002; 47: 1766-1778.

704 Das BK, Roy A, Koschorreck M, Mandal SM, Wendt-Potthoff K, Bhattacharya J. Occurrence
705 and role of algae and fungi in acid mine drainage environment with special reference
706 to metals and sulfate immobilization. Water Research 2009; 43: 883-894.

707 De Bisthoven LJ, Gerhardt A, Soares AMVM. Chironomidae larvae as bioindicators of an
708 acid mine drainage in Portugal. Hydrobiologia 2005; 532: 181-191.

709 Dean AP, Lynch S, Rowland P, Toft BD, Pittman JK, White KN. Natural wetlands are
710 efficient at providing long-term metal remediation of freshwater systems polluted by
711 acid mine drainage. Environmental Science & Technology 2013; 47: 12029-12036.

712 Dean AP, Sigee DC, Estrada B, Pittman JK. Using FTIR spectroscopy for rapid
713 determination of lipid accumulation in response to nitrogen limitation in freshwater
714 microalgae. Bioresource Technology 2010; 101: 4499-4507.

715 DeLong EF. Archaea in coastal marine environments. Proceedings of the National Academy
716 of Sciences, USA 1992; 89: 5685-5689.

717 Deneke R. Review of rotifers and crustaceans in highly acidic environments of pH values ≤ 3 .
718 Hydrobiologia 2000; 433: 167-172.

719 Dessouki TCE, Hudson JJ, Neal BR, Bogard MJ. The effects of phosphorus additions on the
720 sedimentation of contaminants in a uranium mine pit-lake. Water Research 2005; 39:
721 3055-3061.

722 Diez-Ercilla M, Sánchez-España J, Yusta I, Wendt-Potthoff K, Koschorreck M. Formation of
723 biogenic sulphides in the water column of an acidic pit lake: biogeochemical controls
724 and effects on trace metal dynamics. Biogeochemistry 2014; 121: 519-536.

725 Driver T, Bajhaiya AK, Allwood JW, Goodacre R, Pittman JK, Dean AP. Metabolic responses
726 of eukaryotic microalgae to environmental stress limit the ability of FT-IR
727 spectroscopy for species identification. Algal Research 2015; 11: 148-155.

728 Florence K, Sapsford DJ, Johnson DB, Kay CM, Wolkersdorfer C. Iron-mineral accretion
729 from acid mine drainage and its application in passive treatment. Environmental
730 Technology 2016; 37: 1428-1440.

731 Fott B, McCarthy AJ. Three acidophilic Volvocine flagellates in pure culture. The Journal of
732 Protozoology 1964; 11: 116-120.

733 Gao S, Yan R, Cao M, Yang W, Wang S, Chen F. Effects of copper on growth, antioxidant
734 enzymes and phenylalanine ammonia-lyase activities in *Jatropha curcas* L. seedling.
735 Plant Soil Environment 2008; 54: 117-122.

736 Geller W. Case Studies and Regional Surveys. In: Geller W, Schultze M, Kleinmann R,
737 Wolkersdorfer C, editors. Acidic Pit Lakes: The Legacy of Coal and Metal Surface
738 Mines. Springer Berlin Heidelberg, Berlin, Heidelberg, 2013, pp. 265-435.

739 Geller W, Schultze M. Remediation and Management of Acidified Pit Lakes and Outflowing
740 Waters. In: Geller W, Schultze M, Kleinmann R, Wolkersdorfer C, editors. Acidic Pit
741 Lakes: The Legacy of Coal and Metal Surface Mines. Springer Berlin Heidelberg,
742 Berlin, Heidelberg, 2013, pp. 225-264.

743 Gerloff-Elias A, Barua D, Mölich A, Spijkerman E. Temperature- and pH-dependent
744 accumulation of heat-shock proteins in the acidophilic green alga *Chlamydomonas*
745 *acidophila*. FEMS Microbiology Ecology 2006; 56: 345-354.

746 Gerloff-Elias A, Spijkerman E, Proschold T. Effect of external pH on the growth,
747 photosynthesis and photosynthetic electron transport of *Chlamydomonas acidophila*
748 Negoro, isolated from an extremely acidic lake (pH 2.6). Plant, Cell & Environment
749 2005; 28: 1218-1229.

750 Giordano M, Kansiz M, Heraud P, Beardall J, Wood B, McNaughton D. Fourier transform
751 infrared spectroscopy as a novel tool to investigate changes in intracellular
752 macromolecular pools in the marine microalga *Chaetoceros muellerii*
753 (Bacillariophyceae). Journal of Phycology 2001; 37: 271-279.

754 Gray NF, Delaney E. Comparison of benthic macroinvertebrate indices for the assessment
755 of the impact of acid mine drainage on an Irish river below an abandoned Cu–S
756 mine. Environmental Pollution 2008; 155: 31-40.

757 Greenhaigh M, Ovenden D. Freshwater Life: Britain and Northern Europe. London, UK:
758 Collins, 2007.

759 Gustafsson JP. Visual MINTEQ ver. 3.0. Department of Land and Water Resources
760 Engineering, The Royal Institute of Technology, Sweden., 2010.

761 Hargreaves JW, Lloyd EJH, Whitton BA. Chemistry and vegetation of highly acidic streams.
762 Freshwater Biology 1975; 5: 563-576.

763 Hassler CS, Slaveykova VI, Wilkinson KJ. Discriminating between intra- and extracellular
764 metals using chemical extractions. Limnology and Oceanography-Methods 2004; 2:
765 237-247.

766 Hedin R, Weaver T, Wolfe N, Weaver K. Passive treatment of acidic coal mine drainage:
767 The Anna S mine passive treatment complex. Mine Water and the Environment
768 2010; 29: 165-175.

769 Hillebrand H, Durselen CD, Kirschtel D, Pollinger U, Zohary T. Biovolume calculation for
770 pelagic and benthic microalgae. Journal of Phycology 1999; 35: 403-424.

771 Hirooka S, Hirose Y, Kanesaki Y, Higuchi S, Fujiwara T, Onuma R, et al. Acidophilic green
772 algal genome provides insights into adaptation to an acidic environment.
773 Proceedings of the National Academy of Sciences, USA 2017; 114: E8304-E8313.

774 Huss VAR, Frank C, Hartmann EC, Hirmer M, Kloboucek A, Seidel BM, et al. Biochemical
775 taxonomy and molecular phylogeny of the genus *Chlorella* sensu lato (Chlorophyta).
776 Journal of Phycology 1999; 35: 587-598.

777 Jamers A, Blust R, De Coen W, Griffin JL, Jones OAH. Copper toxicity in the microalga
778 *Chlamydomonas reinhardtii*: an integrated approach. BioMetals 2013; 26: 731-740.

779 Jiang Y, Zhu Y, Hu Z, Lei A, Wang J. Towards elucidation of the toxic mechanism of copper
780 on the model green alga *Chlamydomonas reinhardtii*. Ecotoxicology 2016; 25: 1417-
781 1425.

782 John DM, Whitton BA, Brook AJ. The Freshwater Algae of the British Isles. An Identification
783 Guide to Freshwater and Terrestrial Algae. Cambridge: Cambridge University Press,
784 2002.

785 Johnson DB. Chemical and microbiological characteristics of mineral spoils and drainage
786 waters at abandoned coal and metal mines. Water, Air, & Soil Pollution: Focus 2003;
787 3: 47-66.

788 Johnson DB, Hallberg KB. The microbiology of acidic mine waters. Research in Microbiology
789 2003; 154: 466-473.

790 Johnson DB, Hallberg KB. Acid mine drainage remediation options: a review. Science of the
791 Total Environment 2005; 338: 3-14.

792 Küpper H, Šetlík I, Šetliková E, Ferimazova N, Spiller M, Küpper FC. Copper-induced
793 inhibition of photosynthesis: limiting steps of *in vivo* copper chlorophyll formation in
794 *Scenedesmus quadricauda*. Functional Plant Biology 2003; 30: 1187-1196.

795 Levy JL, Angel BM, Stauber JL, Poon WL, Simpson SL, Cheng SH, et al. Uptake and
796 internalisation of copper by three marine microalgae: Comparison of copper-sensitive
797 and copper-tolerant species. Aquatic Toxicology 2008; 89: 82-93.

798 Malmqvist B, Hoffsten P-O. Influence of drainage from old mine deposits on benthic
799 macroinvertebrate communities in central Swedish streams. *Water Research* 1999;
800 33: 2415-2423.

801 Martens H, Stark E. Extended multiplicative signal correction and spectral interference
802 subtraction: new preprocessing methods for near infrared spectroscopy. *Journal of*
803 *Pharmaceutical and Biomedical Analysis* 1991; 9: 625-635.

804 Mehta SK, Gaur JP. Use of algae for removing heavy metal ions from wastewater: progress
805 and prospects. *Critical Reviews in Biotechnology* 2005; 25: 113-152.

806 Mendez-Garcia C, Pelaez AI, Mesa V, Sánchez J, Golyshina OV, Ferrer M. Microbial
807 diversity and metabolic networks in acid mine drainage habitats. *Frontiers in*
808 *Microbiology* 2015; 6: 475.

809 Michailova P, Ilkova J, Kerr R, White KN. Chromosome variability in *Chironomus acidophilus*
810 Keyl, 1960 from the Afon Goch, UK – a river subject to long-term trace metal
811 pollution. *Aquatic Insects* 2009; 31: 213-225.

812 Minoda A, Sawada H, Suzuki S, Miyashita S-i, Inagaki K, Yamamoto T, et al. Recovery of
813 rare earth elements from the sulfothermophilic red alga *Galdieria sulphuraria* using
814 aqueous acid. *Applied Microbiology and Biotechnology* 2015; 99: 1513-1519.

815 Ñancuqueo I, Johnson DB. Acidophilic algae isolated from mine-impacted environments and
816 their roles in sustaining heterotrophic acidophiles. *Frontiers in Microbiology* 2012; 3:
817 325.

818 Ñancuqueo I, Johnson DB. Significance of microbial communities and interactions in
819 safeguarding reactive mine tailings by ecological engineering. *Applied and*
820 *Environmental Microbiology* 2011; 77: 8201-8208.

821 Neculita C-M, Zagury GJ, Bussière B. Passive treatment of acid mine drainage in
822 bioreactors using sulfate-reducing bacteria. *Journal of Environmental Quality* 2007;
823 36: 1-16.

824 Nishikawa K, Tominaga N. Isolation, growth, ultrastructure, and metal tolerance of the green
825 alga, *Chlamydomonas acidophila* (Chlorophyta). *Bioscience, Biotechnology, and*
826 *Biochemistry* 2001; 65: 2650-2656.

827 Nixdorf B, Mischke U, Leßmann D. Chrysophytes and chlamydomonads: pioneer colonists in
828 extremely acidic mining lakes (pH <3) in Lusatia (Germany). *Hydrobiologia* 1998;
829 369: 315-327.

830 Nordstrom DK. Advances in the hydrogeochemistry and microbiology of acid mine waters.
831 *International Geology Review* 2000; 42: 499-515.

832 Olaveson MM, Stokes PM. Responses of the acidophilic alga *Euglena mutabilis*
833 (Euglenophyceae) to carbon enrichment at pH 3. *Journal of Phycology* 1989; 25:
834 529-539.

835 Olsson S, Penacho V, Puente-Sánchez F, Díaz S, Gonzalez-Pastor JE, Aguilera A.
836 Horizontal gene transfer of phytochelatin synthases from bacteria to extremophilic
837 green algae. *Microbial Ecology* 2017; 73: 50-60.

838 Olsson S, Puente-Sánchez F, Gómez MJ, Aguilera A. Transcriptional response to copper
839 excess and identification of genes involved in heavy metal tolerance in the
840 extremophilic microalga *Chlamydomonas acidophila*. *Extremophiles* 2015; 19: 657-
841 672.

842 Osundeko O, Davies H, Pittman JK. Oxidative stress-tolerant microalgae strains are highly
843 efficient for biofuel feedstock production on wastewater. *Biomass and Bioenergy*
844 2013; 56: 284-294.

845 Osundeko O, Dean AP, Davies H, Pittman JK. Acclimation of microalgae to wastewater
846 environments involves increased oxidative stress tolerance activity. *Plant and Cell*
847 *Physiology* 2014; 55: 1848-1857.

848 Pedrozo F, Kelly L, Diaz M, Temporetti P, Baffico G, Kringel R, et al. First results on the
849 water chemistry, algae and trophic status of an Andean acidic lake system of
850 volcanic origin in Patagonia (Lake Caviahue). *Hydrobiologia* 2001; 452: 129-137.

851 Pollio A, Cennamo P, Ciniglia C, De Stefano M, Pinto G, A.R. Huss V. *Chlamydomonas*
852 *pitschmannii* Ettl, a little known species from thermoacidic environments. *Protist*
853 2005; 156: 287-302.

854 Puente-Sánchez F, Díaz S, Penacho V, Aguilera A, Olsson S. Basis of genetic adaptation to
855 heavy metal stress in the acidophilic green alga *Chlamydomonas acidophila*. *Aquatic*
856 *Toxicology* 2018; 200: 62-72.

857 Puente-Sánchez F, Olsson S, Aguilera A. Comparative transcriptomic analysis of the
858 response of *Dunaliella acidophila* (Chlorophyta) to short-term cadmium and chronic
859 natural metal-rich water exposures. *Microbial Ecology* 2016; 72: 595-607.

860 Quigley M. *Invertebrates of Streams and Rivers: A Key to Identification*. London, UK:
861 Edward Arnold, 1977.

862 Raikova S, Smith-Baedorf H, Bransgrove R, Barlow O, Santomauro F, Wagner JL, et al.
863 Assessing hydrothermal liquefaction for the production of bio-oil and enhanced metal
864 recovery from microalgae cultivated on acid mine drainage. *Fuel Processing*
865 *Technology* 2016; 142: 219-227.

866 Rout JR, Ram SS, Das R, Chakraborty A, Sudarshan M, Sahoo SL. Copper-stress induced
867 alterations in protein profile and antioxidant enzymes activities in the in vitro grown
868 *Withania somnifera* L. *Physiology and Molecular Biology of Plants* 2013; 19: 353-361.

869 Sabatini SE, Juárez ÁB, Eppis MR, Bianchi L, Luquet CM, Ríos de Molina MdC. Oxidative
870 stress and antioxidant defenses in two green microalgae exposed to copper.
871 *Ecotoxicology and Environmental Safety* 2009; 72: 1200-1206.

872 Sánchez España J, Pamo EL, Pastor ES, Ercilla MD. The acidic mine pit lakes of the Iberian
873 Pyrite Belt: An approach to their physical limnology and hydrogeochemistry. *Applied*
874 *Geochemistry* 2008; 23: 1260-1287.

875 Schönknecht G, Chen W-H, Ternes CM, Barbier GG, Shrestha RP, Stanke M, et al. Gene
876 transfer from bacteria and archaea facilitated evolution of an extremophilic eukaryote.
877 *Science* 2013; 339: 1207-1210.

878 Schultze M. Limnology of Pit Lakes. In: Geller W, Schultze M, Kleinmann R, Wolkersdorfer
879 C, editors. Acidic Pit Lakes: The Legacy of Coal and Metal Surface Mines. Springer
880 Berlin Heidelberg, Berlin, Heidelberg, 2013, pp. 23-224.

881 Sharma SS, Dietz K-J. The relationship between metal toxicity and cellular redox imbalance.
882 Trends in Plant Science 2009; 14: 43-50.

883 Smucker NJ, Drerup SA, Vis ML. Roles of benthic algae in the structure, function, and
884 assessment of stream ecosystems affected by acid mine drainage. Journal of
885 Phycology 2014; 50: 425-436.

886 Spijkerman E. Phosphorus limitation of algae living in iron-rich, acidic lakes. Aquatic
887 Microbial Ecology 2008; 53: 201-210.

888 Spijkerman E, Barua D, Gerloff-Elias A, Kern J, Gaedke U, Heckathorn SA. Stress
889 responses and metal tolerance of *Chlamydomonas acidophila* in metal-enriched lake
890 water and artificial medium. Extremophiles 2007a; 11: 551-562.

891 Spijkerman E, Bissinger V, Meister A, Gaedke U. Low potassium and inorganic carbon
892 concentrations influence a possible phosphorus limitation in *Chlamydomonas*
893 *acidophila* (Chlorophyceae). European Journal of Phycology 2007b; 42: 327-339.

894 Stamatakis A. RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with
895 thousands of taxa and mixed models. Bioinformatics 2006; 22: 2688-2690.

896 Totsche O, Fyson A, Steinberg CEW. Microbial alkalinity production to prevent reacidification
897 of neutralized mining lakes. Mine Water and the Environment 2006; 25: 204-213.

898 Twiss MR. Copper tolerance of *Chlamydomonas acidophila* (Chlorophyceae) isolated from
899 acidic, copper-contaminated soils. Journal of Phycology 1990; 26: 655-659.

900 Valko M, Morris H, Cronin MTD. Metals, toxicity and oxidative stress. Current Medicinal
901 Chemistry 2005; 12: 1161-1208.

902 Varshney P, Mikulic P, Vonshak A, Beardall J, Wangikar PP. Extremophilic micro-algae and
903 their potential contribution in biotechnology. Bioresource Technology 2015; 184: 363-
904 372.

905 Vrba J, Kopáček J, Fott J, Kohout L, Nedbalová L, Pražáková M, et al. Long-term studies
906 (1871–2000) on acidification and recovery of lakes in the Bohemian Forest (central
907 Europe). *Science of The Total Environment* 2003; 310: 73-85.

908 Webster RE, Dean AP, Pittman JK. Cadmium exposure and phosphorus limitation increases
909 metal content in the freshwater alga *Chlamydomonas reinhardtii*. *Environmental*
910 *Science & Technology* 2011; 45: 7489-7496.

911 Younger PL, Potter HAB. Parys in Springtime: Hazard Management and Steps Towards
912 Remediation of the UKs Most Polluted Acidic Mine Discharge. 9th International
913 Conference on Acid Rock Drainage (ICARD), Ottawa, Canada, 2012.


914 Yun H-S, Lee H, Park Y-T, Ji M-K, Kabra AN, Jeon C, et al. Isolation of novel microalgae
915 from acid mine drainage and its potential application for biodiesel production. *Applied*
916 *Biochemistry and Biotechnology* 2014; 173: 2054-2064.

917 Zalack JT, Smucker NJ, Vis ML. Development of a diatom index of biotic integrity for acid
918 mine drainage impacted streams. *Ecological Indicators* 2010; 10: 287-295.

919


920

921 **Figures**




922

923 Fig. 1. Map and satellite image of Parys Mountain copper mine in Anglesey, North Wales,
924 UK. The sampled AMD ponds are labelled 1–9 and the AMD-polluted Southern Afon Goch
925 river is indicated. The location coordinates of each sampled pond are also shown (A). (B)
926 Satellite image of the mine and the sampled ponds. Source, Google Earth, 2016.
927



928


929 Fig. 2. Water chemistry of the AMD ponds. (A - C) pH and conductivity values (A), and
 930 concentration of dissolved Fe, Cd (B), Cu and Zn (C) for each pond during 2015 (spring,
 931 summer, autumn samples). Boxes show the 25th and 75th percentiles, the line within the
 932 boxes shows the median values, and the whisker bars show minimum and maximum values
 933 ($n = 3 - 18$). (D) PO₄-P, NO₃-N and NH₄-N concentration in each pond (June sample). Pond
 934 7 data is not available. Values are means ($n = 3 - 12$) and error bars correspond to the
 935 standard error of the mean. (E) Mean chlorophyll-a (Chl a) concentration in each pond
 936 during 2015 (spring, summer, autumn samples). Chl a change in Pond 1 during the year and
 937 pond water temperature is shown (inset). Each pond was sampled in triplicate on three
 938 separate occasions (spring, summer, autumn). Values are means ($n = 9$) and error bars
 939 correspond to the standard error of the mean. For all data, bars that do not share a lower
 940 case letter show significant difference ($p < 0.05$) between pond sites.
 941



942


943 Fig. 3. PCA of water chemistry data from each of the AMD ponds. (A) Observation plot for
 944 each pond. The upper site small ponds are shown as blue triangles, the near-neutral pH
 945 pond as a red circle, and the lower site large lagoon ponds as green squares. (B) Variables
 946 plot with selected environmental variables labelled.

947



948


949 Fig. 4. Taxa diversity of eukaryotic organisms in the AMD ponds as determined by numbers
 950 of distinct taxonomic families of microalgae, moss/liverwort, fungi, protozoa, as identified by
 951 18S rRNA gene amplicon sequencing, and families of invertebrates, as identified by visual
 952 observation and use of taxonomic identification keys. Data for Pond 8 is not available.
 953



954


955 Fig. 5. (A) Abundance of *C. acidophila* microalgae and Chironomidae insects in each pond
 956 from summer sampling. Values of *C. acidophila* are means ($n = 3$) and error bars correspond
 957 to the standard error of the mean, while value of Chironomidae are total counts. For *C.*
 958 *acidophila* data, bars that do not share a lower case letter show significant difference ($p <$
 959 0.05) between pond sites. Chironomidae data for Pond 8 is not available. (B) Linear
 960 regression analyses for dissolved phosphate, pH, conductivity and dissolved Cu in relation to
 961 *C. acidophila* cell counts in AMD ponds. Apart from pH values, all data were natural log
 962 transformed.

963




964

965 Fig. 6. 18S rDNA sequence analysis of *C. acidophila* strains. Phylogenetic tree based on
 966 18S nucleotide sequence obtained from the PM01 strain isolated from Pond 1, sequences of
 967 *C. acidophila* CCAP 11/136 and CCAP 11/137 strains, other known or putative *C. acidophila*
 968 strains, and selected unicellular Chlorophyta microalgae of the *Chlamydomonas moewusii*
 969 clade, including three strains originally classified as *C. acidophila*. The sequence lengths
 970 were between 1528 – 1792 nucleotides. *Chlamydomonas reinhardtii* is included as an out-
 971 group. Accession numbers are shown for each sequence. Bootstrap percentage values are
 972 indicated at the tree nodes of branches for 100 replications and indicate confidence in tree
 973 node positions. The branch length scale bar indicates evolutionary distance.
 974



975

976 Fig. 7. Cu tolerance of *C. acidophila* PM01 in comparison to *C. acidophila* strains CCAP
 977 11/136 and 11/137. (A) Growth of algae spot dilutions on MAM pH 3.0 plates with or without
 978 added Cu and photographed after 12 d. Image is representative of 3 independent
 979 experiments. (B) Growth of strains in liquid MAM pH 3.0 determined after 14 d cultivation in
 980 response to a range of Cu concentrations. Values are means (n = 3 - 4) and error bars
 981 correspond to the standard error of the mean. Bars that do not share a lower case letter
 982 show significant difference (p < 0.05) between strains.
 983



984

985 Fig. 8. PCA clustering of FT-IR spectra from *C. acidophila* PM01 in comparison to *C.*
 986 *acidophila* strains CCAP 11/136 and 11/137 in response to a range of Cu concentrations. (A)
 987 PCA scores plot of replicate (n = 4) FT-IR spectra obtained from cells grown after 14 d
 988 cultivation in liquid MAM pH 3.0 with increasing concentrations of added Cu or no added Cu.
 989 Only PM01 cells could grow in 130 mg L⁻¹ Cu. (B) PC1 and PC2 loading plot. Band peaks
 990 which explain most of the variation for each PC are ν C=O of amide I (1655 cm⁻¹) and δ N-H
 991 of amide II (1545 cm⁻¹).

992