



## LJMU Research Online

**Delargy, I, Crowley, D and van Hout, MC**

**Twenty years on, the Methadone Treatment Protocol in Ireland: Reflections on General Practice**

<http://researchonline.ljmu.ac.uk/9780/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Delargy, I, Crowley, D and van Hout, MC Twenty years on, the Methadone Treatment Protocol in Ireland: Reflections on General Practice. Harm Reduction Journal. ISSN 1477-7517 (Accepted)**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

[Click here to view linked References](#)

**Title:** Twenty years of the Methadone Treatment Protocol in Ireland: Reflections on the role  
of General Practice.

### **Authors**

Dr Ide Delargy, Director of the Substance Misuse Programme Irish College of General  
Practitioners, Dublin, Ireland [ide.delargy@icgp.ie](mailto:ide.delargy@icgp.ie)

Dr Des Crowley, Assistant Director of the Substance Misuse Programme, Irish College of  
General Practitioners, Dublin, Ireland. [des.crowley@icgp.ie](mailto:des.crowley@icgp.ie)

Dr Marie Claire Van Hout Professor, Public Health Institute, Liverpool John Moore's  
University, Liverpool, United Kingdom [m.c.vanhout@ljmu.ac.uk](mailto:m.c.vanhout@ljmu.ac.uk) .

**Corresponding Author** Dr Ide Delargy, Director of the Substance Misuse Programme Irish  
College of General Practitioners, Dublin, Ireland [ide.delargy@icgp.ie](mailto:ide.delargy@icgp.ie)

## **Abstract**

### **Background**

Opioid dependence, characterised by socio economic disadvantage and significant morbidity and mortality, remains a major public health problem in Ireland. Through the methadone treatment protocol (MTP), Irish general practice has been a leader in the introduction and expansion of Irish harm reduction services, including opioid substitution treatment (OST), needle and syringe programs (NSP) and naloxone provision. These services have been effective in engaging opiate users in treatment, reducing human deficiency virus (HIV) and hepatitis C virus (HCV) transmission and reducing drug related morbidities. Challenges remain in relation to choice of substitution treatments, timely access to OST services, adequate coverage of NSP, naloxone provision and increasing drug related deaths.

### **Methods**

A narrative review was conducted and designed to present a broad perspective on the Irish MTP, and to describe its history and development in terms of clinical care, stakeholder views and changing trends.

### **Results**

Three themes emerged from the analysis; The History of the Methadone Treatment Protocol, Service User and Provider Views, and Challenges and Developments. Despite initial concern about methadone maintenance treatment (MMT) in Ireland, increased participation by Irish GPs in the treatment of opioid dependence is observed over the last two decades. There are now over 10,000 people on methadone treatment in Ireland, with 40% treated in general practice. The MTP provides structure, remuneration and guidance to GPs and is underpinned by; training, ongoing education and a system of quality assurance provided by the Irish College of General Practice (ICGP). Challenges include the negative views of patients around how methadone services are delivered, the stigma associated with methadone treatment, the

1 lack of choice around substitution medication, waiting lists for treatment in certain areas and  
2 rates of fatal overdose.  
3

#### 4 **Conclusion**

5  
6  
7 Twenty years of the MTP has been the mainstay of harm reduction services in Ireland. It has  
8 provided a network of specially trained GPs who provide methadone to over 10,000 patients  
9 across Ireland within a structured framework of training, quality assurance and remuneration.  
10  
11

12  
13  
14 With the ongoing commitment of Irish specialists in the field of addiction medicine, further  
15 improvements to support and treat patients can be made.  
16  
17  
18  
19  
20  
21

22 **Key Words;** Ireland, general practice, methadone, opioid dependence, harm reduction  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Background

Opiate use in Ireland as elsewhere is characteristic of socio economic disadvantage, low educational attainment and restricted economic opportunity.<sup>1,2</sup> The most recent capture recapture study of opiate users indicates that opiate use in Ireland is now entering a stabilised phase, with a small decrease in use compared to 2006 and the rate of use among young adults aged 15-24 years declining, and with a visible aging cohort effect of those in the 35 to 64 years age group.<sup>3</sup> In 2003, 6,667 individuals were registered on the Central Treatment List (CTL) out of 15,000 opiate users.<sup>4</sup> Kelly, Carvalho and Teljeur<sup>5</sup> estimated that in 2006 there were between 18,136 and 23,576 opioid users resident in Ireland, with the highest rate of heroin use in Europe at just over 7 cases per 1000 population, and small increases in the older drug using population and in Irish females. National prevalence and treatment data indicated at the time that opiate use was no longer confined to the greater urban context in the capital.<sup>1,5,6,7,8</sup>

The overall number of cases treated for problem opiate use (mainly heroin) increased between 2006 and 2015.<sup>9</sup> Most recent trends indicate a concerning shift towards greater levels of poly substance use<sup>5,6,9</sup> with related treatment demand (i.e. benzodiazepines) since 2007, dependence rates on over the counter and prescribed opiates such as codeine since 2008,<sup>10</sup> increase in drug related deaths,<sup>9</sup> and the emergence of potent and potentially fatal synthetic fentanyl in 2018. Treatment data in 2013 reports that incidence of treated problem substance use among Travellers is three times that compared to the general population (523 per hundred versus 173 per 100,000),<sup>11</sup> with reports in 2017 indicating increased vulnerability of this ethnic minority to problem opiate use.<sup>9,12</sup> In 2016, the Health Protection Surveillance Centre reported that the rate of infectious diseases (human immune deficiency virus, (HIV) and hepatitis C (HCV) is declining amongst people who inject drugs (PWID), despite an increase in 2015 due to a reported HIV outbreak.<sup>13</sup>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

In terms of pharmacological options to treat opiate dependence, substitution treatment using methadone is most common in Ireland, with buprenorphine-naloxone currently available on a limited basis.<sup>10</sup> The Irish model of care for delivery of methadone treatment acknowledges the central role of the specialist trained general practitioner (GP) in primary care.<sup>14</sup> The Irish College of General Practitioners (ICGP) has played a central role in developing the Methadone Treatment Protocol (MTP) which provided for the delivery of methadone treatment in primary care in the Irish context.

A network of prescribing GPs work closely with statutory (funded and operated by the Health Services Executive, HSE) and non-statutory (part funded by the HSE through a service level agreement) organisations to optimise methadone treatment delivery. The network includes specialist GPs working in HSE addiction clinics and Level 1 and Level 2 GPs working in primary care. Methadone treatment is commenced by specialist GPs in either addiction clinics or in a general practice setting (Level 2 GP). Level 2 trained doctors are qualified to initiate treatment in general practice, stabilise doses and provide ongoing maintenance treatment.<sup>15</sup> Level 1 trained doctors are encouraged to manage patients registered to their practice with patient stabilisation occurring in specialised clinics or with a Level 2 trained doctor. Once the patient is stabilised on methadone, referral to a Level 1 GPs working in the community for ongoing care is encouraged. All patients on methadone are listed on the confidential Central Treatment List (CTL) with each patient linked to one specific prescriber and a single dispensing site. Both Level 1 and Level 2 contracts attract additional remuneration for GPs caring for opioid dependent patients. Since 1998, greater prescribing of methadone in primary care has occurred, with the number of trained GPs rising steadily each year. There are currently (mid 2018) a total of 1718 trained Level 1 GPs and 188 Level 2 trained GPs on the ICGP database. We present here a narrative review of the history and development of the Irish MTP, 20 years since its initial inception.

## Methods

1  
2 A narrative review was conducted and designed to present a broad perspective on the Irish  
3  
4 MTP, and describe its history and development in terms of clinical care, stakeholder views  
5  
6 and changing trends. A comprehensive search was conducted on the National Documentation  
7  
8 Centre for Drugs, Health Research Board database, with no restriction on date range or types  
9  
10 of records. Key search terms used were methadone treatment, general practice, opioid  
11  
12 dependence and harm reduction combined with Ireland. Databases searched included  
13  
14 PubMed, Science Direct, EMBASE, PsycINFO, Cochrane library and Medline. No limits  
15  
16 were placed on dates. Follow up search strategies included hand searching relevant national  
17  
18 websites including the Health Service Executive (HSE), Irish Prison Service (IPS),  
19  
20 Departments of Health and Justice (DOH and DJ), the Irish Penal Reform Trust (IPRT),  
21  
22 Health Surveillance Protection Centre (HSPC) and EMCDDA. A hand search of reference  
23  
24 lists from published peer reviewed studies was also undertaken. References were managed by  
25  
26 the citation manager Endnote®. National experts and authors of existing papers were  
27  
28 contacted to identify possible sources of unpublished and grey literature. The research team  
29  
30 reviewed the relevant literature and agreed on the structuring of the review under the  
31  
32 following three themes; The History of the Methadone Treatment Protocol, Service User and  
33  
34 Provider Views, and Challenges and Developments.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

### **The History of the Methadone Treatment Protocol (MTP)**

46  
47  
48 The Drug Treatment and Advisory Service was established in Jervis Street hospital in 1969.  
49  
50 In 1971, methadone was introduced as the standardised therapeutic approach for treatment of  
51  
52 opiate dependence, with physeptone treatment available to a small cohort of patients  
53  
54 attending the service. In 1979, the Jervis Street clinic in Dublin treated 55 heroin users; this  
55  
56 figure rose to 213 in 1980.<sup>16</sup> In the early 1980's Dublin experienced what can be described as  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

‘an *opiate epidemic*’ For a 20 year period this was the only medical drug treatment facility in Dublin.<sup>17</sup> In 1988, following the closure of the Jervis Street hospital, the drug treatment clinic was relocated to a central Dublin site named Trinity Court. In 1990, the newly established Dublin Drug Treatment Reporting System<sup>18</sup> had 2037 opiate users treated in its expanding treatment system. Routine voluntary testing of drug users for HIV began in 1985 and over the first 2 years, 19% of those tested were diagnosed as being HIV positive. Needle and syringe provision was first provided in Ireland in 1989 in the former Eastern Health Board (EHB) AIDS Resource Centre in Dublin, in response to the heroin problem at the time.<sup>19</sup> Methadone services, mainly focused on detoxification, have been available since 1992<sup>20</sup> and was initially restricted to the capital, Dublin.<sup>17</sup> The *‘Report of the Expert Group on the Establishment of a Protocol for the Prescribing of Methadone’* was conducted in 1993<sup>20</sup>. It recommended that services should be developed using this protocol to make methadone available free of charge and national in coverage to all persons undergoing methadone treatment for opiate dependence. It also included recommendations on the type and concentration of methadone to be used, the roles of general practitioners and pharmacists, and the relationships between treatment centres, general practitioners and pharmacists. The CTL for the prescribing of methadone was established in 1993, and by June 1997, 2232 people were registered as on methadone treatment. See **Figure One**, MTP Timeline.

**Insert Figure One ‘MTP Timeline’ about here**

The Methadone Treatment Services Review Group was set up by the Department of Health and Children in 1997 to assess methadone use in the treatment of heroin dependence. In 1998 the report of the Methadone Treatment Services Review Group was published (Methadone Treatment Services Review Group, 1998).<sup>21</sup> In October 1998, legislation was



1 introduced as the ‘Misuse of Drugs (Supervision of Prescription and Supply of Methadone)  
2 *Regulations*’ which stipulated the regulatory structures for treating opioid dependent patients  
3  
4 with methadone. It was also intended to encourage GPs to become involved in the treatment  
5  
6 of drug dependence. The 1998 MTP was written to guide opioid dependent treatment delivery  
7  
8 in primary care, with detailed protocols for methadone prescribing, guidance and standards  
9  
10 for patient management and care, specialist training requirements for GPs, and protocols for  
11  
12 clinical audit. As GPs were subsequently contractually required to have specific training if  
13  
14 they wished to provide methadone treatment, the ICGP developed and delivered a number of  
15  
16 training programmes to enable GPs to fulfil these contractual requirements: Level 1 GPs were  
17  
18 provided with a daylong training programme while more experienced GPs who had been  
19  
20 working with opiate dependent patients prior to the introduction of the protocol, were offered  
21  
22 Level 2 status via a “grandfather clause”. Best practice guidelines, Working with Opiate  
23  
24 Users in Community Based Primary Care were published by the ICGP<sup>22</sup> and were issued to  
25  
26 all GPs participating in the MTP. Ongoing audit formed part of the contractual obligations  
27  
28 and the ICGP convened an audit review committee in conjunction with the EHB with the  
29  
30 purpose of developing an audit process for GPs participating in the MTP. This Joint Audit  
31  
32 Review Group (ARG)<sup>23</sup> developed a model of external audit of adherence to the MTP, which  
33  
34 was the first of its kind in Irish general practice. In addition to overseeing the audit process,  
35  
36 the ARG was given the remit of overseeing training and continuing medical education (CME)  
37  
38 for GPs participating in the MTP delivered by the ICGP. The audit process underwent a  
39  
40 number of changes since then and, since 2017, utilises a blended method of online self-audit  
41  
42 supplemented by random external audit.<sup>24</sup> The aim of this novel audit process is to assess the  
43  
44 quality of care provided by GPs to patients on methadone maintenance treatment (MMT) so  
45  
46 as to ensure that patient care meets national and international best practice standards.  
47  
48 Furthermore, it aims to enhance practice based learning and reflective practice in order to  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

improve patient care and safety, minimise methadone diversion, reduce drug overdose rates, address associated health conditions and optimise patient rehabilitative outcomes within the practice.<sup>24</sup>

Reviews of the MTP were conducted internally in 2005 by the *'Methadone Prescribing Implementation Committee'* itself and externally in 2010.<sup>25</sup> The 2010 review was an external review of the MTP undertaken to inform and maximise treatment provision and assess clinical governance and audit, referral pathways, doctor enrolment, training (Level 1 and 2) and GP coordination.<sup>25</sup> The review commented on improved prescribing and quality of independent practitioner practice, and advised the need to maximize treatment provision and referral pathways. It also commented on a number of other issues; timely responses to requests for detoxification to be reviewed as part of a service audit process, (see *'National Drugs Rehabilitation Framework'* Working Group on Drugs Rehabilitation, 2007),<sup>26</sup> rural service development, improved integration between and among services, improved clinical governance and audit, a need to review benzodiazepine prescribing (see *'Report of the Benzodiazepine Committee'* Department of Health and Children 2002),<sup>27</sup> changing urine analysis regimes, prescribing of methadone in police stations and the expansion of the number of Level 2 doctors with greater emphasis on transfer of patients from Level 2 to Level 1 doctors. Farrell and Barry<sup>25</sup> also commented on the inclusion of buprenorphine and naloxone treatment modalities, and the need to revise the title to “The Opioid Treatment Protocol”. See **Table One** and **Figure Two**.

**Insert Table One. ‘Number of patients on MMT , by year, CTL 2007-2017’ about here**

**Insert Figure Two. ‘Number on MMT and new HIV Diagnosis related to IDU’ about here**

## Service User and Provider Views

1  
2 Early studies on Irish treatment provider experiences of methadone were mixed.<sup>28</sup> Service  
3  
4 users described negative aspects centring on the patient lack of choice, humiliating  
5  
6 experiences in consuming methadone in a public space, difficulties complying with punitive  
7  
8 contracts and urine screening, and engaging with uncaring service providers.<sup>29,30,31,32</sup> Service  
9  
10 user views were reported to centre on methadone as vital to initial stages of stabilisation, and  
11  
12 that, due to the lack of alternatives to methadone, treatment regimens and patient experiences  
13  
14 became restrictive and confining, and seen as the only solution representative of the 'one size  
15  
16 fits all' phenomenon.<sup>32,33</sup> A 2012 study highlighted the influence of the GP in supporting  
17  
18 recovery.<sup>34</sup> Studies conducted in 2013 with GPs and service stakeholders<sup>35,36</sup> and in 2016  
19  
20 indicated a generally positive attitude of prescribing GPs toward methadone treatment.<sup>14</sup>  
21  
22  
23  
24  
25  
26

27 The outcomes of a national treatment and rehabilitation outcome study were reported  
28  
29 in 2009 and showed high retention rates in treatment, high levels of completion of  
30  
31 detoxification programs, significant reduction in all illicit drug use (heroin, cocaine,  
32  
33 benzodiazepine, cannabis, and illicit methadone), reduction in criminal activity, improved  
34  
35 physical and health, reduction in risky drug taking behaviour and injecting drug use (IDU),  
36  
37 improved uptake in training and employment and accommodation. Of note was that the  
38  
39 incidence of non-fatal overdose remained constant.<sup>6,37</sup> See **Table Two** and **Figure Three**.  
40  
41  
42  
43  
44  
45

46 **Insert Table Two 'Number of deaths, by year, NDRDI 2004 to 2015' here**

47  
48 **Insert Figure Three 'Drug related deaths in Ireland 2004-2015 2004 to 2015' here**

## Challenges and Developments

54  
55 Complexities around methadone provision in Ireland in the initial years centred on negative  
56  
57 attitudes toward drug users and differences of opinions around abstinence as an ultimate  
58  
59 goal.<sup>38,39,40</sup> The MTP is restricted to the medico-pharmacological treatment of stabilised drug  
60  
61

1 dependent patients, with minor reference to psychotherapeutic treatment (Kenny & O'  
2 Carroll, 2012).<sup>41</sup> The absence of a framework for the use of psychotherapeutic interventions  
3  
4 is deemed by Kenny and O Carroll<sup>41</sup> to warrant improvement.  
5  
6

7 A number of non-statutory agencies have a national brief and have in recent years  
8  
9 advocated for the expansion of treatment, the decriminalisation of drug use, and the setting up  
10  
11 of drug consumption rooms. Many provide support and advocacy groups and a number of the  
12  
13 larger agencies provide residential detoxification facilities.<sup>24</sup> A need for a widespread  
14  
15 increase of detoxification services at both community and residential levels continues, along  
16  
17 with a strengthened focus on family member support, dual diagnosis and mental health.  
18  
19

20  
21 26,32,42,43,44,45,46,47,48,49,50,51  
22  
23

24 Regional treatment structures continue to struggle to meet the demand outside of the  
25  
26 greater Dublin area, amid increasing drug related deaths where methadone is implicated  
27  
28 particularly within poly substance use, and changes in type of opioid abuse.<sup>24,52</sup> Irish studies  
29  
30 have reported on the elevated risk of drug related mortality in methadone maintenance  
31  
32 treatment during treatment initiation, transition, post drop out or discharge.<sup>53,54</sup> Rates of  
33  
34 benzodiazepine use implicated in over dose deaths are concerning for patients on  
35  
36 methadone.<sup>54,55</sup> Improved monitoring, risk assessment and methadone treatment retention  
37  
38 strategies are needed to inform national drug overdose plans and overdose prevention.<sup>52,56</sup>  
39  
40  
41  
42

43 Given the increased trend in problem drug use among ethnic groups, current Irish  
44  
45 opioid substitution guidelines are mainstream and do not refer specifically to the cultural  
46  
47 considerations of ethnic minorities such as the Traveller community.<sup>57</sup> Given the  
48  
49 complexities of methadone maintained patients in terms of multi morbidity and chronic  
50  
51 illness, O'Toole, Hambly, Cox, O' Shea and Darker<sup>58</sup> recommended a more holistic approach  
52  
53 to integrating MMT with general medical needs, the increased role of the GP Practice nurse,  
54  
55 GP incentives to train in MMT, that any formal review of MMT GP remuneration should take  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 into account the additional workload of taking care of MMT patients, and the promotion of  
2 better record-keeping with regard to chronic diseases and virology. Roy, McCarthy, Kiernan,  
3  
4 McGorrian, Keenan, Mahon, & Sweeney<sup>59</sup> more recently have recommended an increased  
5 awareness of cardiac safety guidelines, including relevant clinical and family history,  
6  
7 baseline, and trough dose ECG monitoring, should be incorporated into methadone  
8  
9 maintenance therapy protocols.  
10  
11  
12  
13

14 Several additional key developments have occurred. Enhanced multi sited data  
15 modelling will be utilised to track and monitor outcomes for policy makers, service providers  
16 and service users.<sup>60</sup> Van Hout & McElrath<sup>61</sup> have reported on the development of service user  
17  
18 fora, designed to advocate and empower service users of MMT services in Ireland within the  
19  
20 treatment care pathways and beyond. The National Poisons Information centre reported on 16  
21  
22 paediatric admissions for methadone toxicity in children under four years old in the period  
23  
24 2005-2014 and in response, a safe storage of methadone protocol has been rolled out  
25  
26 nationally.<sup>62</sup> With regard to choice, at the time of writing in August 2018, buprenorphine-  
27  
28 naloxone (Suboxone®) is prescribed to approximately 100 patients predominantly with  
29  
30 codeine dependence or other opiate based medications and with very little general practice  
31  
32 involvement.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

## 44 **Discussion**

45 The MTP in Ireland represents a model of care which uses existing primary care resources for  
46  
47 maximum benefit. As was the case in other jurisdictions, the drivers for devising the MTP in  
48  
49 Ireland, included the emergence of HIV amongst intravenous drug users and the unregulated  
50  
51 prescribing of methadone by untrained GPs and pharmacists.<sup>63,64</sup> The lack of co-ordination  
52  
53 around the delivery of services prior to the introduction of the MTP contributed to poor  
54  
55 practices, increased risk for patients and the refusal of GPs and others to look after drug  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 users. Attitudes amongst health care professionals twenty years ago prior to the inception of  
2 the Irish MTP, were that the opioid dependent patient group were problematic, difficult to  
3 deal with and would create chaos in the surgery setting. Despite evidence to the contrary,  
4 there was also a large degree of scepticism amongst health care professionals about the  
5 benefits of methadone, particularly methadone maintenance as abstinence based treatments  
6 and models of care were the norm at the time. The development of a new model of care in the  
7 form of the MTP represented recognition of harm reduction as a treatment option and  
8 provided a regulated structure for delivering MMT services. It also provided a legitimate “*opt*  
9 *out*” for GPs who did not wish to participate in the MTP; GPs could choose not to participate  
10 in the required addiction training and were consequently not eligible to provide MMT. The  
11 rationale for empowering GPs to become involved in treating drug users emerged from the  
12 UK and wider European experience.<sup>65,66</sup> A national retention in treatment study also  
13 supported the view that the treatment of patients in primary care was a protective factor for  
14 retention in treatment.<sup>67</sup>

15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34 Official endorsement for the MTP from the ICGP as the professional body for GPs  
35 was crucial in making it acceptable for GPs to get involved in this work. Prior to the  
36 introduction of the MTP, some of the messages emanating from the professional body were  
37 urging caution about the safety aspects of GP involvement in provision of MMT. The design  
38 of the protocol and regulatory controls around delivery of services gave GPs a greater sense  
39 of professional assurances and satisfaction in delivering services to drug users in their own  
40 practice.<sup>14</sup> GPs also played a key role in the delivery of services in specialist addiction centres  
41 with most of Health Services Executive (HSE) specialist services delivered by experienced  
42 Level 2 GPs. Current negotiations for a new general medical services (GMS) GP contract in  
43 Ireland include recommendations that all GPs should have completed the Level 1 training  
44 programme and be eligible to take a Level 1 contract. The ICGP continues to encourage all  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 GP training schemes to avail of the Foundation Training in Substance Misuse (Level 1  
2 training) which is available free of charge to all GP registrars and members of the ICGP.  
3

4 The evidence from patients on the MTP suggests that it has contributed significantly  
5 to the improved health and wellbeing of many drug using patients.<sup>6</sup> One of the major  
6 achievements on the MTP over the past 20 years is the containment of HIV transmission  
7 among Irish people who inject drugs (PWID). At the time of the design and introduction of  
8 the MTP there was major concern around rates of HIV infection and AIDS among small  
9 communities in the Dublin Inner City.<sup>68</sup> Over the past decade the rates of new infections  
10 have been modest and reflect the impact of the overall harm reduction strategy which  
11 combined the provision of OST and the development and expansion of NSP. The number of  
12 new HIV diagnoses amongst injecting drug users in Ireland more than halved over the period  
13 from 2004 to 2009, and only 12 new infections were notified in 2017 among PWID<sup>57</sup> An  
14 unexpected increase in HIV notifications which occurred in 2015 were related to increased  
15 injecting use of new psychoactive substances (NSP) among homeless and chaotic PWID.<sup>69</sup>  
16 This highlights the need for vigilance and lack of complacency with regard to ongoing  
17 support of harm reduction services in Ireland.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 The success with Hepatitis C infection has not been so marked and there is a  
40 substantial sub population of current and former injectors who have chronic HCV related  
41 liver disease.<sup>70,71,72</sup> The challenge of treating and managing such individuals with the new  
42 direct acting anti-retroviral (DAA) therapies will demand substantial medical resources  
43 including the transfer of HCV assessment and treatment into community addiction services  
44 and primary care.<sup>73,74</sup> The need to scale up HCV treatment to levels where it provides for  
45 “treatment as prevention” will be a major challenge over the next decade.<sup>57,73,74,75</sup> The  
46 provision of MMT in the Irish Prison Service (IPS) was of further benefit to the introduction  
47 of the MTP. Until treatment became more widespread and continuity of care could be  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

guaranteed on entry or exit from prison, opiate users who were serving sentences were offered detoxification only. This resulted in increased risk of HIV and Hepatitis C transmission as well as the increased risk of fatal overdose on release. Policy within the IPS reflected the benefits identified with OST and prisoners are now either commenced on OST where appropriate or continued on their existing treatment regime when incarcerated. Re-engagement with their community-based OST treatment service is now an important risk management element of the prison discharge policy.<sup>76</sup>

Despite the benefits of the MTP in Ireland over the past 20 years, significant challenges remain today. Access to treatment can be difficult specifically in areas outside of Dublin where waiting lists to access treatment continue to exist. It remains a concern that some PWID's can wait long periods of time to access treatment which may reflect a rigidity within the existing protocol and how it is being interpreted in certain areas. The lack of choice in terms of which substitution medication is available is another constraint as buprenorphine based medication is currently available on a limited basis only. Lack of flexibility with regard to geographical location of treatment is a factor which can become a barrier to treatment access. In some rural areas patients may have long journeys to travel to access treatment and in areas where availability of public transport is an issue this can significantly influence both access and retention in treatment.<sup>24</sup>

The single biggest challenge related to the MTP in Ireland today is the level of drug related deaths.<sup>52</sup> The ICGP convened a working group to discuss strategies to minimise drug related deaths as far back as 2007 however<sup>77</sup> drug overdose death rates have remained stubbornly high over the past decade with 224 people dying in 2015.<sup>78</sup> Most of those who died were male and were in their late 30s. The mean age of victims in 2015 was 39 years, the highest ever recorded, mainly due to the increase in deaths in those aged 55 years or older compared with 2014. The reason for this increase is not yet known, although more than half



1 of deaths among those aged 55 years or older were among women. Opioids were most  
2 commonly associated with drug-induced deaths, although they were frequently found  
3 together with other psychoactive substances, such as alcohol and prescription medicines.  
4  
5 Drug users not on OST are at greater risk from opioid and methadone related deaths. The  
6  
7 drug-induced mortality rate among adults (aged 15-64 years) in Ireland was 70 deaths per  
8  
9 million in 2015, which is more than three times the most recent European average of 21.8  
10  
11 deaths per million.  
12  
13  
14  
15  
16  
17  
18

## 19 **Conclusion**

21 Recent trends suggest a stabilisation or reduction in Irish opioid dependence rates yet opioid  
22 use remains a major public health problem. The MTP was put in place 20 years ago to enable  
23 the expansion of harm reduction services in Ireland in order to combat the high rates of HIV  
24 infections and the deaths associated with an increasing heroin epidemic in socially deprived  
25 areas of Dublin. Its introduction was controversial with mixed support from politicians,  
26 policy makers and medical professionals. The evidence based harm reduction model  
27 challenged the existing traditional abstinence model of addiction treatment. The MTP is  
28 effective in supporting the retention of patients in MMT, reducing HIV and HCV  
29 transmission and improving the health and social functioning of opioid users. Despite the  
30 successes of the MTP, many challenges remain, including; negative views of patients around  
31 how services are delivered, the stigma associated with methadone treatment, the lack of  
32 choice around substitution medication, waiting lists for treatment in certain areas and the  
33 rates of fatal overdose. The first twenty years has taken the Irish leaders in this field of  
34 medicine on a journey of improvement, advocacy and political lobbying in order to get the  
35 best possible services for patients in need of OST.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

The ICGP has been a leader in the expansion of MMT services and has been instrumental in mainstreaming the care of this marginalised patient group in general practice. It has developed accessible training and has continuously made efforts to make substance misuse a mainstream issue in primary care rather than a niche area for a small number of GP's. It has been instrumental in developing a quality assurance audit with an educational focus at its core. This structured framework of training and quality assurance along with increased remuneration, helped to allay the fears of many GPs who were reluctant to get involved in methadone treatment. Today a network of specially trained GPs provides methadone to over 10,000 patients across Ireland. Reducing stigma and normalising OST in primary care is a challenge particularly in the context of ongoing criticisms of long term maintenance treatment.<sup>79,80,81</sup> The MTP has provided a structure and support for GPs prepared to provide holistic care to drug users. As the range and type of substance misuse continues to evolve, GPs have a key role in identifying patients who may have a problem. Over the counter (OTC) codeine abuse as well as opiate analgesic dependency are a growing cause for concern and general practice is a less stigmatising environment to offer care.<sup>79</sup> Raising awareness and equipping GPs to meet the needs of their patients who may have substance use issues is of paramount importance if lives are to be saved. This area of medicine should no longer remain an '*opt in*' option; it is time to ensure that every GP is adequately trained to service their patients.

## Declarations

**Ethics Approval and consent to participate:** Not applicable, no human subjects involved.

**Consent for Publication:** Not applicable, no human subjects involved.

**Availability of Data and material:** On request from first Author.

**Competing Interests:** Authors declare no competing interests

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Funding:** Self funded.

**Authors Contributions:** MCVH was responsible for designing the study and writing the paper. ID was involved in the study design, contributed to writing, critically reviewed the manuscript, and had full control, including final responsibility for the decision to submit the paper for publication. DC was involved in the study design, contributed to writing, critically reviewed the manuscript, and had full control, including final responsibility for the decision to submit the paper for publication. All authors read and approved the final manuscript.

## References

1. Carew AM, Bellerose D, Lyons S, Long, J. Trends in treated problem opioid use in Ireland, 2002 to 2007. HRB Trends Series 7. Dublin: Health Research Board, 2009.
2. Davies HR, Nazareth I, Petersen I. Trends of People Using Drugs and Opioid Substitute Treatment Recorded in England and Wales General Practice (1994-2012). PLoS One. 2015;10:e0122626.
3. Hay G, Jaddoa A, Oyston J, Webster J, Van Hout MC, Rael dos Santos A. Estimating the prevalence of problematic opiate use in Ireland using indirect statistical methods. Dublin: National Advisory Committee on Drugs and Alcohol, 2017. Retrieved from <http://www.drugsandalcohol.ie/27233/>.
4. Kelly A, Carvalho M, Teljeur C. Prevalence of Opiate Use in Ireland 2006: A 3-Source Capture Recapture Study. Dublin: National Advisory Committee on Drugs, Stationery Office, 2003.
5. Kelly A, Carvalho M, Teljeur C. Prevalence of opioid use in Ireland 2006: a 3-source capture recapture study. Dublin: National Advisory Committee on Drugs, 2009.
6. Cox G, Comiskey C, Kelly P. ROSIE Findings 4: Summary of 1-year outcomes: Methadone modality. Dublin: National Advisory Committee on Drugs, 2007.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
7. Lyons S, Lynn E, Walsh S, Long, J. Trends in drug related deaths and deaths among drug users in Ireland, 1998 to 2005. HRB Trends Series 4. Dublin: Health Research Board, 2008.
8. National Advisory Committee on Drugs. Drug Use in Ireland and Northern Ireland 2006/2007 Drug Prevalence Survey. Regional Drugs Task Force (Ireland) and Health and Social Services Board (Northern Ireland). Dublin and Belfast: NACD & PHIRB, 2008.
9. Bates G. The drugs situation in Ireland: an overview of trends from 2005 to 2015. Liverpool: Centre for Public Health at Liverpool John Moores University, 2017.
10. Van Hout MC, Delargy I, Ryan G, Flanagan S, Gallagher H. 2015 Dependence on over the counter (OTC) codeine containing analgesics: treatment and recovery with buprenorphine naloxone. *Int J Ment Health Addict*. 2015;14:873-883.
11. Carew AM, Cafferty S, Long J, Bellerose D, Lyons S. Travellers accessing addiction services in Ireland (2007 to 2010): analysis of routine surveillance data. *J Ethn Subst Abuse*. 2013;12:339–355.
12. Claffey C, MacLachlan M, Crowley D, Van Hout MC. 2017 Exploring Irish Travellers’ experiences of Opioid Substitute Treatment: A Phenomenological Study. *Heroin Addiction and Related Clinical Problems* 2017;19:73-80.
13. Health Service Executive, Health Protection Surveillance Centre. HIV in Ireland, 2016. Dublin: Health Protection Surveillance Centre, 2016.
14. Delargy I, O’Shea M, Van Hout MC, Collins C. General Practitioners perspectives on and attitudes toward the Methadone Treatment Protocol in Ireland. *Heroin Addiction and Related Clinical Problems* 2016;18:43-50.
15. Delargy I. Survey of general practitioners participating in the methadone treatment programme. Dublin: Irish College of General Practitioners, 2008.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
16. Butler S.. Drug problems and drug policies in Ireland: a quarter of a century reviewed. Administration. 1991;39:210-233.
  17. Butler S. Alcohol, drugs and health promotion in modern Ireland. Dublin: IPA, 2002.
  18. O'Hare A. O'Brien M. Treated Drug Misuse in the Greater Dublin Area 1990. Dublin: Health Research Board, 1992.
  19. McVeigh J, Bates G, Hearne E, Van Hout MC. Community pharmacist experiences of providing needle and syringe programmes in Ireland. Res Social Adm Pharm. 2017;13:767-777.
  20. Department of Health. Report on the Expert Group on the Establishment of Protocol for the Prescribing of methadone. Dublin: Department of Health, 1993.
  21. Methadone Treatment Services Review Group. Report of the Methadone Treatment Services Review Group. Dublin: Department of Health and Children, 1998.
  22. ICGP. 2008 Publication Working with Opiate Users in Community Based Primary Care DOHC Hep C Guidelines. <http://health.gov.ie/national-patient-safety-office/ncec/national-clinical-guidelines/>. Accessed June 20<sup>th</sup> 2018.
  23. Irish College of General Practitioners. Terms of reference for Joint HSE/ICGP Audit Review Group (ARG) – Methadone treatment protocol – (Revised and approved December 2011. Dublin: Irish College of General Practitioners, 2012.
  24. Van Hout MC, Crowley D, McBride A, Delargy I. Piloting online self-audit of methadone treatment in Irish general practice: results, reflections and educational outcomes. BMC Med Educ. 2018;18:153.
  25. Farrell M, Barry J. The Introduction of the Opioid Treatment Protocol. Dublin: Health Service Executive, 2010. [www.drugsandalcohol.ie/14458](http://www.drugsandalcohol.ie/14458) Accessed June 20<sup>th</sup> 2018.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
26. Department of Community, Rural and Gaeltacht Affairs. Report of the Working Group on Drugs Rehabilitation Dublin: Department of Community, Rural and Gaeltacht Affairs, 2007.
  27. Benzodiazepine Committee. Report of the Benzodiazepine Committee. Dublin: Department of Health and Children, 2002. [www.drugsandalcohol.ie/5348](http://www.drugsandalcohol.ie/5348) Accessed June 20<sup>th</sup> 2018.
  28. Langton D, Hickey A, Bury G, Smith M, O'Kelly F, Barry J, Sweeney B, Bourke M. Methadone maintenance in general practice: impact on staff attitudes. *Ir J Med Sci.* 2000;169:133-136.
  29. O'Connor S. Pharmaceutical Services for Patients in Methadone Treatment in Ireland and the Introduction of the Methadone Protocol Scheme. University of Dublin, Thesis for the degree of PhD. Available at the Library in Trinity College Dublin, 2002.
  30. Cox G, Lawless M. Maintaining or enabling? Evaluation of a methadone prescribing service in Dublin City. In *Pieces of the jigsaw: six reports addressing homelessness and drug use in Ireland.* Dublin: Merchants Quay Ireland, 2004 p. 79-140.
  31. UISCE. Methadone: What's the Story? Dublin: Union for Improved Services, Communication and Education (Internal Publication), 2003.
  32. Van Hout MC, Bingham T. Methadone Maintenance and Special Community Employment schemes: A study of Irish participants' views. *J Vocat Rehabil.* 2012;37:63-73.
  33. Lawless K. Listening and learning: evaluation of Special Community Employment programmes in Dublin North East. Dublin: Dublin North East Drugs Task Force, 2006.
  34. Latham L. Methadone treatment in Irish general practice: voices of service users. *Ir J Psychol Med.* 2012;29:147-156.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
35. Van Hout MC, Bingham T. A qualitative study of prescribing doctor experiences of methadone maintenance treatment. *Int J Ment Health Addict*. 2013;12:227-242.
  36. Van Hout MC, Bingham T. Service stakeholders' perspectives on methadone maintenance treatment, Special Community Employment schemes and client recovery pathways. *J Vocat Behav*. 2014;40:49-58.
  37. Comiskey CM, Kelly P, Leckey Y, McCulloch L, O'Duill B, Stapleton RD, White E. Research Outcome Study In Ireland Evaluating Drug Treatment Effectiveness (Rosie). Maynooth: National University Of Ireland, 2009.  
<https://www.drugsandalcohol.ie/11542/1/ROSIE3-YearReport.pdf> Accessed June 20<sup>th</sup> 2018.
  38. MacGreil M, Prejudice in Ireland Revisited: based on a national survey of inter group attitudes in the Republic of Ireland. Dublin: Survey and Research Unit, St. Patrick's College Maynooth, 1996.
  39. Bryan A, Moran R, Farrell E, O'Brien M. Drug-related Knowledge, Attitudes and Beliefs in Ireland: report of a nationwide survey. Dublin: Health Research Board, 2000.
  40. Carlin T. A therapeutic response or 'a free fix'? The Irish experience of methadone maintenance treatment. *Ir Prob J*. 2005;2:87-94.
  41. Kenny K, O'Carroll A. The use of psychotherapeutic interventions by primary care GPs in Ireland in the treatment of their methadone patients: a grounded theory study. *Ir J Med Sci*. 2012;181:43-48.
  42. Mannix M. Drug and Alcohol Detoxification Services A Needs Assessment for Cork and Kerry 2005. Dublin: Dept. of Public Health HSE South, 2006.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
43. Corrigan D, O'Gorman A. Report of the HSE Working Group on Residential Treatment and Rehabilitation Substance Users. Dublin: Health Service Executive, 2008.
  44. Department of Community, Rural and Gaeltacht Affairs. National Drugs Strategy 2009–2016. Dublin: Department of Community, Rural and Gaeltacht Affairs, 2009.
  45. Doyle J, Ivanovic J. National Drugs Rehabilitation Framework Document, National Drugs Rehabilitation Implementation Committee. Dublin: Health Services Executive, 2010.
  46. McDonnell A, Van Hout MC. Maze and minefield: a grounded theory of opiate self-detoxification in rural Ireland. *Drugs and Alcohol Today*. 2010;10:24–31.
  47. McDonnell A, Van Hout M. Heroin detoxification-seeking, a grounded theory of process and practicalities. *The Grounded Theory Review*. 2011;10:17-14.
  48. Progression Routes Initiative. Community detoxification protocols: benzodiazepines. Dublin: Progression Routes Initiative, 2011a.
  49. Progression Routes Initiative. Community detoxification protocols: methadone. Dublin: Progression Routes Initiative, 2011b.
  50. Ana Liffey Drug Project. Community detoxification protocols: methadone. Dublin: Ana Liffey Drug Project, 2016a. <https://www.drugsandalcohol.ie/26888/1/Community-Detox-Guidelines-Methadone-2017.pdf> Accessed June 20<sup>th</sup> 2018.
  51. Ana Liffey Drug Project. National community detoxification: methadone guidelines. Dublin: Ana Liffey Drug Project, 2016b. <https://www.drugsandalcohol.ie/26888/> Accessed June 20<sup>th</sup> 2018.
  52. Van Hout MC, Crowley D, Barry A, Collins C, Delargy I. Characteristics of methadone related overdose deaths and comparisons between those dying on and off



1 methadone substitution treatment: a national cohort study. *Heroin Addiction and*  
2 *Related Clinical Problems* 2018;20:37-44.  
3

4  
5 53. Truskowska E, McCarron P, Konovalov P, Galander T, Lyons S, Keenan E, Smyth B.  
6  
7 Case-Control Study of Risks and Causes of Death Amongst Opioid Dependent  
8  
9 Patients on Methadone Maintenance Treatment. *Can J Addict.* 2015;6:18-26.  
10

11  
12 54. Cousins G, Boland F, Courtney B, Barry J, Lyons S, Fahey T. Risk of mortality on  
13  
14 and off methadone substitution treatment in primary care: a national cohort study.  
15  
16 *Addiction.* 2016;111:73-82.  
17

18  
19 55. Gilroy D, O'Brien S, Barry J, Ivers J, Whiston L, Keenan E, Darker C.  
20  
21 Benzodiazepine use in a methadone maintained opiate dependent cohort in Ireland.  
22  
23 *Heroin Addiction and Related Clinical Problems* 2014;16:23-30.  
24

25  
26 56. Darker C, Ho J, Kelly G, Whiston L, Barry J. Demographic and clinical factors  
27  
28 predicting retention in methadone maintenance: results from an Irish cohort. *Ir J Med*  
29  
30 *Sci.* 2016;185:433-441.  
31

32  
33 57. Health Service Executive. Clinical guidelines for opioid substitution treatment.  
34  
35 Dublin: Health Service Executive, 2017.  
36  
37 [https://www.hse.ie/eng/services/publications/Primary/clinical-guidelines-for-](https://www.hse.ie/eng/services/publications/Primary/clinical-guidelines-for-opioidsubstitution-treatment.pdf)  
38  
39 [opioidsubstitution-treatment.pdf](https://www.hse.ie/eng/services/publications/Primary/clinical-guidelines-for-opioidsubstitution-treatment.pdf). Accessed June 20<sup>th</sup> 2018.  
40  
41

42  
43 58. O'Toole J, Hambly R, Cox AM, O'Shea B, Darker C. Methadone-maintained patients  
44  
45 in primary care have higher rates of chronic disease and multimorbidity, and use  
46  
47 health services more intensively than matched controls. *Eur Gen Pract.* 2014;20:275–  
48  
49 320.  
50  
51

52  
53 59. Roy AK, McCarthy C, Kiernan G, McGorrian C, Keenan E, Mahon NG, Sweeney B.  
54  
55 Increased incidence of QT interval prolongation in a population receiving lower doses  
56  
57 of methadone maintenance therapy. *Addiction.* 2012;107:1132-1139.  
58  
59  
60  
61

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
60. Comiskey CM, Snel A. Using clients' routine urinalysis records from multiple treatment systems to model five-year opioid substitution treatment outcomes. *Subst Use Misuse*. 2016;51:498–507.
  61. Van Hout MC, McElrath K. Service user involvement in drug treatment programs: Barriers to implementation and potential benefits for client recovery. *Drugs*. 2012;19:474-483.
  62. Horan A, Gooney M, Van Hout MC. Safe storage of Methadone in the home: results from an Irish audit of service users and pharmacists. *Heroin Addiction and Related Clinical Problems*. 2016;18:31-40.
  63. Bruce RD. Methadone as HIV prevention: high volume methadone sites to decrease HIV incidence rates in resource limited settings. *Int J Drug Pol*. 2010;21:122–124.
  64. Zaric GS, Brandeau ML, Barnett PG. Methadone Maintenance and HIV Prevention: A Cost-Effectiveness Analysis. *Manag Sci*. 2000;46:1013–1031.
  65. Keenan E, Barry J. Managing drug misuse in general practice. Republic of Ireland has set up scheme to regulate methadone prescribing by GPs. *BMJ*. 1999;319:1497.
  66. Wilson P, Watson R, Ralston GE. Methadone maintenance in general practice: patients, workload, and outcomes. *BMJ*. 1994;309:641–644.
  67. Mullen L, Barry J, Long J, Keenan E, Mulholland D, Grogan L, Delargy I. A national study of the retention of Irish opiate users in methadone substitution treatment. *Am J Drug Alcohol Abuse*. 2012;38:551-558.
  68. O'Kelly FD, & O'Kelly CM. The natural history of injecting drug use: A 25-year longitudinal study of a cohort of injecting drug users in inner city Dublin. *Ir J Med Sci*. 2012;181:541–548.
  69. Giese C, Igoe D, Gibbons Z, Hurley C, Stokes S, McNamara S., Ennis O, O'Donnell K, Keenan E, De Gascun C, Lyons F, Ward M, Danis K, Glynn R, Waters A,

1 Fitzgerald M, outbreak control team. on behalf of the outbreak control. Injection of  
2 new psychoactive substance snow blow associated with recently acquired HIV  
3 infections among homeless people who inject drugs in Dublin, Ireland, 2015. Euro  
4 Surveill. 2015;20:30036.  
5  
6  
7  
8

9  
10 70. Keegan D, Crowley D, Laird E, Van Hout, MC. Prevalence and risk factors for  
11 hepatitis C viral infection amongst a cohort of Irish drug users attending a drug  
12 treatment centre for agonist opioid treatment (AOT). Heroin Addiction and Related  
13 Clinical Problems, 2017;19:47–56. <http://www.drugsandalcohol.ie/25815/>  
14  
15  
16  
17

18  
19 71. Smyth BP, Keenan E, O'Connor JJ. Bloodborne viral infection in Irish injecting drug  
20 users. Addiction. 1998;93:1649–1656. [https://doi.org/10.1046/j.1360-](https://doi.org/10.1046/j.1360-0443.1998.931116493.x)  
21  
22  
23  
24  
25  
26

27 72. Thornton L, Murphy N, Jones L, Connell J, Dooley S, Gavin S, Hunter K, Brennan A.  
28 Determination of the burden of hepatitis C virus infection in Ireland. Epidemiol  
29 Infect. 2012;140:1461-1468.  
30  
31  
32  
33

34 73. Department of Health. Public Health Plan for the Pharmaceutical Treatment of  
35 Hepatitis C. Dublin: Department of Health, 2014. [http://health.gov.ie/wp-](http://health.gov.ie/wp-content/uploads/2015/07/Final-Public-Health-Plan-for-the-Pharmaceutical-Treatment-of-Hep-C-Final-Copy-Circulated-July-2015.pdf)  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

54 74. Health Service Executive. National Hepatitis C Strategy 2011-2014. Dublin: Health  
55 Service Executive, 2012.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

66 75. Department of Health. Hepatitis C Screening (NCEC National Clinical Guideline No.  
67 15). Dublin: Department of Health, 2017. [http://health.gov.ie/national-patient-safety-](http://health.gov.ie/national-patient-safety-office/ncec/national-clinical-guidelines)  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
76. Irish Prison Service. Irish Prison Service Health Care Standards. Dublin: Irish Prison Service, 2009.
77. Drugnet Ireland. Drug-related deaths and strategies for prevention. Drugnet Ireland. 2007;21:2-3.
78. Health Research Board. National Drug-Related Deaths Index 2004 to 2015 data. Dublin: Health Research Board, 2017.
79. Latowsky M, Kallen E. Mainstreaming methadone maintenance treatment: the role of the family physician. CMAJ. 1997;157:395-398.
80. McPhee I, Brown A, Martin C. Stigma and perceptions of recovery in Scotland: a qualitative study of injecting drug users attending methadone treatment. Drugs and Alcohol Today. 2013;13:244–257. <https://doi.org/10.1108/DAT-05-2013-0022>.
81. Woo J, Bhalerao A, Bawor M, Bhatt M, Dennis B, Mouravska N, Zielinski L, Samaan Z. “Don’t Judge a Book by Its Cover”: A Qualitative Study of Methadone Patients’ Experiences of Stigma. Subst Abuse. 2017;11:1-12.


**Table One. 'Number of patients on MMT by year, CTL 2007-2017'**

<b>Year</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>2014</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>
<b>Total numbers on MMT</b>	8537	8718	9047	9266	9251	9419	9655	9764	9940	10087	10316
<b>Numbers of MMT with GPs</b>	3047	3085	3199	3360	3477	3589	3812	3951	4097	4184	4220
<b>New HIV Cases related to IDU</b>	55	40	30	22	16	17	21	28	49	20	17


**Table Two ‘Number of deaths, by year, NDRDI 2004 to 2015’**

<b>Year</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>2014</b>	<b>2015</b>
<b>All Deaths</b>	431	503	554	620	628	656	607	645	660	704	719	695
<b>Poisonings</b>	266	301	326	387	386	373	339	377	356	400	364	348

(N = 7,422)



**Figure One 'MTP Timeline'**



**Figure Two. ‘Number on MMT and new HIV Diagnosis related to IDU’**



**Figure Three ‘Drug related deaths in Ireland 2004-2015 2004 to 2015’**

