

*Citation for published version:*

del Barrio, L., Martin-de-Saavedra, MD, Romero, A., Parada, E., Egea, J., Avila, J., McIntosh, JM, Wonnacott, S & Lopez, MG 2011, 'Neurotoxicity Induced by Okadaic Acid in the Human Neuroblastoma SH-SY5Y Line Can Be Differentially Prevented by alpha 7 and beta 2* Nicotinic Stimulation', *Toxicological Sciences*, vol. 123, no. 1, pp. 193-205. <https://doi.org/10.1093/toxsci/kfr163>

DOI:

[10.1093/toxsci/kfr163](https://doi.org/10.1093/toxsci/kfr163)

Publication date:

2011

Document Version

Peer reviewed version

[Link to publication](#)

This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *Toxicological Sciences* following peer review. The definitive publisher-authenticated version del Barrio, L., Martin-de-Saavedra, M. D., Romero, A., Parada, E., Egea, J., Avila, J., McIntosh, J. M., Wonnacott, S. and Lopez, M. G., 2011. Neurotoxicity Induced by Okadaic Acid in the Human Neuroblastoma SH-SY5Y Line Can Be Differentially Prevented by alpha 7 and beta 2* Nicotinic Stimulation. *Toxicological Sciences*, 123 (1), pp. 193-205 is available online at: <http://dx.doi.org/10.1093/toxsci/kfr163>

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1
2
3 **Neurotoxicity induced by okadaic acid in the human neuroblastoma SH-SY5Y line can**
4 **be differentially prevented by $\alpha 7$ and $\beta 2^*$ nicotinic stimulation**
5
6
7
8
9

10
11 ^{*,†,¶}Laura del Barrio¹, ^{*,#}María Dolores Martín-de-Saavedra¹, ^{*,†,#}Alejandro Romero, ^{*,†}Esther
12 Parada, ^{*,†,#}Javier Egea, [‡]Jesús Avila, [§]J Michael McIntosh, [¶]Susan Wonnacott and
13
14
15
16 ^{*,†,#}Manuela G. López
17
18
19
20

21 ^{*}Instituto Teófilo Hernando, [†]Departamento de Farmacología y Terapéutica; Facultad de
22 Medicina, Universidad Autónoma de Madrid, Spain.
23
24
25

26 [‡]Centro de Biología Molecular Severo Ochoa. Universidad Autónoma de Madrid. Spain
27

28 [§]Department of Biology and Psychiatry, University of Utah, Salt Lake City, Utah, USA,
29

30 [¶]Department of Biology and Biochemistry, University of Bath, Bath, United Kingdom.
31
32

33 [#]HIV Unit. Hospital La Paz/Autónoma University School of Medicine. IdiPAZ
34
35
36
37

38 ¹*These authors contributed equally*
39

40 **Running title:** nAChR reduce okadaic acid neurotoxicity
41
42

43
44 **Corresponding author:** Manuela G. López
45

46 Departamento de Farmacología, Facultad de Medicina,
47

48 Universidad Autónoma de Madrid
49

50 C/ Arzobispo Morcillo 4 E-28029 Madrid. Spain
51

52 E-mail: manuela.garcia@uam.es
53

54 Phone: +34-914975386 Fax: +34-914973120
55
56
57
58
59
60

ABSTRACT

A good model of neuronal death that reproduces the characteristic tau (τ) hyperphosphorylation of Alzheimer's disease is the use of okadaic acid (OA). The aim of this study was to determine the contribution of $\alpha 7$ and $\beta 2^*$ nicotinic acetylcholine receptor (nAChR) subtypes to neuroprotection against OA in the SH-SY5Y cell line, by using the selective $\alpha 7$ and $\beta 2^*$ nAChR agonists PNU 282987 and 5Iodo-A85380, respectively. The results of this study show that both $\alpha 7$ and $\beta 2^*$ nAChR can afford neuroprotection against OA-induced neurotoxicity. Protection mediated by $\alpha 7$ nAChRs was independent of Ca^{2+} and involved the intracellular signaling pathway JAK2/PI3K/Akt. When Ca^{2+} entry was promoted through the $\alpha 7$ nAChR by using the $\alpha 7$ -selective positive allosteric modulator PNU 120596, protection was lost. By contrast, protection mediated by $\beta 2^*$ nAChRs was Ca^{2+} -dependent and implicated the signaling pathways PI3K/Akt and ERK1/2. Both $\alpha 7$ and $\beta 2^*$ nAChR activation converged on down-regulation of GSK3 β and reduction of τ phosphorylation in cells undergoing cell death induced by OA. Therefore, targeting nAChR could offer a strategy for reducing neurodegeneration secondary to hyperphosphorylation of protein τ .

Key words: Okadaic acid, SH-SY5Y neuroblastoma, nicotinic receptors, hyperphosphorylation of Tau, PNU 282987, 5IA 85380.

INTRODUCTION

Alzheimer Disease (AD) is characterized by a progressive loss of memory, often with deterioration of language as well as deficits in visual and motor coordination. At the cellular level, it has been shown that cell death (Cummings 2004) occurs especially in cholinergic neurons (Whitehouse *et al.* 1982) and that this neuronal death is related to the loss of memory in AD patients (Haass and Selkoe 2007). At the molecular level, AD is characterized by the presence of tangles rich in the intracellular microtubule-associated protein tau (τ) (Lee and Trojanowski 1992) and extracellular deposits of amyloid peptides that form senile plaques (Price *et al.* 1991). A link between these two proteins has been demonstrated since beta-amyloid can trigger tau phosphorylation and the subsequent degeneration of the affected neurons (Busciglio *et al.* 1995; Alvarez *et al.*, 2002; Liu *et al.*, 2004). It has also been shown that beta-amyloid fibrils can induce neurotoxicity and tau hyperphosphorylation without plaque formation; it therefore seems that beta-amyloid pathology could precede hyperphosphorylation of tau and the formation of neurofibrillary tangles and neurodegeneration (Busciglio *et al.*, 1995)

In this study we have focused on τ hyperphosphorylation as a mechanism for neuronal death (Irizarry *et al.* 1997; Pizzi *et al.* 1995). In the brains of AD patients, there is a high percentage of hyperphosphorylated τ protein (Avila 2000) and an accumulation, in certain neurons, of paired helical filaments (PHF) in which hyperphosphorylated τ is the major protein present (Grundke-Iqbal *et al.* 1986; Grundke-Iqbal *et al.* 1988). Destabilization of microtubules and the consequent decrease in axonal transport is believed to be responsible for retrograde degeneration of neurons in AD (Alonso *et al.* 1994; Alonso *et al.* 1997; De Vos *et al.* 2008) and also for cell death in “*in vitro*” models of τ hyperphosphorylation (Cowan *et al.*; Perez *et al.* 2002; Tanaka *et al.* 1998). Inhibition of phosphatases that act on some abnormally phosphorylated sites characteristic of AD (Gong *et al.* 1994) may also lead to τ

1
2
3 protein hyperphosphorylation (Tanaka *et al.* 1998). These phosphatases include the Ser/Thr
4 PP-1, PP-2A and PP-2B (Gong *et al.* 1994; Wang *et al.* 1995) and their activity has been
5
6 shown to be diminished in brain tissue from Alzheimer's disease patients (Gong *et al.* 1995;
7
8 Gong *et al.* 1993).

9
10
11
12 Okadaic acid, a seaweed toxin, inhibits phosphatases with the preference PP2A > PP1
13 > PP2B. A consequence of its action is τ hyperphosphorylation "*in vivo*" (Tian *et al.* 2004)
14
15 and "*in vitro*" (Tanaka *et al.* 1998; Uberti *et al.* 1997). It therefore provides a good model that
16
17 reproduces τ hyperphosphorylation and cell death. In this study we have used OA in the
18
19 human neuroblastoma cell line SH-SY5Y to model aspects of cell damage relevant to
20
21 Alzheimer's disease (Arias *et al.* 2005; Cowan *et al.*; Perez *et al.* 2002; Tanaka *et al.* 1998), in
22
23 order to investigate neuroprotective strategies.

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
There is a body of evidence for nAChRs exerting neuroprotective effects against a
variety of insults (Dajas-Bailador and Wonnacott 2004; Picciotto and Zoli 2008). With
respect to Alzheimer's disease, nicotinic agonists have been reported to be effective against
 $A\beta$ -induced toxicity "*in vivo*" and "*in vitro*" (Arias *et al.* 2005; Kihara *et al.* 1998). Their
efficacy in τ hyperphosphorylation models is less well documented.

The majority of studies have highlighted the $\alpha 7$ nAChR as the main nAChR subtype
responsible for neuroprotective effects mediated by nicotinic agonists (Arias *et al.* 2005;
Bitner *et al.* 2009; Haydar and Dunlop). However, non $\alpha 7$ nAChR subtypes have also been
found to contribute to neuroprotection (Tizabi *et al.* 2004), depending on the cell type and
toxic stimulus. Because the roles of different nAChR subtypes are not fully understood, we
have exploited subtype-selective ligands, namely the $\alpha 7$ nAChR agonist PNU 282987
(Bodnar *et al.* 2005), the β^* -selective agonist 5-Iodo-A-85380 (5IA 85380) (Mukhin *et al.*
2000) and the $\alpha 7$ nAChR positive allosteric modulator PNU 120596 (Hurst *et al.* 2005) in this
study

1
2
3 The results demonstrated nAChR-mediated neuroprotection against OA. Protection
4 mediated by $\alpha 7$ nAChRs was independent of Ca^{2+} and involved the intracellular signaling
5 pathway JAK2/PI3K/Akt. Sustained activation of $\alpha 7$ nAChR to promote Ca^{2+} entry was
6 achieved with the positive allosteric modulator but under these conditions protection was lost.
7
8 In contrast, protection mediated by $\beta 2^*$ nAChRs was Ca^{2+} -dependent and implicated the
9 signaling pathways PI3K/Akt and ERK1/2. Both $\alpha 7$ and $\beta 2^*$ nAChR activation converged on
10 the inhibition of GSK-3 β , and consequently τ phosphorylation, to afford neuroprotection.
11
12
13
14
15
16
17
18
19
20

21 MATERIALS AND METHODS

22 *Materials*

23
24
25
26
27
28 PNU 282987, 5IA 85380, PNU 120596, AG 490, LY 294002, PD 98059 were
29 purchased from Tocris Cookson Inc. (Bristol, UK). Okadaic acid, α -bungarotoxin (BGT),
30 dihydro- β -erythroidine (DH β E), methyllycaconitine (MLA), mecamylamine and AR-A01448
31 were purchased from Sigma-Aldrich (St. Louis, MO). α -conotoxin-MII and α -conotoxin-
32 AuIB were synthesized as previously described (Luo et al., 1998). Alpha-conotoxin AuIB
33 selectively blocks $\alpha 3\beta 4$ nicotinic acetylcholine receptors and nicotine-evoked norepinephrine
34 release (Lou et al, 1998).
35
36
37
38
39
40
41
42
43
44
45
46
47

48 *Culture of SH-SY5Y Cells*

49
50 SH-SY5Y cells were maintained in a 1:1 mixture of F-12 Nutrient Mixture (Ham12)
51 (Sigma Aldrich, Madrid, Spain) and Eagle's minimum essential medium (EMEM)
52 supplemented with 15 non-essential amino acids, 1 mM sodium pyruvate, 10% heat-
53 inactivated fetal bovine serum (FBS), 100 units/ml penicillin, and 100 $\mu\text{g}/\text{ml}$ streptomycin
54 (reagents from Invitrogen, Madrid, Spain). Cultures were seeded into flasks containing
55 supplemented medium and maintained at 37°C in a humidified atmosphere of 5% CO_2 and
56
57
58
59
60

1
2
3 95% air. For assays, SH-SY5Y cells were sub-cultured in 48-well plates at a seeding density
4 of 1×10^5 cells per well. Cells were treated with the drugs before confluence in F-12/EMEM
5 with 1% FBS. These cells, when undifferentiated, express functional nicotinic receptors
6 (Dajas-Bailador *et al.* 2002a). All cells used in this study were used at a low passage number
7 (<13).
8
9

10 11 12 13 14 15 16 17 ***Measurement of lactate dehydrogenase activity***

18
19
20 Extracellular and intracellular lactate dehydrogenase (LDH) activity was measured
21 spectrophotometrically using a cytotoxicity cell death kit (Roche Diagnostics, Mannheim,
22 Germany) according to the manufacturer's instructions. Total LDH activity was defined as the
23 sum of intracellular and extracellular LDH activity; released LDH was defined as the
24 percentage of extracellular compared with total LDH activity.
25
26
27
28
29
30
31
32
33

34 ***Calcium fluorimetry.***

35
36 Increases in intracellular Ca^{2+} in confluent cultures of SH-SY5Y cells grown in 96-
37 well plates were monitored as described by Dajas-Bailador *et al.* (2002a). In brief, SH-SY5Y
38 cells were washed twice with Tyrode's salt solution (TSS: in mM 137 NaCl, 2.7 KCl, 1.0
39 MgCl_2 , 2.5 CaCl_2 , 0.2 NaH_2PO_4 , 12 NaHCO_3 , 5.5 glucose; pH 7.4) and incubated with the
40 membrane-permeable Ca^{2+} -sensitive dye fluo-3 AM (10 μM) and 0.02% pluronic F127 for 1 h
41 at room temperature in the dark. Cells were then washed twice with TSS, before adding 80 μL
42 buffer, with or without nAChR antagonists or PNU 120596, per well. After 10 min, basal
43 fluorescence (excitation 485 nm, emission 520 nm) was recorded for 4 s using a Fluoroskan
44 Ascent fluorescence plate reader (Labsystems, Helsinki, Finland). nAChR agonists (20 μL)
45 were added using an automatic dispenser and changes in fluorescence were monitored for a
46 further 10 s. To normalize fluo-3 AM signals, responses from each well were calibrated by
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 determination of the maximum and minimum fluorescence values by addition of 0.2% Triton-
4 X100 (F_{\max}) followed by 40 mM MnCl_2 (F_{\min}). Maximum fluorescence responses were
5
6
7
8 calculated as a percentage of $F_{\max} - F_{\min}$.
9

10 11 12 ***Immunoblotting***

13
14
15 SH-SY5Y cells were washed once with cold phosphate-buffered saline and lysed in
16
17 100 μL ice-cold lysis buffer (1% Nonidet P-40, 10% glycerol, 137 mmol/L NaCl, 20 mmol/L
18
19 Tris-HCl, pH 7.5, 1 $\mu\text{g}/\text{mL}$ leupeptin, 1 mmol/L phenylmethylsulfonyl fluoride, 20 mmol/L
20
21 NaF, 1 mmol/L sodium pyrophosphate, and 1 mmol/L Na_3VO_4). Protein (30 μg) from the cell
22
23
24 lysates was resolved by sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–
25
26 PAGE) and transferred to PVDF membranes (Amersham). Membranes were incubated with
27
28 the following antibodies: anti-extracellular regulated kinase 1/2 (ERK 1/2) and anti-phospho-
29
30 ERK1/2 at 1:1000 (Santa Cruz Biotechnology Inc., Heidelberg, Germany); anti-JAK-2 , anti-
31
32 phospho-Tyr-211-JAK-2, Anti-phospho-Thr-308-Akt, Anti-Akt, Anti-phospho-Ser-9-GSK3-
33
34 β , Anti-phospho-Tyr-216-GSK3- β and Anti-GSK-3 β at 1:1000 (Cell Signalling, Izasa S.A.,
35
36 Barcelona, Spain); anti- β -actin at 1:100000 (Sigma, Madrid, Spain). Membranes were also
37
38
39 incubated with Anti-phospho-Ser-396/404-Tau and Anti-Tau at 1:1000 which were
40
41
42 previously used by Lucas et al. (2001). Appropriate peroxidase-conjugated secondary
43
44
45 antibodies at 1:10000 were used to detect proteins by enhanced chemiluminescence.
46
47
48
49

50 51 ***Cytoskeleton staining***

52
53 SH-SY5Y cells were plated on coverslips: 24-48 h after plating the cells, they were
54
55 separated into 4 groups: two were treated with EMEM, one with 10 μM PNU282987 and
56
57 another one with 100 μM 5IA 85380 for 24 h. Then, the control group was maintained in
58
59 EMEM and the rest of the groups were treated with okadaic acid for 5 h. After exposure of
60

1
2
3 the cells to the different treatments, they were rinsed three times with PBS (NaCl 9g/L,
4 NaH₂PO₄ 10 mM, K₂HPO₄ 10 mM). Then, they were fixed in 2% paraformaldehyde in PBS
5
6 for 15 min and washed three more times with PBS. Cells were permeabilized by exposure to
7
8 0.1% triton in PBS for 1 min. Preparations were washed several times with PBS and then
9
10 incubated at room temperature with rhodamine-labeled phalloidin (Sigma-Aldrich, Madrid,
11
12 Spain) 1:1000 in PBS. Coverslips were then thoroughly washed with PBS. In the second
13
14 rinsing, Hoechst (Invitrogen, Madrid, Spain) was added (1 μL/mL) in order to mark nuclei
15
16 and after the third rinsing the coverslips were mounted in glycerol-PBS (1:1 ; vol/vol). Images
17
18 were taken with a confocal microscope (TCS SPE; Leica, Wetzlar, Germany).
19
20
21
22
23
24
25
26

27 *Statistical Analysis*

28
29 Statistically significant differences between groups were determined by an analysis of
30
31 variance followed by a Newman-Keuls post hoc analysis. The level of statistical significance
32
33 was taken at $p < 0.05$.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

Effect of different nicotinic agonists and the $\alpha 7$ allosteric modulator PNU 120596 on OA-induced toxicity

Exposure of SH-SY5Y cells to 30 nM OA for 16 h increased LDH release from 7-10 % (basal conditions) to 36-42 % (OA treatment), consistent with the induction of modest levels of cell death (Fig 1). To determine if nAChRs are able to protect against OA-induced toxicity we incubated SH-SY5Y cells with either the $\alpha 7$ or $\beta 2^*$ selective agonist PNU 282987 (Parada *et al.*) or 5IA 85380, respectively, or with the $\alpha 7$ allosteric modulator PNU 120596. With pre-incubation increasing concentrations of PNU 282987, for 24 h before the addition of OA (see protocol at the top of fig. 1), we observed a gradual reduction in cell death measured as LDH released into the extracellular medium. PNU 282987 exerted a significant but partial protection against OA-induced toxicity at 10 and 30 μM (fig. 1 A). 5IA 85380 also produced significant protection at 100 μM (fig. 1 B). The $\alpha 7$ positive allosteric modulator PNU 120596 alone showed no protection at any of the concentration assayed (0.3 to 10 μM) (fig. 1 C)

Intracellular Ca^{2+} increases mediated by PNU 282987, 5IA 85380 and PNU 120596 plus PNU 282987.

SH-SY5Y cells express $\alpha 3$, $\alpha 5$, $\alpha 7$, $\beta 2$, and $\beta 4$ nAChR subunits (Gentry and Lukas 2002; Peng *et al.* 1994). In order to determine the effects of PNU 282987 and 5IA 85380 on intracellular Ca^{2+} levels and their selectivity on nAChR subtypes, we measured Ca^{2+} signals in SH-SY5Y cells loaded with the dye Fluo-3. For these experiments we employed concentrations of the agonists that showed protection in the previous section in order to gather information about the possible neuroprotective mechanisms of action.

PNU 282987 alone did not induce a significant increase in $[\text{Ca}^{2+}]_c$ at the neuroprotective concentration of 10 μM . However, when the agonist was added to cells pre-

1
2
3 incubated with the allosteric modulator PNU 120596, a significant increase in fluorescence
4 was observed, consistent with increased cytosolic Ca^{2+} . These $[\text{Ca}^{2+}]_c$ increases were
5 mediated by activation of nAChR since the signal was significantly attenuated by the non-
6 selective nAChR antagonist mecamylamine. In particular, these Ca^{2+} signals could be
7 attributed to stimulation of the $\alpha 7$ subtype nAChR since the selective $\alpha 7$ nAChR antagonists
8 α BGT and MLA completely inhibited the increases in fluorescence (fig. 2 A).
9

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

5IA 85380, at the protective concentration of 100 μM , produced a significant intracellular Ca^{2+} signal by itself (fig. 2B). Selective $\beta 2^*$ antagonists such as DH β E and α -conotoxin MII or mecamylamine were able to significantly block Ca^{2+} signals induced by 5IA 85380. However, $\alpha 3\beta 4$ (α -conotoxin AuIB) or $\alpha 7$ (BGT) antagonists did not modify the $[\text{Ca}^{2+}]_c$ responses mediated by 5IA 85380.

Therefore, under these experimental conditions, 5IA 85380 appears to be activating heteromeric $\beta 2$ -containing nAChRs while PNU 282987 is activating $\alpha 7$ nAChRs.

Ca^{2+} -dependence in the protective effects of 5IA 85380 and PNU 282987

To determine the influence of Ca^{2+} entry mediated by $\alpha 7$ and $\beta 2^*$ nAChRs in the neuroprotective effects afforded by PNU 282987 and 5IA 85380, we performed experiments in the absence or presence of extracellular Ca^{2+} . Since prolonged lack of extracellular Ca^{2+} can be a toxic stimulus *per se*, we first performed a time course to determine the minimum time of agonist pre-incubation required to achieve the maximal protective effect (see protocol on top of Fig. 3).

As figure 3 illustrates, PNU 282987 reduced LDH release in cells injured with OA in a time-dependent manner; cell death was significantly reduced following only 5 min incubation with the agonist, 24 hours before exposing the cells to 16 h to OA (fig. 3 A). In the case of 5IA 85380, protection was significant following 1 h pre-incubation (fig. 3 B). For

1
2
3 comparability we selected a 4 h preincubation period for both agonists since this length of
4 exposure produced a level of protection similar to that seen following 24 h pre-incubation.
5
6 The presence or absence of extracellular Ca^{2+} was maintained only during the 4 h exposure to
7
8 the nicotinic agonist; thereafter, the medium was replaced with fresh EMEM containing Ca^{2+}
9
10 for 20 h, and OA was then added for 16 h. The results show that protection mediated by 5IA
11
12 85380 was dependent on extracellular Ca^{2+} (fig. 3 D) while that induced by PNU 282987 was
13
14 not (fig. 3 C). Moreover, when cells were pre- and co-incubated with the $\alpha 7$ allosteric
15
16 modulator PNU 120596 in the presence of Ca^{2+} , the neuroprotective effect of PNU 282987
17
18 was lost (fig. 3C). Finally, in the absence of extracellular Ca^{2+} , PNU 120596 did not reverse
19
20 the neuroprotection mediated by PNU 282987 (fig. 3 C). These results suggest that Ca^{2+} entry
21
22 via $\beta 2^*$ nAChR activation is required to afford a neuroprotective effect. However, in the case
23
24 of the $\alpha 7$ nAChR, PNU 282987 exerted a Ca^{2+} -independent neuroprotective effect.
25
26 Furthermore, as shown by the experiments with the $\alpha 7$ agonist in the presence of the $\alpha 7$
27
28 nAChR allosteric modulator that promotes Ca^{2+} entry through the $\alpha 7$ nAChR subtype (Fig.
29
30 2A), protection was lost in a Ca^{2+} dependent way (Fig. 3 C).
31
32
33
34
35
36
37
38
39
40

41 ***JAK-2 is involved in protection mediated by $\alpha 7$ but not by $\beta 2^*$ nAChRs***

42
43 JAK-2 is a protein that can be physically associated with $\alpha 7$ nAChRs and activated
44 when nicotine binds to this receptor subtype (Shaw *et al.* 2002). To determine if $\alpha 7$ and/or
45
46 $\beta 2^*$ nAChR subtype stimulation could activate this kinase under our experimental conditions,
47
48 SH-SY5Y cells were incubated with PNU 282987 10 μM , PNU 282987 10 μM plus PNU
49
50 120596 10 μM or 5IA 85380 100 μM for different times (5, 15, 30 min and 1 or 24 h) and
51
52 JAK-2 phosphorylation was then analyzed by western blotting. Cells treated with PNU
53
54 282987 10 μM showed maximally phosphorylated JAK-2 after 15 min incubation; thereafter,
55
56 phospho-JAK-2 decreased to basal levels (fig. 4 A). In contrast, 5IA 85380 failed to promote
57
58
59
60

1
2
3 the activation of JAK-2 at any of the times tested (fig. 4 B). Also, no alteration in phospho-
4
5 JAK-2 levels was observed when the $\alpha 7$ agonist was incubated in the presence of the $\alpha 7$
6
7 allosteric modulator (fig. 4 C).
8
9

10 JAK-2 has been previously shown to be involved in neuroprotection induced by
11
12 nicotine via $\alpha 7$ nAChR stimulation in neurons injured with $A\beta_{1-42}$ (Shaw *et al.* 2002). For this
13
14 reason, and because the $\alpha 7$ -selective agonist increased JAK-2 phosphorylation, we evaluated
15
16 if the protective effect of PNU 282987 against OA would be lost in the presence of a JAK-2
17
18 inhibitor, AG-490. To corroborate that JAK-2 activation was linked with its action on the $\alpha 7$
19
20 and not with the $\beta 2^*$ nAChRs, we also included 5IA 85380 in these experiments. As shown in
21
22 figure 4 D, the protective effect of PNU 282987 was completely prevented by pre- (15
23
24 minutes before) and co-incubation with AG-490, but this compound did not affect protection
25
26 afforded by 5IA 85380. Therefore, these results indicate that JAK-2 is linked to $\alpha 7$ nAChR-
27
28 mediated survival. Moreover, the presence of the $\alpha 7$ allosteric modulator prevented PNU
29
30 282987-induced JAK phosphorylation (fig. 4 C), and this result correlates with the loss of
31
32 protection under the same drug treatment conditions (see Fig.1C).
33
34
35
36
37
38
39
40

41 ***Implication of PI3K/Akt in the neuroprotective effect mediated by $\alpha 7$ and $\beta 2^*$ nAChR*** 42 ***stimulation*** 43 44

45
46 Multiple nicotinic agonists cause activation of PI3K/Akt (Kihara *et al.* 2001).
47
48 PI3K/Akt is known to be one of the targets of JAK-2 (Shaw *et al.* 2002). To investigate if
49
50 exposure to $\alpha 7$ and $\beta 2^*$ nAChR agonists led to Akt activation, we measured Akt
51
52 phosphorylation by western blot analysis. Incubation with PNU 282987 or 5IA 85380
53
54 produced a statistically significant activation of Akt at 5 and 15 minutes for both agonists (fig.
55
56 5 A and B). To determine if protection afforded by $\alpha 7$ and $\beta 2^*$ nAChR was dependent on the
57
58 PI3K/Akt pathway, cells were pre-(15 minutes before) and co-incubated with a PI3K
59
60

1
2
3 inhibitor, LY 294002 (3 μ M). Protection elicited by PNU 282987 and 5IA 85380 against OA-
4
5 toxicity was prevented by the inhibitor LY 294002 (fig. 5 C). Taken together, these results
6
7 indicate that protection obtained with both nicotinic agonists is dependent on activation of
8
9 PI3K/Akt.
10
11

12
13
14
15 ***ERK 1/2 is involved in the neuroprotective effect mediated by 5IA 85380 but not by PNU***
16
17 ***282987.***
18

19
20 The MAPK intracellular pathway has also been implicated in neuroprotection
21
22 processes mediated by nAChR activation (Egea *et al.* 2007). To assess if PNU 282987 and/or
23
24 5IA 85380 were able to activate ERK 1/2, we measured the phosphorylation of these kinases
25
26 by western blot. As shown in figure 6 A, PNU 282987 10 μ M by itself did not produce ERK
27
28 1/2 phosphorylation at any of the time intervals studied. However, 5IA 85380 significantly
29
30 increased the level of P-ERK 1/2 after 15 min (fig. 6 B). To corroborate the involvement of
31
32 MAPK activation in the neuroprotective effect mediated by 5IA 85380 and not by PNU
33
34 282987, we employed a MEK inhibitor, PD 98059. As shown in figure 6 C, the presence of
35
36 PD 98059 completely inhibited the protection induced by 5IA 85380 but had no effect on that
37
38 of PNU 282987. From these results, it can be concluded that MEK participates in the
39
40 neuroprotective effect mediated by β 2* but not by α 7 nAChR stimulation.
41
42
43
44
45
46
47

48 ***Involvement of GSK-3 β in neuroprotection induced by α 7 and β 2* nAChR.***
49

50
51 GSK-3 β is a substrate of Akt. Akt phosphorylates GSK-3 β at position Ser-9 and this
52
53 phosphorylation inactivates GSK-3 β (Stambolic and Woodgett 1994). On the other hand,
54
55 phosphorylation at Tyr-216 has the opposite effect and activates GSK-3 β ; however the
56
57 kinase(s) responsible for this phosphorylation are not well characterized (Cole *et al.* 2004;
58
59 Hughes *et al.* 1993; Wang *et al.* 1994). We therefore measured nAChR agonist-induced GSK-
60

1
2
3 3 β phosphorylation by western blot. As shown in figure 7, both PNU 282987 and 5IA 85380
4
5 were able to increase Ser 9 phosphorylation after 15 and 5 minutes incubation, respectively
6
7 (fig. 7 A and B). Subsequently, this phosphorylation decreased over time and after 24 hours it
8
9 was not statistically significantly different from OA control. P-Tyr 216-GSK-3 β and total
10
11 GSK-3 β levels were not altered at any of the incubation times (data not shown).
12
13
14

15 To determinate if GSK-3 β inhibition could be involved in the protection mediated by
16
17 PNU 282987 and 5IA 85380, we conducted a neuroprotection experiment in the presence or
18
19 absence of a GSK-3 β inhibitor, AR-A014418. Figure 7 C shows how AR-A014418 alone
20
21 produced protection against OA and such protection was not statistically different from that
22
23 obtained with the nicotinic agonists alone. Since the combination of nAChR agonists plus
24
25 AR-A014418 did not afford a greater neuroprotective effect, we deduce that both agonists
26
27 were acting through a common mechanism which involves a transient inactivation of GSK-
28
29 3 β .
30
31
32
33

34 To further investigate how nicotinic agonists and the GSK-3 β inhibitor, pre-incubated
35
36 for 24 h before adding OA, induced neuroprotection we evaluated the activation and the
37
38 expression of GSK-3 β at the end of the experiment, i.e. after OA treatment. Western blot
39
40 analysis of GSK-3 β was carried out after OA exposure for 16 hours alone or in the presence
41
42 of the neuroprotective compounds. As shown in figure 8 A, the expression of total GSK-3 β
43
44 increased after OA treatment and nicotinic agonists prevented this increased expression to
45
46 almost basal levels (fig. 8 A); a similar effect was obtained with the GSK-3 β antagonist AR-
47
48 A014418. Interestingly, OA produced a significant increase of total P-Tyr 216 that was also
49
50 significantly reduced by the nicotinic agonists and AR-A014418 (fig. 8 B).
51
52
53
54
55
56
57
58
59
60

1
2
3 ***$\alpha 7$ and $\beta 2^*$ nicotinic agonists reduced τ hyperphosphorylation***
4
5

6 As previously mentioned, OA is an inhibitor of phosphatases that augments τ
7 hyperphosphorylation. In addition, one of the kinases involved in τ phosphorylation is GSK-
8 3β , whose target in the τ substrate is Ser396/404 (Cho and Johnson 2004). The question of
9 how GSK- 3β activity affected the phosphorylation on τ Ser396/404 residue was raised. After
10 OA treatment for 16 h in the presence or absence of PNU 282987, 5IA85380 and AR-
11 A014418, we measured Ser-396/404 τ phosphorylation by western blot. As shown in figure 9,
12 PNU 282987, 5IA 85380 and AR-A014418 significantly reduced τ phosphorylation induced
13 by OA on its Ser396/404 residue.
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 ***Effect of $\alpha 7$ and $\beta 2^*$ nicotinic agonists on the cytoskeleton alterations caused by okadaic***
29 ***acid***
30
31

32 AO, besides causing hyperphosphorylation of tau, induces changes in the cytoskeleton
33 that are identified with strong retraction and rounding in cell morphology (Vale and Botana,
34 2008; Yoon et al., 2008; Cabado et al., 2004). Therefore, we also analyzed how treatment of
35 the cells with nicotinic agonists could modify the changes in the cytoskeleton induced by OA
36 treatment.
37
38
39
40
41
42
43

44 As shown in figure 10, control cells exposed only to EMEM (Fig. 10A) showed a
45 fibrous cytoarchitecture with a well extended and branched morphology. OA-treated cells
46 showed a retracted morphology with almost no neurites (Fig. 10B). When cells were pre-
47 incubated for 24 h with PNU282987 (10 μ M) (Fig. 10C) or 5IA 85380 (100 μ M) (Fig. 10D)
48 before adding OA, we observed a partial recovery of the fibrous and branched
49 cytoarchitecture as observed in intact cells. This partial recovery of the cytoarchitecture
50 correlates with the partial neuroprotective effects observed when measured as LDH release
51 (Fig. 1 A and B).
52
53
54
55
56
57
58
59
60

DISCUSSION

Central to this study is the observation that both $\alpha 7$ and $\beta 2^*$ nAChR can transduce signals through different mechanisms to provide protection of cells that undergo hyperphosphorylation of τ . Ultimately, these distinct signalling pathways converge on the regulation of GSK3 β and reduction of τ phosphorylation.

To analyse how Ca^{2+} could be involved in the protective effects of PNU 282987 and 5IA 85380, we first measured cytosolic Ca^{2+} levels in Fluo-3-loaded SH-SY5Y cells. Interestingly, PNU 282987 at the protective concentration of 10 μM did not induce a measureable intracellular Ca^{2+} increase, but in the presence of the allosteric modulator PNU 120596, it gave a robust intracellular Ca^{2+} rise which was blocked by the $\alpha 7$ nAChR-selective antagonists BGT and MLA. These results are in agreement with those previously reported in bovine chromaffin cells (del Barrio *et al.*) or PC12 cells (Dickinson *et al.* 2007). Unlike PNU 282987, 5IA 85380 alone, at a protective concentration, was able to increase intracellular Ca^{2+} levels; this response was blocked by selective $\beta 2^*$ antagonists such as DH β E and α -conotoxin MII but not by $\alpha 7$ (Dickinson *et al.* 2007) or $\alpha 3\beta 4$ selective antagonists, which corroborates the selectivity of 5IA 85380 for $\beta 2^*$ nAChR subtype (Dajas-Bailador *et al.* 2002a; Mukhin *et al.* 2000).

The observation that the $\alpha 7$ agonist exerted its protective effect without apparently inducing intracellular Ca^{2+} increases is against the belief that Ca^{2+} is the second messenger responsible for kinases intracellular activation involved in neuroprotection mediated by $\alpha 7$ nAChR (Dajas-Bailador *et al.* 2002b; Donnelly-Roberts *et al.* 1996; Ferchmin *et al.* 2003). However, examination of the protective efficacy of PNU 282987 in a medium without Ca^{2+} corroborated the Ca^{2+} -independence of the protective effect. In fact, when Ca^{2+} was forced to increase via $\alpha 7$ nAChR activation, with the aid of the positive allosteric modulator, protection

1
2
3 was lost in a Ca^{2+} dependent way. In contrast, the protective effect of 5IA 85380 was
4
5 dependent on extracellular Ca^{2+} which corroborates a previous study in which this agonist
6
7 offered Ca^{2+} -dependent neuroprotection against glutamate-induced toxicity (Ueda *et al.*
8
9 2008). Taken together, these results indicate that $\alpha 7$ nAChR protection is Ca^{2+} -independent
10
11 while $\beta 2^*$ nAChR protection is Ca^{2+} -dependent.
12
13

14
15 Regarding the intracellular signalling pathways involved in the neuroprotective
16
17 mechanism associated with the nAChR activation, there are several intracellular kinases
18
19 involved in this process (Buckingham *et al.* 2009). In the case of the $\alpha 7$ nAChR, JAK-2
20
21 kinase is able to bind to this nAChR subtype under nicotinic stimulation, to transduce signals
22
23 to PI3-kinase and Akt (Shaw *et al.* 2002), which results in neuroprotection. Our results show
24
25 that only stimulation of the $\alpha 7$ and not the $\beta 2^*$ nAChR can induce JAK-2 phosphorylation.
26
27 Activation of the JAK-2/PI3K/Akt route can be independent of Ca^{2+} (Chernyavsky *et al.*
28
29 2009) which agrees with the results of this study. The implication of this kinase in protection
30
31 mediated by $\alpha 7$ nAChR was confirmed when protection induced by PNU 282987, but not by
32
33 5IA 85380, was lost in cells treated with the JAK-2 inhibitor AG-490. Furthermore, the loss
34
35 of protection by the $\alpha 7$ agonist in the presence of the allosteric modulator seems to be related
36
37 to the loss of phosphorylation of JAK-2 under similar conditions, as determined by the
38
39 western blot experiments (Fig. 4C).
40
41
42
43
44
45

46 Continuing with the intracellular signalling pathways, data from other groups
47
48 (Buckingham *et al.* 2009; Quesada *et al.* 2008) as well as our own (Arias *et al.* 2005; Canas *et*
49
50 *al.* 2007) have shown that activation of PI3K/Akt is related to cell survival. In line with these
51
52 observations, we have seen that both PNU 282987 and 5IA 85380 increased Akt
53
54 phosphorylation. Furthermore, the PI3K inhibitor LY 294002 blocked the protective effects of
55
56 the $\alpha 7$ and $\beta 2^*$ agonist. Therefore, this signalling pathway is shared by both nAChR subtypes.
57
58 In the case of $\alpha 7$ nAChR, it has been described that one of the targets of JAK-2 is precisely
59
60

1
2
3 PI3K/Akt (Shaw *et al.* 2002); these kinases can be activated in a Ca²⁺-dependent (Soletti *et*
4
5 *al.*) or independent manner (Chernyavsky *et al.* 2009).
6
7

8 Based on a previous study that showed that non $\alpha 7$ nAChRs are able to activate
9
10 MAPK/ERK in a Ca²⁺-dependent way (Nakayama *et al.* 2001) we evaluated if 5IA 85380
11
12 could induce ERK 1/2 phosphorylation. In our experimental conditions, ERK1/2 activation
13
14 was only induced by 5IA 85380. Participation of MAPK in $\beta 2^*$ nAChR-mediated protection
15
16 was supported with the experiments with the inhibitor PD98059 (Fig. 6C).
17
18

19
20 Since both agonists activated Akt, we tried to identify Akt targets which could be
21
22 related to the mechanism of action of OA, which induces τ hyperphosphorylation. One of the
23
24 kinases that phosphorylates τ is GSK-3 β (Grimes and Jope 2001), and it is known that P-Akt
25
26 inactivates GSK-3 β by phosphorylation of its Ser-9 (Cross *et al.* 1995; Srivastava and Pandey
27
28 1998). We observed that $\alpha 7$ and $\beta 2^*$ nAChR stimulation increased phosphorylation of Ser-9-
29
30 GSK-3 β after 15 min but it decreased after 24h exposure to the agonist. This temporal pattern
31
32 excludes sustained inactivation of GSK-3 β as a prerequisite for the observed neuroprotection.
33
34 However, protection mediated by both nAChR subtypes does seem to be related to inhibition
35
36 of GSK-3 β since the combination of AR-A014418 (a GSK-3 β inhibitor) with the $\alpha 7$ or $\beta 2^*$
37
38 nAChR agonist did not afford additional protective effects (fig. 7G).
39
40
41
42

43
44 After exposing the cells for 16 h to OA, total GSK-3 β expression was increased by
45
46 almost two-fold. Increases in the expression of GSK-3 β have been reported in glial progenitor
47
48 cells of Alzheimer's disease patients treated with A β (He and Shen 2009) as well as in post-
49
50 mortem brain of Alzheimer patients (Lau *et al.* 1999). In fact, there are several transgenic
51
52 models that over-express GSK-3 β and reproduce many pathophysiological features of
53
54 Alzheimer disease (Engel *et al.* 2006). Taken together, OA seems to increase the expression
55
56 of GSK-3 β as observed in Alzheimer's disease patients. Therefore, reduction of total GSK-3 β
57
58 expression induced by nicotinic agonists and the GSK-3 β inhibitor could be an interesting
59
60

1
2
3 therapeutic approach. Recent studies have shown that lithium, a GSK-3 β inhibitor, and
4 tobacco extracts are able to reduce mRNA and GSK-3 β protein levels respectively (Mendes *et*
5 *al.* 2009; Tian *et al.* 2009). Moreover, total Tyr-216-GSK-3 β phosphorylation (active levels)
6 was also increased by OA alone and this increase was significantly reduced by nAChR
7 agonists and AR- A014418.
8
9

10
11
12
13
14
15 GSK-3 β specifically phosphorylates Ser-396/404 of τ (PHF-1, Paired Helicoidal
16 Filament-1) both in cell cultures (Lovestone *et al.* 1994; Tanaka *et al.* 1998) and “*in vivo*”
17 models (Spittaels *et al.* 2000). τ hyperphosphorylation by PP2A inhibition induces a
18 disruption of microtubules causing cell death in SH-SY5Y cells treated with OA (Tanaka *et*
19 *al.* 1998). This microtubule destabilisation compromises axonal transport and induces
20 neurodegeneration of neurons with tangles, a widely accepted characteristic in brains of
21 Alzheimer’s disease patients (Alonso *et al.* 1994; Alonso *et al.* 1997). We have observed that
22 OA almost doubled τ phosphorylation on its Ser-396/404 and that the $\alpha 7$ and $\beta 2^*$ nicotinic
23 agonists, together with the GSK-3 β inhibitor, reduced Ser-396/404- τ -hyperphosphorylation.
24 Furthermore, cytoskeleton alterations caused by okadaic acid were partially reverted with
25 nAChR agonist treatment (Fig. 10).
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41
42 Apart from reducing tau phosphorylation, nicotinic stimulation also confers
43 neuroprotection and anti-inflammatory actions by activation of some of the intracellular
44 pathways here described (Marrero and Bencherif, 2009; Kawamata and Shimohama, 2011;
45 Parada *et al.*, 2010; Egea *et al.*, 2007), which could also be contributing to the protective
46 effects observed by both $\alpha 7$ and $\beta 2$ nAChR activation.
47
48
49
50
51
52

53
54 In conclusion, both $\alpha 7$ and $\beta 2^*$ nAChR activation can mediate protection against OA.
55 $\alpha 7$ nAChR mediated protection involves a Ca²⁺-independent-JAK-2/PI3K/Akt/GSK-3 β
56 signalling pathway. In contrast, the Ca²⁺-dependent protective mechanism mediated by $\beta 2^*$
57 nAChR relates to MEK/ERK1/2, PI3K/Akt activation and GSK-3 β inactivation. However,
58
59
60

1
2
3 both signalling pathways converge on GSK-3 β down regulation and reduction of
4 phosphorylation of τ in cells exposed to okadaic acid which represents a good model of
5 neuronal death related to the characteristic tau hyperphosphorylation found in Alzheimer's
6 disease patients.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FUNDING INFORMATION

This work was supported in part by grants from the Spanish Ministry of Education and [SAF2009-12150], the Spanish Ministry of Health-Instituto de Salud Carlos III- [RETICS-RD06/0026], Comunidad Autónoma de Madrid [SAL2006/0275] to MGL and MH63631 and GM48677 to JMM.

ACKNOWLEDGEMENTS

We thank Ms Vanessa Gómez for technical assistance and the support of Fundación Teófilo Hernando.

LEGENDS TO FIGURES

Figure 1. Protective effects mediated by the nicotinic agonists PNU 282987 and 5IA 85380 and the $\alpha 7$ allosteric modulator PNU 120596 against okadaic acid-induced toxicity. Cells were pre-treated for 24 hours with increasing concentrations of the $\alpha 7$ nicotinic agonist PNU282987, the $\beta 2^*$ nicotinic agonist 5IA 85380 or the $\alpha 7$ positive allosteric modulator PNU 120596. Thereafter, SH-SY5Y were incubated for 16 h with okadaic acid (OA) 30 nM and at the end of the experiment cell death was evaluated by measuring the release of LDH release to the extracellular medium (see top of the figure). PNU 282987 (**A**) and 5IA 85380 (**B**) offered protection against OA-induced toxicity, however the $\alpha 7$ allosteric modulator PNU 120596 did not (**C**). The data represent means \pm S.E.M. of triplicates of at least 4 different batches of cells. * $p < 0.05$, ** $p < 0.01$ with respect to OA alone.

Figure 2. Intracellular Ca^{2+} increases mediated by the nicotinic agonists PNU 282987 and 5IA 85380 in SH-SY5Y cells loaded with Fluo-3. (A) SH-SY5Y cells were stimulated with 10 μ M PNU 282987 alone or in cells pre-incubated during 15 min with the $\alpha 7$ allosteric modulator PNU 120596 (10 μ M); PNU 120596 was also co-incubated with various nicotinic antagonist such as mecamylamine (“Meca” 30 μ M), α -bungarotoxin (“BGT” 100 nM) or methyllycaconitine (“MLA” 100 nM). (B) Cytosolic Ca^{2+} increases induced by 5IA 85380 at 100 μ M alone or in the presence of the nicotinic antagonist Meca 30 μ M, BGT 100 nM, α -conotoxin-AuIB (“AuIB” 10 μ M), dihydro- β -eritroidine (“DH β E” 10 μ M) or α -conotoxin-MII (“MII” 200 nM). The data represent the average of maximum fluorescence increases during 10 seconds after the implementation of PNU 282987 10 μ M (**A**) or 5IA 85380 100 μ M (**B**). Experiments were conducted in triplicate on at least 6 different batches of cells. *** $p < 0.001$ with respect to PNU 282987 alone (**A**) or with respect to 5IA 85380 (**B**);

1
2
3 $###p < 0.001$ with respect to PNU 282987+PNU 120596 (**B**); $**p < 0.01$ with respect to 5IA
4
5 85380 (**B**).
6
7
8
9

10 **Figure 3. Time and Ca²⁺ dependence of cytoprotection mediated by PNU 282987 and**
11 **5IA 85380.** As shown in the upper panel, cells were incubated with PNU 282987 (10 μ M) (**A**)
12 or 5IA 85380 (100 μ M) (**B**) during 5 min, 1 h, 4 h or 24 h, then the agonist was removed and
13 fresh EMEM was added until to complete 24 h; after this period, EMEM was replaced by OA
14 (30 nM) for 16 h. At the end of the experiment, cell death was evaluated by LDH. Figures (**C**)
15 and (**D**) show the mean of LDH values when agonists were pre-incubated for 4 h in the
16 presence or absence of extracellular Ca²⁺ using the same protocol described above. PNU
17 120596 (“PNU 12” 10 μ M) was pre-incubated for 15 min and co-incubated during the
18 application of PNU 282987 (**C**). Data represents the mean of LDH values normalized with
19 respect to OA alone \pm S.E.M. of triplicates of at least 6 different cultures. $*p < 0.05$,
20 $**p < 0.01$ and $***p < 0.001$ with respect to OA; $^{\#}p < 0.05$ and $###p < 0.001$ with respect to
21 the agonists plus OA.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 **Figure 4. Differential phosphorylation and implication of JAK-2 in neuroprotection**
42 **mediated by PNU 282987 and 5IA 85380.** PNU282987 (**A**), 5IA 85380 (**B**) and PNU
43 282987 plus the allosteric modulator PNU 120596 were incubated during different time
44 periods (5 min, 15 min, 30 min, 1 h and 24 h); at the end, P-JAK-2 (immunoblots on the top)
45 and JAK-2 total (immunoblots at the bottom) were quantified by western blot. (**D**) Represents
46 the effect of the JAK-2 inhibitor, AG-490, on the neuroprotective action of PNU 282987 10
47 μ M (“PNU 28”) and 5IA 85380 100 μ M (“5IA”). Data are the mean \pm S.E.M. of at least 4
48 different cultures; $*p < 0.05$ with respect to the basal (**A**); $***p < 0.001$ with respect to OA
49 alone and $^{\#}p < 0.05$ with respect to PNU 282987 (**B**).
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 5. Phosphorylation of Akt and implication of PI3K in neuroprotection mediated by PNU 282987 and 5IA 85380. Akt phosphorylation by PNU 282987 10 μ M (A) and 5IA 85380 100 μ M (B) at different incubation periods. (C) Shows the effect of the PI3K inhibitor 3 μ M LY 294002 on the neuroprotection mediated by PNU 282987 10 μ M (“PNU 28”) and 5IA 85380 100 μ M (“5IA”) against OA-induced toxicity. Each value is a mean of at least 4 different cell cultures \pm S.E.M. $*p < 0.05$ and $**p < 0.01$ with respect to the basal in figures (A) and (B); $***p < 0.001$ with respect to OA alone, $^{\#}p < 0.05$ with respect to PNU 282987, $^{\$}p < 0.01$ with respect to 5IA 85380 (C).

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 6. Phosphorylation of ERK 1/2 and implication of MEK in neuroprotection mediated by PNU 282987 and 5IA 85380. ERK 1/2 phosphorylation by PNU 282987 10 μ M (A) and 5IA 85380 100 μ M (B) at different incubation periods. (C) Shows the effect of the MEK inhibitor PD 98059 1 μ M on neuroprotection mediated by PNU 282987 10 μ M (“PNU 28”) and 5IA 85380 100 μ M (“5IA”) against OA-induced toxicity. Each value is a mean of at least 4 different cell cultures \pm S.E.M. $*p < 0.05$ with respect to the basal in (B); $***p < 0.001$ with respect to OA alone, $^{\#\#}p < 0.01$ with respect to 5IA 85380 (C).

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 7. GSK-3 β phosphorylation and implication in neuroprotection mediated by PNU 282987 and 5IA 85380. Time course of Ser-9 were analysed by western blot in protein extracts obtained from cells treated with PNU 282987 10 μ M (A) and 5IA 85380 100 μ M (B) during different time periods. (C) Shows the effect of the GSK-3 β inhibitor AR-A014418 1 μ M on the neuroprotection mediated by PNU 282987 10 μ M (“PNU 28”) and 5IA 85380 100 μ M (“5IA”) against OA-induced toxicity. Each value is a mean of at least 4 different cultures \pm S.E.M. $*p < 0.05$ with respect to the basal in (A and B); $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$ with respect to OA alone (C).

1
2
3
4
5
6 **Figure 8. GSK-3 β expression/phosphorylation under the treatment of OA.** Protein
7
8 samples were collected from cells that were pre-incubated for 24 h with PNU 282987 10 μ M,
9
10 5IA 85380 100 μ M or AR-A014418 1 μ M and, subsequently this pre-treatment was replaced
11
12 by medium with OA 30 nM for 16 hours. Histograms represent the densitometric
13
14 quantification of total GSK-3 β (A) and total Tyr-216 (B) phosphorylation, using β -actin for
15
16 normalization. Each value is a mean of 4 different cultures \pm S.E.M. * p < 0.05, ** p < 0.01
17
18 with respect to basal levels and, # p < 0.05, ## p < 0.01 and ### p < 0.001 with respect to OA
19
20 alone.
21
22
23
24
25
26

27 **Figure 9. Effect of nicotinic agonists and AR-A014418 on Ser-396/404 τ phosphorylation**
28
29 **in cells treated with AO.** Protein samples were collected from cells that were pre-incubated
30
31 for 24 h with PNU 282987 10 μ M, 5IA 85380 100 μ M or AR-A014418 1 μ M and
32
33 subsequently medium was replaced by OA 30 nM for 16 hours. Histograms represent the
34
35 densitometric quantification of P-Ser-396/404- τ , using τ total for normalization. Each value is
36
37 a mean of 5 different cultures \pm S.E.M. *** p < 0.001 with respect to basal levels and
38
39 # p < 0.05 with respect to OA alone.
40
41
42
43
44
45
46

47 **Figure 10. PNU 282987 and 5IA 85380 partially prevent cytoarchitecture disruption**
48
49 **elicited by OA.** Photomicrographs of double staining of SH-SY5Y marked with Rhodamine-
50
51 phalloidin (filamentous actin marker, red) and Hoechst (nuclear marker, blue). Cells were
52
53 incubated with PNU 282987 (10 μ M) (C) or 5IA 85380 (100 μ M) (D) during 24 h, then OA
54
55 30 nM was added for 5h. A control group without any treatment (A) and another just treated
56
57 with OA 30 nM for 5 h were run in parallel (B). As shown in the photomicrographs, PNU
58
59 282987 and 5IA 85380 prevented partially the cytoarchitecture alterations caused by OA.
60

REFERENCES

- Alonso, A. C., Zaidi, T., Grundke-Iqbal, I., and Iqbal, K. (1994). Role of abnormally phosphorylated tau in the breakdown of microtubules in Alzheimer disease. *Proc Natl Acad Sci U S A* **91**, 5562-6.
- Alonso, A. D., Grundke-Iqbal, I., Barra, H. S., and Iqbal, K. (1997). Abnormal phosphorylation of tau and the mechanism of Alzheimer neurofibrillary degeneration: sequestration of microtubule-associated proteins 1 and 2 and the disassembly of microtubules by the abnormal tau. *Proc Natl Acad Sci U S A* **94**, 298-303.
- Arias, E., Gallego-Sandin, S., Villarroya, M., Garcia, A. G., and Lopez, M. G. (2005). Unequal neuroprotection afforded by the acetylcholinesterase inhibitors galantamine, donepezil, and rivastigmine in SH-SY5Y neuroblastoma cells: role of nicotinic receptors. *J Pharmacol Exp Ther* **315**, 1346-53.
- Avila, J. (2000). Tau aggregation into fibrillar polymers: taupathies. *FEBS Lett* **476**, 89-92.
- Bitner, R. S., Nikkel, A. L., Markosyan, S., Otte, S., Puttfarcken, P., and Gopalakrishnan, M. (2009). Selective alpha7 nicotinic acetylcholine receptor activation regulates glycogen synthase kinase3beta and decreases tau phosphorylation in vivo. *Brain Res* **1265**, 65-74.
- Bodnar, A. L., Cortes-Burgos, L. A., Cook, K. K., Dinh, D. M., Groppi, V. E., Hajos, M., Higdon, N. R., Hoffmann, W. E., Hurst, R. S., Myers, J. K., Rogers, B. N., Wall, T. M., Wolfe, M. L., and Wong, E. (2005). Discovery and structure-activity relationship of quinuclidine benzamides as agonists of alpha7 nicotinic acetylcholine receptors. *J Med Chem* **48**, 905-8.
- Buckingham, S. D., Jones, A. K., Brown, L. A., and Sattelle, D. B. (2009). Nicotinic acetylcholine receptor signalling: roles in Alzheimer's disease and amyloid neuroprotection. *Pharmacol Rev* **61**, 39-61.
- Canas, N., Valero, T., Villarroya, M., Montell, E., Verges, J., Garcia, A. G., and Lopez, M. G. (2007). Chondroitin sulfate protects SH-SY5Y cells from oxidative stress by inducing heme oxygenase-1 via phosphatidylinositol 3-kinase/Akt. *J Pharmacol Exp Ther* **323**, 946-53.
- Cole, A., Frame, S., and Cohen, P. (2004). Further evidence that the tyrosine phosphorylation of glycogen synthase kinase-3 (GSK3) in mammalian cells is an autophosphorylation event. *Biochem J* **377**, 249-55.
- Cowan, C. M., Chee, F., Shepherd, D., and Mudher, A. Disruption of neuronal function by soluble hyperphosphorylated tau in a Drosophila model of tauopathy. *Biochem Soc Trans* **38**, 564-70.
- Cross, D. A., Alessi, D. R., Cohen, P., Andjelkovich, M., and Hemmings, B. A. (1995). Inhibition of glycogen synthase kinase-3 by insulin mediated by protein kinase B. *Nature* **378**, 785-9.
- Cummings, J. L. (2004). Treatment of Alzheimer's disease: current and future therapeutic approaches. *Rev Neurol Dis* **1**, 60-9.
- Chernyavsky, A. I., Arredondo, J., Qian, J., Galitovskiy, V., and Grando, S. A. (2009). Coupling of ionic events to protein kinase signaling cascades upon activation of alpha7 nicotinic receptor: cooperative regulation of alpha2-integrin expression and Rho kinase activity. *J Biol Chem* **284**, 22140-8.
- Cho, J. H., and Johnson, G. V. (2004). Glycogen synthase kinase 3 beta induces caspase-cleaved tau aggregation in situ. *J Biol Chem* **279**, 54716-23.
- Dajas-Bailador, F., and Wonnacott, S. (2004). Nicotinic acetylcholine receptors and the regulation of neuronal signalling. *Trends Pharmacol Sci* **25**, 317-24.

- 1
2
3 Dajas-Bailador, F. A., Mogg, A. J., and Wonnacott, S. (2002a). Intracellular Ca²⁺ signals
4 evoked by stimulation of nicotinic acetylcholine receptors in SH-SY5Y cells: contribution of
5 voltage-operated Ca²⁺ channels and Ca²⁺ stores. *J Neurochem* **81**, 606-14.
- 6 Dajas-Bailador, F. A., Soliakov, L., and Wonnacott, S. (2002b). Nicotine activates the
7 extracellular signal-regulated kinase 1/2 via the alpha7 nicotinic acetylcholine receptor and
8 protein kinase A, in SH-SY5Y cells and hippocampal neurones. *J Neurochem* **80**, 520-30.
- 9 De Vos, K. J., Grierson, A. J., Ackerley, S., and Miller, C. C. (2008). Role of axonal transport
10 in neurodegenerative diseases. *Annu Rev Neurosci* **31**, 151-73.
- 11 del Barrio, L., Egea, J., Leon, R., Romero, A., Ruiz, A., Montero, M., Alvarez, J., and Lopez,
12 M. G. Calcium signalling mediated through alpha7 and non-alpha7 nAChR stimulation is
13 differentially regulated in bovine chromaffin cells to induce catecholamine release. *Br J*
14 *Pharmacol* **162**, 94-110.
- 15 Dickinson, J. A., Hanrott, K. E., Mok, M. H., Kew, J. N., and Wonnacott, S. (2007).
16 Differential coupling of alpha7 and non-alpha7 nicotinic acetylcholine receptors to calcium-
17 induced calcium release and voltage-operated calcium channels in PC12 cells. *J Neurochem*
18 **100**, 1089-96.
- 19 Donnelly-Roberts, D. L., Xue, I. C., Arneric, S. P., and Sullivan, J. P. (1996). In vitro
20 neuroprotective properties of the novel cholinergic channel activator (ChCA), ABT-418.
21 *Brain Res* **719**, 36-44.
- 22 Egea, J., Rosa, A. O., Cuadrado, A., Garcia, A. G., and Lopez, M. G. (2007). Nicotinic
23 receptor activation by epibatidine induces heme oxygenase-1 and protects chromaffin cells
24 against oxidative stress. *J Neurochem* **102**, 1842-52.
- 25 Engel, T., Hernandez, F., Avila, J., and Lucas, J. J. (2006). Full reversal of Alzheimer's
26 disease-like phenotype in a mouse model with conditional overexpression of glycogen
27 synthase kinase-3. *J Neurosci* **26**, 5083-90.
- 28 Ferchmin, P. A., Perez, D., Eterovic, V. A., and de Vellis, J. (2003). Nicotinic receptors
29 differentially regulate N-methyl-D-aspartate damage in acute hippocampal slices. *J*
30 *Pharmacol Exp Ther* **305**, 1071-8.
- 31 Gentry, C. L., and Lukas, R. J. (2002). Regulation of nicotinic acetylcholine receptor numbers
32 and function by chronic nicotine exposure. *Curr Drug Targets CNS Neurol Disord* **1**, 359-85.
- 33 Gong, C. X., Grundke-Iqbal, I., and Iqbal, K. (1994). Dephosphorylation of Alzheimer's
34 disease abnormally phosphorylated tau by protein phosphatase-2A. *Neuroscience* **61**, 765-72.
- 35 Gong, C. X., Shaikh, S., Wang, J. Z., Zaidi, T., Grundke-Iqbal, I., and Iqbal, K. (1995).
36 Phosphatase activity toward abnormally phosphorylated tau: decrease in Alzheimer disease
37 brain. *J Neurochem* **65**, 732-8.
- 38 Gong, C. X., Singh, T. J., Grundke-Iqbal, I., and Iqbal, K. (1993). Phosphoprotein
39 phosphatase activities in Alzheimer disease brain. *J Neurochem* **61**, 921-7.
- 40 Grimes, C. A., and Jope, R. S. (2001). CREB DNA binding activity is inhibited by glycogen
41 synthase kinase-3 beta and facilitated by lithium. *J Neurochem* **78**, 1219-32.
- 42 Grundke-Iqbal, I., Iqbal, K., Tung, Y. C., Quinlan, M., Wisniewski, H. M., and Binder, L. I.
43 (1986). Abnormal phosphorylation of the microtubule-associated protein tau (tau) in
44 Alzheimer cytoskeletal pathology. *Proc Natl Acad Sci U S A* **83**, 4913-7.
- 45 Grundke-Iqbal, I., Vorbrodt, A. W., Iqbal, K., Tung, Y. C., Wang, G. P., and Wisniewski, H.
46 M. (1988). Microtubule-associated polypeptides tau are altered in Alzheimer paired helical
47 filaments. *Brain Res* **464**, 43-52.
- 48 Haass, C., and Selkoe, D. J. (2007). Soluble protein oligomers in neurodegeneration: lessons
49 from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* **8**, 101-12.
- 50 Haydar, S. N., and Dunlop, J. Neuronal Nicotinic Acetylcholine Receptors - Targets for the
51 Development of Drugs to Treat Cognitive Impairment Associated with Schizophrenia and
52 Alzheimer's Disease. *Curr Top Med Chem*.
- 53
54
55
56
57
58
59
60

- 1
2
3 He, P., and Shen, Y. (2009). Interruption of beta-catenin signaling reduces neurogenesis in
4 Alzheimer's disease. *J Neurosci* **29**, 6545-57.
- 5 Hughes, K., Nikolakaki, E., Plyte, S. E., Totty, N. F., and Woodgett, J. R. (1993). Modulation
6 of the glycogen synthase kinase-3 family by tyrosine phosphorylation. *Embo J* **12**, 803-8.
- 7 Hurst, R. S., Hajos, M., Raggenbass, M., Wall, T. M., Higdon, N. R., Lawson, J. A.,
8 Rutherford-Root, K. L., Berkenpas, M. B., Hoffmann, W. E., Piotrowski, D. W., Groppi, V.
9 E., Allaman, G., Ogier, R., Bertrand, S., Bertrand, D., and Arneric, S. P. (2005). A novel
10 positive allosteric modulator of the alpha7 neuronal nicotinic acetylcholine receptor: in vitro
11 and in vivo characterization. *J Neurosci* **25**, 4396-405.
- 12 Irizarry, M. C., Soriano, F., McNamara, M., Page, K. J., Schenk, D., Games, D., and Hyman,
13 B. T. (1997). Abeta deposition is associated with neuropil changes, but not with overt
14 neuronal loss in the human amyloid precursor protein V717F (PDAPP) transgenic mouse. *J*
15 *Neurosci* **17**, 7053-9.
- 16 Kihara, T., Shimohama, S., Sawada, H., Honda, K., Nakamizo, T., Shibasaki, H., Kume, T.,
17 and Akaike, A. (2001). alpha 7 nicotinic receptor transduces signals to phosphatidylinositol 3-
18 kinase to block A beta-amyloid-induced neurotoxicity. *J Biol Chem* **276**, 13541-6.
- 19 Kihara, T., Shimohama, S., Urushitani, M., Sawada, H., Kimura, J., Kume, T., Maeda, T., and
20 Akaike, A. (1998). Stimulation of alpha4beta2 nicotinic acetylcholine receptors inhibits beta-
21 amyloid toxicity. *Brain Res* **792**, 331-4.
- 22 Lau, K. F., Miller, C. C., Anderton, B. H., and Shaw, P. C. (1999). Molecular cloning and
23 characterization of the human glycogen synthase kinase-3beta promoter. *Genomics* **60**, 121-8.
- 24 Lee, V. M., and Trojanowski, J. Q. (1992). The disordered neuronal cytoskeleton in
25 Alzheimer's disease. *Curr Opin Neurobiol* **2**, 653-6.
- 26 Lovestone, S., Reynolds, C. H., Latimer, D., Davis, D. R., Anderton, B. H., Gallo, J. M.,
27 Hanger, D., Mulot, S., Marquardt, B., Stabel, S., and et al. (1994). Alzheimer's disease-like
28 phosphorylation of the microtubule-associated protein tau by glycogen synthase kinase-3 in
29 transfected mammalian cells. *Curr Biol* **4**, 1077-86.
- 30 Lucas, J. J., Hernandez, F., Gomez-Ramos, P., Moran, M. A., Hen, R., and Avila, J. (2001).
31 Decreased nuclear beta-catenin, tau hyperphosphorylation and neurodegeneration in GSK-
32 3beta conditional transgenic mice. *Embo J* **20**, 27-39.
- 33 Mendes, C. T., Mury, F. B., de Sa Moreira, E., Alberto, F. L., Forlenza, O. V., Dias-Neto, E.,
34 and Gattaz, W. F. (2009). Lithium reduces Gsk3b mRNA levels: implications for Alzheimer
35 Disease. *Eur Arch Psychiatry Clin Neurosci* **259**, 16-22.
- 36 Mukhin, A. G., Gundisch, D., Horti, A. G., Koren, A. O., Tamagnan, G., Kimes, A. S.,
37 Chambers, J., Vaupel, D. B., King, S. L., Picciotto, M. R., Innis, R. B., and London, E. D.
38 (2000). 5-Iodo-A-85380, an alpha4beta2 subtype-selective ligand for nicotinic acetylcholine
39 receptors. *Mol Pharmacol* **57**, 642-9.
- 40 Nakayama, H., Numakawa, T., Ikeuchi, T., and Hatanaka, H. (2001). Nicotine-induced
41 phosphorylation of extracellular signal-regulated protein kinase and CREB in PC12h cells. *J*
42 *Neurochem* **79**, 489-98.
- 43 Parada, E., Egea, J., Romero, A., del Barrio, L., Garcia, A. G., and Lopez, M. G. Poststress
44 treatment with PNU282987 can rescue SH-SY5Y cells undergoing apoptosis via alpha7
45 nicotinic receptors linked to a Jak2/Akt/HO-1 signaling pathway. *Free Radic Biol Med* **49**,
46 1815-21.
- 47 Peng, X., Katz, M., Gerzanich, V., Anand, R., and Lindstrom, J. (1994). Human alpha 7
48 acetylcholine receptor: cloning of the alpha 7 subunit from the SH-SY5Y cell line and
49 determination of pharmacological properties of native receptors and functional alpha 7
50 homomers expressed in Xenopus oocytes. *Mol Pharmacol* **45**, 546-54.
- 51
52
53
54
55
56
57
58
59
60

- 1
2
3 Perez, M., Hernandez, F., Gomez-Ramos, A., Smith, M., Perry, G., and Avila, J. (2002).
4 Formation of aberrant phosphotau fibrillar polymers in neural cultured cells. *Eur J Biochem*
5 **269**, 1484-9.
6
7 Picciotto, M. R., and Zoli, M. (2008). Neuroprotection via nAChRs: the role of nAChRs in
8 neurodegenerative disorders such as Alzheimer's and Parkinson's disease. *Front Biosci* **13**,
9 492-504.
10 Pizzi, M., Valerio, A., Arrighi, V., Galli, P., Belloni, M., Ribola, M., Alberici, A., Spano, P.,
11 and Memo, M. (1995). Inhibition of glutamate-induced neurotoxicity by a tau antisense
12 oligonucleotide in primary culture of rat cerebellar granule cells. *Eur J Neurosci* **7**, 1603-13.
13 Price, J. L., Davis, P. B., Morris, J. C., and White, D. L. (1991). The distribution of tangles,
14 plaques and related immunohistochemical markers in healthy aging and Alzheimer's disease.
15 *Neurobiol Aging* **12**, 295-312.
16
17 Quesada, A., Lee, B. Y., and Micevych, P. E. (2008). PI3 kinase/Akt activation mediates
18 estrogen and IGF-1 nigral DA neuronal neuroprotection against a unilateral rat model of
19 Parkinson's disease. *Dev Neurobiol* **68**, 632-44.
20
21 Shaw, S., Bencherif, M., and Marrero, M. B. (2002). Janus kinase 2, an early target of alpha 7
22 nicotinic acetylcholine receptor-mediated neuroprotection against Abeta-(1-42) amyloid. *J*
23 *Biol Chem* **277**, 44920-4.
24
25 Soletti, R. C., del Barrio, L., Daffre, S., Miranda, A., Borges, H. L., Moura-Neto, V., Lopez,
26 M. G., and Gabilan, N. H. Peptide gomesin triggers cell death through L-type channel calcium
27 influx, MAPK/ERK, PKC and PI3K signaling and generation of reactive oxygen species.
28 *Chem Biol Interact* **186**, 135-43.
29
30 Spittaels, K., Van den Haute, C., Van Dorpe, J., Geerts, H., Mercken, M., Bruynseels, K.,
31 Lasrado, R., Vandezande, K., Laenen, I., Boon, T., Van Lint, J., Vandenheede, J., Moechars,
32 D., Loos, R., and Van Leuven, F. (2000). Glycogen synthase kinase-3beta phosphorylates
33 protein tau and rescues the axonopathy in the central nervous system of human four-repeat tau
34 transgenic mice. *J Biol Chem* **275**, 41340-9.
35
36 Srivastava, A. K., and Pandey, S. K. (1998). Potential mechanism(s) involved in the
37 regulation of glycogen synthesis by insulin. *Mol Cell Biochem* **182**, 135-41.
38
39 Stambolic, V., and Woodgett, J. R. (1994). Mitogen inactivation of glycogen synthase kinase-
40 3 beta in intact cells via serine 9 phosphorylation. *Biochem J* **303** (Pt 3), 701-4.
41
42 Tanaka, T., Zhong, J., Iqbal, K., Trenkner, E., and Grundke-Iqbal, I. (1998). The regulation of
43 phosphorylation of tau in SY5Y neuroblastoma cells: the role of protein phosphatases. *FEBS*
44 *Lett* **426**, 248-54.
45
46 Tian, D., Zhu, M., Li, J., Ma, Y., and Wu, R. (2009). Cigarette smoke extract induces
47 activation of beta-catenin/TCF signaling through inhibiting GSK3beta in human alveolar
48 epithelial cell line. *Toxicol Lett* **187**, 58-62.
49
50 Tian, Q., Lin, Z. Q., Wang, X. C., Chen, J., Wang, Q., Gong, C. X., and Wang, J. Z. (2004).
51 Injection of okadaic acid into the meynert nucleus basalis of rat brain induces decreased
52 acetylcholine level and spatial memory deficit. *Neuroscience* **126**, 277-84.
53
54 Tizabi, Y., Manaye, K. F., Smoot, D. T., and Taylor, R. E. (2004). Nicotine inhibits ethanol-
55 induced toxicity in cultured cerebral cortical cells. *Neurotox Res* **6**, 311-6.
56
57 Uberti, D., Rizzini, C., Spano, P. F., and Memo, M. (1997). Characterization of tau proteins in
58 human neuroblastoma SH-SY5Y cell line. *Neurosci Lett* **235**, 149-53.
59
60 Ueda, M., Iida, Y., Kitamura, Y., Kawashima, H., Ogawa, M., Magata, Y., and Saji, H.
(2008). 5-Iodo-A-85380, a specific ligand for alpha 4 beta 2 nicotinic acetylcholine receptors,
prevents glutamate neurotoxicity in rat cortical cultured neurons. *Brain Res* **1199**, 46-52.
Wang, J. Z., Gong, C. X., Zaidi, T., Grundke-Iqbal, I., and Iqbal, K. (1995).
Dephosphorylation of Alzheimer paired helical filaments by protein phosphatase-2A and -2B.
J Biol Chem **270**, 4854-60.

1
2
3 Wang, Q. M., Fiol, C. J., DePaoli-Roach, A. A., and Roach, P. J. (1994). Glycogen synthase
4 kinase-3 beta is a dual specificity kinase differentially regulated by tyrosine and
5 serine/threonine phosphorylation. *J Biol Chem* **269**, 14566-74.

6
7 Whitehouse, P. J., Price, D. L., Struble, R. G., Clark, A. W., Coyle, J. T., and Delon, M. R.
8 (1982). Alzheimer's disease and senile dementia: loss of neurons in the basal forebrain.
9 *Science* **215**, 1237-9.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60





Fig. 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49




Fig. 2

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Experimental Protocol		
5 min	Fresh medium	OA 30nM 16h
1 h	Fresh medium	OA 30nM 16h
4h	Fresh medium	OA 30nM 16h
24 h		OA 30nM 16h ↑
Agonists pre-incubation time		LDH measurement









Fig. 3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49




Fig. 5












Fig. 6

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49





1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49






Fig. 10