



Citation for published version:

Margos, G, Castillo-Ramírez, S & Hoen, AG 2012, 'Phylogeography of Lyme borreliosis-group spirochetes and methicillin-resistant *Staphylococcus aureus*', *Parasitology*, vol. 139, no. 14, pp. 1952-1965.
<https://doi.org/10.1017/S0031182012000741>

DOI:

[10.1017/S0031182012000741](https://doi.org/10.1017/S0031182012000741)

Publication date:

2012

Document Version

Publisher's PDF, also known as Version of record

[Link to publication](#)

© Cambridge University Press 2012

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Phylogeography of Lyme borreliosis-group spirochetes and methicillin-resistant *Staphylococcus aureus*

GABRIELE MARGOS¹, SANTIAGO CASTILLO-RAMÍREZ¹ and ANNE GATEWOOD HOEN²

¹Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, UK

²Department of Community and Family Medicine, Dartmouth Medical School, Hanover, NH 03756, USA

(Received 6 December 2011; revised 30 March 2012; accepted 14 April 2012; first published online 23 May 2012)

SUMMARY

Multilocus sequence typing (MLST) and multilocus sequence analysis (MLSA) have revolutionized understanding the global epidemiology of many medically relevant bacteria utilizing a number, mostly seven, of housekeeping genes. A more recent introduction, single nucleotide polymorphisms (SNPs), constitutes an even more powerful tool for bacterial typing, population genetic studies and phylogeography. The introduction of massive parallel sequencing has made genome re-sequencing and SNP discovery more economical for investigations of microbial organisms. In this paper we review phylogeographic studies on Lyme borreliosis (LB)-group spirochetes and methicillin-resistant *Staphylococcus aureus* (MRSA). Members of the LB-group spirochetes are tick-transmitted zoonotic bacteria that have many hosts and differ in their degree of host specialism, constituting a highly complex system. MRSA is a directly transmitted pathogen that may be acquired by contact with infected people, animals or MRSA-contaminated objects. For the LB-group spirochetes, MLSA has proved a powerful tool for species assignment and phylogeographic investigations while for *S. aureus*, genome-wide SNP data have been used to study the very short-term evolution of two important MRSA lineages, ST239 and ST225. These data are detailed in this review.

Key words: Lyme borreliosis-group spirochetes, MLST, methicillin-resistant *Staphylococcus aureus*, MRSA, SNPs, phylogeography.

INTRODUCTION

The term ‘phylogeography’ was coined by Avise and colleagues in 1987 (Avise *et al.* 1987). Originally, phylogeography was introduced to bridge the gap between population genetics and systematics in eukaryotic organisms. Molecular markers were used to explore microevolutionary processes and the phylogenetic relationship in populations to explain the evolution of species. Initial studies in phylogeography relied on mitochondrial DNA (mtDNA) due to its fast evolution, no or limited recombination and haploidy which makes these loci suitable for phylogenetic inferences. The properties of mtDNA in animals yielded observations around geographic structuring of genetic lineages. This correspondence between genealogy and geographic origin, in turn, allows for inferences related to population history and demography. Since the introduction of phylogeography as a discipline much progress has been made in terms of (1) development of a statistical framework in which phylogeographic hypothesis can be contrasted (e.g. Knowles and Maddison, 2002; Garrick *et al.* 2010); (2) the amount of genetic information used

(multilocus analyses, sequencing technology and single nucleotide polymorphism [SNPs]) (e.g. Aanensen and Spratt, 2005; Chan, 2005; Hall, 2007) and (3) the incorporation of geographic elements through geographic information systems (e.g. Kidd and Ritchie, 2006).

Due to similarities in genome characteristics (haploidy and fast mutation rates) between mitochondria and bacterial species, there has been a clear application of the phylogeographic techniques developed for animals using mtDNA to bacterial taxa, including pathogenic bacteria. However, the use of phylogenetics to infer the very recent past is generally limited by the type or number of loci used. While it was initially suggested that homoplasy in fast evolving loci may limit phylogenetic inferences to the very recent past such as single hospital outbreaks (Spratt, 1999), a more recent view is that the use of such loci is hampered by processes like horizontal gene transfer or gene loss. This complicates their use for applications such as tracing the source of hospital outbreaks because these loci very likely do not reflect the phylogenetic relationships between the strains considered. On the other hand, genes that are highly conserved, such as 16S RNA, are usually not able to provide a resolved picture of intra-species bacterial lineages. The same is true for genes under balancing selection, because these loci might reflect the

Corresponding author: Gabriele Margos, University of Bath, Department of Biology and Biochemistry, 3 South, Claverton Down, Bath BA2 7AY, U.K. Tel: 0044-1225-385116. Email: gmargos1@gmail.com

evolutionary history of the locus rather than that of the population or species. Slowly evolving loci (such as housekeeping genes) can be useful for medically important bacteria for explaining long-term historic patterns of emergence or co-evolution with hosts, but their limited genetic variation may make it difficult to estimate processes over shorter time-spans when only few loci are used. Until recently, power was limited for investigating finer scale landscape ecological processes that have shaped contemporary populations, partially as a result of complex demographic processes such as sequential population bottlenecks, population expansions, founder effects, hybridization or gene flow.

A more recent introduction for microevolutionary studies, i.e. at the population level in a spatial context, is landscape genetics. As it combines landscape ecology with molecular population genetics, landscape genetics aims to explain local adaptation and gene flow or barriers to it, providing a powerful approach for determining populations or their substructure and for quantifying their connectivity. An advantage of landscape genetics over conventional population genetics is that populations and their boundaries do not need to be defined *a priori*. It has been developed to investigate genetic variation under consideration of landscape heterogeneity potentially at finer spatial and temporal scales than phylogeography (Manel *et al.* 2003; Storfer *et al.* 2007; Reisen, 2010). It has been predicted that an increase in the number of genetic markers used, including the use of genome-wide SNPs, will allow the optimal choice of genetic marker for the study in question, and this will greatly enhance the power to identify the underlying processes of population heterogeneity. Indeed, as shown by recent studies on bacterial pathogens, interrogation of genome-wide SNPs and their phylogenetic inferences can inform about bacterial relationships at ultra-fine temporal and spatial scales, i.e. they might bridge the gap between phylogeography and landscape genetics for these systems.

Phylogeography aims to untangle microevolutionary processes that lead to diversification and ultimately speciation. However, multilocus analyses and the availability of fully sequenced bacterial genomes have revealed that horizontal gene transfer between and within bacterial species may be more common than previously thought, which raised questions about the nature of bacterial species and whether bacterial species do exist (Lan and Reeves, 2001; Cohan, 2002; Konstantinidis and Tiedje, 2005; Achtman and Wagner, 2008). Although a theory-based prokaryotic species concept has not been agreed on, it has been suggested repeatedly that cohesive forces cause genetic clustering of bacterial populations (Rossello-Mora and Amann, 2001; Cohan, 2002; Gevers *et al.* 2006; Konstantinidis *et al.* 2006) and that such clusters may have quintessential properties to be classified as species. As a pragmatic approach

most bacteriologists define bacterial species through genetic or phenotypic cut-offs (Goris *et al.* 2007), and multilocus sequence analysis (MLSA) has been suggested as a convenient tool for bacterial species assignment (Gevers *et al.* 2005; Bishop *et al.* 2009). It has been shown for several bacteria that sequence clusters are remarkably congruent with bacterial ecotypes (Koeppel *et al.* 2008; Margos *et al.* 2009) lending support to such an approach.

The availability of many fully sequenced genomes has led to the concept that bacterial genomes consist of a conserved 'core' genome which basically codes for housekeeping functions, and an 'accessory' genome which is susceptible to horizontal gene transfer and gene loss and where more mobile genetic elements (plasmids, phage, transposons, etc.) are located. While the core genome may be used to ascertain the evolutionary relationship among bacteria and to generate species trees, horizontal gene transfer of genetic elements may occasionally occur between (un)related prokaryotes and can lead to 'evolutionary jumps' (e.g. resistance to antibiotics or host switches). Such events may enable new strains to emerge and to occupy a different ecological niche. These ecotypic changes may not be observed immediately at the level of core genes (by means of MLST) as suggested for *Yersinia pestis* (Achtman *et al.* 1999; Kotetishvili *et al.* 2005), *Bacillus anthracis* (Keim *et al.* 2004) or other so-called monomorphic bacterial pathogens (Achtman and Wagner, 2008; Pearson *et al.* 2009). Regardless of such concerns, it appears that bacterial lineages do exist which differ considerably in their geno- and phenotypes, at least in some instances even if this cannot be applied to all bacterial taxa (Hanage *et al.* 2005; Corander *et al.* 2011).

BACTERIAL TYPING TOOLS

Multilocus sequence typing (MLST) and multilocus sequence analysis (MLSA)

For many bacterial genera, multilocus sequence typing (MLST) or multilocus sequence analyses (MLSA) have considerably advanced the knowledge of intraspecies bacterial relationships (see mlst.net or pubmlst.org). These techniques utilize slowly evolving loci, such as enzyme encoding genes that are under purifying selection, which have usually only few mutational changes that may not give sufficient phylogenetic resolution to distinguish closely related bacteria strains when studied individually. Therefore, to increase the phylogenetic signal and to understand the longer term relationship of bacteria, multiple loci are included for population genetic analyses (Maiden *et al.* 1998; Maiden, 2006). Compelling reasons for this choice of marker type were the assumption of nearly neutral variation, ability to detect or buffer against recombination, and high discriminatory power (reviewed by Maiden, 2006). However, the

phylogenetic signal obtained, even from several loci, may be limited for the inference of very short-term or deep evolutionary relationships in 'monomorphic' species such as *Yersinia pestis* or *Bacillus anthracis* (reviewed by Achtman and Wagner, 2008) or in species with elevated levels of homologous recombination (Pearson *et al.* 2009).

Genome-wide single nucleotide polymorphisms (SNPs)

SNPs have been shown to be a powerful instrument for understanding bacterial evolution (even for genetically monomorphic microbes) (Van Ert *et al.* 2007; Achtman and Wagner, 2008; Holt *et al.* 2008; Harris *et al.* 2010). The introduction of massive parallel sequencing and improvements such as increased read length and depth, or reduced error rates have made genome re-sequencing using next generation platforms more economical for investigations of microbial organisms (Hall, 2007). The large number of SNPs that can be found in genomes of many prokaryotes offers unprecedented power to infer bacterial relationships at temporal scales that cover deep and very fine evolutionary relationships. SNPs can be filtered according to whether they are from coding or non-coding regions, from core or accessory genome, permitting enormous scope for evolutionary inferences.

The large number of software tools which are available to analyse MLSA and SNP data facilitate such approaches (Excoffier and Heckel, 2006). Such data will not only permit much more precise insights into the history of species, they will reveal complications such as incomplete lineage sorting, branch length heterogeneity or recombination.

Here, we review phylogeographic data for two different types of bacterial pathogens: one vector-borne pathogen, *Borrelia burgdorferi* sensu lato, also known as Lyme borreliosis (LB)-group spirochetes and one directly transmitted human pathogen, methicillin-resistant *Staphylococcus aureus* (MRSA). While members of the LB-group spirochetes are tick-transmitted zoonotic bacteria that have many hosts and differ in their degree of host specialism, MRSA was initially thought to occur only in humans, but now it has been found in horses, dogs, cast, cattle and pigs. MRSA is transmitted by direct contact with an infected or colonized person or animal, and also by contact with MRSA-contaminated objects. In both cases, pathogen dispersal is host dependent. In the case of LB-group spirochetes, MLSA data proved to be very useful in ecotype and species determination (Richter *et al.* 2006; Postic *et al.* 2007; Chu *et al.* 2008; Margos *et al.* 2009, 2010; Rudenko *et al.* 2009a,b; Crowder *et al.* 2010) and to distinguish between long-term and more recent demographic and evolutionary processes (Margos *et al.* unpublished). In *S. aureus* MLST was used to study the global population structure as well as the dispersal of particular clonal

complexes (Enright *et al.* 2000). In MLST analyses, clonal complexes are defined by using a model of clonal expansion and divergence. The eBURST algorithm was developed to provide a convenient method to establish relationships of bacterial populations via allelic multilocus sequence profiles. It clusters STs as disjointed tree collections and uses a set of hierarchical rules to determine single locus (SLV) and double locus variants (DLV) and founder stains of complexes that consist of SLV and DLVs. Furthermore, very recently genome-wide SNP data have been used to study the very short-term evolution of two important MRSA lineages, ST239 and ST225, which is described in detail below.

EXAMPLES OF BACTERIAL PHYLOGEOGRAPHIC STUDIES

LB-group spirochetes: a vector-transmitted bacterial species complex

The LB-group spirochetes form a species complex which now comprises 18 named species (Table 1) (Margos *et al.* 2011) but more species are likely to be described in the near future as highly diverse LB strains have been found in environmental samples (e.g. Scott *et al.* 2010, Lane personal communication).

LB-group spirochetes are transmitted by ticks belonging to the *Ixodes ricinus-persulcatus* species complex. Vector competence has been demonstrated for a number of *Ixodes* species (reviewed by Eisen and Lane, 2002) but the extent to which vector competence for LB-group spirochete species varies is by no means fully understood. The main vector species transmitting LB-group spirochetes to humans are *I. pacificus* in Western North America, *I. scapularis* in Central and Eastern North America, *I. ricinus* in Europe and *I. persulcatus* in Eastern Europe and Asia (Ogden *et al.* 2011a).

A hallmark of LB-group spirochetes is the pronounced host associations of some of its members, which means that these hosts support completion of the whole transmission cycle, i.e. tick to host and host to tick (Kurtenbach *et al.* 1998a, 2002; Humair and Gern, 2000). These host specialisations match the ability of spirochete species to deflect complement-mediated lysis of the respective host species (Kurtenbach *et al.* 1998a, 2002). Furthermore, they have been shown to play an important role in the geographic distribution and spread of LB-group spirochetes (Vollmer *et al.* 2011).

Although less well studied, the compatibility of LB-group spirochetes and vector associations may also impact the distribution range of LB-group spirochetes (Fig. 1) and both host and vector associations contribute to niche formation which may lead to speciation of LB-group spirochetes (reviewed by Randolph, 2004; Tsao, 2009; Kurtenbach *et al.* 2010; Margos *et al.* 2011).

Table 1. Lyme borreliosis group spirochetes

| Species name | Reservoir hosts | Late disease symptoms |
|-------------------------------------|-----------------------|---|
| <i>B. burgdorferi</i> sensu stricto | generalist | arthritis |
| <i>B. afzelii</i> | rodents, insectivores | acrodermatitis |
| <i>B. garinii</i> | birds | neuroborreliosis |
| <i>B. bavariensis</i> (p) | rodents | neuroborreliosis |
| <i>B. spielmanii</i> | door mice, hedgehogs | unknown |
| <i>B. lusitaniae</i> | lizards | unknown |
| <i>B. valaisiana</i> | birds | unknown pathogenicity in humans and animals |
| <i>B. japonica</i> | rodents | |
| <i>B. tanukii</i> | rodents | |
| <i>B. turdi</i> | birds | |
| <i>B. sinica</i> | rodents | |
| <i>B. yangtze</i> (p) | rodents | |
| <i>B. andersonii</i> | rabbits, birds | |
| <i>B. bissettii</i> | rodents | |
| <i>B. californiensis</i> | rodents | |
| <i>B. carolinensis</i> | rodents | |
| <i>B. americana</i> (p) | birds | |
| <i>B. kurtenbachii</i> (p) | rodents | |

p = proposed species.

Members of the LB-group spirochetes are distributed across the northern temperate zones of the globe but the species and strains are not evenly distributed globally, across continents or regions (Kurtenbach *et al.* 2006; Margos *et al.* 2011). This is due to complex multifactorial processes that govern the ecology and evolution of LB-group spirochetes including host and vector associations, climate and landscape features and long-term evolutionary and demographic processes (Randolph, 2000, 2004; Kurtenbach *et al.* 2010). LB-group spirochetes reside either in their hosts or in their vector ticks. It is conceivable that the strong impact that Pleistocene climate variations had on vertebrate and invertebrate species and their present day geographic ranges (Waters, 1963; Hewitt, 1999) will also have impacted the distribution of LB spirochete species and sequence types. Therefore, the post-glacial recolonization by vertebrate and vector hosts of northern regions in Europe and North America might be reflected in the sequence type and species distribution of LB-group spirochetes (Vollmer *et al.* unpublished).

Of the 18 species recognized in the LB group of spirochetes to date, seven occur in North America, eight in Europe and eight in Asia. There is some overlap between the regions in that three species, *B. burgdorferi* sensu stricto, *B. bissettii*, and possibly *B. kurtenbachii*, occur in Europe and North America (Picken *et al.* 1995, 1996, 1997; Picken and Picken 2000) and three species, *B. afzelii*, *B. garinii*, and *B. bavariensis*, occur in Europe and Asia but remarkably no overlap of species from North America and Asia has been described so far (Masuzawa, 2004; Margos *et al.* 2010). This is interesting in view of the origin of LB-group spirochetes, their spread and the global colonisation of this group of spirochetes.

North America. The picture of LB-group spirochetes in North America has recently become more complex. As far as LB-group spirochetes and their vectors are concerned, there are several distinctly different regions on the North American continent: coastal regions west of the Rocky Mountains (California, Oregon, Washington, British Columbia), the southeastern States (Florida, Georgia, South Carolina, North Carolina, Virginia, Maryland), north-eastern States (Pennsylvania, New Jersey, New York, Connecticut, Massachusetts, New Hampshire, Maine, Vermont) and midwestern states (Wisconsin, Minnesota, Missouri, Illinois). The main vector species maintaining the bacterial species in natural transmission cycles in these regions are *I. pacificus* and *I. spinipalpis* in the west, *I. affinis* and *I. minor* in the southeast and *Ixodes scapularis* in northeast and mid-west (Brown and Lane, 1992; Maupin *et al.* 1994; Lin *et al.* 2004; Brown *et al.* 2006; Diuk-Wasser *et al.* 2006, 2010; Hamer *et al.* 2010; Maggi *et al.* 2010).

The distribution of species and genotypes suggests a turbulent evolutionary history of LB spirochetes in North America with long-term (millions of years ago) and short-term (500 years ago until present) demographic events (Hoen *et al.* 2009; Margos *et al.* unpublished). The long-term events may have led to separation of strains, isolation and speciation. It has been suggested that the present day distribution of the species may be the result of southward migration during glacial periods and northward migration during interglacials (Humphrey *et al.* 2010; Margos *et al.* 2010). But so far, this is speculative as there are no data to estimate mutation rates or speciation events for LB spirochetes and not all *Borrelia* species that are likely to reside on this continent are currently known (Scott *et al.* 2010) and can be included into phylogenetic analysis.

Geographic distribution and vector associations of Lyme borreliosis group spirochetes




Fig. 1. Geographic distribution and vector associations of Lyme borreliosis group spirochetes. The figure shows combinations of main ixodid vectors and Lyme borreliosis group spirochetes they transmit. While for many Lyme borreliosis group spirochetes vector associations have been established, this is not the case for all species, e.g. *B. californiensis*, *B. carolinensis*, *B. kurtenbachii* in North America. ^aFor *B. bissettii* the vector in Europe is uncertain. ^b*B. burgdorferi*, *B. valaisiana* and *B. lusitaniae* are transmitted by *I. ricinus* in Eastern Europe but their prevalence is low. These species do not appear to be transmitted by *I. persulcatus*. Reproduced with modifications and with permission from the author.

B. americana, *B. bissettii* and *B. burgdorferi* appear in eastern and western coastal regions (Rudenko *et al.* 2009b; Margos *et al.* 2010) while species such as *B. californiensis* or *B. carolinensis* seem to have limited distribution ranges and were described from the west or southeast, respectively (Postic *et al.* 2007; Rudenko *et al.* 2009a). *B. andersonii* seems to be distributed through the eastern range of LB spirochete occurrence (Lin *et al.* 2004; Hamer *et al.* 2010). The distribution range of *B. kurtenbachii* needs to be investigated further as it was previously subsumed in *B. bissettii* and only recently proposed as a separate species (Margos *et al.* 2010). Several samples previously described as *B. bissettii* in the various regions need to be re-evaluated as they showed RFLP pattern similar to *B. kurtenbachii* (Lin *et al.* 2001, 2005; Oliver *et al.* 2003). While *B. bissettii* was associated with *I. spinipalpis* and *I. pacificus* and the woodrats *Neotoma mexicana* or *Neotoma fuscipes* in Colorado and California, respectively, *B. kurtenbachii* were isolated from the vole *Microtus pennsylvanicus* in Illinois (Picken and Picken, 2000) and the vector is unknown. Other *B. kurtenbachii* strains were isolated from host-attached *I. scapularis*, which raised the possibility that *B. kurtenbachii* transmission cycles differ ecologically from that of *B. bissettii*. These data underpin the importance of using appropriate typing tools in order to understand the evolutionary history of this species complex. This needs to be particularly emphasized as a relatively new species, *B. miyamotoi*, which represents a relapsing-fever spirochete that

occurs sympatrically with LB-group spirochetes (Barbour *et al.* 2010).

B. burgdorferi and *B. bissettii* are without doubt the species with the widest distributions as both occur not only in most regions of the USA where LB-group spirochetes are found but also in Europe. Based on analyses of the genetic diversity of the outer surface protein C (*ospC*) it has been suggested that *B. burgdorferi* originated in North America (Marti Ras *et al.* 1997), and that migration between Europe and North America occurred in the past 200 years (Qiu *et al.* 2008). In contrast, data based on MLST of housekeeping genes suggested that *B. burgdorferi* originated in Europe (Margos *et al.* 2008). Analysis of the pairwise mismatch distribution of variation of *B. burgdorferi* samples from northeastern and mid-western USA suggested a population expansions that took place thousands or million of years ago (Hoen *et al.* 2009). This time estimate makes the presence of *B. burgdorferi* in North America likely during the glacial-interglacial periods of the Pleistocene (2.5 million to about 10,000 years ago).

Evidence for more recent demographic processes of *B. burgdorferi* populations comes from the analysis of allele frequencies in the various regions. *B. burgdorferi* was isolated and the participation of ticks as transmission vectors was demonstrated in the early 1980s (Burgdorfer *et al.* 1982). However, museum specimens of *Peromyscus* from Massachusetts collected in 1894 were PCR positive for *B. burgdorferi* DNA (Marshall *et al.* 1994). In addition, reports suggesting

LB cases (and presence of *B. burgdorferi*) were from Cape Cod in 1962 and Wisconsin in 1969 (Scrimanti, 1970; Steere *et al.* 1986) suggesting that the bacterium was widely distributed in North America.

The epidemic increase of human LB in parts of the USA in the past 40–50 years has been attributed to range expansions of vector and pathogen and this has been linked to reforestation and the exponential increase in numbers of white tail deer (*Odocoileus virginianus* Zimmermann) since 1900 (Barbour and Fish, 1993; Falco *et al.* 1995; McCabe and McCabe, 1997). Before 1500, white-tailed deer populations covered a large area, probably ranging from Florida to New England in the east to British Columbia in the west of North America. The distribution of deer does not necessarily match the distribution range of *Ixodes* spp. tick vectors because other factors such as climate play an important role. It is, however, highly likely that *Ixodes* spp. populations were also widely distributed (Brownstein *et al.* 2003). Due to over-hunting, woodland clearance and land-use changes, by 1900 *O. virginianus* had been driven to almost extinction (Christensen, 1959; Cronon, 1983; Halls, 1984; McCabe and McCabe, 1997). The lack of the main hosts of adult ticks most certainly led to a decline of *Ixodes* spp. populations (Wilson *et al.* 1985) which in the 1920s were only known to occur on some islands off the coast of Massachusetts, on Long Island and Wisconsin (Jackson and DeFoliart, 1970; Spielman, 1994). It would be expected that a decline in *Ixodes* spp. produced a severe bottleneck for *B. burgdorferi* populations. Presently, in the northeast USA *B. burgdorferi* appears to be the only LB species transmitted by *I. scapularis*.

MLST was used to analyse geo-referenced samples from the Northeast (New England and southern Canada), the upper midwest, southern Canada (Hoen *et al.* 2009; Ogden *et al.* 2010, 2011b) and California (Margos *et al.* unpublished) and to investigate potential population subdivisions in a spatial context. These analyses suggest that the contemporary populations of *B. burgdorferi* are geographically separated with no or only limited gene flow between the regions. The determination of boundaries between sub-populations strongly supported suggestions of local expansions of regional *B. burgdorferi* populations since the middle of the last century and was consistent with hypotheses of bottlenecks occurring in the recent evolutionary history (Margos *et al.* unpublished). Phylogenetic and other analyses indicated that the populations from all three regions are genetically closely related and revealed signatures of admixture, particularly between STs found in northeast and midwest. It was hypothesized that these signals reflect an ancestral population structure before the recent population bottleneck. Interestingly, some strains from California showed levels of divergence that suggest longer-term separation from STs clustering in

neighbouring phylogenetic clades. These data revealed genetic signatures of contemporary and ancestral population structure of *B. burgdorferi* in North America and these were in agreement with current knowledge of the recent historical events of major ecological components impacting the transmission cycle of this parasitic microorganism.

These studies also suggested that gene-flow occurred in an east-to-west direction as most founders of clonal complexes were located in the northeast (Hoen *et al.* 2009; Ogden *et al.* 2011b). Whilst this is consistent with suggestions that *B. burgdorferi* originated in Europe (Margos *et al.* 2008) and absent in Asia (Masuzawa, 2004) (therefore, not colonising North America from the west), bootstrap support of clonal complex founders was not particularly strong and more samples need to be analysed to confirm these hypotheses. Furthermore, genetic analyses of the present day sub-populations would benefit from use of tools providing higher resolution. As exemplified by the *S. aureus* data, genome-wide SNP analyses may provide the discriminatory power required for such an enterprise.

Europe and Asia. In Europe, eight LB spirochete species occur (*B. afzelii*, *B. garinii*, *B. bavariensis*, *B. lusitanae*, *B. spielmanii*, *B. valaisiana*, *B. burgdorferi*, and *B. bissetii*) of which three, namely *B. afzelii*, *B. garinii*, and *B. bavariensis* (or *B. bavariensis*-like strains), also occur in Asia. Whether or not *B. kurtenbachii* occurs in Europe needs to be evaluated (Picken *et al.* 1996). It is interesting to note that some species have been described as occurring more regionally in Asia and Europe. These include *B. japonica*, *B. turdi*, *B. tanukii*, which have only been described from Japan while *B. sinica* and *B. yangtze* were found in southeastern China. Several of these species are transmitted by specialised vectors (*B. turdi* by *I. turdus*; *B. tanukii* by *I. tanuki*; *B. sinica* and *B. japonica* by *I. ovatus*), which may limit the geographic distribution range of these species. The notion that vector associations may determine the geographic range that a species can occupy is supported by the following observations: (1) *B. valaisiana* is widely distributed in Europe in the distribution range of *I. ricinus* but whether it occurs in Asia is uncertain. It is not commonly reported in *I. persulcatus*-dominated habitats and a single report in *I. columnae* exists from Japan (Korenberg *et al.* 2002; Bormane *et al.* 2004; Masuzawa, 2004). Phylogenetically the most closely related species is the rodent-associated species *B. yangtze* (previously known as *B. valaisiana*-like strains) which is transmitted by *I. granulatus* in China and *I. nipponensis* in Korea but not by *I. persulcatus* (Masuzawa, 2004; Chu *et al.* 2008). (2) Similarly, *B. burgdorferi* has not been reported from Asia. (3) *B. garinii* is the most widespread species in Eurasia and the Arctic and

Antarctic Circles where *I. ricinus*, *I. persulcatus* and *I. urea* serve as vectors, respectively. It has also been reported to be present in sea bird colonies at the east coast of the American continent (Smith *et al.* 2006) but it has not (yet) spread inland of the American continent which may have to do with vector incompetence of local *Ixodes* or reservoir incompetence of vertebrate hosts.

Rodent-adapted variants of *B. garinii* were described from Europe where they were named OspA serotype 4 and in Asia where they were designated NT29. In western Europe these strains have a focal distribution and are highly invasive for humans (Wilske *et al.* 1996; Fingerle *et al.* 2002). NT29 strains (also known as ribotype IV or Asian-type *B. garinii* (Nakao *et al.* 1994)), were known to occur in Russia and Asia. MLSA revealed that OspA serotype 4 and NT29 are genetically closely related but are distinct from bird-associated *B. garinii* and a new species, *B. bavariensis*, has been proposed (Margos *et al.* 2009). MLSA revealed that the majority of human cases in Japan was caused by strains closely related to *B. bavariensis* (Takano *et al.* 2011). Strains from Asia show much more genetic heterogeneity suggesting that the species likely originated in Asia. However, it has been reported that NT29 strains are only transmitted by *I. persulcatus* while OspA serotype 4 strains are transmitted by *I. ricinus*. This and the genetic homogeneity of *B. bavariensis* in Europe suggest that *B. bavariensis* made its way into a new vector and it is possible that this currently leads to speciation. This hypothesis requires testing by transmission assays.

In Europe all of the known LB species (*B. burgdorferi*, *B. garinii*, *B. afzelii*, *B. bavariensis*, *B. spielmannii*, *B. lusitaniae*, and *B. valaisiana*) are transmitted by the generalist vector *I. ricinus* with the exception of perhaps *B. bissettii*. This tick species feeds on birds as well as on reptiles or small-to-medium sized mammals. It renders the tick an ideal place for mixing different strains and species and, therefore, the host associations described in Europe are not driven by adaptation to an endophilic tick with a narrow host preference but are truly host driven (Kurtenbach *et al.* 1998b). The host associations described for LB-group species in Europe have provided an ideal system to compare and contrast the impact of hosts on the population structure, dispersal or spread of the different species. The phylogeography of several LB species has been analysed using MLST/MLSA.

The first species to be analysed was *B. lusitaniae* using samples collected in two ecologically different locations north (Mafra) and south (Grandola) of Lisbon (Vitorino *et al.* 2008). While Mafra resembles a European deciduous forest habitat and harbours several *Borrelia* species, Grandola is much drier and hotter, and *B. lusitaniae* is the sole *Borrelia* species

described from here. Lizards of the family Lacertidae are one of the main hosts for *I. ricinus* and adult ticks can be found to have very high infection prevalences (up to 70%) (De Michelis *et al.* 2000). MLST on samples of this *Borrelia* species revealed a pronounced fine-scale geographic structure of populations from Mafra and Grandola suggesting limited migration between the two localities. It was further suggested that the population structure displayed by the *Borrelia* populations reflected the highly parapatric population structure of their lizard hosts underpinning the importance of host associations in the geographic dispersal of parasitic microorganisms. In view of several other studies on LB spirochetes which show strong spatial structuring but appear polyphyletic in phylogenies (such as *B. afzelii*) the (almost) monophyletic clustering of the two regional *B. lusitaniae* populations warrants further investigations (Vitorino *et al.* 2008).

Of the eight LB species occurring in Europe, *B. afzelii*, *B. garinii* and *B. valaisiana* are the most common. Vollmer *et al.* (2011) used an MLSA system to compare the population structure of the species *B. valaisiana* and *B. garinii* which are associated with highly mobile hosts (passerine birds) (Hanincova *et al.* 2003b; Taragel'ova *et al.* 2008; Dubska *et al.* 2009), and the rodent-associated species *B. afzelii* (Gern *et al.* 1998; Hanincova *et al.* 2003a; Skuballa *et al.* 2007). As predicted, on a European scale, the two bird-associated species showed spatial mixing (although this was more pronounced in *B. garinii*) while *B. afzelii* showed spatial structuring. However, when bird associated *B. garinii* strains from Asia were included into the analyses, the majority of European strains clustered independently from Asian strains indicating spatial structuring at this continental scale even for this highly mobile species.

Spatial structuring of *B. garinii* was also reported when samples from marine and terrestrial transmission cycles were compared. Gomez and colleagues (Gomez-Diaz *et al.* 2011) investigated the population structure of *B. garinii* maintained in the different transmission cycles using MLSA of chromosomally located loci and two plasmid-encoded genes (*ospA* and *ospC*). The samples included in the study consisted of strains from the terrestrial transmission cycle and two distinct marine transmission cycles, Pacific and Atlantic. Analyses of strains from the Pacific, Atlantic and terrestrial *B. garinii* populations using only the chromosomal loci suggested population structuring although clades containing strains from the various regions appeared polyphyletic. Population structuring between the two marine cycles and the terrestrial cycles was highly significant but no evidence of host associated genetic structure was observed. This is interesting in view of the described host races of *I. uriae* which suggested some form of adaptation between vector and host (McCoy

et al. 2003, 2005). Interestingly, the *B. garinii* type strain 20047 clustered separately from the other terrestrial strains seemingly closer related to *Borrelia* strains detected in Atlantic seabirds. Although the plasmid-encoded loci showed higher genetic variation and evidence for recombination (particularly when both marine and terrestrial strains were considered) they did not reveal population structure. Based on their results, the authors suggested that contact zones exist where exchange between the different transmission cycles occurs.

B. afzelii populations showed pronounced structuring at the European scale (Vollmer *et al.* 2011). When Asian *B. afzelii* were included into the analyses, the majority of these samples formed a distinct cluster in phylogenetic trees and isolation-by-distance between European and Asian STs was significant at the continental scale but not at a European scale (Vollmer *et al.* unpublished). Interestingly, the European STs formed two phylogenetic clades, one consisting mostly of STs found in eastern Europe (Latvia) while the other one carried mostly STs found in western Europe such as England and France. Strains from Germany were scattered inside and outside these clades. This clustering of STs evoked some extent of similarity to geographic structuring of mammalian species due to post-glacial recolonisation of northern Europe (Hewitt, 1999). The last of the Pleistocene glacial maxima is dated to about 18,000 years before present (BP) although there might be continental differences of temporal advance and retreat of the ice sheets. In Europe, ice sheets started to retreat from 16,000 BP and by 13,000 BP Britain had been colonised by some Mediterranean species (Atkinson *et al.* 1987) before the re-formation of the English Channel (Coles, 1998). In Europe, the formation of three refuge populations for many vertebrate and invertebrate species including host species of LB spirochetes (e.g. voles, hedgehogs, shrews) has been proposed: a western refuge on the Iberian peninsula, an eastern refuge in the Baltics, and a potential refuge on the Italian peninsula. It was also suggested that the migration of these species may have led to a pattern of hybrid zones in the region of France, Germany or Czech Republic (Hewitt, 1999). It would be interesting to investigate additional samples from the Iberian Peninsula and Baltic states to confirm this hypothesis.

Methicillin-resistant S. aureus (MRSA) clones

S. aureus colonises the skin and is also found in the anterior nares in around 30% of healthy people (Wertheim *et al.* 2005). However, *S. aureus* can cause infections that vary from minor skin infections to pneumonia all over the world (Grundmann *et al.* 2006). From an epidemiological perspective, there has been great interest in monitoring the geographical

distribution and the spread of *S. aureus* in general and MRSA in particular. For this purpose, several typing methods have been developed and used, including MLST, genomic DNA fingerprinting by pulse-field electrophoresis, sequence analysis of the 'X-region' in the staphylococcal protein A that has a highly polymorphic succession of short and variable tandem repeats (spa typing), multilocus variable tandem repeat analysis (VNTR) and, most recently, genome-wide SNP data (Nubel *et al.* 2008, 2011). Since the turn of the century, the most widely used typing method for *S. aureus* was MLST. Using this approach, the *S. aureus* global population structure has been classified into some major and many minor clonal complexes (Feil *et al.* 2003). Many of these clonal complexes consist of particular type of strains, the so-called MRSA, which are resistant to many classes of antibiotics. Being a very common cause of hospital-acquired infections, MRSA is a considerable burden to hospitals all over the world. The methicillin-resistant phenotype is encoded by the *mecA* gene that is located on the staphylococcal cassette chromosome (SCCmec), which can integrate into the genome of *S. aureus* at a particular site near the origin of replication (Kuroda *et al.* 2001). It has been shown that MRSA strains do not form a monophyletic group suggesting that this phenotype has evolved many times by acquisition of SCCmec elements by diverse lineages of *S. aureus* (Robinson and Enright, 2003). In what follows we will consider two very recently emerged MRSA clones, namely ST239 and ST225. These two MLST sequence types have been shown to be restricted to hospital settings and for these STs genome-wide SNP data sets, generated either by a second-generation sequencing platform (Harris *et al.* 2010) or a dHPLC mutation discovery approach (Nubel *et al.* 2010), have been used to study their evolutionary history considering very confined spatial and very recent temporal scales.

The pandemic ST239 shows resistance against multiple antibiotics and has been found in hospitals in Asia, central and eastern Europe, South America and Australia. In a recent study, Harris and colleagues have unveiled the global geographic structure and the intercontinental spread of this clone over four decades (Harris *et al.* 2010). The authors first mapped reads for 62 globally collected isolates against a reference genome and identified 6714 high-quality SNPs. After removing all SNPs found in the non-core genome they generated a maximum likelihood phylogeny based on 4310 SNPs mapping to the core genome. The rationale behind this strategy was to disregard the genomic regions that might have been affected by recombination. The authors found that the phylogenetic clustering presented a remarkable consistency with the geographic origin of the samples (Fig. 2). Three groups were observed corresponding to three geographic areas, i.e., Asia, Europe, and South America (blue, red and green




Fig. 2. Maximum likelihood phylogeny using core genome SNPs to analyze the intercontinental transmission of MRSA ST239 isolates. The label colours reflect the continental origin for each isolate: Asia, blue; Australia, yellow; Europe, red; North America, black; South America, green. Bootstrap values are shown below the branches, stars represent a bootstrap value of 100%. The scale bar represents the number of substitutions per SNP. The inset shows the subset from Thailand, here the bootstrap values are above the branches and, below the branch, the number of distinguishing SNPs for each isolate. Original Fig. from Harris *et al.* 2010, *Science* 327 (5964). Reprinted with permission from AAAS.

name labels, respectively, in Fig. 2). While the South American and the Asian groups were monophyletic the European clade was not. They also noted some exceptions to this geographical pattern and these clearly allude to intercontinental transmissions. For example, Portuguese strains and even a strain from the Czech Republic were found within the South American clade. Remarkably, the study was able to resolve fully the phylogenetic relationship of a subset of 20 isolates, all of which were obtained from patients from a single hospital in northeast Thailand over a period of 7 months (see the higher resolution panel in Fig. 2). This clearly demonstrates the resolving power of SNP interrogation as the authors were able to resolve very confined geographical (one hospital) and temporal scales. Additionally, having the date of isolation of each strain, the authors were not only able to estimate the mutation rate but also to date the origin of this clone. It has been dated between the mid and late 1960s, which, interestingly, is few years after the introduction of methicillin in 1960.

Nubel and colleagues, in a slightly more recent study analysed the MLST ST225, which is a highly prevalent ST in Central European Hospitals (Nubel *et al.* 2010). The study considered 269 loci using

an international collection of 73 isolates (including samples from Europe and the USA). The observation of very low diversity in European isolates suggested that a recent population bottleneck preceded the expansion of this ST in Europe. The authors suggested that isolates collected in the USA comprised an ancestral population, being more divergent than the European population. As the sampling dates were known, the authors were able to estimate the emergence of this ST. Using Bayesian approaches, they established that ST225 had diverged since ~1990 and that the expansion of the European clade began in ~1995. Through a demographic analysis, the authors inferred a drastic increase of population size in Europe from 2001 to 2004 and this is consistent with the known prevalence of this ST in European countries. Furthermore, a pattern of frequent transmission of ST225 between hospitals within central Europe was suggested by a spatio-temporal dynamics analysis. This study exemplifies how the recent demographic history of a particular ST can be studied in great detail by means of genome-wide SNPs in combination with biologically appropriate statistical techniques.

Notably, a similar phylogeographic pattern was identified in both studies. Both STs, 239 and 225,

appear to be relatively low-dispersal lineages where very closely related isolates are geographically proximate, forming tight groups, while basal long branches distinguish these groups. This makes sense when the ecology of these STs is taken into account. Importantly, MRSA mainly colonize humans in hospital settings and these particular settings can be regarded as 'islands' on which these lineages thrive. These STs seem to be dispersing among hospitals within the same geographical regions. Nonetheless, the intercontinental transmission events detected in the first study are clear exceptions to this pattern. Only by utilizing genome-wide SNP data sets were these studies able to explore the short-term evolution of these STs at an ultra-fine scale. Actually, the common motivation of both studies was the lack of power of MLST to grasp microevolutionary processes within individual clonal complexes. It is important to emphasize that these studies explored either 40 or 20 years in time and, remarkably, in the first study no two isolates had the same haplotype. Hence, these data provide such high resolution that each individual strain can be distinguished from other strains. The unprecedented resolving power of this genome-wide SNP approaches is exemplified in the study by Harris and colleagues (2010), which has estimated that a mutation event occurs every six weeks. From an epidemiological point of view, this total discrimination between isolates is quite remarkable and offers the possibility of tracking the spread of individual clones over very short periods of time.

COMPARING AND CONTRASTING THE SYSTEMS REVIEWED

In this review we intended to give a comprehensive overview of the phylogeography for the LB-group spirochetes. The idea of including the discussion about two very recently evolved MRSA clones was not to give a full account of the MRSA clones but to exemplify how the use of genome-wide SNPs can be of tremendous use in uncovering very recent phylogeographic patterns for pathogenic bacteria with very restrictive niches. Nonetheless, there are similarities between the systems discussed here, LB-group spirochetes and MRSA, that we wish to highlight. First, the phylogeography of both bacteria is dependent on host species and, in the case of LB-group spirochetes, on the vector as well. While the LB-group spirochetes can infect either their hosts or their tick vectors, the two MRSA clones have been found only in humans and specifically only in hospitalized patients. This has major implications - for example, recurrent population bottlenecks may affect these two groups of bacteria whenever they are transmitted from one host to another. Due to this consistent reduction in population size, these bacteria would be predicted to show considerable levels of genetic drift.

Indeed, a recent study reanalyzing ST239 data has shown increased levels of drift on the core genomes of this clone (Castillo-Ramirez *et al.* 2011). As a consequence of the smaller populations of these bacteria, one would expect that more slightly deleterious mutations would go to fixation by drift and thus these bacteria might have faster substitution rates than their close relatives with larger population sizes. It remains to be seen whether LB-group spirochetes also show high levels of genetic drift and corroborate the idea that these bacteria will have faster substitution rates than free-living relatives, which should have larger population sizes. Because the complete transmission cycle of LB-group spirochetes involves two passages (from tick to host and from host to tick), the levels of genetic drift could be more pronounced for this group.

Another issue of host dependence is that the mobility of hosts will inevitably influence the spread of these bacteria and the bacterial populations will be considerably affected by the demographic process shaping the populations of their host. A major difference between these systems is that MRSA is directly transmitted from human to human (or via contaminated equipment) while LB-group spirochetes represent a zoonotic pathogen with a highly complex life cycle. This difference is clearly reflected in their distribution range. Although both pathogens may utilise highly mobile hosts (humans and birds, respectively), the spread of LB-group spirochetes was likely limited by availability and abundance of vector species and populations and the environmental factors that act upon them underpinning the complexity of such systems (Kurtenbach *et al.* 2006; Margos *et al.* 2011).

Finally, in spite of being initially described as rather clonal species, strains from both of these groups undergo recombination. Actually, very recent studies using whole genome sequences have found considerable amounts of recombination in *S. aureus* strains (Chan *et al.* 2009; Takuno *et al.* 2012). For instance, Chan and colleagues noted that ~ 27% of the genes within the *Staphylococcus* genus have been affected by recombination and also found elevated rates of recombination, particularly in *S. aureus* genomes (Chan *et al.* 2011). Interestingly, we found compelling evidence that recombinations have also affected *B. burgdorferi* sensu stricto, one of the emblematic species of the LB-group (Margos *et al.* unpublished). However, more systematic studies are required to assay the impact of recombination on the various species of the LB-group spirochetes quantitatively. This implies that caution should be exercised when dealing with even supposedly clonal species and the presence of recombination should be tested in the loci under consideration before inferring phylogenetic relationships. On the other hand, moderate levels of recombination have the advantage of breaking the linkage between genes and in this

regard different realisations of the history of the species would be available as the genes may be unlinked. Statistical methods are now at hand for this to be taken into consideration when inferring species trees (Brito and Edwards, 2009).

OUTLOOK

In the foreseeable future, approaches using genome-wide SNP data will be very useful for conducting more informed surveillance over very short periods of time (weeks), at local and global scales, for a variety of bacterial pathogens. Furthermore, these genome-wide SNP data sets could not only be used to study the phylogeography of these microorganisms but also the evolutionary forces that have been shaping the genetic variation in these clones. For example, by reanalyzing the data set generated by Harris and colleagues (2010), it was possible to determine the effect of recombination on recently emerged clones and to show that the non-core regions of ST239 had been enriched in synonymous changes through recombination (Castillo-Ramirez *et al.* 2011).

Additionally, SNP data are likely to provide a sufficient signal to investigate deeper evolutionary relationships such as speciation events and to infer population genetic parameters such as mutation rates or effective population sizes. Furthermore, these data may also allow disentangling locus-specific effects (mutation, recombination) from genome-wide effects owing to demographic processes and, in this regard, both demographic processes and single loci could be studied in great detail. The ever decreasing costs of genome sequencing suggest that eventually such studies will be carried out routinely for many bacterial pathogens and this would allow a much better understanding not only about the spread and population structure of any given bacterium but also about the evolutionary forces that impact such bacteria.

However, for many of bacteria *ad hoc* MLST/MLSA have proved to be very revealing in terms of population structure and evolutionary relationship. Major advantages of these approaches are the employment of targeted PCR, which confers high specificity and permits the analysis of field collected samples without the need to culture the species/strains of interest (particularly appealing to species that are difficult to culture). In addition, the accumulative nature of database, their accessibility, the link to Google Maps (permitting generation of global distribution maps for STs) and the potential to use EpiCollect (Aanensen *et al.* 2009) makes this a convenient system for epidemiologists, ecologists and/or clinicians (Aanensen and Spratt, 2005). Another virtue of MLSA is its usefulness for taxonomic purposes (Gevers *et al.* 2005; Bishop *et al.* 2009).

The enormous speed of scientific advance using sequencing methods in the last few years and

additional technological and methodological developments that are likely to be made in the near future will most certainly accelerate progress in this exciting field of research.

ACKNOWLEDGEMENTS

The authors acknowledge Klaus Kurtenbach (who passed away in March 2009) for his many contributions to Lyme borreliosis research, his enthusiasm and the initiation of phylogeographic studies on *Borrelia* and S.A. Vollmer for sharing unpublished data. SCR is funded by the TROCAR consortium (EU FP7-HEALTH #223031).

REFERENCES

- Aanensen, D. M., Huntley, D. M., Feil, E. J., Al-Own, F. and Spratt, B. G. (2009). EpiCollect: linking smartphones to web applications for epidemiology, ecology and community data collection. *PLoS ONE* **4**, e6968.
- Aanensen, D. M. and Spratt, B. G. (2005). The multilocus sequence typing network: mlst.net. *Nucleic Acids Research* **33**, W728–733.
- Achtman, M. and Wagner, M. (2008). Microbial diversity and the genetic nature of microbial species. *Nature Reviews Microbiology* **6**, 431–440.
- Achtman, M., Zurth, K., Morelli, G., Torrea, G., Guiyoule, A. and Carniel, E. (1999). *Yersinia pestis*, the cause of plague, is a recently emerged clone of *Yersinia pseudotuberculosis*. *Proceedings of the National Academy of Sciences, USA* **96**, 14043–14048.
- Atkinson, T. C., Briffa, K. R. and Coope, G. R. (1987). Seasonal temperatures in Britain during the past 22,000 years reconstructed using beetle remains. *Nature* **325**, 587–592.
- Avise, J. C., Arnold, J., Ball, R. M., Bermingham, E., Lamb, T., Neigel, J. E., Reeb, C. A. and Saunders, N. C. (1987). Intraspecific phylogeography: The mitochondrial DNA bridge between population genetics and systematics. *Annual Reviews of Ecological Systems* **18**, 489–522.
- Barbour, A. G., Bunikis, J., Travinsky, B., Hoen, A. G., Diuk-Wasser, M. A., Fish, D. and Tsao, J. I. (2010). Niche partitioning of *Borrelia burgdorferi* and *Borrelia miyamotoi* in the same tick vector and mammalian reservoir species. *American Journal of Tropical Medicine and Hygiene* **81**, 1120–1131.
- Barbour, A. G. and Fish, D. (1993). The biological and social phenomenon of Lyme disease. *Science* **260**, 1610–1616.
- Bishop, C. J., Aanensen, D. M., Jordan, G. E., Kilian, M., Hanage, W. P. and Spratt, B. G. (2009). Assigning strains to bacterial species via the internet. *BMC Biology* **7**, 3.
- Bormane, A., Lucenko, I., Duks, A., Mavtchoutko, V., Ranka, R., Salmina, K. and Baumanis, V. (2004). Vectors of tick-borne diseases and epidemiological situation in Latvia in 1993–2002. *International Journal of Medical Microbiology* **293** Suppl 37, 36–47.
- Brito, P. H. and Edwards, S. V. (2009). Multilocus phylogeography and phylogenetics using sequence-based markers. *Genetica* **135**, 439–455.
- Brown, R. N. and Lane, R. S. (1992). Lyme disease in California: a novel enzootic transmission cycle of *Borrelia burgdorferi*. *Science* **256**, 1439–1442.
- Brown, R. N., Peot, M. A. and Lane, R. S. (2006). Sylvatic maintenance of *Borrelia burgdorferi* (Spirochaetales) in Northern California: untangling the web of transmission. *Journal of Medical Entomology* **43**, 743–751.
- Brownstein, J. S., Holford, T. R. and Fish, D. (2003). A climate-based model predicts the spatial distribution of the Lyme disease vector *Ixodes scapularis* in the United States. *Environmental Health Perspectives* **111**, 1152–1157.
- Burgdorfer, W., Barbour, A. G., Hayes, S. F., Benach, J. L., Grunwaldt, E. and Davis, J. P. (1982). Lyme disease—a tick-borne spirochetosis? *Science* **216**, 1317–1319.
- Castillo-Ramirez, S., Harris, S. R., Holden, M. T., He, M., Parkhill, J., Bentley, S. D. and Feil, E. J. (2011). The impact of recombination on dN/dS within recently emerged bacterial clones. *PLoS Pathogens* **7**, e1002129.
- Chan, C. X., Beiko, R. G., Darling, A. E. and Ragan, M. A. (2009). Lateral transfer of genes and gene fragments in prokaryotes. *Genome Biology and Evolution* **1**, 429–438.
- Chan, C. X., Beiko, R. G. and Ragan, M. A. (2011). Lateral transfer of genes and gene fragments in *Staphylococcus* extends beyond mobile elements. *Journal of Bacteriology* **193**, 3964–3977.
- Chan, E. Y. (2005). Advances in sequencing technology. *Mutation Research* **573**, 13–40.

- Christensen, E. M.** (1959). A historical view of the ranges of the white-tailed deer in northern Wisconsin forests. *American Midland Naturalist* **61**, 230–238.
- Chu, C. Y., Liu, W., Jiang, B. G., Wang, D. M., Jiang, W. J., Zhao, Q. M., Zhang, P. H., Wang, Z. X., Tang, G. P., Yang, H. and Cao, W. C.** (2008). Novel genospecies of *Borrelia burgdorferi* sensu lato from rodents and ticks in Southwestern China. *Journal of Clinical Microbiology* **46**, 3130–3133.
- Cohan, F. M.** (2002). What are bacterial species? *Annual Reviews in Microbiology* **56**, 457–487.
- Coles, B. J.** (1998). Doggerland: a speculative survey *Proceedings of the Prehistoric Society* **64** 45–81.
- Corander, J., Connor, T. R., O'dwyer, C. A., Kroll, J. S. and Hanage, W. P.** (2011). Population structure in the *Neisseria*, and the biological significance of fuzzy species. *Journal of the Royal Society Interface* **71**, 1208–1215. doi: 10.1098/rsif.2011.0601.
- Cronon, W.** (1983). *Changes in the Land: Indians, Colonists, and the Ecology of New England*, 1st edition. edn. Hill and Wang, New York.
- Crowder, C. D., Matthews, H. E., Schutzer, S., Rounds, M. A., Luft, B. J., Nolte, O., Campbell, S. R., Phillipson, C. A., Li, F., Sampath, R., Ecker, D. J. and Eshoo, M. W.** (2010). Genotypic variation and mixtures of Lyme *Borrelia* in *Ixodes* ticks from North America and Europe. *PLoS ONE* **5**, e10650.
- De Michelis, S., Sewell, H. S., Collares-Pereira, M., Santos-Reis, M., Schouls, L. M., Benes, V., Holmes, E. C. and Kurtenbach, K.** (2000). Genetic diversity of *Borrelia burgdorferi* sensu lato in ticks from mainland Portugal. *Journal of Clinical Microbiology* **38**, 2128–2133.
- Diuk-Wasser, M. A., Gatewood, A. G., Cortinas, M. R., Yaremych-Hamer, S., Tsao, J., Kitron, U., Hickling, G., Brownstein, J. S., Walker, E., Piesman, J. and Fish, D.** (2006). Spatiotemporal patterns of host-seeking *Ixodes scapularis* nymphs (Acari: Ixodidae) in the United States. *Journal of Medical Entomology* **43**, 166–176.
- Diuk-Wasser, M. A., Vourc'h, G., Cislo, P., Hoen, A. G., Melton, F., Hamer, S. A., Rowland, M., Cortinas, R., Hickling, G. J., Tsao, J., Barbour, A. G., Kitron, U., Piesman, J. and Fish, D.** (2010). Field and climate-based model for predicting the density of host-seeking nymphal *Ixodes scapularis*, an important vector of tick-borne disease agents in the eastern United States. *Global Ecology and Biogeography* **19** 504–514.
- Dubská, L., Literák, I., Kocianová, E., Taragelova, V. and Sychra, O.** (2009). Differential role of passerine birds in distribution of *Borrelia* spirochetes, based on data from ticks collected from birds during the postbreeding migration period in Central Europe. *Applied and Environmental Microbiology* **75**, 596–602.
- Eisen, L. and Lane, R. S.** (2002). Vectors of *Borrelia burgdorferi* sensu lato. In *Lyme Borreliosis: Biology, Epidemiology and Control* (eds. Gray, J., Kahl, O., Lane, R. S. and Stanek, G.), pp. 91–115. CABI Publishing, Wallingford.
- Enright, M. C., Day, N. P., Davies, C. E., Peacock, S. J. and Spratt, B. G.** (2000). Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *Journal of Clinical Microbiology* **38**, 1008–1015.
- Excoffier, L. and Heckel, G.** (2006). Computer programs for population genetics data analysis: a survival guide. *Nature Reviews Genetics* **7**, 745–758.
- Falco, R. C., Daniels, T. J. and Fish, D.** (1995). Increase in abundance of immature *Ixodes scapularis* (Acari: Ixodidae) in an emergent Lyme disease endemic area. *Journal of Medical Entomology* **32**, 522–526.
- Feil, E. J., Cooper, J. E., Grundmann, H., Robinson, D. A., Enright, M. C., Berendt, T., Peacock, S. J., Smith, J. M., Murphy, M., Spratt, B. G., Moore, C. E. and Day, N. P.** (2003). How clonal is *Staphylococcus aureus*? *Journal of Bacteriology* **185**, 3307–3316.
- Fingerle, V., Rauser, S., Hammer, B., Kahl, O., Heimerl, C., Schulte-Spechtel, U., Gern, L. and Wilske, B.** (2002). Dynamics of dissemination and outer surface protein expression of different European *Borrelia burgdorferi* sensu lato strains in artificially infected *Ixodes ricinus* nymphs. *Journal of Clinical Microbiology* **40**, 1456–1463.
- Garrick, R. C., Caccone, A. and Sunnucks, P.** (2010). Inference of population history by coupling exploratory and model-driven phylogeographic analyses. *International Journal of Molecular Science* **11**, 1190–1227.
- Gern, L., Estrada-Pena, A., Frandsen, F., Gray, J. S., Jaenson, T. G., Jongejan, F., Kahl, O., Korenberg, E., Mehl, R. and Nuttall, P. A.** (1998). European reservoir hosts of *Borrelia burgdorferi* sensu lato. *Zentralblatt für Bakteriologie* **287**, 196–204.
- Gevers, D., Cohan, F. M., Lawrence, J. G., Spratt, B. G., Coenye, T., Feil, E. J., Stackebrandt, E., Van De Peer, Y., Vandamme, P., Thompson, F. L. and Swings, J.** (2005). Opinion: Re-evaluating prokaryotic species. *Nature Reviews Microbiology* **3**, 733–739.
- Gevers, D., Dawyndt, P., Vandamme, P., Willems, A., Vancanneyt, M., Swings, J. and De Vos, P.** (2006). Stepping stones towards a new prokaryotic taxonomy. *Philosophical Transactions of the Royal Society London B Biological Sciences* **361**, 1911–1916.
- Gomez-Diaz, E., Boulmier, T., Sertour, N., Cornet, M., Ferquel, E. and McCoy, K. D.** (2011). Genetic structure of marine *Borrelia garinii* and population admixture with the terrestrial cycle of Lyme borreliosis. *Environmental Microbiology* **9**, 2453–2467.
- Goris, J., Konstantinidis, K. T., Klappenbach, J. A., Coenye, T., Vandamme, P. and Tiedje, J. M.** (2007). DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *International Journal of Systematic and Evolutionary Microbiology* **57**, 81–91.
- Grundmann, H., Aires-De-Sousa, M., Boyce, J. and Tiemersma, E.** (2006). Emergence and resurgence of methicillin-resistant *Staphylococcus aureus* as a public-health threat. *Lancet* **368**, 874–885.
- Hall, N.** (2007). Advanced sequencing technologies and their wider impact in microbiology. *Journal of Experimental Biology* **210**, 1518–1525.
- Halls, L. K.** (1984). *White-tailed Deer: Ecology and Management*, Stackpole Books, Harrisburg, PA.
- Hamer, S. A., Tsao, J. I., Walker, E. D. and Hickling, G. J.** (2010). Invasion of the Lyme Disease vector *Ixodes scapularis*: Implications for *Borrelia burgdorferi* endemicity. *EcoHealth* DOI: 10.1007/s10393-010-0287-0.
- Hanage, W. P., Fraser, C. and Spratt, B. G.** (2005). Fuzzy species among recombinogenic bacteria. *BMC Biology* **3**, 6.
- Hanincova, K., Schafer, S. M., Etti, S., Sewell, H. S., Taragelova, V., Ziak, D., Labuda, M. and Kurtenbach, K.** (2003a). Association of *Borrelia afzelii* with rodents in Europe. *Parasitology* **126**, 11–20.
- Hanincova, K., Taragelova, V., Koci, J., Schafer, S. M., Hails, R., Ullmann, A. J., Piesman, J., Labuda, M. and Kurtenbach, K.** (2003b). Association of *Borrelia garinii* and *B. valaisiana* with songbirds in Slovakia. *Applied and Environmental Microbiology* **69**, 2825–2830.
- Harris, S. R., Feil, E. J., Holden, M. T. G., Quail, M. A., Nickerson, E. K., Chantraita, N., Gardete, S., Tavares, A., Day, N., Lindsay, J. A., Edgeworth, J. D., Lencastre, D. H., Parkhill, J., Peacock, S. J. and Bentley, S. D.** (2010). Evolution of MRSA During Hospital Transmission and Intercontinental Spread. *Science* **327**, 469–474.
- Hewitt, G. M.** (1999). Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society* **68**, 87–112.
- Hoen, A. G., Margos, G., Bent, S. J., Kurtenbach, K. and Fish, D.** (2009). Phylogeography of *Borrelia burgdorferi* in the eastern United States reflects multiple independent Lyme disease emergence events. *Proceedings of the National Academy of Sciences, USA* **106**, 15013–15018.
- Holt, K. E., Parkhill, J., Mazzoni, C. J., Roumagnac, P., Weill, F. X., Goodhead, I., Rance, R., Baker, S., Maskell, D. J., Wain, J., Dolecek, C., Achtman, M. and Dougan, G.** (2008). High-throughput sequencing provides insights into genome variation and evolution in *Salmonella Typhi*. *Nature Genetics* **40**, 987–993.
- Humair, P. and Gern, L.** (2000). The wild hidden face of Lyme borreliosis in Europe. *Microbes and Infection* **2**, 915–922.
- Humphrey, P. T., Caporale, D. A. and Brisson, D.** (2010). Uncoordinated Phylogeography of *Borrelia burgdorferi* and Its tick vector, *Ixodes scapularis*. *Evolution* **64**, 2653–2663.
- Jackson, J. O. and Defoliart, G. R.** (1970). *Ixodes scapularis* Say in northern Wisconsin. *Journal of Medical Entomology* **7**, 124–125.
- Keim, P., Van Ert, M. N., Pearson, T., Vogler, A. J., Huynh, L. Y. and Wagner, D. M.** (2004). Anthrax molecular epidemiology and forensics: using the appropriate marker for different evolutionary scales. *Infection, Genetics and Evolution* **4**, 205–213.
- Kidd, D. M. and Ritchie, M. G.** (2006). Phylogeographic information systems: putting the geography into phylogeography. *Journal of Biogeography* **33**, 1851–1865.
- Knowles, L. L. and Maddison, W. P.** (2002). Statistical phylogeography. *Molecular Ecology* **11**, 2623–2635.
- Koepfel, A., Perry, E. B., Sikorski, J., Krizanc, D., Warner, A., Ward, D. M., Rooney, A. P., Brambilla, E., Connor, N., Ratcliff, R. M., Nevo, E. and Cohan, F. M.** (2008). Identifying the fundamental units of bacterial diversity: a paradigm shift to incorporate ecology into bacterial systematics. *Proceedings of the National Academy of Sciences, USA* **105**, 2504–2509.
- Konstantinidis, K. T., Ramette, A. and Tiedje, J. M.** (2006). The bacterial species definition in the genomic era. *Philosophical Transactions of the Royal Society London B Biological Sciences* **361**, 1929–1940.
- Konstantinidis, K. T. and Tiedje, J. M.** (2005). Genomic insights that advance the species definition for prokaryotes. *Proceedings of the National Academy of Sciences, USA* **102**, 2567–2572.
- Korenberg, E. I., Gorelova, N. B. and Kovalevskii, Y. V.** (2002). Ecology of *Borrelia burgdorferi* sensu lato in Russia. In *Lyme*

- Borreliosis: Biology, Epidemiology and Control* (eds. Gray, J., Kahl, O., Lane, R. S. and Stanek, G.), CABI Publishing, Wallingford.
- Kotetishvili, M., Kreger, A., Wauters, G., Morris, J. G., Jr., Sulakvelidze, A. and Stine, O. C. (2005). Multilocus sequence typing for studying genetic relationships among *Yersinia* species. *Journal of Clinical Microbiology* **43**, 2674–2684.
- Kuroda, M., Ohta, T., Uchiyama, I., Baba, T., Yuzawa, H., Kobayashi, I., Cui, L., Oguchi, A., Aoki, K. and Nagai, Y. (2001). Whole genome sequencing of methicillin-resistant *Staphylococcus aureus*. *Lancet* **357**, 1225–1240.
- Kurtenbach, K., De Michelis, S., Etti, S., Schafer, S. M., Sewell, H. S., Brade, V. and Kraiczy, P. (2002). Host association of *Borrelia burgdorferi* sensu lato—the key role of host complement. *Trends in Microbiology* **10**, 74–79.
- Kurtenbach, K., Hanincova, K., Tsao, J. I., Margos, G., Fish, D. and Ogden, N. H. (2006). Fundamental processes in the evolutionary ecology of Lyme borreliosis. *Nature Reviews Microbiology* **4**, 660–669.
- Kurtenbach, K., Hoen, A. G., Bent, S. J., Vollmer, S. A., Ogden, N. H. and Margos, G. (2010). Population biology of Lyme borreliosis spirochetes. In *Bacterial Population Genetics in Infectious Disease* (eds. Robinson, D. A., Falush, D. and Feil, E. J.), John Wiley & Sons, Inc.
- Kurtenbach, K., Peacey, M., Rijpkema, S. G., Hoodless, A. N., Nuttall, P. A. and Randolph, S. E. (1998a). Differential transmission of the genospecies of *Borrelia burgdorferi* sensu lato by game birds and small rodents in England. *Applied and Environmental Microbiology* **64**, 1169–1174.
- Kurtenbach, K., Sewell, H. S., Ogden, N. H., Randolph, S. E. and Nuttall, P. A. (1998b). Serum complement sensitivity as a key factor in Lyme disease ecology. *Infection and Immunity* **66**, 1248–1251.
- Lan, R. and Reeves, P. R. (2001). When does a clone deserve a name? A perspective on bacterial species based on population genetics. *Trends in Microbiology* **9**, 419–424.
- Lin, T., Gao, L., Seyfang, A. and Oliver, J. H., Jr. (2005). ‘Candidatus *Borrelia texasensis*’, from the American dog tick *Dermacentor variabilis*. *International Journal of Systematic and Evolutionary Microbiology* **55**, 685–693.
- Lin, T., Oliver, J. H., Jr. and Gao, L. (2004). Molecular characterization of *Borrelia* isolates from ticks and mammals from the southern United States. *Journal of Parasitology* **90**, 1298–1307.
- Lin, T., Oliver, J. H., Jr., Gao, L., Kollars, T. M., Jr. and Clark, K. L. (2001). Genetic heterogeneity of *Borrelia burgdorferi* sensu lato in the southern United States based on restriction fragment length polymorphism and sequence analysis. *Journal of Clinical Microbiology* **39**, 2500–2507.
- Maggi, R. G., Reichelt, S., Toliver, M. and Engber, B. (2010). *Borrelia* species in *Ixodes affinis* and *Ixodes scapularis* ticks collected from the coastal plain of North Carolina. *Ticks and Tick Borne Diseases* **1**, 168–171.
- Maiden, M. C. (2006). Multilocus sequence typing of bacteria. *Annual Reviews of Microbiology* **60**, 561–588.
- Maiden, M. C., Bygraves, J. A., Feil, E., Morelli, G., Russell, J. E., Urwin, R., Zhang, Q., Zhou, J., Zurth, K., Caugant, D. A., Feavers, I. M., Achtman, M. and Spratt, B. G. (1998). Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proceedings of the National Academy of Sciences, USA* **95**, 3140–3145.
- Manel, S., Schwartz, M. K., Luikart, G. and Roberlet, P. (2003). Landscape genetics: combining landscape ecology and population genetics. *Trends in Ecology and Evolution* **18**, 189–197.
- Margos, G., Gatewood, A. G., Aanensen, D. M., Hanincova, K., Terekhova, D., Vollmer, S. A., Cornet, M., Piesman, J., Donaghy, M., Bormane, A., Hurn, M. A., Feil, E. J., Fish, D., Casjens, S., Wormser, G. P., Schwartz, I. and Kurtenbach, K. (2008). MLST of housekeeping genes captures geographic population structure and suggests a European origin of *Borrelia burgdorferi*. *Proceedings of the National Academy of Sciences, USA* **105**, 8730–8735.
- Margos, G., Hojgaard, A., Lane, R. S., Cornet, M., Fingerle, V., Rudenko, N., Ogden, N., Aanensen, D. M., Fish, D. and Piesman, J. (2010). Multilocus sequence analysis of *Borrelia bisettii* strains from North America reveals a new *Borrelia* species, *Borrelia kurtenbachii*. *Ticks and Tick Borne Diseases* **1**, 151–158.
- Margos, G., Vollmer, S. A., Cornet, M., Garnier, M., Fingerle, V., Wilske, B., Bormane, A., Vitorino, L., Collares-Pereira, M., Drancourt, M. and Kurtenbach, K. (2009). A new *Borrelia* species defined by Multilocus Sequence Analysis of housekeeping genes. *Applied and Environmental Microbiology* **75**, 5410–5416.
- Margos, G., Vollmer, S. A., Ogden, N. H. and Fish, D. (2011). Population genetics, taxonomy, phylogeny and evolution of *Borrelia burgdorferi* sensu lato. *Infection, Genetics and Evolution* **11**, 1545–1563.
- Marshall, W. F., 3rd, Telford, S. R., 3rd, Rys, P. N., Rutledge, B. J., Mathiesen, D., Malawista, S. E., Spielman, A. and Persing, D. H. (1994). Detection of *Borrelia burgdorferi* DNA in museum specimens of *Peromyscus leucopus*. *Journal of Infectious Diseases* **170**, 1027–1032.
- Marti Ras, N., Postic, D., Foretz, M. and Baranton, G. (1997). *Borrelia burgdorferi* sensu stricto, a bacterial species “made in the U.S.A.”? *International Journal for Systematic Bacteriology* **47**, 1112–1117.
- Masuzawa, T. (2004). Terrestrial distribution of the Lyme borreliosis agent *Borrelia burgdorferi* sensu lato in East Asia. *Japanese Journal of Infectious Diseases* **57**, 229–235.
- Maupin, G. O., Gage, K. L., Piesman, J., Monteneri, J., Sviat, S. L., Vanderzanden, L., Happ, C. M., Dolan, M. and Johnson, B. J. (1994). Discovery of an enzootic cycle of *Borrelia burgdorferi* in *Neotoma mexicana* and *Ixodes spinipalpis* from northern Colorado, an area where Lyme disease is nonendemic. *Journal of Infectious Diseases* **170**, 636–643.
- Mccabe, T. R. and McCabe, R. E. (1997). Recounting whitetails past. In *The Science of Overabundance: Deer Ecology and Population Management* (eds. McShea, W. J., Underwood, H. B. and Rappole, J. H.), pp. 11–26. Smithsonian Institution Press, Washington DC.
- Mccooy, K. D., Bouludier, T., Tirard, C. and Michalakakis, Y. (2003). Host-dependent genetic structure of parasite populations: differential dispersal of seabird tick host races. *Evolution* **57**, 288–296.
- Mccooy, K. D., Chapuis, E., Tirard, C., Bouludier, T., Michalakakis, Y., Bohec, C. L., Maho, Y. L. and Gauthier-Clerc, M. (2005). Recurrent evolution of host-specialized races in a globally distributed parasite. *Proceedings of Biological Sciences* **272**, 2389–2395.
- Nakao, M., Miyamoto, K. and Fukunaga, M. (1994). Lyme disease spirochetes in Japan: enzootic transmission cycles in birds, rodents, and *Ixodes persulcatus* ticks. *Journal of Infectious Diseases* **170**, 878–882.
- Nubel, U., Dordel, J., Kurt, K., Strommenger, B., Westh, H., Shukla, S. K., Zemlickova, H., Leblois, R., Wirth, T., Jombart, T. and Al, E. (2010). A timescale for evolution, population expansion, and spatial spread of an emerging clone of methicillin-resistant *Staphylococcus aureus*. *PLoS Pathogens* **6**, e1000855.
- Nubel, U., Roumagnac, P., Feldkamp, M., Song, J. H., Ko, K. S., Huang, Y. C., Coombs, G., Ip, M., Westh, H., Skov, R., Struelens, M. J., Goering, R. V., Strommenger, B., Weller, A., Witte, W. and Achtman, M. (2008). Frequent emergence and limited geographic dispersal of methicillin-resistant *Staphylococcus aureus*. *Proceedings of the National Academy of Sciences, USA* **105**, 14130–14135.
- Nubel, U., Strommenger, B., Layer, F. and Witte, W. (2011). From types to trees: Reconstructing the spatial spread of *Staphylococcus aureus* based on DNA variation. *International Journal of Medical Microbiology* **301**, 614–618.
- Ogden, N. H., Artsob, H., Margos, G. and Tsao, J. (2011a). Non-rickettsial tick-borne bacteria and the diseases they cause. In *Biology of Ticks* (eds. Sonenshine, D. E. and Roe, M.), Oxford University Press, USA.
- Ogden, N. H., Bouchard, C., Lindsay, L. R., Margos, G., Kurtenbach, K., Trudel, L., Nguon, S. and Milord, F. (2011b). Investigation of genotypes of *Borrelia burgdorferi* in *Ixodes scapularis* ticks collected in surveillance in Canada. *Applied and Environmental Microbiology* **77**, 3244–3254.
- Ogden, N. H., Bouchard, C., Kurtenbach, K., Margos, G., Lindsay, L. R., Trudel, L., Nguon, S. and Milord, F. (2010). Active and passive surveillance and phylogenetic analysis of *Borrelia burgdorferi* elucidate the process of Lyme disease risk emergence in Canada. *Environmental Health Perspectives* **118**, 909–914.
- Oliver, J. H., Jr., Lin, T., Gao, L., Clark, K. L., Banks, C. W., Durden, L. A., James, A. M. and Chandler, F. W., Jr. (2003). An enzootic transmission cycle of Lyme borreliosis spirochetes in the south-eastern United States. *Proceedings of the National Academy of Sciences, USA* **100**, 11642–11645.
- Pearson, T., Okinaka, R. T., Foster, J. T. and Keim, P. (2009). Phylogenetic understanding of clonal populations in an era of whole genome sequencing. *Infection, Genetics and Evolution* **9**, 1010–1019.
- Picken, R. N., Cheng, Y., Han, D., Nelson, J. A., Reddy, A. G., Hayden, M. K., Picken, M. M., Strle, F., Bouseman, J. K. and Trenholme, G. M. (1995). Genotypic and phenotypic characterization of *Borrelia burgdorferi* isolated from ticks and small animals in Illinois. *Journal of Clinical Microbiology* **33**, 2304–2315.
- Picken, R. N., Cheng, Y., Strle, F. and Picken, M. M. (1996). Patient isolates of *Borrelia burgdorferi* sensu lato with genotypic and phenotypic similarities of strain 25015. *Journal of Infectious Diseases* **174**, 1112–1115.
- Picken, R. N. and Picken, M. M. (2000). Molecular characterization of *Borrelia* spp. isolates from greater metropolitan Chicago reveals the presence of *Borrelia bisettii*. Preliminary report. *Journal of Molecular Microbiology and Biotechnology* **2**, 505–507.

- Picken, R. N., Strle, F., Ruzic-Sabljić, E., Maraspin, V., Lotric-Furlan, S., Cimperman, J., Cheng, Y. and Picken, M. M. (1997). Molecular subtyping of *Borrelia burgdorferi* sensu lato isolates from five patients with solitary lymphocytoma. *Journal of Investigative Dermatology* **108**, 92–97.
- Postic, D., Garnier, M. and Baranton, G. (2007). Multilocus sequence analysis of atypical *Borrelia burgdorferi* sensu lato isolates - description of *Borrelia californiensis* sp. nov., and genomospecies 1 and 2. *International Journal of Medical Microbiology* **297**, 263–271.
- Qiu, W. G., Bruno, J. F., McCaig, W. D., Xu, Y., Livey, I., Schriefer, M. E. and Luft, B. J. (2008). Wide distribution of a high-virulence *Borrelia burgdorferi* clone in Europe and North America. *Emerging Infectious Diseases* **14**, 1097–1104.
- Randolph, S. E. (2000). Ticks and tick-borne disease systems in space and from space. *Advances in Parasitology* **47**, 217–243.
- Randolph, S. E. (2004). Tick ecology: processes and patterns behind the epidemiological risk posed by ixodid ticks as vectors. *Parasitology* **129** Suppl, S37–S65.
- Reisen, W. K. (2010). Landscape epidemiology of vector-borne diseases. *Annual Reviews in Entomology* **55**, 461–483.
- Richter, D., Postic, D., Sertour, N., Livey, I., Matuschka, F. R. and Baranton, G. (2006). Delineation of *Borrelia burgdorferi* sensu lato species by multilocus sequence analysis and confirmation of the delineation of *Borrelia spielmanii* sp. nov. *International Journal of Systematic and Evolutionary Microbiology* **56**, 873–881.
- Robinson, D. A. and Enright, M. C. (2003). Evolutionary models of the emergence of methicillin-resistant *Staphylococcus aureus*. *Antimicrobial Agents and Chemotherapy* **47**, 3926–3934.
- Rossello-Mora, R. and Amann, R. (2001). The species concept for prokaryotes. *FEMS Microbiology Reviews* **25**, 39–67.
- Rudenko, N., Golovchenko, M., Grubhoffer, L. and Oliver, J. H., Jr. (2009a). *Borrelia carolinensis* sp. nov. – a new (14th) member of *Borrelia burgdorferi* sensu lato complex from the southeastern United States. *Journal of Clinical Microbiology* **47**, 134–141.
- Rudenko, N., Golovchenko, M., Lin, T., Gao, L., Grubhoffer, L. and Oliver, J. H., Jr. (2009b). Delineation of a new species of the *Borrelia burgdorferi* sensu lato complex, *Borrelia americana* sp. nov. *Journal of Clinical Microbiology* **47**, 3875–3880.
- Scott, J. D., Lee, M. K., Fernando, K., Durden, L. A., Jorgensen, D. R., Mak, S. and Morshed, M. G. (2010). Detection of Lyme disease spirochete, *Borrelia burgdorferi* sensu lato, including three novel genotypes in ticks (Acari: Ixodidae) collected from songbirds (Passeriformes) across Canada. *Journal of Vector Ecology* **35**, 124–139.
- Scrimanti, R. J. (1970). Erythema chronicum migrans. *Archives of Dermatology* **102**, 104–105.
- Skuballa, J., Oehme, R., Hartelt, K., Petney, T., Bucher, T., Kimmig, P. and Taraschewski, H. (2007). European Hedgehogs as Hosts for *Borrelia* spp., Germany. *Emerging Infectious Diseases* **13**, 952–953.
- Smith, R. P., Jr., Muzaffar, S. B., Lavers, J., Lacombe, E. H., Cahill, B. K., Lubelczyk, C. B., Kinsler, A., Mathers, A. J. and Rand, P. W. (2006). *Borrelia garinii* in seabird ticks (*Ixodes uriae*), Atlantic Coast, North America. *Emerging Infectious Diseases* **12**, 1909–1912.
- Spielman, A. (1994). The emergence of Lyme disease and human babesiosis in a changing environment. *Annals of the New York Academy of Science* **740**, 146–156.
- Spratt, B. G. (1999). Multilocus sequence typing: molecular typing of bacterial pathogens in an era of rapid DNA sequencing and the Internet. *Current Opinion in Microbiology* **2**, 312–316.
- Steere, A. C., Bartenhagen, N. H., Craft, J. E., Hutchinson, G. J., Newman, J. H., Pachner, A. R., Rahn, D. W., Sigal, L. H., Taylor, E. and Malawista, S. E. (1986). Clinical manifestations of Lyme disease. *Zentralblatt für Bakteriologie, Mikrobiologie und Hygiene [A]* **263**, 201–205.
- Storfer, A., Murphy, M. A., Evans, J. S., Goldberg, C. S., Robinson, S., Spear, S. F., Dezzani, R., Delmelle, E., Vierling, L. and Waits, L. P. (2007). Putting the “landscape” in landscape genetics. *Heredity* **98**, 128–142.
- Takano, A., Nakao, M., Masuzawa, T., Takada, N., Yano, Y., Ishiguro, F., Fujita, H., Ito, T., Ma, X., Oikawa, Y., Kawamori, F., Kumagai, K., Mikami, T., Hanaoka, N., Ando, S., Honda, N., Taylor, K., Tsubota, T., Konnai, S., Watanabe, H., Ohnishi, M. and Kawabata, H. (2011). Multilocus Sequence Typing Implicates Rodents as the Main Reservoir Host of Human-Pathogenic *Borrelia garinii* in Japan. *Journal of Clinical Microbiology* **49**, 2035–2039.
- Takuno, S., Kado, T., Sugino, R. P., Nakhleh, L. and Innan, H. (2012). Population Genomics in Bacteria: A Case Study of *Staphylococcus aureus*. *Molecular Biology and Evolution* **29**, 797–809.
- Taragel'ova, V., Koci, J., Hanincova, K., Kurtenbach, K., Derdakova, M., Ogden, N. H., Literak, I., Kocianova, E. and Labuda, M. (2008). Blackbirds and song thrushes constitute a key reservoir of *Borrelia garinii*, the causative agent of borreliosis in Central Europe. *Applied and Environmental Microbiology* **74**, 1289–1293.
- Tsao, J. I. (2009). Reviewing molecular adaptations of Lyme borreliosis spirochetes in the context of reproductive fitness in natural transmission cycles. *Veterinary Research* **40**, 36.
- Van Ert, M. N., Easterday, W. R., Simonson, T. S., U'ren, J. M., Pearson, T., Kenefic, L. J., Busch, J. D., Huynh, L. Y., Dukerich, M., Trim, C. B., Beaudry, J., Welty-Bernard, A., Read, T., Fraser, C. M., Ravel, J. and Keim, P. (2007). Strain-specific single-nucleotide polymorphism assays for the *Bacillus anthracis* Ames strain. *Journal of Clinical Microbiology* **45**, 47–53.
- Vitorino, L. R., Margos, G., Feil, E. J., Collares-Pereira, M., Ze-Ze, L. and Kurtenbach, K. (2008). Fine-scale phylogeographic structure of *Borrelia lusitaniae* revealed by Multilocus Sequence Typing. *PLoS ONE* **3**, e4002.
- Vollmer, S. A., Margos, G., Donaghy, M., Bormane, A., Drancourt, M., Garnier, M., Cornet, M. and Kurtenbach, K. (2011). Phylogeographic structuring and evolutionary relationships of Lyme Borreliosis spirochetes in Europe as revealed by MLSA. *Environmental Microbiology* **13**, 184–192.
- Waters, J. H. (1963). Biochemical Relationships of the Mouse *Peromyscus* in New England. *Systematic Zoology* **12**, 122–133.
- Wertheim, H. F., Melles, D. C., Vos, M. C., Van Leeuwen, W., Van Belkum, A., Verbrugh, H. A. and Nouwen, J. L. (2005). The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infectious Diseases* **5**, 751–762.
- Wilske, B., Busch, U., Eiffert, H., Fingerle, V., Pfister, H. W., Rossler, D. and Preac-Mursic, V. (1996). Diversity of OspA and OspC among cerebrospinal fluid isolates of *Borrelia burgdorferi* sensu lato from patients with neuroborreliosis in Germany. *Medical Microbiology and Immunology* **184**, 195–201.
- Wilson, M. L., Adler, G. H. and Spielman, A. (1985). Correlation between abundance of deer and that of the deer tick, *Ixodes dammini* (Acari, Ixodidae). *Annals of the Entomological Society of America* **78**, 172–176.