



Citation for published version:

Acerra, N, Kad, NM, Griffith, DA, Ott, S, Crowther, DC & Mason, JM 2014, 'Retro-inversal of intracellular selected -amyloid-interacting peptides: Implications for a novel Alzheimer's disease treatment', *Biochemistry*, vol. 53, no. 13, pp. 2101-2111. <https://doi.org/10.1021/bi5001257>

DOI:

[10.1021/bi5001257](https://doi.org/10.1021/bi5001257)

Publication date:

2014

Document Version

Peer reviewed version

[Link to publication](#)

This document is the Accepted Manuscript version of a Published Work that appeared in final form in *Biochemistry*, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <http://pubs.acs.org/doi/abs/10.1021/bi5001257>

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

Retro-inversal of Intracellular Selected A β Interacting Peptides: Implications for a Novel Alzheimer's Disease Treatment.

Journal:	<i>Biochemistry</i>
Manuscript ID:	bi-2014-001257.R1
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Acerra, Nicola; University Of Essex, Biological Sciences Kad, Neil; University Of Essex, Biological Sciences Griffith, Douglas; University of Cambridge, Department of Genetic Ott, Stanislav; University of Cambridge, Department of Genetic Crowther, Damian; University of Cambridge, Department of Genetic Mason, Jody; University Of Essex, Biological Sciences

SCHOLARONE™
Manuscripts

1
2
3 **Retro-inversal of Intracellular Selected A β Interacting Peptides:**
4
5
6 **Implications for a Novel Alzheimer's Disease Treatment.**
7
8

9 Nicola Acerra¹, Neil. M. Kad¹, Douglas A. Griffith², Stanislav Ott², Damian C. Crowther² and
10
11 Jody M. Mason¹
12

13
14
15 ¹*Department of Biological Sciences, University of Essex, Wivenhoe Park, Colchester,*
16
17 *CO4 3SQ, UK Tel +44 1206 873010; Fax +44 1206 872592*
18

19 ²*Department of Genetics, University of Cambridge, Downing Site, Cambridge, CB2 3EH*
20

21 ¹*To whom correspondence should be addressed: jmason@essex.ac.uk*
22
23
24
25
26

27 Running title: Retro-inversed intracellular selected A β binders
28
29
30

31 Abbreviations: A β ₁₋₄₂, β -amyloid 1-42 variant; CD, circular dichroism; PPI, Protein-protein
32 Interaction; PCA, Protein-fragment Complementation Assay; MTT, (3-(4,5-Dimethylthiazol-2-
33 yl)-2,5-diphenyltetrazolium bromide; ThT, Thioflavin-T; HFIP, hexafluoroisopropanol; TFA,
34 trifluoroacetic acid; CPP, cell-penetrating peptide; TAT, trans-activating transcriptional
35 activator
36
37
38
39
40
41
42
43

44 Funding: JMM thanks Alzheimer's Research UK for a pilot project grant (ART/PPG2008B/2)
45 and AgeUK for a New Investigator Award (340). JMM is the recipient of a Cancer Research
46 UK Career Establishment Award (A11738) and a Wellcome Trust Project Grant
47 (WT090184MA). NA is funded by a University of Essex Departmental Studentship. JMM and
48 NMK also thank Parkinson's UK for awarding a PhD Studentship (H-1001). DCC is supported
49 by Wellcome Trust (082604/2/07/Z), Medical Research Council UK (G0700990) and an
50 Alzheimer's Research UK Senior Research Fellow (ART-SRF2010-2).
51
52
53
54
55
56
57
58
59
60

ABSTRACT:

The aggregation of β -amyloid ($A\beta$) into toxic oligomers is a hallmark of Alzheimer's disease pathology. Here we present a novel approach for the development of peptides capable of preventing amyloid aggregation based upon the previous selection of natural all-L peptides that bind $A\beta_{1-42}$. Using an intracellular selection system, successful library members were further screened via competition selection to identify the most effective peptides capable of reducing amyloid levels. In order to circumvent potential issues arising from stability and protease action for these structures we have replaced all L-residues with D-residues and inverted the sequence. These retro-inverso (RI) peptide analogues therefore encompass reversed sequences that maintain the overall topological order of the native peptides. Our results demonstrate that efficacy in blocking and reversing amyloid formation is maintained while introducing desirable properties to the peptides. Thioflavin-T assays, circular dichroism, and oblique angle fluorescence microscopy collectively indicate that RI-peptides can reduce amyloid load while MTT assays demonstrate modest reductions in cell toxicity. These conclusions are reinforced using *Drosophila melanogaster* studies to monitor pupal hatching rates and fly locomotor activity in the presence of RI-peptides delivered via RI-TAT peptide fusions. We demonstrate that the RI-PCA approach can be used as a generalised method for deriving $A\beta$ -interacting peptides. This approach has subsequently led to several peptide candidates to be further explored as potential treatments for Alzheimer's disease.

INTRODUCTION

A vast body of evidence implicates $A\beta$ as a major player in Alzheimer's Disease (AD) (1), with genetic studies revealing abnormal production in cell culture and animal models. $A\beta$ aggregation is implicated in neuronal death and impaired memory (2), with a wealth of evidence suggesting that amyloid load in AD sufferers does not correlate with disease severity (2, 3). The insoluble fibrils that are characteristic of the disease may serve as reservoirs that sequester a number of more toxic and soluble oligomeric species. Consequently, targeting $A\beta$ via therapeutic intervention has led to numerous inhibition strategies. Rare familial mutations that increase $A\beta$ oligomer concentration (e.g. the E22G arctic mutation (4)) have been shown to accelerate the onset of AD as a result (4). Many β -sheet breaker (BSB) molecules that block or breakdown amyloid fibres have had limited success and in certain cases have been counteractive, owing either to increased oligomer production or a failure to accelerate their removal (5-11). Therefore, since oligomers as small as dimers have been shown to exhibit cytotoxicity (12, 13) molecules effective at lowering amyloid levels may need to either sequester fibrils in the insoluble state, or preferably bind $A\beta$ as a small but ultimately non-toxic oligomer, possibly even the monomer (14).

We have previously used intracellular library screening and selection to identify $A\beta$ interacting peptides (15). In this approach, residues 25-35 of $A\beta$ are used as a design scaffold. This sequence, along with residues 15-20, is known to form amyloid in isolation (5, 6) and is thought to be responsible for instigating $A\beta$ self-association in the parental protein. Many amyloid inhibitors have therefore been based upon these regions (5-11) (see also (16) and references therein), with many strategies focused on simple modifications such that they retain the ability to bind $A\beta$ but prevent amyloid formation by introducing blocking or charged groups to these short sequences. For example, Tjernberg *et al* demonstrated that $A\beta_{16-20}$, despite forming fibrils itself, binds residues 25-35 of $A\beta$ and prevents fibril formation (5).

1
2
3 Soto and co-workers rationally designed proline-containing peptides based on $A\beta_{17-21}$ (10, 11).
4
5 It was subsequently shown that clearance of large amyloid fibrils can lead to the population of
6
7 smaller more cytotoxic intermediates (17). Therefore, the search for BSB peptides has been
8
9 hampered by the fact that successful molecules must be capable of preventing the population of
10
11 amyloid oligomers while avoiding the generation of cytotoxic species. In our current approach
12
13 we use an intracellular protein-fragment complementation assay (PCA) selection, which works
14
15 by recombining a split enzyme that is essential for cell survival (15, 18-20). We apply a
16
17 variation on this system whereby the $A\beta$ target is fused at the genetic level to one half of the
18
19 essential enzyme murine dihydrofolate reductase (mDHFR), with a peptide library fused to the
20
21 other half. Following protein expression in the cytoplasm, binding of a library member to $A\beta$
22
23 reconstitutes mDHFR, leading to bacterial cell growth and colony formation under selective
24
25 conditions. Since the entire process is intracellular, no assumptions are made regarding the
26
27 mechanism of antagonist action or which amyloid states become populated during selection.
28
29 The only prerequisite for success of any given library member is therefore that it must *i*) bind
30
31 to $A\beta$ and reconstitute mDHFR and *ii*) prevent $A\beta$ aggregation that has been shown to slow cell
32
33 growth, with *E.coli* rescued from the toxic effects of aggregation. In addition, the PCA
34
35 approach is predicted to select peptides that are resistant to degradation by bacterial proteases,
36
37 be soluble in solution, and target-specific in the presence of other cytoplasmic proteins. Using
38
39 PCA with an 8000 member library based on $A\beta_{29-35}$ led to an initial interacting sequence. A
40
41 second library of 160,000 members using this first hit as a design scaffold yielded two further
42
43 novel interacting sequences. These peptides consequently shared no homology to the original
44
45 $A\beta_{29-35}$ design template. All selected peptides were found to be capable of binding $A\beta$,
46
47 inhibiting amyloid formation and breaking down preformed fibrils.
48
49
50
51
52

53
54 A potential limitation of this previous study was the use of natural unmodified L-
55
56 peptides which are susceptible to degradation by mammalian proteases. This was partially
57
58
59
60

1
2
3 addressed by undertaking library selection inside the cytoplasm of *E.coli*. However, while
4
5 bacterial growth experiments indicated that toxicity was lowered, cytotoxicity experiments
6
7 using mammalian cells suggested that protease protection was not observed when transferring
8
9 to the PC12 cell environment (15). In a continuation of the above study we present data on
10
11 retro-inversed (RI) analogues of these previously selected peptides whereby we have
12
13 substituted the L-amino acids for their D counterparts and reversed the sequence. The D-amino
14
15 acid sequences give a mirror conformation, while the retro-peptides, consisting of the same
16
17 sequence of L-amino acids, leads to a reversal in order. Sequence retro-inversal therefore leads
18
19 to a mimic of the original peptide owing to inversion of the peptide bonds (21, 22). Using this
20
21 approach peptides have been shown to retain the same inhibitory aggregation qualities while
22
23 displaying vastly increased protease resistance (23). For example, rationally designed RI
24
25 peptide inhibitors of A β aggregation have previously been derived by appending arginine
26
27 residues to the A β ₁₆₋₂₀ (KLVFF) sequence (24, 25). These peptides were also shown to be
28
29 serum stable and with the addition of a RI cell penetrating peptide fusion were also able to
30
31 cross the blood brain barrier and display significant activity in transgenic mouse models (26).
32
33 In addition, although not an RI approach, mirror image phage display has been used to screen a
34
35 12-mer library against an immobilised D-enantiomer of the A β ₁₋₄₂ peptide, with the most
36
37 successful peptide binder replaced by D-amino acids to yield a protease-stable peptide capable
38
39 of binding to the natural L-form of A β ₁₋₄₂ (27). The success of these studies further highlights
40
41 the potential for peptide-based therapeutic strategies.
42
43
44
45
46

47 Here we report upon a novel approach that uses *intracellular* library screening and
48
49 selection followed by sequence retro-inversal (RI-PCA) as a generalised strategy for creating
50
51 stable peptides capable of antagonising protein-protein interactions (PPI). We have tested the
52
53 effectiveness of peptides using a range of *in vitro* techniques that report on the amount of fibril
54
55 present. These experiments have been coupled with PC12 neuronal cell-based assays that
56
57
58
59
60

1
2
3 report on the overall effect of toxicity in the presence of various peptides, as well as *in vivo*
4
5 studies using *drosophila melanogaster*, to verify peptide efficacy in the context of a whole cell
6
7 Alzheimer's disease model organism.
8
9

10 11 12 **MATERIALS AND METHODS**

13
14 ***PCA and expression vector cloning*** - PCA has been extensively used to derive PPI antagonists
15
16 of activator protein-1 (18, 19, 28, 29). More recently this has been extended to $A\beta$ interacting
17
18 peptides, where a full description of the methods can be found (15). Briefly, mDHFR was split
19
20 and one half fused to an $A\beta_{25-35}$ target peptide, and the other half to the library (15). Only target
21
22 binding library members bring two halves of mDHFR into close proximity, render it active,
23
24 and lead to colony formation on M9 selective plates. Trimethoprim is used to selectively
25
26 inhibit bacterial DHFR, thereby ensuring that colonies can only arise as the result of an
27
28 interaction between $A\beta$ and a peptide-library member. The $A\beta_{25-35}$ gene was synthesized using
29
30 overlap extension PCR and cloned into the pES300d-DHFR2 vector system using NheI and
31
32 AscI restriction sites.
33
34
35
36
37

38
39 ***PCA Library construction*** - Library construction and cloning has been described previously
40
41 (15, 18). Briefly, in the first library, positions 31-33 of $A\beta_{29-35}$ were completely randomized
42
43 using degenerate oligonucleotides containing NNK codons. NNK was used to encode all
44
45 twenty residues while removing two of three stop codons (30) to create an 8000 member
46
47 library. The second library was designed using the first PCA winner ('KAT') as a design
48
49 scaffold. In this case residues 29-30 and 34-35 of KAT were randomized, again using the
50
51 codon NNK, this time to generate a library of 160,000 members. Thus in starting with $A\beta_{29-35}$
52
53 as an initial design scaffold a completely unrelated sequence was subsequently derived.
54
55
56
57
58
59
60

1
2
3 Competition selection during PCA means that only the most effective 1–2 sequences are
4
5 isolated from the 50–100 A β binders that are initially identified during single-step selection.
6
7

8
9
10 **A β Peptide preparation** - A β_{1-42} was purchased as a pure recombinant peptide from rPeptide
11 (Stratech) and used for all experiments described. Prior to use, the peptide was treated to three
12 rounds of dissolution in hexafluoro-2-propanol (HFIP), sonication, drying, dissolution in
13 trifluoroacetic acid (TFA), followed by sonication and drying, according to the Zagorski
14 protocol (31). Peptide was then aliquoted into appropriately sized batches for subsequent
15 assays and dried via lyophilisation before being dissolved in 10mM potassium phosphate
16 buffer (pH 7.4) to generate a final concentration of 50 μ M. Rounds of TFA/HFIP treatment
17 were used to ensure that amyloid growth always proceeded from the same monomeric state,
18 thus reducing errors in amyloid growth and consequent assay measurements. For the second
19 library winners, two versions of each peptide was synthesised: (i) the parental sequence
20 including selected residues (L2P1a-RI / L2P2a-RI), and (ii) including additional amino acids
21 from restriction sites during cloning into the pES230d vector (L2P1b-RI / L2P2b-RI) (15). See
22 Table 1 for all peptide sequences.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40
41 **Peptide preparation** – KAT-RI, L2P1a-RI, L2P1b-RI, L2P2a-RI, L2P2b-RI and both a positive
42 control from the literature, iA β 5 (10), and a negative control peptide, TAT-dummy-RI, were
43 obtained by Peptide Protein Research (Fareham, UK) as pure lyophilised peptides. In addition,
44 for *drosophila melanogaster* experiments, TAT-KAT-RI, TAT-L2P1-RI and TAT-L2P2-RI
45 were also used. Peptides were weighed using an analytical balance and stock solutions of 1mM
46 concentration were subsequently dissolved in ultrapure water. Prior to assay peptides were
47 either aliquoted and lyophilised or diluted from stock as required.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Thioflavin T Assays** - ThT inhibition assays were performed using 50 μM Zagorski treated
4 (31) monomeric $A\beta_{1-42}$ in 100 μl of 10 mM potassium phosphate buffer, pH 7.4, with or
5 without each peptide at a concentration of 5 μM (for 1:0.1 molar ratio), 50 μM (for 1:1 molar
6 ratio), 100 μM (for 1:2 molar ratio) and 200 μM (for 1:4 molar ratio). In addition, for sub-
7 stoichiometric experiments 0.5 μM (1:0.01 molar ratio), 50 nM (1:0.001 molar ratio) and 5 nM
8 (1:0.0001 molar ratio) were also included to demonstrate progressively reduced activity as the
9 peptide dose is increasingly lowered, thus demonstrating a trend of dose dependency. To
10 achieve this, sufficient $A\beta_{1-42}$ was lyophilized, dissolved, and thoroughly vortexed as one
11 single batch (for immediate use in all $A\beta_{1-42}$ target-peptide mixes) to a concentration of 50 μM
12 in potassium phosphate buffer. Each inhibitor was lyophilized and redissolved in an Eppendorf
13 tube to a concentration of 5 μM (1:0.1), 50 μM (1:1), 100 μM (1:2), and 200 μM (1:4). Finally,
14 a 100 μL aliquot of the target solution was added to each inhibitor to give a total assay volume
15 of 100 μL containing 50 μM target and the appropriate inhibitor. The assay mixture was
16 vortexed and stored at 37°C for three days to induce aggregation in the presence of each
17 inhibitor. The ThT assay solution was prepared from a 25x stock containing 500 μM ThT. The
18 stock was aliquoted and kept frozen until required. It was then allowed to thaw at room
19 temperature for 10min before dilution into 10mM Tris buffer pH 7.4, giving the required
20 freshly prepared ThT assay solution containing 20 μM ThT in 10mM Tris and buffer at pH 7.4.
21 A total of 197.1 μL of the ThT assay solution was then added into 2.9 μL of each
22 inhibition/reversal assay mixture, thoroughly vortexed and transferred into an appropriate well
23 of the multiplate. The fluorescence of amyloid-bound ThT was measured by fluorescence
24 spectroscopy using a Cary Eclipse fluorescence spectrophotometer; bound ThT exhibits a new
25 excitation maxima at 450nm and an enhanced emission maxima at 482nm (32). For the
26 inhibition assays, the $A\beta_{1-42}$ target-peptide mixtures were incubated together on day zero at
27 37°C and single ThT readings were taken on day three. For the reversal assays, the target was
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 incubated alone at 37°C for three days before the addition of 100 µL to each lyophilized
4
5 inhibitor. The vortexed A β ₁₋₄₂ target-peptide solutions were then incubated at 37°C for a further
6
7 three days, at which single ThT readings were taken.
8
9

10
11 **Circular Dichroism (CD)** - Far-UV circular dichroism (CD) spectra were recorded on an
12
13 Applied Photophysics Chirascan CD spectrometer at 20°C. Peptide (10µM in 10mM Potassium
14
15 Phosphate buffer pH 7.4) was added to a 1mm CD cell (Hellma) and spectra recorded over the
16
17 200-300 nm range at a scan rate of 10nm/min with step size of 1nm. Spectra were recorded as
18
19 the average of two scans as raw ellipticity. Spectra for RI-peptides alone were subtracted from
20
21 A β ₁₋₄₂ target + peptide spectra, to leave normalised CD spectra accounting for the effect of the
22
23 peptide upon the A β ₁₋₄₂.
24
25
26
27

28
29 **3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium Bromide (MTT) Cell-Toxicity Assay -**

30
31 MTT experiments were undertaken using Rat phaeochromocytoma (PC12) cells to assess the
32
33 effect of the toxicity of A β ₁₋₄₂. PC12 cells are known to be particularly sensitive and their use
34
35 in this assay is well established (33). The MTT Vybrant® MTT Cell Proliferation Assay Kit
36
37 (Invitrogen) was used to measure the conversion of the water soluble MTT dye to formazan,
38
39 which is then solubilized, and the concentration determined by a colour change monitored via
40
41 absorbance measurement at 570 nm. The change in absorbance can then be converted to a
42
43 percentage MTT reduction which can be used as an indicator of the PC12 cell health in the
44
45 assay. The assay was performed with 10 µM A β ₁₋₄₂ and varying molar ratios of peptide
46
47 corresponding to 1:0.1 (0.1 µM), 1:1 (10 µM), 1:2 (20 µM), 1:4 (40 µM). PC12 cells were
48
49 maintained in RPMI 1640 +2mM glutamine medium mixed with 10% horse serum, 5% foetal
50
51 bovine serum, supplemented with a 20mg/mL gentamycin. Cells were transferred to a sterile
52
53 96-well plate with 30,000 cells per well and experiments performed in triplicate. Briefly,
54
55
56
57
58
59
60

1
2
3 different concentrations of peptides were screened in the presence of 10 μM $\text{A}\beta_{1-42}$. The
4
5 required volume from peptide and target stock solutions was freeze-dried overnight. The
6
7 freeze-dried peptide and $\text{A}\beta_{1-42}$ target were resuspended in 100% dimethyl sulfoxide (DMSO),
8
9 each at 100x stock concentration (i.e. 1 mM, 2 mM, 4 mM or 10 mM). For example for the
10
11 molar ratio 1:1 a total of 5 μL from each of the resuspended peptide/DMSO and target/DMSO
12
13 was mixed in a well of a 96-well preparation plate, thus giving 10 μL of 1:1 mM peptide/ $\text{A}\beta_{1-42}$
14
15 target concentration ratio in 100% DMSO. A total of 90 μL of RPMI media was added to the
16
17 10 μL peptide/ $\text{A}\beta_{1-42}$ target mixture (100:100 μM peptide/ $\text{A}\beta_{1-42}$ target ratio in 10% DMSO). A
18
19 total of 10 μL of the 50:50 μM peptide/ $\text{A}\beta_{1-42}$ target mixture in 10% DMSO was then dispensed
20
21 into 90 μL of media/PC12 cells, at final peptide and $\text{A}\beta_{1-42}$ target concentrations of 10
22
23 μM . These were incubated for 24 h at 37 $^{\circ}\text{C}$, 5% CO_2 , prior to the addition of the MTT dye. A
24
25 total of 10 μL of the dye was added to each well and incubated for a further 4h at 37 $^{\circ}\text{C}$, 5%
26
27 CO_2 . A total of 100 μL of the DMSO (stop/solubilisation solution) was then added to each well
28
29 and was allowed to stand for 10 minutes. The absorbance was measured at 570 nm using a 96-
30
31 well Versamax tuneable microplate reader.
32
33
34
35
36
37

38 ***Oblique Angle Fluorescence Microscopy Experiments*** - Samples were imaged on a custom
39
40 built oblique angle fluorescence system as described previously (OAF, (15, 34)). The
41
42 excitation and emission wavelengths were 488 nm and 500-605 nm respectively. Although
43
44 both the excitation and emission wavelengths were off peak for ThT, the image quality and
45
46 photobleaching characteristics were excellent. All samples were pre-stained with 10 μM ThT,
47
48 pipetted onto a clean glass slide, air dried and then imaged in KPP buffer supplemented with
49
50 100 mM DTT to further minimize photobleaching. For consistency and cross-correlation, the
51
52 same samples were used for inhibition/reversal imaging as those in ThT and CD experiments.
53
54 In addition, all samples provided for OAF imaging were supplied blind.
55
56
57
58
59
60

1
2
3
4
5 *Drosophila Melanogaster Assays* – The effect of RI peptides upon *drosophila melanogaster*
6
7 was assayed by fusing sequences to the (also retro-inversed) nine residue cell-penetrating
8 peptide (CPP), TAT (35-37). In these experiments, flies expressing $A\beta_{1-42}$ were fed nutrient
9 containing either RI-peptides or RI-TAT-peptide fusions (see Table 1) and two key effects
10 monitored; *i*) the speed of pupae hatching in which the cumulative hatching fractions for each
11 treatment was monitored relative to a negative control group *ii*) fly motility using an automated
12 fly tracking system to monitor the walking speed of the transgenic flies. In the control group,
13 flies typically become immobile as they age over the first 10-15 days of life.
14
15
16
17
18
19
20
21
22
23
24

25 RESULTS

26
27 We have previously undertaken PCA screening on libraries to derive peptides capable of
28 clearing $A\beta_{1-42}$ aggregates and generating significant enhancements in bacterial growth rates
29 (15). To produce peptides that retain the ability to bind $A\beta_{1-42}$ while bringing additional
30 properties to the molecule such as stability and protease resistance (23, 38) we have now retro-
31 inversed these sequences. The RI-PCA derived peptides (Table 1) have been synthesized and
32 characterised using a number of methods including ThT dye fluorescence to report on amyloid
33 formation, CD to measure global changes in β -sheet content associated with fibril formation,
34 and direct imaging using OAF microscopy. These techniques have demonstrated that RI-
35 peptides retain the ability of the PCA selected parental peptides to prevent aggregation and
36 remove preformed fibrils. In addition, MTT cytotoxicity assays undertaken using a neuronal
37 PC12 cell line were used to demonstrate that $A\beta_{1-42}$ toxicity is lowered when bacterially
38 selected peptides (which were not effective in MTT experiments (15)) are retro-inversed in
39 sequence. To explore cell-based experiments further, we have undertaken studies in which
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *drosophila melanogaster* expressing wild-type $A\beta_{1-42}$ were fed RI-peptides as well as RI-TAT-
4 peptide fusions to bring cell permeability to these $A\beta_{1-42}$ interacting compounds.
5
6
7

8
9
10 ***ThT binding indicates reduced fibril load*** - To determine the ability of PCA derived peptides
11 to either prevent fibril assembly (inhibition) and/or breakdown preformed fibrils (reversal),
12 ThT was used as an indicator of the degree to which $A\beta_{1-42}$ had aggregated into amyloid fibrils.
13 In this assay $A\beta_{1-42}$ was rendered monomeric (31) and redissolved at a concentration of 50 μ M
14 before being aggregated by incubating without agitation at 37°C. For the inhibition assay,
15 peptides were added on day zero, whereas for the reversal assay the peptides were added after
16 three days of $A\beta_{1-42}$ fibril growth. Once incubated together, $A\beta_{1-42}$ target-peptide solutions were
17 tested after three days. In addition to the five RI-PCA derived peptides, the positive control L-
18 peptide iA β 5 was also included as it is known to perform well in ThT assays and lead to a
19 reduction in fibril load (11). Finally a negative control peptide TAT-dummy-RI was included
20 to demonstrate no effect on ThT binding and therefore specificity for the PCA-derived
21 peptides. Figure 1 shows the results of these experiments at a number of $A\beta_{1-42}$:peptide ratios
22 for each peptide. Experiments were undertaken at stoichiometries of 1:0.0001, 1:0.001, 1:0.01,
23 1:0.1, 1:1, 1:2 and 1:4.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 Inhibition experiments demonstrate that peptides are able to prevent aggregation
41 (Figure 1a) with reductions in ThT bound relative to the $A\beta_{1-42}$ control of up to 80%. During
42 these experiments, a concentration dependence was observed as the $A\beta$:peptide ratio was
43 lowered (Figure 1). At increasingly higher stoichiometries this trend was less apparent. For
44 example, L2P1a-RI lowered bound ThT by ~30-40% at $A\beta$:peptide stoichiometries of 1:0.01 or
45 greater. Similarly, L2P2b-RI was also able to reduce bound ThT by >40%. RI-peptides
46 incubated in the absence of $A\beta$ did not bind ThT, displayed weakly helical CD spectra, and
47 were soluble to high concentrations in stock solutions. The lack of concentration dependence
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 for ThT assays at higher molar ratios is likely to be due to signal-to-noise difficulties, as is
4 shown by consistently large errors that are typical of this assay. In addition, ThT is known to
5 exhibit different fluorescence levels according to the oligomeric state populated (39). Given
6 that there are a number of potential intermediates in the growth of A β , different fluorescence
7 intensities could result. Reassuringly, the all L-residue positive control peptide, iA β 5 (11)
8 performed well, providing a reduction of 60-80%, that was lost in a dose dependent manner at
9 lower A β :iA β 5 ratios. In addition the negative control TAT-dummy-RI peptide had no effect
10 over a range of molar ratios, again demonstrating specificity for the PCA-derived sequences.
11 RI peptides performed favourably relative to their PCA-derived L-peptide counterparts.
12
13
14
15
16
17
18
19
20
21
22

23 In the case of reversal experiments (Figure 1b), as was observed previously for the L-
24 peptide counterparts, more pronounced reductions are observed relative to inhibition
25 experiments; over 60% for all peptides tested in all but six instances. At molar ratios of 1:0.1
26 reductions of >60% were observed for all peptides. Increasing the molar ratio did not lead to
27 improvements in reversal of amyloid in most cases, indicating that a ten-fold sub-
28 stoichiometric concentration of peptide may be sufficient to reverse fibril formation. As we
29 previously reported for parental L-peptides, a progressively reduced activity was observed as
30 the peptide dose tended towards zero, demonstrating a dose dependency (15). Experiments on
31 RI-peptides in isolation indicate that peptides do not aggregate into fibrils. They do not bind
32 ThT and do not generate CD spectra consistent with a β -sheet structure. Again, as predicted
33 from the literature, iA β 5 performed well at higher molar ratios, and this effect is lost at lower
34 ratios. These results compare favourably with PCA-derived L-peptide counterparts in being
35 able to reduce the ThT bound by ~60% (15). Again, the negative control TAT-dummy-RI
36 peptide also had no effect over a range of molar ratios. Errors associated with ThT
37 experiments preclude more detailed interpretations from being drawn.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **CD studies indicate reduced β -sheet content** – Owing to the fact that amyloid fibrils are
4 predominantly β -sheet, we have used CD as a measure of the global signal upon incubation
5 with peptides in inhibition and reversal assays. In these experiments the same aggregating
6 samples as used in ThT experiments have been measured to allow for direct comparison
7 between experiments. However, CD experiments in which the aggregating all L-residue $A\beta_{1-42}$
8 target is mixed with D-residue containing RI-peptides must be approached with caution; in
9 incubating the peptides together with $A\beta_{1-42}$ it was unknown if a signal at 218 nm caused by the
10 RI-peptides might obscure the signal at 218 nm arising from the $A\beta_{1-42}$ target. Therefore, the
11 CD signal of $A\beta_{1-42}$ and RI-peptides were both measured in isolation. The CD spectra arising
12 from these controls could then be accounted for (i.e. CD spectra of $A\beta_{1-42}$ against [$A\beta_{1-42}$ + RI-
13 peptide] – [RI-peptide alone]; Figure 2), allowing the overall loss or gain in β -sheet signal
14 exerted by the action of RI-peptide upon $A\beta$ aggregation to be established. In this assay we
15 observe impressive reductions in β -sheet content for all peptides at various molar ratios,
16 supporting the ThT data by demonstrating that RI-peptides reduce the global β -sheet content of
17 the sample and therefore the amyloid content. In agreement with the ThT data the negative
18 control peptide TAT-dummy-RI had little effect on the CD signal demonstrating $A\beta$ specificity
19 for the PCA-derived sequences.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 **ThT and CD experiments demonstrate that RI peptides do not aggregate in isolation** - ThT
44 experiments and Circular Dichroism spectroscopy experiments undertaken on RI-peptides in
45 isolation that have been incubated at 50 μ M for three days under conditions identical to
46 aggregation assays using $A\beta_{1-42}$ demonstrate that peptides do not bind significant amounts of
47 ThT, and that the CD signal for all peptides (at 0:1) is consistent with that of a random coil or
48 weakly helical conformation. They therefore indicate along with computational aggregation
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 prediction programs on the L-residue parent peptides (e.g. Waltz (40), Amylpred (41), Pasta
4 (42), Zyggregator (43), and Tango (44) that peptides do not form amyloid in isolation.
5
6
7

8
9
10 ***MTT experiments indicate reduced amyloid toxicity to cells*** - MTT ((3-(4,5-Dimethylthiazol-
11 2-yl)-2,5-diphenyltetrazolium bromide)) cell toxicity experiments were performed using Rat
12 phaeochromocytoma (PC12) neuronal-like cells to assess toxicity of $A\beta_{1-42}$ and the
13 preventative effects of the peptides generated in this study. MTT assays (Fig. 4) were
14 performed across $A\beta_{1-42}$ target:RI-peptide ratios and normalised relative to cells in isolation
15 (normalised as 0% death) and cells incubated with $A\beta_{1-42}$ alone (normalised as 100% death).
16 Peptides did not improve cell viability when incubated at 1:0.1. At the increased ratio of 1:1 a
17 decrease in toxicity of between ~30 % was observed for all RI-peptides studied. At increased
18 molar ratios of 1:2 and 1:4 a decrease in toxicity of approximately 10-20% was observed for all
19 RI-peptides studied. This experiment demonstrates a modest but significant reduction in $A\beta_{1-42}$
20 induced toxicity in the presence of RI-peptides. This was more promising than for the PCA
21 derived L-peptide parent molecules (15) and more impressive than the $iA\beta 5$ peptide which has
22 been previously shown to perform poorly in MTT cytotoxicity experiments using the PC12
23 cell-line (7). Reassuringly, the TAT-dummy-RI peptide had minimal effect of the viability of
24 PC12 cells incubated with $A\beta$.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 ***Oblique Angle Fluorescence (OAF) Microscopy indicates a reduction in amyloid levels*** – To
46 allow direct comparison, samples used in ThT and CD experiments were also imaged using
47 OAF microscopy (34) for both inhibition and reversal experiments. To prevent bias toward any
48 one sample the experiment was carried out blind. This technique allows for surface associated
49 and stacked aggregates of amyloid fibres to be imaged directly. OAF also permits assessment
50 of the amount of protein deposited as amyloid and its morphology. To further quantify the
51
52
53
54
55
56
57
58
59
60

1
2
3 amount of amyloid deposited we analysed the mean fluorescence value for each condition
4
5 using ImageJ (NIH, USA) over a randomly chosen 160x160 pixel² area and found a similar
6
7 correlation. In studying the peptides' effects on reversal we found KAT-RI, L2P1b-RI and
8
9 L2P2a-RI were the most potent, resulting in the fewest observed fibrils on the surface. Also
10
11 consistent with the reduction in ThT bound, L2P1a-RI and L2P1b-RI appear to have removed
12
13 the vast majority of fibrils present in the solution. Finally L2P2b-RI displays a number of
14
15 deposits that are smaller and of different morphology to the A β ₁₋₄₂ sample.
16
17

18
19 OAF microscopy data for inhibition experiments demonstrated that fibrils are present in
20
21 iA β ₅, L2P1a-RI, L2P1b-RI, L2P2a-RI and L2P2b-RI samples. However the average intensity
22
23 of these fibrils was reduced relative to A β ₁₋₄₂. KAT-RI stood out as the one sample with
24
25 reduced fluorescence intensity and no appearance of fibrils on the surface indicating that KAT-
26
27 RI reduced the deposition of amyloid. Collectively these data suggest that KAT-RI is most
28
29 effective overall, while L2P1b-RI and L2P2a-RI are also effective at reversal.
30
31
32
33

34 ***Retro-Inverso Peptides Rescue Developmental Delay in a Drosophila Model of A β Toxicity -***

36
37 Our initial experiments used retinal expression of the Arctic (E22G) variant of A β ₁₋₄₂ to screen
38
39 for amelioration of the expected rough eye phenotype. The flies were treated with RI peptides
40
41 in the food throughout larval development. Upon eclosion the severity of the phenotype was
42
43 scored by investigators blind to the treatment history. No differences in the rough eye
44
45 phenotype, as compared to untreated flies, could be detected for any of the peptides (data not
46
47 shown). We did however notice that some treated flies appeared to hatch either earlier or later
48
49 than normal. When we formally assessed the hatching times of flies, expressing wild type A β ₁₋₄₂,
50
51 we found that two RI peptides lacking a TAT sequence led to an increase in the hatching
52
53 time as compared to untreated control flies expressing A β ₁₋₄₂ (Fig. 6A, left panel). In addition
54
55 we saw three significant differences for TAT-containing peptides. Considering the A β ₁₋₄₂-
56
57
58
59
60

1
2
3 expressing flies we found that those treated with the TAT-L2P2-RI peptide hatched
4 significantly earlier while those treated with TAT-KAT-RI and TAT-L2P1-RI, which were
5 comparable to water-fed controls. Surprisingly treatment with a poly-gly dummy peptide
6 linked to the TAT sequence resulted in a more severe developmental delay as compared to
7 untreated flies (Fig. 6B, right panel). This sequence reassuringly demonstrates that the effect of
8 improved hatching is sequence specific, with this control sequence having the opposite effect.
9 As an additional control, non-transgenic flies were treated with TAT-L2P2-RI and TAT-
10 dummy-RI peptide; these flies hatched at the same time as control flies treated with water (data
11 not shown) suggesting that the effects on development are A β -specific.
12
13
14
15
16
17
18
19
20
21
22
23
24

25 ***Retro-Inverso Peptides Promote Locomotor Activity in a Drosophila Model of A β Toxicity-***

26
27 The expression of A β peptides in the nervous system of the fly is known to reduce the walking
28 velocity of flies with increasing age. To determine whether A β_{1-42} -expressing flies treated
29 with RI peptides might retain their locomotor function for longer. As we can see from Fig. 7A,
30 left panel, when we treated flies with RI peptides lacking a TAT sequence, there were only
31 small effects on the walking velocity, particularly for flies younger than 5 days. Those
32 peptides that carry a TAT sequence (Fig 7B, right panel) also increased the walking velocity of
33 young A β_{1-42} -expressing flies. In particular flies treated with the TAT-L2P1-RI peptide
34 exhibited markedly increased walking velocities.
35
36
37
38
39
40
41
42
43
44
45
46
47

48 ***Retro-inversed peptides have minimal effect on the presence of plaques -*** Oral administration

49 of RI peptides did not alter the pattern of A β_{42} deposits in the *Drosophila* brain (Figure 8).
50 Flies treated with KAT-RI, L2P1-RI, L2P2-RI, TAT-KAT-RI, TAT-L2P1-RI, TAT-L2P2-RI,
51 TAT-Dummy-RI and water all appeared to exhibit similar deposits of A β -reactive material
52
53
54
55
56
57
58
59
60

1
2
3 (arrow, stained with 6E10 antibody), suggesting that although amyloid deposits recognised by
4
5 this antibody have not been abolished, the production of toxic species has been modulated.
6
7

8 9 10 **DISCUSSION**

11
12 The PPI field has long been considered undruggable using conventional small
13
14 molecules, while peptide-based approaches have gained considerable traction in recent years
15
16 (23, 38). This owes to the fact that PPIs feature extended interacting surfaces with many
17
18 points of contact that are too shallow to accommodate traditional small molecule inhibitors,
19
20 making peptides and their mimetics promising candidates for intervention. Peptides often
21
22 form highly specific interactions with their target and many associated barriers (e.g. cell
23
24 permeability and bioavailability) can now be addressed via a number of modification options.
25
26 To develop peptide inhibitors of A β amyloidosis, we have combined a Protein-fragment
27
28 Complementation Assay (PCA) approach with semi-rational library design, and PCA
29
30 screening. This has been followed by retro-inversion of the selected peptide sequences to
31
32 identify molecules capable of binding A β_{1-42} and lowering toxicity. The process has been
33
34 iterative; the initial library search used the A β_{29-35} sequence as a design scaffold, and the
35
36 second library design was based on the initial winner, to yield a completely unrelated
37
38 sequence to the parent template. By retro-inverting these PCA library-selected sequences we
39
40 aimed to increase peptide bioavailability while retaining desirable inhibitory properties. Our
41
42 results demonstrate that the inhibitory properties found in the L-peptide templates are retained
43
44 in the RI sequences. For both inhibition and reversal ThT experiments we observed a
45
46 reduction in the ThT-bound to A β_{1-42} upon incubation with RI-sequences. Reassuringly the
47
48 reduction in ThT bound is less at increasingly sub-stoichiometric peptide concentrations,
49
50 demonstrating a dose dependency. ThT data is supported by a large decrease in the global β -
51
52 sheet CD signal upon examination of the same samples. These experiments therefore suggest
53
54
55
56
57
58
59
60

1
2
3 that RI-peptides can bind $A\beta_{1-42}$ and exert their effect by reducing the amount of the protein
4 in amyloid form. The reduction in CD signal was more pronounced than for ThT reflecting
5 that the assays measure different outputs. Thus while fibrils are being broken down into
6 smaller structures with significantly less β -sheet content, the species that become populated
7 are still able to bind ThT, although to a lesser extent than for $A\beta_{1-42}$ in isolation. In addition,
8 as was observed for L-peptides, the reduction in ThT bound is more pronounced for reversal
9 experiments, suggesting that RI-peptides are more effective at reversing preformed fibrils
10 than preventing amyloid assembly. RI-Peptides in isolation displayed either random coil or
11 weakly helical spectra, perhaps indicating a mechanism of binding that involves a structural
12 change in the target protein. MTT cytotoxicity data additionally indicate that although at sub-
13 stoichiometric ratios RI-peptides are effective at reducing β -sheet content and ThT-binding,
14 they are ineffective at causing a reduction in $A\beta_{1-42}$ toxicity. For this to be achieved molar
15 ratios of 1:1 or greater are required. This reduction in toxicity is modest (~30% at most),
16 suggesting that although populated amyloid species are not toxic to the bacteria in which they
17 were selected, the reduction in toxicity is however reduced when transferred to the context of
18 a mammalian line. Since bacterial cells harbouring parental L-peptide-DHFR2 fusions have
19 demonstrated an improvement in DHFR activity as well as increased bacterial doubling rates,
20 this suggests that toxicity reduction is successful within the confines of the selection system.
21 One possibility to circumvent this issue and improve outputs from the MTT assay would be to
22 use libraries based on existing sequences while transferring the PCA selection to a
23 therapeutically relevant neuronal cell line (20). However, testing these RI compounds in
24 *Drosophila melanogaster* has demonstrated that several sequences can go on to improve
25 either hatching rates or locomotor activity, with TAT-L2P2-RI being the most striking for the
26 former and TAT-L2P1-RI for the latter. Lastly, the presence of comparable brain plaque
27 levels for each of the peptides suggests that amyloid deposits recognised by this antibody
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 have not been abolished. Interpreting this result alongside fly hatching times and locomotor
4
5 activity data suggests that the production of the toxic A β species has been modulated. These
6
7 results compare favourably with ThT, CD, OAF microscopy and MTT studies in
8
9 demonstrating that peptides function by modulating amyloid levels and consequently
10
11 reducing associated cytotoxicity. The RI-PCA approach therefore offers the potential to
12
13 derive protease resistant A β -interacting peptides that are capable of lowering the toxicity
14
15 associated with amyloid, which may in turn serve as potential precursors for an Alzheimer's
16
17 disease treatment.
18
19

20 21 22 23 **ACKNOWLEDGEMENTS**

24
25 The authors wish to thank Dr Miao Yu and Dr Victoria Allen-Baume for excellent technical
26
27 assistance throughout the project.
28
29

30 31 32 33 **REFERENCES**

- 34
35
36 1. Karran, E., Mercken, M., and De Strooper, B. (2011) The amyloid cascade hypothesis
37
38 for Alzheimer's disease: an appraisal for the development of therapeutics, *Nat Rev Drug*
39
40 *Discov 10*, 698-712.
41
42 2. Haass, C., and Selkoe, D. J. (2007) Soluble protein oligomers in neurodegeneration:
43
44 lessons from the Alzheimer's amyloid beta-peptide, *Nat Rev Mol Cell Biol 8*, 101-112.
45
46 3. McLean, C. A., Cherny, R. A., Fraser, F. W., Fuller, S. J., Smith, M. J., Beyreuther, K.,
47
48 Bush, A. I., and Masters, C. L. (1999) Soluble pool of Abeta amyloid as a determinant
49
50 of severity of neurodegeneration in Alzheimer's disease, *Ann Neurol 46*, 860-866.
51
52 4. Nilsberth, C., Westlind-Danielsson, A., Eckman, C. B., Condron, M. M., Axelman, K.,
53
54 Forsell, C., Stenh, C., Luthman, J., Teplow, D. B., Younkin, S. G., Naslund, J., and
55
56
57
58
59
60

- 1
2
3 Lannfelt, L. (2001) The 'Arctic' APP mutation (E693G) causes Alzheimer's disease by
4 enhanced Abeta protofibril formation, *Nat Neurosci* 4, 887-893.
5
6
7
8 5. Tjernberg, L. O., Naslund, J., Lindqvist, F., Johansson, J., Karlstrom, A. R., Thyberg,
9 J., Terenius, L., and Nordstedt, C. (1996) Arrest of beta-amyloid fibril formation by a
10 pentapeptide ligand, *J Biol Chem* 271, 8545-8548.
11
12
13
14 6. Hughes, E., Burke, R. M., and Doig, A. J. (2000) Inhibition of toxicity in the beta-
15 amyloid peptide fragment beta -(25-35) using N-methylated derivatives: a general
16 strategy to prevent amyloid formation, *J Biol Chem* 275, 25109-25115.
17
18
19
20
21 7. Kokkoni, N., Stott, K., Amijee, H., Mason, J. M., and Doig, A. J. (2006) N-Methylated
22 peptide inhibitors of beta-amyloid aggregation and toxicity. Optimization of the
23 inhibitor structure, *Biochemistry* 45, 9906-9918.
24
25
26
27
28 8. Tjernberg, L. O., Callaway, D. J., Tjernberg, A., Hahne, S., Lilliehook, C., Terenius, L.,
29 Thyberg, J., and Nordstedt, C. (1999) A molecular model of Alzheimer amyloid beta-
30 peptide fibril formation, *J Biol Chem* 274, 12619-12625.
31
32
33
34 9. Gordon, D. J., Sciarretta, K. L., and Meredith, S. C. (2001) Inhibition of beta-
35 amyloid(40) fibrillogenesis and disassembly of beta-amyloid(40) fibrils by short beta-
36 amyloid congeners containing N-methyl amino acids at alternate residues, *Biochemistry*
37 40, 8237-8245.
38
39
40
41
42
43 10. Soto, C., Kindy, M. S., Baumann, M., and Frangione, B. (1996) Inhibition of
44 Alzheimer's amyloidosis by peptides that prevent beta-sheet conformation, *Biochem*
45 *Biophys Res Commun* 226, 672-680.
46
47
48
49
50 11. Soto, C., Sigurdsson, E. M., Morelli, L., Kumar, R. A., Castano, E. M., and Frangione,
51 B. (1998) Beta-sheet breaker peptides inhibit fibrillogenesis in a rat brain model of
52 amyloidosis: implications for Alzheimer's therapy, *Nat Med* 4, 822-826.
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
12. O'Nuallain, B., Freir, D. B., Nicoll, A. J., Risse, E., Ferguson, N., Herron, C. E., Collinge, J., and Walsh, D. M. (2010) Amyloid beta-protein dimers rapidly form stable synaptotoxic protofibrils, *J Neurosci* 30, 14411-14419.
 13. O'Nuallain, B., Klyubin, I., Mc Donald, J. M., Foster, J. S., Welzel, A., Barry, A., Dykoski, R. K., Cleary, J. P., Gebbink, M. F., Rowan, M. J., and Walsh, D. M. (2011) A monoclonal antibody against synthetic Abeta dimer assemblies neutralizes brain-derived synaptic plasticity-disrupting Abeta, *J Neurochem* 119, 189-201.
 14. Giuffrida, M. L., Caraci, F., Pignataro, B., Cataldo, S., De Bona, P., Bruno, V., Molinaro, G., Pappalardo, G., Messina, A., Palmigiano, A., Garozzo, D., Nicoletti, F., Rizzarelli, E., and Copani, A. (2009) Beta-amyloid monomers are neuroprotective, *J Neurosci* 29, 10582-10587.
 15. Acerra, N., Kad, N. M., and Mason, J. M. (2013) Combining Intracellular Selection with Protein-fragment Complementation to Derive A β interacting Peptides, *Protein Eng Des Sel* 26, 463-470.
 16. Doig, A. J. (2007) Peptide inhibitors of beta-amyloid aggregation, *Curr Opin Drug Discov Devel* 10, 533-539.
 17. Lowe, T. L., Strzelec, A., Kiessling, L. L., and Murphy, R. M. (2001) Structure-function relationships for inhibitors of beta-amyloid toxicity containing the recognition sequence KLVFF, *Biochemistry* 40, 7882-7889.
 18. Pelletier, J. N., Campbell-Valois, F. X., and Michnick, S. W. (1998) Oligomerization domain-directed reassembly of active dihydrofolate reductase from rationally designed fragments, *Proc Natl Acad Sci U S A* 95, 12141-12146.
 19. Mason, J. M., Schmitz, M. A., Muller, K. M., and Arndt, K. M. (2006) Semirational design of Jun-Fos coiled coils with increased affinity: Universal implications for leucine zipper prediction and design, *Proc Natl Acad Sci U S A* 103, 8989-8994.

- 1
2
3 20. Remy, I., Campbell-Valois, F. X., and Michnick, S. W. (2007) Detection of protein-
4 protein interactions using a simple survival protein-fragment complementation assay
5 based on the enzyme dihydrofolate reductase, *Nat Protoc* 2, 2120-2125.
6
7
8
9
10 21. Chorev, M., and Goodman, M. (1995) Recent developments in retro peptides and
11 proteins--an ongoing topochemical exploration, *Trends Biotechnol* 13, 438-445.
12
13
14 22. Fletcher, M. D., and Campbell, M. M. (1998) Partially Modified Retro-Inverso
15 Peptides: Development, Synthesis, and Conformational Behavior, *Chem Rev* 98, 763-
16 796.
17
18
19
20
21 23. Mason, J. M. (2010) Design and development of peptides and peptide mimetics as
22 antagonists for therapeutic intervention, *Future Med Chem* 2, 1813-1822.
23
24
25 24. Matharu, B., El-Agnaf, O., Razvi, A., and Austen, B. M. (2010) Development of retro-
26 inverso peptides as anti-aggregation drugs for beta-amyloid in Alzheimer's disease,
27 *Peptides* 31, 1866-1872.
28
29
30
31
32 25. Taylor, M., Moore, S., Mayes, J., Parkin, E., Beeg, M., Canovi, M., Gobbi, M., Mann,
33 D. M., and Allsop, D. (2010) Development of a proteolytically stable retro-inverso
34 peptide inhibitor of beta-amyloid oligomerization as a potential novel treatment for
35 Alzheimer's disease, *Biochemistry* 49, 3261-3272.
36
37
38
39
40
41 26. Parthasarathy, V., McClean, P. L., Holscher, C., Taylor, M., Tinker, C., Jones, G.,
42 Kolosov, O., Salvati, E., Gregori, M., Masserini, M., and Allsop, D. (2013) A novel
43 retro-inverso peptide inhibitor reduces amyloid deposition, oxidation and inflammation
44 and stimulates neurogenesis in the APP^{swe}/PS1^{DeltaE9} mouse model of Alzheimer's
45 disease, *PLoS One* 8, e54769.
46
47
48
49
50
51
52 27. Wiesehan, K., Buder, K., Linke, R. P., Patt, S., Stoldt, M., Unger, E., Schmitt, B.,
53 Bucci, E., and Willbold, D. (2003) Selection of D-Amino-Acid peptides that bind to
54
55
56
57
58
59
60

- 1
2
3 Alzheimer's disease amyloid peptide A beta(1-42) by mirror image phage display,
4
5 *Chembiochem* 4, 748-753.
6
- 7 28. Mason, J. M., Muller, K. M., and Arndt, K. M. (2007) Positive aspects of negative
8 design: simultaneous selection of specificity and interaction stability, *Biochemistry* 46,
9 4804-4814.
10
- 11 29. Mason, J. M., Hagemann, U. B., and Arndt, K. M. (2009) Role of hydrophobic and
12 electrostatic interactions in coiled coil stability and specificity, *Biochemistry* 48, 10380-
13 10388.
14
- 15 30. Virnekas, B., Ge, L., Pluckthun, A., Schneider, K. C., Wellnhofer, G., and Moroney, S.
16 E. (1994) Trinucleotide phosphoramidites: ideal reagents for the synthesis of mixed
17 oligonucleotides for random mutagenesis, *Nucleic Acids Res* 22, 5600-5607.
18
- 19 31. Zagorski, M. G., Yang, J., Shao, H., Ma, K., Zeng, H., and Hong, A. (1999)
20 Methodological and chemical factors affecting amyloid beta peptide amyloidogenicity,
21 *Methods Enzymol* 309, 189-204.
22
- 23 32. LeVine, H., 3rd. (1993) Thioflavine T interaction with synthetic Alzheimer's disease
24 beta-amyloid peptides: detection of amyloid aggregation in solution, *Protein Sci* 2, 404-
25 410.
26
- 27 33. Shearman, M. S., Ragan, C. I., and Iversen, L. L. (1994) Inhibition of PC12 cell redox
28 activity is a specific, early indicator of the mechanism of beta-amyloid-mediated cell
29 death, *Proc Natl Acad Sci U S A* 91, 1470-1474.
30
- 31 34. Kad, N. M., Wang, H., Kennedy, G. G., Warshaw, D. M., and Van Houten, B. (2010)
32 Collaborative dynamic DNA scanning by nucleotide excision repair proteins
33 investigated by single- molecule imaging of quantum-dot-labeled proteins, *Mol Cell* 37,
34 702-713.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 35. Frankel, A. D., and Pabo, C. O. (1988) Cellular Uptake of the Tat Protein from Human
4 Immunodeficiency Virus, *Cell* 55, 1189-1193.
5
6
7 36. Fawell, S., Seery, J., Daikh, Y., Moore, C., Chen, L. L., Pepinsky, B., and Barsoum, J.
8 (1994) Tat-Mediated Delivery of Heterologous Proteins into Cells, *P Natl Acad Sci*
9 *USA* 91, 664-668.
10
11
12
13
14 37. Morris, M. C., Deshayes, S., Heitz, F., and Divita, G. (2008) Cell-penetrating peptides:
15 from molecular mechanisms to therapeutics, *Biology of the Cell* 100, 201-217.
16
17
18 38. Craik, D. J., Fairlie, D. P., Liras, S., and Price, D. (2013) The future of peptide-based
19 drugs, *Chem Biol Drug Des* 81, 136-147.
20
21
22
23 39. Biancalana, M., and Koide, S. (2010) Molecular mechanism of Thioflavin-T binding to
24 amyloid fibrils, *Biochim Biophys Acta* 1804, 1405-1412.
25
26
27 40. Maurer-Stroh, S., Debulpaep, M., Kuemmerer, N., Lopez de la Paz, M., Martins, I. C.,
28 Reumers, J., Morris, K. L., Copland, A., Serpell, L., Serrano, L., Schymkowitz, J. W.,
29 and Rousseau, F. (2010) Exploring the sequence determinants of amyloid structure
30 using position-specific scoring matrices, *Nat Methods* 7, 237-242.
31
32
33
34 41. Frousios, K. K., Iconomidou, V. A., Karletidi, C. M., and Hamodrakas, S. J. (2009)
35 Amyloidogenic determinants are usually not buried, *BMC Struct Biol* 9, 44.
36
37
38
39 42. Trovato, A., Seno, F., and Tosatto, S. C. (2007) The PASTA server for protein
40 aggregation prediction, *Protein Eng Des Sel* 20, 521-523.
41
42
43
44 43. Tartaglia, G. G., and Vendruscolo, M. (2008) The Zyggregator method for predicting
45 protein aggregation propensities, *Chem Soc Rev* 37, 1395-1401.
46
47
48
49 44. Fernandez-Escamilla, A. M., Rousseau, F., Schymkowitz, J., and Serrano, L. (2004)
50 Prediction of sequence-dependent and mutational effects on the aggregation of peptides
51 and proteins, *Nat Biotechnol* 22, 1302-1306.
52
53
54
55
56
57
58
59
60

FIGURE LEGENDS

Figure 1: ThT Inhibition and Reversal Data. The data in a) and b) show the effects of different stoichiometries of the peptides KAT-RI, L2P1a-RI, L2P1b-RI, L2P2a-RI, and L2P2b-RI on the aggregation of 50 μ M A β ₁₋₄₂ (at three days for the inhibition assay and at six days for reversal assay). Molar ratios (A β :peptide) are shown at 1:0.0001, 1:0.001, 1:0.01, 1:0.1, 1:1, 1:2 and 1:4 for each peptide. iA β 5 and TAT-dummy-RI are included as positive and negative controls respectively. Errors are given as the standard deviation of all errors at each molar ratio. The data show that for the three lowest molar ratios (all sub-stoichiometric; 1:0.01, 1:0.001, and 1:0.0001) the average reduction in ThT bound was minimal (106% for Inhibition and 89% for reversal). In contrast at the three highest molar ratios the reduction in ThT bound was significantly greater (58% for inhibition, 38% for reversal). The most effective average molar ratio for peptides was 1:0.1 which displayed ThT bound values of 35% and 26% for inhibition and reversal, approximating to 71% and 63% less than the average of the three lowest stoichiometries respectively.

Figure 2: Circular Dichroism spectra show i) β -sheet content of RI peptides with A β ii) RI peptides alone and iii) with RI-peptides subtracted from the combined signal (i.e. ii) – i)). This is shown at all molar ratios studied for inhibition and reversal conditions. Both experiments are undertaken at three days postmix and performed at a concentration 10 μ M A β ₁₋₄₂.

Figure 3: Circular Dichroism spectroscopy and ThT experiments undertaken on RI-peptides in isolation that have been incubated at 50 μ M for three days under conditions identical to aggregation assays using A β ₁₋₄₂.

1
2
3 **Figure 4:** MTT toxicity assay using $A\beta_{1-42}$ and selected RI-peptides using different molar ratios
4 after 24 hours of incubation. The assay was performed with 10 μM $A\beta_{1-42}$ and different
5 concentrations of inhibitor, for example, 1:0.1 (1 μM), 1:1 (10 μM), 1:2 (20 μM), 1:4 (40 μM).
6
7
8
9

10 **Figure 5:** Oblique Angle Fluorescence (OAF) Microscopy data. During reversal experiments,
11 $A\beta_{1-42}$ was grown alone for three days, after which RI-peptide was added at a stoichiometry of
12 1:4 and followed by a further three day incubation to assay for peptide induced reversal of
13 amyloid deposition. Each sample was then imaged by fluorescence microscopy and panels
14 showing representative images obtained. To quantify amyloid deposition the mean grey value
15 over a 160x160 pixel area randomly chosen for five separate images is plotted as fluorescence
16 intensity. Each data point is normalized to the control iA β_5 peptide by subtraction. Shown are
17 a) Inhibition data and b) Reversal data. It can be clearly seen that both KAT and L2P1B are
18 strongly inhibitory for this reversal assay. The scale bars represent a distance of 2 μm . Each
19 data point was then scaled to overcome the 'background noise' by taking $A\beta$ (1:0) as the
20 maximum.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 **Figure 6:** Oral administration of RI peptides suppresses the delayed eclosion associated with
37 pan-neuronal expression of $A\beta_{42}$. A) As compared to $A\beta_{42}$ treated with water (control), those
38 flies treated with KAT-RI and L2P1-RI showed small but significant suppression of the
39 developmental delay ($p < 0.05$). B) Upon fusion with the TAT peptide the potency of the RI
40 peptides increased with TAT-KAT-RI ($p < 0.05$) and TAT-L2P2-RI ($p < 0.001$) exhibiting
41 significant suppression of delayed eclosion as compared to water (control). Oral administration
42 of a dummy RI peptide resulted in significant increase in the delay in eclosion ($p < 0.05$).
43 Significance determined by the log-rank test.
44
45
46
47
48
49
50
51
52
53

54 **Figure 7:** Oral administration of RI peptides suppresses the locomotor deficits associated with
55 pan-neuronal expression of $A\beta_{42}$. A) As compared to flies treated with water (control), flies
56
57
58
59
60


1
2
3 receiving RI peptides recorded higher walking velocities in the first days of adult life. The
4
5 rank order of velocities was L2P1>L2P2>KAT>control. B) Upon fusion with the TAT peptide
6
7 the potency of the TAT-L2P1-RI peptide was particularly enhanced. The rank order of the
8
9 velocities was similar: L2P1>L2P2>(Dummy>)KAT>control.
10
11

12 **Figure 8:** Oral administration of RI peptides did not alter the pattern of A β ₄₂ deposits in the
13 *Drosophila* brain. Flies treated with KAT-RI (A), L2P1-RI (B), L2P2-RI (C), TAT-KAT-RI
14
15 (D), TAT-L2P1-RI (E), TAT-L2P2-RI (F), TAT-Dummy-RI (G) and water (H) all exhibited
16
17 similar deposits of A β -reactive material (arrow, stained with 6E10 antibody).
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Name	Sequence
KAT-RI	mltakag-NH ₂
L2P1a-RI	nstaksf-NH ₂
L2P1b-RI	pagnstaksfsa-NH ₂
L2P2a-RI	attakvp-NH ₂
L2P2b-RI	pagattakvpsa-NH ₂
Tat-KAT-RI	rrrqrrkkrmItakag-NH ₂
Tat-L2P1-RI	rrrqrrkkrnstaksf-NH ₂
Tat-L2P2-RI	rrrqrrkkrrattakvp-NH ₂
TAT-dummy-RI	rrrqrrkkrrggggggg-NH ₂
iAβ5	LPFFD-NH ₂

Table 1: PCA derived retro-inversed sequences and related sequences with retro-inversed TAT-fusions to promote intracellular uptake for drosophila melanogaster studies.

a) Inhibition




b) Reversal




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58


Inhibition 1:0.1




Reversal 1:0.1




Inhibition 1:1




Reversal 1:1




Inhibition 1:2




Reversal 1:2



Inhibition 1:4



Reversal 1:4




- $A\beta_{1-42}$
- KAT-RI
- L2P1a-RI
- L2P1b-RI
- L2P2a-RI
- L2P2b-RI


Wavelength (nm)

Wavelength (nm)


ThT RI Inhibition 0:1 Day 3




CD RI Inhibition 0:1 Day 3




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43





Inhibition experiments at 1:4 Stoichiometry






Reversal experiments at 1:4 Stoichiometry





1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43





1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43