

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a preprint version which may differ from the publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/106928>

Please be advised that this information was generated on 2017-12-06 and may be subject to change.

Search for Neutral Higgs Bosons in Events with Multiple Bottom Quarks at the Tevatron

T. Aaltonen^{†,12} V.M. Abazov^{†,48} B. Abbott^{†,112} B.S. Acharya^{†,31} M. Adams^{†,78} T. Adams^{†,74} G.D. Alexeev^{†,48}
 G. Alkhazov^{†,52} A. Alton^{†a,96} B. Álvarez González^{†a,57} G. Alverson^{†,92} S. Amerio^{†,35} D. Amidei^{†,96}
 A. Anastassov^{†b,76} A. Annovi^{†,34} J. Antos^{†,53} G. Apollinari^{†,76} J.A. Appel^{†,76} T. Arisawa^{†,41} A. Artikov^{†,48}
 J. Asaadi^{†,119} W. Ashmanskas^{†,76} A. Askew^{†,74} S. Atkins^{†,89} B. Auerbach^{†,72} K. Augsten^{†,9} A. Aurisano^{†,119}
 C. Avila^{†,7} F. Azfar^{†,66} F. Badaud^{†,13} W. Badgett^{†,76} T. Bae^{†,43} L. Bagby^{†,76} B. Baldin^{†,76} D.V. Bandurin^{†,74}
 S. Banerjee^{†,31} A. Barbaro-Galtieri^{†,68} E. Barberis^{†,92} P. Baringer^{†,87} V.E. Barnes^{†,85} B.A. Barnett^{†,90}
 P. Barria^{†#d,36} J.F. Bartlett^{†,76} P. Bartos^{†,53} U. Bassler^{†,18} M. Bauc^{†#b,35} V. Bazterra^{†,78} A. Bean^{†,87}
 F. Bedeschi^{†,36} M. Begalli^{†,2} S. Behari^{†,90} L. Bellantoni^{†,76} G. Bellettini^{†#c,36} J. Bellinger^{†,125} D. Benjamin^{†,109}
 A. Beretvas^{†,76} S.B. Beri^{†,29} G. Bernardi^{†,17} R. Bernhard^{†,22} I. Bertram^{†,61} M. Besançon^{†,18} R. Beuselinck^{†,63}
 P.C. Bhat^{†,76} S. Bhatia^{†,99} V. Bhatnagar^{†,29} A. Bhatti^{†,105} D. Bisello^{†#b,35} I. Bizjak^{†,64} K.R. Bland^{†,122}
 G. Blazey^{†,79} S. Blessing^{†,74} K. Bloom^{†,100} B. Blumenfeld^{†,90} A. Bocci^{†,109} A. Bodek^{†,106} A. Boehnlein^{†,76}
 D. Boline^{†,107} E.E. Boos^{†,50} G. Borissov^{†,61} D. Bortoletto^{†,85} T. Bose^{†,91} J. Boudreau^{†,116} A. Boveia^{†,77}
 A. Brandt^{†,118} O. Brandt^{†,23} L. Brigliadori^{†#a,33} R. Brock^{†,98} C. Bromberg^{†,98} A. Bross^{†,76} D. Brown^{†,17}
 J. Brown^{†,17} E. Brucken^{†,12} J. Budagov^{†,48} X.B. Bu^{†,76} H.S. Budd^{†,106} M. Buehler^{†,76} V. Buescher^{†,25}
 V. Bunichev^{†,50} S. Burdin^{†b,61} K. Burkett^{†,76} G. Busetto^{†#b,35} P. Bussey^{†,60} C.P. Buszello^{†,58} A. Buzatu^{†,4}
 A. Calamba^{†,115} C. Calancha^{†,56} E. Camacho-Pérez^{†,45} S. Camarda^{†,54} M. Campanelli^{†,64} M. Campbell^{†,96}
 F. Canelli^{†,77} B. Carls^{†,81} D. Carlsmith^{†,125} R. Carosi^{†,36} S. Carrillo^{†c,73} S. Carron^{†,76} B. Casal^{†d,57} M. Casarsa^{†,38}
 B.C.K. Casey^{†,76} H. Castilla-Valdez^{†,45} A. Castro^{†#a,33} P. Catastini^{†,93} S. Caughron^{†,98} D. Cauz^{†,38} V. Cavaliere^{†,81}
 M. Cavalli-Sforza^{†,54} A. Cerri^{†e,68} L. Cerrito^{†f,64} S. Chakrabarti^{†,107} D. Chakraborty^{†,79} K.M. Chan^{†,84}
 A. Chandra^{†,121} E. Chapon^{†,18} G. Chen^{†,87} Y.C. Chen^{†,6} M. Chertok^{†,69} S. Chevalier-Théry^{†,18} G. Chiarelli^{†,36}
 G. Chlachidze^{†,76} F. Chlebana^{†,76} D.K. Cho^{†,117} K. Cho^{†,43} S.W. Cho^{†,44} S. Choi^{†,44} D. Chokheli^{†,48}
 B. Choudhary^{†,30} W.H. Chung^{†,125} Y.S. Chung^{†,106} S. Cihangir^{†,76} M.A. Ciocci^{†#d,36} D. Claes^{†,100} A. Clark^{†,59}
 C. Clarke^{†,97} J. Clutter^{†,87} G. Compostella^{†#b,35} M.E. Convery^{†,76} J. Conway^{†,69} M. Cooke^{†,76} W.E. Cooper^{†,76}
 M. Corbo^{†,76} M. Corcoran^{†,121} M. Cordelli^{†,34} F. Couderc^{†,18} M.-C. Cousinou^{†,15} C.A. Cox^{†,69} D.J. Cox^{†,69}
 F. Crescioli^{†#c,36} A. Croc^{†,18} J. Cuevas^{†a,57} R. Culbertson^{†,76} D. Cutts^{†,117} D. Dagenhart^{†,76} N. d'Ascenzo^{†g,76}
 A. Das^{†,67} M. Datta^{†,76} G. Davies^{†,63} P. de Barbaro^{†,106} S.J. de Jong^{†,46,47} E. De La Cruz-Burelo^{†,45} F. Déliot^{†,18}
 M. Dell'Orso^{†#c,36} R. Demina^{†,106} L. Demortier^{†,105} M. Deninno^{†,33} D. Denisov^{†,76} S.P. Denisov^{†,51} M. d'Errico^{†#b,35}
 S. Desai^{†,76} C. Deterre^{†,18} K. DeVaughan^{†,100} F. Devoto^{†,12} A. Di Canto^{†#c,36} B. Di Ruzza^{†,76} H.T. Diehl^{†,76}
 M. Diesburg^{†,76} P.F. Ding^{†,65} J.R. Dittmann^{†,122} A. Dominguez^{†,100} S. Donati^{†#c,36} P. Dong^{†,76} M. D'Onofrio^{†,62}
 M. Dorigo^{†,38} T. Dorigo^{†,35} A. Dubey^{†,30} L.V. Dudko^{†,50} D. Duggan^{†,101} A. Duperrin^{†,15} S. Dutt^{†,29}
 A. Dyshkant^{†,79} M. Eads^{†,100} K. Ebina^{†,41} D. Edmunds^{†,98} A. Elagin^{†,119} J. Ellison^{†,71} V.D. Elvira^{†,76} Y. Enari^{†,17}
 A. Eppig^{†,96} R. Erbacher^{†,69} S. Errede^{†,81} N. Ershaidat^{†h,76} R. Eusebi^{†,119} H. Evans^{†,82} A. Evdokimov^{†,108}
 V.N. Evdokimov^{†,51} G. Facini^{†,92} S. Farrington^{†,66} M. Feindt^{†,24} L. Feng^{†,79} T. Ferbel^{†,106} J.P. Fernandez^{†,56}
 F. Fiedler^{†,25} R. Field^{†,73} F. Filthaut^{†,46,47} W. Fisher^{†,98} H.E. Fisk^{†,76} G. Flanagan^{†i,76} R. Forrest^{†,69}
 M. Fortner^{†,79} H. Fox^{†,61} M.J. Frank^{†,122} M. Franklin^{†,93} J.C. Freeman^{†,76} S. Fuess^{†,76} Y. Funakoshi^{†,41} I. Furic^{†,73}
 M. Gallinaro^{†,105} A. Garcia-Bellido^{†,106} J.E. Garcia^{†,59} J.A. García-González^{†,45} G.A. García-Guerra^{†c,45}
 A.F. Garfinkel^{†,85} P. Garosi^{†#d,36} V. Gavrilov^{†,49} P. Gay^{†,13} W. Geng^{†,15,98} D. Gerbaudo^{†,102} C.E. Gerber^{†,78}
 H. Gerberich^{†,81} E. Gerchtein^{†,76} Y. Gershtein^{†,101} S. Giagu^{†,37} V. Giakoumopoulou^{†,28} P. Giannetti^{†,36}
 K. Gibson^{†,116} C.M. Ginsburg^{†,76} G. Ginter^{†,76,106} N. Giokaris^{†,28} P. Giromini^{†,34} G. Giurgiu^{†,90} V. Glagolev^{†,48}
 D. Glenzinski^{†,76} M. Gold^{†,103} D. Goldin^{†,119} N. Goldschmidt^{†,73} A. Golossanov^{†,76} G. Golovanov^{†,48}
 G. Gomez-Ceballos^{†,94} G. Gomez^{†,57} M. Goncharov^{†,94} O. González^{†,56} I. Gorelov^{†,103} A.T. Goshaw^{†,109}
 K. Goulianos^{†,105} A. Goussiou^{†,124} P.D. Grannis^{†,107} S. Greder^{†,19} H. Greenlee^{†,76} G. Grenier^{†,20} S. Grinstein^{†,54}
 Ph. Gris^{†,13} J.-F. Grivaz^{†,16} A. Grohsjean^{†#d,18} C. Grosso-Pilcher^{†,77} R.C. Group^{†,123,76} S. Grünendahl^{†,76}
 M.W. Grünewald^{†,32} T. Guillemain^{†,16} J. Guimaraes da Costa^{†,93} G. Gutierrez^{†,76} P. Gutierrez^{†,112} S. Hagopian^{†,74}
 S.R. Hahn^{†,76} J. Haley^{†,92} E. Halkiadakis^{†,101} A. Hamaguchi^{†,40} J.Y. Han^{†,106} L. Han^{†,5} F. Happacher^{†,34}
 K. Hara^{†,42} K. Harder^{†,65} D. Hare^{†,101} M. Hare^{†,95} A. Harel^{†,106} R.F. Harr^{†,97} K. Hatakeyama^{†,122}
 J.M. Hauptman^{†,86} C. Hays^{†,66} J. Hays^{†,63} T. Head^{†,65} T. Hebbeker^{†,21} M. Heck^{†,24} D. Hedin^{†,79} H. Hegab^{†,113}

J. Heinrich[†],¹¹⁴ A.P. Heinson[†],⁷¹ U. Heintz[†],¹¹⁷ C. Hensel[†],²³ I. Heredia-De La Cruz[†],⁴⁵ M. Herndon[†],¹²⁵
 K. Herner[†],⁹⁶ G. Hesketh[†],⁶⁵ S. Hewamanage[†],¹²² M.D. Hildreth[†],⁸⁴ R. Hirosky[†],¹²³ T. Hoang[†],⁷⁴ J.D. Hobbs[†],¹⁰⁷
 A. Hocker[†],⁷⁶ B. Hoeneisen[†],¹¹ J. Hogan[†],¹²¹ M. Hohlfeld[†],²⁵ W. Hopkins[†],⁷⁶ D. Horn[†],²⁴ S. Hou[†],⁶ I. Howley[†],¹¹⁸
 Z. Hubacek[†],^{9,18} R.E. Hughes[†],¹¹⁰ M. Hurwitz[†],⁷⁷ U. Husemann[†],⁷² N. Hussain[†],⁴ M. Hussein[†],⁹⁸ J. Huston[†],⁹⁸
 V. Hynek[†],⁹ I. Iashvili[†],¹⁰⁴ Y. Ilchenko[†],¹²⁰ R. Illingworth[†],⁷⁶ G. Introzzi[†],³⁶ M. Iori[†],³⁷ A.S. Ito[†],⁷⁶ A. Ivanov[†],⁶⁹
 S. Jabeen[†],¹¹⁷ M. Jaffré[†],¹⁶ E. James[†],⁷⁶ D. Jang[†],¹¹⁵ A. Jayasinghe[†],¹¹² B. Jayatilaka[†],¹⁰⁹ E.J. Jeon[†],⁴³
 M.S. Jeong[†],⁴⁴ R. Jesik[†],⁶³ S. Jindariani[†],⁷⁶ K. Johns[†],⁶⁷ E. Johnson[†],⁹⁸ M. Johnson[†],⁷⁶ A. Jonckheere[†],⁷⁶
 M. Jones[†],⁸⁵ P. Jonsson[†],⁶³ K.K. Joo[†],⁴³ J. Joshi[†],⁷¹ S.Y. Jun[†],¹¹⁵ A.W. Jung[†],⁷⁶ T.R. Junk[†],⁷⁶ A. Juste[†],⁵⁵
 K. Kaadze[†],⁸⁸ E. Kajfasz[†],¹⁵ T. Kamon[†],¹¹⁹ P.E. Karchin[†],⁹⁷ D. Karmanov[†],⁵⁰ A. Kasmi[†],¹²² P.A. Kasper[†],⁷⁶
 Y. Kato[†],⁴⁰ I. Katsanos[†],¹⁰⁰ R. Kehoe[†],¹²⁰ S. Kermiche[†],¹⁵ W. Ketchum[†],⁷⁷ J. Keung[†],¹¹⁴ N. Khalatyan[†],⁷⁶
 A. Khanov[†],¹¹³ A. Kharchilava[†],¹⁰⁴ Y.N. Kharzhev[†],⁴⁸ V. Khotilovich[†],¹¹⁹ B. Kilminster[†],⁷⁶ D.H. Kim[†],⁴³
 H.S. Kim[†],⁴³ J.E. Kim[†],⁴³ M.J. Kim[†],³⁴ S.B. Kim[†],⁴³ S.H. Kim[†],⁴² Y.J. Kim[†],⁴³ Y.K. Kim[†],⁷⁷ N. Kimura[†],⁴¹
 M. Kirby[†],⁷⁶ I. Kiselevich[†],⁴⁹ S. Klimenko[†],⁷³ K. Knoepfel[†],⁷⁶ J.M. Kohli[†],²⁹ K. Kondo[†],⁴¹ D.J. Kong[†],⁴³
 J. Konigsberg[†],⁷³ A.V. Kotwal[†],¹⁰⁹ A.V. Kozelov[†],⁵¹ J. Kraus[†],⁹⁹ M. Kreps[†],²⁴ J. Kroll[†],¹¹⁴ D. Krop[†],⁷⁷
 M. Kruse[†],¹⁰⁹ V. Krutelyov[†],¹¹⁹ T. Kuhr[†],²⁴ S. Kulikov[†],⁵¹ A. Kumar[†],¹⁰⁴ A. Kupco[†],¹⁰ M. Kurata[†],⁴²
 T. Kurča[†],²⁰ V.A. Kuzmin[†],⁵⁰ S. Kwang[†],⁷⁷ A.T. Laasanen[†],⁸⁵ S. Lami[†],³⁶ S. Lammel[†],⁷⁶ S. Lammers[†],⁸²
 M. Lancaster[†],⁶⁴ R.L. Lander[†],⁶⁹ G. Landsberg[†],¹¹⁷ K. Lannon[†],¹¹⁰ A. Lath[†],¹⁰¹ G. Latino[†],³⁶ P. Lebrun[†],²⁰
 T. LeCompte[†],⁷⁵ E. Lee[†],¹¹⁹ H.S. Lee[†],⁴⁴ H.S. Lee[†],⁷⁷ J.S. Lee[†],⁴³ S.W. Lee[†],¹¹⁹ S.W. Lee[†],⁸⁶ W.M. Lee[†],⁷⁶
 X. Lei[†],⁶⁷ J. Lellouch[†],¹⁷ S. Leo[†],³⁶ S. Leone[†],³⁶ J.D. Lewis[†],⁷⁶ H. Li[†],¹⁴ L. Li[†],⁷¹ Q.Z. Li[†],⁷⁶ J.K. Lim[†],⁴⁴
 A. Limosani[†],¹⁰⁹ D. Lincoln[†],⁷⁶ C.-J. Lin[†],⁶⁸ M. Lindgren[†],⁷⁶ J. Linnemann[†],⁹⁸ V.V. Lipaev[†],⁵¹ E. Lipeles[†],¹¹⁴
 R. Lipton[†],⁷⁶ A. Lister[†],⁵⁹ D.O. Litvintsev[†],⁷⁶ C. Liu[†],¹¹⁶ H. Liu[†],¹²⁶ H. Liu[†],¹²⁰ Q. Liu[†],⁸⁵ T. Liu[†],⁷⁶ Y. Liu[†],⁵
 A. Lobodenko[†],⁵² S. Lockwitz[†],⁷² A. Loginov[†],⁷² M. Lokajicek[†],¹⁰ R. Lopes de Sa[†],¹⁰⁷ H.J. Lubatti[†],¹²⁴
 D. Lucchesi[†],³⁵ J. Lueck[†],²⁴ P. Lujan[†],⁶⁸ P. Lukens[†],⁷⁶ R. Luna-Garcia[†],⁴⁵ G. Lungu[†],¹⁰⁵ A.L. Lyon[†],⁷⁶
 R. Lysak[†],⁵³ J. Lys[†],⁶⁸ A.K.A. Maciel[†],¹ R. Madar[†],¹⁸ R. Madrak[†],⁷⁶ K. Maeshima[†],⁷⁶ P. Maestro[†],³⁶
 R. Magaña-Villalba[†],⁴⁵ S. Malik[†],¹⁰⁵ S. Malik[†],¹⁰⁰ V.L. Malyshev[†],⁴⁸ G. Manca[†],⁶² A. Manousakis-Katsikakis[†],²⁸
 Y. Maravin[†],⁸⁸ F. Margaroli[†],³⁷ C. Marino[†],²⁴ M. Martínez[†],⁵⁴ J. Martínez-Ortega[†],⁴⁵ P. Mastrandrea[†],³⁷
 K. Matera[†],⁸¹ M.E. Mattson[†],⁹⁷ A. Mazzacane[†],⁷⁶ P. Mazzanti[†],³³ R. McCarthy[†],¹⁰⁷ K.S. McFarland[†],¹⁰⁶
 C.L. McGivern[†],⁶⁵ P. McIntyre[†],¹¹⁹ R. McNulty[†],⁶² A. Mehta[†],⁶² P. Mehtala[†],¹² M.M. Meijer[†],^{46,47}
 A. Melnitchouk[†],⁹⁹ D. Menezes[†],⁷⁹ P.G. Mercadante[†],³ M. Merkin[†],⁵⁰ A. Meyer[†],²¹ J. Meyer[†],²³ T. Miao[†],⁷⁶
 F. Miconi[†],¹⁹ D. Mietlicki[†],⁹⁶ A. Mitra[†],⁶ H. Miyake[†],⁴² S. Moed[†],⁷⁶ N. Moggi[†],³³ N.K. Mondal[†],³¹
 M.N. Mondragon[†],⁷⁶ C.S. Moon[†],⁴³ R. Moore[†],⁷⁶ M.J. Morello[†],³⁶ J. Morlock[†],²⁴ P. Movilla Fernandez[†],⁷⁶
 A. Mukherjee[†],⁷⁶ M. Mulhearn[†],¹²³ Th. Muller[†],²⁴ P. Murat[†],⁷⁶ M. Mussini[†],³³ J. Nachtman[†],⁷⁶ Y. Nagai[†],⁴²
 J. Naganoma[†],⁴¹ E. Nagy[†],¹⁵ M. Naimuddin[†],³⁰ I. Nakano[†],³⁹ A. Napier[†],⁹⁵ M. Narain[†],¹¹⁷ R. Nayyar[†],⁶⁷
 H.A. Neal[†],⁹⁶ J.P. Negret[†],⁷ J. Nett[†],¹¹⁹ M.S. Neubauer[†],⁸¹ C. Neu[†],¹²⁶ P. Neustroev[†],⁵² J. Nielsen[†],⁶⁸
 L. Nodulman[†],⁷⁵ S.Y. Noh[†],⁴³ O. Norriella[†],⁸¹ T. Nunnemann[†],²⁶ L. Oakes[†],⁶⁶ S.H. Oh[†],¹⁰⁹ Y.D. Oh[†],⁴³
 I. Oksuzian[†],¹²⁶ T. Okusawa[†],⁴⁰ R. Orava[†],¹² J. Orduna[†],¹²¹ L. Ortolan[†],⁵⁴ N. Osman[†],¹⁵ J. Osta[†],⁸⁴ M. Padilla[†],⁷¹
 S. Pagan Griso[†],³⁵ C. Pagliarone[†],³⁸ A. Pal[†],¹¹⁸ E. Palencia[†],⁵⁷ V. Papadimitriou[†],⁷⁶ A.A. Paramonov[†],⁷⁵
 N. Parashar[†],⁸³ V. Parihar[†],¹¹⁷ S.K. Park[†],⁴⁴ R. Partridge[†],¹¹⁷ N. Parua[†],⁸² J. Patrick[†],⁷⁶ A. Patwa[†],¹⁰⁸
 G. Pauletta[†],³⁸ M. Paulini[†],¹¹⁵ C. Paus[†],⁹⁴ D.E. Pellett[†],⁶⁹ B. Penning[†],⁷⁶ A. Penzo[†],³⁸ M. Perfilov[†],⁵⁰
 Y. Peters[†],⁶⁵ K. Petridis[†],⁶⁵ G. Petrillo[†],¹⁰⁶ P. Pétroff[†],¹⁶ T.J. Phillips[†],¹⁰⁹ G. Piacentino[†],³⁶ E. Pianori[†],¹¹⁴
 J. Pilot[†],¹¹⁰ K. Pitts[†],⁸¹ C. Plager[†],⁷⁰ M.-A. Pleier[†],¹⁰⁸ P.L.M. Podesta-Lerma[†],⁴⁵ V.M. Podstavkov[†],⁷⁶
 L. Pondrom[†],¹²⁵ A.V. Popov[†],⁵¹ S. Poprocki[†],⁷⁶ K. Potamianos[†],⁸⁵ A. Pranko[†],⁶⁸ M. Prewitt[†],¹²¹ D. Price[†],⁸²
 N. Prokopenko[†],⁵¹ F. Prokoshin[†],⁴⁸ F. Ptohos[†],³⁴ G. Punzi[†],³⁶ J. Qian[†],⁹⁶ A. Quadri[†],²³ B. Quinn[†],⁹⁹
 A. Rahaman[†],¹¹⁶ V. Ramakrishnan[†],¹²⁵ M.S. Rangel[†],¹ K. Ranjan[†],³⁰ N. Ranjan[†],⁸⁵ P.N. Ratoff[†],⁶¹ I. Razumov[†],⁵¹
 I. Redondo[†],⁵⁶ P. Renkel[†],¹²⁰ P. Renton[†],⁶⁶ M. Rescigno[†],³⁷ T. Riddick[†],⁶⁴ F. Rimondi[†],³³ I. Ripp-Baudot[†],¹⁹
 L. Ristori[†],^{116,76} F. Rizatdinova[†],¹¹³ A. Robson[†],⁶⁰ T. Rodrigo[†],⁵⁷ T. Rodriguez[†],¹¹⁴ E. Rogers[†],⁸¹ S. Rolli[†],⁹⁵
 M. Rominsky[†],⁷⁶ R. Roser[†],⁷⁶ A. Ross[†],⁶¹ C. Royon[†],¹⁸ P. Rubinov[†],⁷⁶ R. Ruchti[†],⁸⁴ F. Ruffini[†],³⁶ A. Ruiz[†],⁵⁷
 J. Russ[†],¹¹⁵ V. Rusu[†],⁷⁶ A. Safonov[†],¹¹⁹ G. Sajot[†],¹⁴ W.K. Sakumoto[†],¹⁰⁶ Y. Sakurai[†],⁴¹ P. Salcido[†],⁷⁹
 A. Sánchez-Hernández[†],⁴⁵ M.P. Sanders[†],²⁶ L. Santi[†],³⁸ A.S. Santos[†],¹ K. Sato[†],⁴² G. Savage[†],⁷⁶ V. Saveliev[†],⁷⁶
 A. Savoy-Navarro[†],⁷⁶ L. Sawyer[†],⁸⁹ T. Scanlon[†],⁶³ R.D. Schamberger[†],¹⁰⁷ Y. Scheglov[†],⁵² H. Schellman[†],⁸⁰
 P. Schlabach[†],⁷⁶ S. Schlobohm[†],¹²⁴ A. Schmidt[†],²⁴ E.E. Schmidt[†],⁷⁶ C. Schwanenberger[†],⁶⁵ T. Schwarz[†],⁷⁶

R. Schwienhorst^{‡,98} L. Scodellaro^{†,57} A. Scribano^{†#d,36} F. Scuri^{†,36} S. Seidel^{†,103} Y. Seiya^{†,40} J. Sekaric^{‡,87}
 A. Semenov^{†,48} H. Severini^{‡,112} F. Sforza^{†#d,36} E. Shabalina^{‡,23} S.Z. Shalhout^{†,69} V. Shary^{‡,18} S. Shaw^{‡,98}
 A.A. Shchukin^{‡,51} T. Shears^{†,62} P.F. Shepard^{†,116} M. Shimojima^{†aa,42} R.K. Shivpuri^{†,30} M. Shochet^{†,77}
 I. Shreyber-Tecker^{†,127} V. Simak^{‡,9} A. Simonenko^{†,48} P. Sinervo^{†,4} P. Skubic^{‡,112} P. Slattery^{‡,106} K. Sliwa^{†,95}
 D. Smirnov^{‡,84} J.R. Smith^{†,69} K.J. Smith^{‡,104} F.D. Snider^{†,76} G.R. Snow^{†,100} J. Snow^{†,111} S. Snyder^{†,108}
 A. Soha^{†,76} S. Söldner-Rembold^{‡,65} H. Song^{†,116} L. Sonnenschein^{‡,21} V. Sorin^{†,54} K. Soustruznik^{‡,8}
 P. Squillacioti^{†#d,36} R. St. Denis^{†,60} M. Stancari^{†,76} J. Stark^{‡,14} O. Stelzer-Chilton^{†,4} B. Stelzer^{†,4} D. Stentz^{†b,76}
 D.A. Stoyanova^{‡,51} M. Strauss^{‡,112} J. Strologas^{†,103} G.L. Strycker^{†,96} Y. Sudo^{†,42} A. Sukhanov^{†,76} I. Suslov^{†,48}
 L. Suter^{‡,65} P. Svoisky^{‡,112} M. Takahashi^{‡,65} K. Takemasa^{†,42} Y. Takeuchi^{†,42} J. Tang^{†,77} M. Tecchio^{†,96}
 P.K. Teng^{†,6} J. Thom^{†j,76} J. Thome^{†,115} G.A. Thompson^{†,81} E. Thomson^{†,114} M. Titov^{‡,18} D. Toback^{†,119}
 S. Tokar^{†,53} V.V. Tokmenin^{†,48} K. Tollefson^{†,98} T. Tomura^{†,42} D. Tonelli^{†,76} S. Torre^{†,34} D. Torretta^{†,76}
 P. Totaro^{†,35} M. Trovato^{†#e,36} Y.-T. Tsai^{†,106} K. Tschann-Grimm^{‡,107} D. Tsybychev^{‡,107} B. Tuchming^{‡,18}
 C. Tully^{‡,102} F. Ukegawa^{†,42} S. Uozumi^{†,43} L. Uvarov^{‡,52} S. Uvarov^{‡,52} S. Uzunyan^{†,79} R. Van Kooten^{‡,82}
 W.M. van Leeuwen^{‡,46} N. Varelas^{‡,78} A. Varganov^{†,96} E.W. Varnes^{‡,67} I.A. Vasilyev^{‡,51} F. Vázquez^{†c,73} G. Velev^{†,76}
 C. Vellidis^{†,76} P. Verdier^{‡,20} A.Y. Verkheev^{†,48} L.S. Vertogradov^{†,48} M. Verzocchi^{‡,76} M. Vesterinen^{‡,65} M. Vidal^{†,85}
 I. Vila^{†,57} D. Vilanova^{‡,18} R. Vilar^{†,57} J. Vizán^{†,57} M. Vogel^{†,103} P. Vokac^{‡,9} G. Volpi^{†,34} P. Wagner^{†,114}
 R.L. Wagner^{†,76} H.D. Wahl^{†,74} T. Wakisaka^{†,40} R. Wallny^{†,70} S.M. Wang^{†,6} M.H.L.S. Wang^{†,76} A. Warburton^{†,4}
 J. Warchol^{‡,84} D. Waters^{†,64} G. Watts^{‡,124} M. Wayne^{‡,84} J. Weichert^{‡,25} L. Welty-Rieger^{‡,80} W.C. Wester III^{†,76}
 A. White^{‡,118} D. Whiteson^{†bb,114} F. Wick^{†,24} D. Wicke^{‡,27} A.B. Wicklund^{†,75} E. Wicklund^{†,76} S. Wilbur^{†,77}
 H.H. Williams^{†,114} M.R.J. Williams^{‡,61} G.W. Wilson^{‡,87} J.S. Wilson^{†,110} P. Wilson^{†,76} B.L. Winer^{†,110}
 P. Wittich^{†j,76} M. Wobisch^{‡,89} S. Wolbers^{†,76} H. Wolfe^{†,110} D.R. Wood^{‡,92} T. Wright^{†,96} X. Wu^{†,59} Z. Wu^{†,122}
 T.R. Wyatt^{‡,65} Y. Xie^{‡,76} R. Yamada^{‡,76} K. Yamamoto^{†,40} D. Yamato^{†,40} S. Yang^{‡,5} T. Yang^{†,76} U.K. Yang^{†cc,77}
 W.-C. Yang^{†,65} Y.C. Yang^{†,43} W.-M. Yao^{†,68} T. Yasuda^{‡,76} Y.A. Yatsunenkov^{†,48} W. Ye^{‡,107} Z. Ye^{‡,76} G.P. Yeh^{†,76}
 K. Yi^{†u,76} H. Yin^{†,76} K. Yip^{†,108} J. Yoh^{†,76} K. Yorita^{†,41} T. Yoshida^{†dd,40} S.W. Youn^{†,76} G.B. Yu^{†,109} I. Yu^{†,43}
 J.M. Yu^{‡,96} S.S. Yu^{†,76} J.C. Yun^{†,76} A. Zanetti^{†,38} Y. Zeng^{†,109} J. Zennaro^{‡,104} T. Zhao^{†,124} T.G. Zhao^{‡,65}
 B. Zhou^{‡,96} C. Zhou^{†,109} J. Zhu^{‡,96} M. Zielinski^{‡,106} D. Zieminska^{‡,82} L. Zivkovic^{‡,117} and S. Zucchelli^{†#a33}

(CDF[†] and D0[‡] Collaborations)

¹LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

²Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

³Universidade Federal do ABC, Santo André, Brazil

⁴Institute of Particle Physics: McGill University, Montréal, Québec,

Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia,

Canada V5A 1S6; University of Toronto, Toronto, Ontario,

Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3

⁵University of Science and Technology of China, Hefei, People's Republic of China

⁶Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China

⁷Universidad de los Andes, Bogotá, Colombia

⁸Charles University, Faculty of Mathematics and Physics,

Center for Particle Physics, Prague, Czech Republic

⁹Czech Technical University in Prague, Prague, Czech Republic

¹⁰Center for Particle Physics, Institute of Physics,

Academy of Sciences of the Czech Republic, Prague, Czech Republic

¹¹Universidad San Francisco de Quito, Quito, Ecuador

¹²Division of High Energy Physics, Department of Physics,

University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland

¹³LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France

¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3,

Institut National Polytechnique de Grenoble, Grenoble, France

¹⁵CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

¹⁶LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

¹⁷LPNHE, Universités Paris VI and VII, CNRS/IN2P3, Paris, France

¹⁸CEA, Irfu, SPP, Saclay, France

¹⁹IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France

²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France

²¹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany

²²Physikalisches Institut, Universität Freiburg, Freiburg, Germany

²³II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany

- ²⁴*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*
- ²⁵*Institut für Physik, Universität Mainz, Mainz, Germany*
- ²⁶*Ludwig-Maximilians-Universität München, München, Germany*
- ²⁷*Fachbereich Physik, Bergische Universität Wuppertal, Wuppertal, Germany*
- ²⁸*University of Athens, 157 71 Athens, Greece*
- ²⁹*Panjab University, Chandigarh, India*
- ³⁰*Delhi University, Delhi, India*
- ³¹*Tata Institute of Fundamental Research, Mumbai, India*
- ³²*University College Dublin, Dublin, Ireland*
- ³³*Istituto Nazionale di Fisica Nucleare Bologna, ^{3a}University of Bologna, I-40127 Bologna, Italy*
- ³⁴*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*
- ³⁵*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, ^{3b}University of Padova, I-35131 Padova, Italy*
- ³⁶*Istituto Nazionale di Fisica Nucleare Pisa, ^{3c}University of Pisa,*
- ³⁷*University of Siena and ^{3d}Scuola Normale Superiore, I-56127 Pisa, Italy*
- ³⁸*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1,*
- ³⁹*Sapienza Università di Roma, I-00185 Roma, Italy*
- ⁴⁰*Istituto Nazionale di Fisica Nucleare Trieste/Udine,*
- ⁴¹*I-34100 Trieste, ⁴⁹University of Udine, I-33100 Udine, Italy*
- ⁴²*Okayama University, Okayama 700-8530, Japan*
- ⁴³*Osaka City University, Osaka 588, Japan*
- ⁴⁴*Waseda University, Tokyo 169, Japan*
- ⁴⁵*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- ⁴⁶*Center for High Energy Physics: Kyungpook National University,*
- ⁴⁷*Daegu 702-701, Korea; Seoul National University, Seoul 151-742,*
- ⁴⁸*Korea; Sungkyunkwan University, Suwon 440-746,*
- ⁴⁹*Korea; Institute of Science and Technology Information,*
- ⁵⁰*Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757,*
- ⁵¹*Korea; Chonbuk National University, Jeonju 561-756, Korea*
- ⁵²*Korea Detector Laboratory, Korea University, Seoul, Korea*
- ⁵³*CINVESTAV, Mexico City, Mexico*
- ⁵⁴*Nikhef, Science Park, Amsterdam, the Netherlands*
- ⁵⁵*Radboud University Nijmegen, Nijmegen, the Netherlands*
- ⁵⁶*Joint Institute for Nuclear Research, Dubna, Russia*
- ⁵⁷*Institute for Theoretical and Experimental Physics, Moscow, Russia*
- ⁵⁸*Moscow State University, Moscow, Russia*
- ⁵⁹*Institute for High Energy Physics, Protvino, Russia*
- ⁶⁰*Petersburg Nuclear Physics Institute, St. Petersburg, Russia*
- ⁶¹*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*
- ⁶²*Institut de Física d'Altes Energies, ICREA, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*
- ⁶³*Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Física d'Altes Energies (IFAE), Barcelona, Spain*
- ⁶⁴*Centro de Investigaciones Energeticas Medioambientales y Tecnológicas, E-28040 Madrid, Spain*
- ⁶⁵*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*
- ⁶⁶*Uppsala University, Uppsala, Sweden*
- ⁶⁷*University of Geneva, CH-1211 Geneva 4, Switzerland*
- ⁶⁸*Glasgow University, Glasgow G12 8QQ, United Kingdom*
- ⁶⁹*Lancaster University, Lancaster LA1 4YB, United Kingdom*
- ⁷⁰*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ⁷¹*Imperial College London, London SW7 2AZ, United Kingdom*
- ⁷²*University College London, London WC1E 6BT, United Kingdom*
- ⁷³*The University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁷⁴*University of Oxford, Oxford OX1 3RH, United Kingdom*
- ⁷⁵*University of Arizona, Tucson, Arizona 85721, USA*
- ⁷⁶*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*
- ⁷⁷*University of California, Davis, Davis, California 95616, USA*
- ⁷⁸*University of California, Los Angeles, Los Angeles, California 90024, USA*
- ⁷⁹*University of California Riverside, Riverside, California 92521, USA*
- ⁸⁰*Yale University, New Haven, Connecticut 06520, USA*
- ⁸¹*University of Florida, Gainesville, Florida 32611, USA*
- ⁸²*Florida State University, Tallahassee, Florida 32306, USA*
- ⁸³*Argonne National Laboratory, Argonne, Illinois 60439, USA*
- ⁸⁴*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*
- ⁸⁵*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*
- ⁸⁶*University of Illinois at Chicago, Chicago, Illinois 60607, USA*
- ⁸⁷*Northern Illinois University, DeKalb, Illinois 60115, USA*

- ⁸⁰Northwestern University, Evanston, Illinois 60208, USA
⁸¹University of Illinois, Urbana, Illinois 61801, USA
⁸²Indiana University, Bloomington, Indiana 47405, USA
⁸³Purdue University Calumet, Hammond, Indiana 46323, USA
⁸⁴University of Notre Dame, Notre Dame, Indiana 46556, USA
⁸⁵Purdue University, West Lafayette, Indiana 47907, USA
⁸⁶Iowa State University, Ames, Iowa 50011, USA
⁸⁷University of Kansas, Lawrence, Kansas 66045, USA
⁸⁸Kansas State University, Manhattan, Kansas 66506, USA
⁸⁹Louisiana Tech University, Ruston, Louisiana 71272, USA
⁹⁰The Johns Hopkins University, Baltimore, Maryland 21218, USA
⁹¹Boston University, Boston, Massachusetts 02215, USA
⁹²Northeastern University, Boston, Massachusetts 02115, USA
⁹³Harvard University, Cambridge, Massachusetts 02138, USA
⁹⁴Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
⁹⁵Tufts University, Medford, Massachusetts 02155, USA
⁹⁶University of Michigan, Ann Arbor, Michigan 48109, USA
⁹⁷Wayne State University, Detroit, Michigan 48201, USA
⁹⁸Michigan State University, East Lansing, Michigan 48824, USA
⁹⁹University of Mississippi, University, Mississippi 38677, USA
¹⁰⁰University of Nebraska, Lincoln, Nebraska 68588, USA
¹⁰¹Rutgers University, Piscataway, New Jersey 08855, USA
¹⁰²Princeton University, Princeton, New Jersey 08544, USA
¹⁰³University of New Mexico, Albuquerque, New Mexico 87131, USA
¹⁰⁴State University of New York, Buffalo, New York 14260, USA
¹⁰⁵The Rockefeller University, New York, New York 10065, USA
¹⁰⁶University of Rochester, Rochester, New York 14627, USA
¹⁰⁷State University of New York, Stony Brook, New York 11794, USA
¹⁰⁸Brookhaven National Laboratory, Upton, New York 11973, USA
¹⁰⁹Duke University, Durham, North Carolina 27708, USA
¹¹⁰The Ohio State University, Columbus, Ohio 43210, USA
¹¹¹Langston University, Langston, Oklahoma 73050, USA
¹¹²University of Oklahoma, Norman, Oklahoma 73019, USA
¹¹³Oklahoma State University, Stillwater, Oklahoma 74078, USA
¹¹⁴University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
¹¹⁵Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA
¹¹⁶University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
¹¹⁷Brown University, Providence, Rhode Island 02912, USA
¹¹⁸University of Texas, Arlington, Texas 76019, USA
¹¹⁹Texas A&M University, College Station, Texas 77843, USA
¹²⁰Southern Methodist University, Dallas, Texas 75275, USA
¹²¹Rice University, Houston, Texas 77005, USA
¹²²Baylor University, Waco, Texas 76798, USA
¹²³University of Virginia, Charlottesville, Virginia 22904, USA
¹²⁴University of Washington, Seattle, Washington 98195, USA
¹²⁵University of Wisconsin, Madison, Wisconsin 53706, USA
¹²⁶University of Virginia, Charlottesville, Virginia 22906, USA
¹²⁷Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- (Dated: July 12, 2012)

The combination of searches performed by the CDF and D0 collaborations at the Fermilab Tevatron Collider for neutral Higgs bosons produced in association with b quarks is reported. The data, corresponding to 2.6 fb^{-1} of integrated luminosity at CDF and 5.2 fb^{-1} at D0, have been collected in final states containing three or more b jets. Upper limits are set on the cross section multiplied by the branching ratio varying between 44 pb and 0.7 pb in the Higgs boson mass range 90 to 300 GeV, assuming production of a narrow scalar boson. Significant enhancements to the production of Higgs bosons can be found in theories beyond the standard model, for example in supersymmetry. The results are interpreted as upper limits in the parameter space of the minimal supersymmetric standard model in a benchmark scenario favoring this decay mode.

PACS numbers: 14.80.Da, 12.38.Qk, 12.60.Fr, 13.85.Rm

*Deceased

†With CDF visitors from †^aUniversidad de Oviedo, E-33007

In the standard model (SM), electroweak symmetry breaking is achieved through the introduction of a single scalar doublet field and the existence of a single neutral scalar boson [1–6] is predicted. However, extensions of the SM exist with richer structure. Introducing a second Higgs doublet, such as in Type II 2-Higgs Doublet Models (2HDM) [7], leads to multiple scalar bosons and can give scenarios with enhanced couplings to down-type fermions. Supersymmetry is a plausible extension to the SM that introduces an additional symmetry between fermions and bosons. The two Higgs boson doublets in the minimal supersymmetric standard model (MSSM) [8, 9] lead to five physical Higgs bosons: three neutral (collectively denoted as ϕ): h , H , and A ; and two charged: H^+ and H^- . At leading order the MSSM is a Type II 2HDM model, and two parameters are sufficient to describe the Higgs sector. These are conventionally chosen as the ratio of the two Higgs doublet vacuum expectation values, $\tan\beta$, and, M_A , the mass of the pseudoscalar boson, A . The couplings to the down-type fermions are enhanced by a factor of $\tan\beta$ relative to those in the SM. Thus, the main decay mode is $\phi \rightarrow \bar{b}$, with branching fractions near 90% at leading order (the remainder being mostly $\phi \rightarrow \tau^+\tau^-$). While searches

for τ -lepton-pair final states at hadron colliders are relatively insensitive to higher-order radiative corrections, due to cancellations between the production and decay processes, this is not the case for decays to \bar{b} . Thus, information from decays to $b\bar{b}$ together with stringent limits from decays to $\tau^+\tau^-$ can constrain higher-order effects and yield additional information about electroweak symmetry breaking, supersymmetry or other new physics beyond the SM with similar final states such as pair production of color octet scalars [12–14].

Since an inclusive search for $\phi \rightarrow \bar{b}$ is difficult due to large multijet backgrounds, these searches rely on the case where the ϕ boson is produced in association with one or more b quarks. This final state with at least three b quarks represents a powerful search channel, with the third b jet providing additional suppression of the large multijet background at a hadron collider.

MSSM Higgs boson production has been studied at the CERN LEP e^+e^- collider excluding $M_{h,A} < 93$ GeV for all $\tan\beta$ values [15]. The CDF [16–18] and D0 [19–28] collaborations extended such searches to higher masses and for large $\tan\beta$. The most stringent upper limits on $\tan\beta$ in searches for neutral Higgs production for masses above the LEP bounds come from searches in final states with τ leptons pair produced at the Large Hadron Collider [29–31].

The results reported in this Article make use of $p\bar{p}$ collisions at a center of mass energy of 1.96 TeV. The data, corresponding to integrated luminosities of 2.6 fb^{-1} and 5.2 fb^{-1} , were collected during Run II at the Fermilab Tevatron Collider by the CDF and D0 collaborations, respectively. The combination of searches for neutral Higgs bosons in final states with three or more b jets [18, 25] is presented.

The CDF and D0 detectors are described in Ref. [32–35]. A brief outline of the reconstruction of the final states and event selection used in these searches follows. A cone algorithm [36] with a radius of $\mathcal{R} = \sqrt{(\Delta y)^2 + (\Delta\varphi)^2} = 0.7$ (CDF) and 0.5 (D0), where y is the rapidity and φ the azimuthal angle, is used to reconstruct jets from energy deposition in the calorimeters. Identification of jets arising from b -quark fragmentation (b -tagging) uses an algorithm based on reconstructing secondary vertices of charged particles displaced from the $p\bar{p}$ interaction vertex at CDF [37] and a neural network combining lifetime information from secondary vertices and the minimum distance of approach of charged particle trajectories to the primary interaction at D0 [38].

Events are selected at CDF with dedicated online event selections (triggers) requiring at least two jets and using b -tagging information, while at least three jets are required at D0. For most of the data collected at D0, b -tagging information is also used in the trigger decision. Events are selected offline with at least three jets within the fiducial region. This is defined by requirements on the momentum transverse to the proton beam direction,

Oviedo, Spain, ^{†b}Northwestern University, Evanston, IL 60208, USA, ^{†c}Universidad Iberoamericana, Mexico D.F., Mexico, ^{†d}ETH, 8092 Zurich, Switzerland, ^{†e}CERN, CH-1211 Geneva, Switzerland, ^{†f}Queen Mary, University of London, London, E1 4NS, United Kingdom, ^{†g}National Research Nuclear University, Moscow, Russia, ^{†h}Yarmouk University, Irbid 211-63, Jordan, ^{†i}Muons, Inc., Batavia, IL 60510, USA, ^{†j}Cornell University, Ithaca, NY 14853, USA, ^{†k}Kansas State University, Manhattan, KS 66506, USA, ^{†l}Kinki University, Higashi-Osaka City, Japan 577-8502, ^{†m}University of CA Santa Barbara, Santa Barbara, CA 93106, USA, ^{†n}University of Notre Dame, Notre Dame, IN 46556, USA, ^{†o}Ewha Womans University, Seoul, 120-750, Korea, ^{†p}Texas Tech University, Lubbock, TX 79609, USA, ^{†q}University of Melbourne, Victoria 3010, Australia, ^{†r}Institute of Physics, Academy of Sciences of the Czech Republic, Czech Republic, ^{†s}Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, ^{†t}University College Dublin, Dublin 4, Ireland, ^{†u}University of Iowa, Iowa City, IA 52242, USA, ^{†v}University of CA Santa Cruz, Santa Cruz, CA 95064, USA, ^{†w}Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile, ^{†x}University of Cyprus, Nicosia CY-1678, Cyprus, ^{†y}Office of Science, U.S. Department of Energy, Washington, DC 20585, USA, ^{†z}CNRS-IN2P3, Paris, F-75205 France, ^{†aa}Nagasaki Institute of Applied Science, Nagasaki, Japan, ^{†bb}University of CA Irvine, Irvine, CA 92697, USA, ^{†cc}University of Manchester, Manchester M13 9PL, United Kingdom, and ^{†dd}University of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017.

[‡]and D0 visitors from ^{†a}Augustana College, Sioux Falls, SD, USA, ^{†b}The University of Liverpool, Liverpool, UK, ^{†c}UPIITA-IPN, Mexico City, Mexico, ^{†d}DESY, Hamburg, Germany, ^{†e}University College London, London, UK, ^{†f}Centro de Investigacion en Computacion - IPN, Mexico City, Mexico, ^{†g}SLAC, Menlo Park, CA, USA, ^{†h}ECFM, Universidad Autonoma de Sinaloa, Culiacán, Mexico and ^{†i}Universidade Estadual Paulista, São Paulo, Brazil.

p_T , and detector pseudorapidity, $\eta_{\text{det}} = -\ln(\tan(\theta/2))$, of jets, where θ is the polar angle with respect to the proton beam direction and an origin at the center of the detector: three jets with $p_T > 20$ GeV and $|\eta_{\text{det}}| < 2.0$ at CDF and three jets with $p_T > 20$ GeV and at least two jets with $p_T > 25$ GeV, $|\eta_{\text{det}}| < 2.5$ at D0. Events with more jets are accepted at CDF but only the leading (in p_T) four jets are used in the analysis. Exclusive channels with exactly three or exactly four jets are used at D0. The signal sample is defined by both experiments requiring at least three b -tagged jets. In the CDF inclusive and the D0 4-jet channel the two leading jets and at least one of the third and fourth jets are required to be b tagged. In addition, a large sample requiring only two b -tagged jets is used by both experiments to build models of the multijet backgrounds.

At CDF, the background is modeled using a sum of templates of the invariant mass distribution of the leading two jets, representing contributions from different background modes categorized according to kinematics and flavor content. The templates are constructed from events in the double-tagged data sample where at least one of the leading two jets is b -tagged. The events are then weighted according to the probability for at least one of the jets with no b tag to pass the tagging requirements under three different assumptions as to whether it arises from a b quark, a c quark or a light quark (or gluon). This results in six different mass templates. The separation between the different components is enhanced by using an additional jet-flavor-sensitive discriminant, x_{tags} , that makes use of kinematic properties of the charged particles from secondary vertices associated with each b -tagged jet.

In the D0 analysis the background model is formed by correcting the shape of the dijet invariant mass distribution of a data sample with two b -tagged jets using the ratio of simulated multijet samples with three b tags and two b tags. The simulated samples are generated using ALPGEN [39] with showering and hadronization carried out using PYTHIA [40] and detailed simulation of the detector using GEANT [41]. Their flavor composition, in terms of the relative numbers of b , c and light jets per event, is determined from a simultaneous fit to the data across samples with differing numbers of tagged jets, different b -tagger operating points, and in small intervals of the scalar sum of the transverse momenta of the jets. The shape correction is applied as a function of the dijet invariant mass and the value of a likelihood-ratio discriminant, \mathcal{L} , designed to select signal-like events in preference to background-like events. Only the two possible jet pairings from the leading jet plus either the second or third leading jet are considered when forming Higgs candidates, and the choice that gives the highest value of $\mathcal{L} > 0.65$ is selected. If neither pairing in the event is above this threshold then the event is discarded. Systematic uncertainties are assessed to take into account

the imperfect modeling of the background simulation. However, the definition of the correction as a ratio of distributions significantly reduces the sensitivity of the final model to these imperfections. Small contributions to the background from top-quark-pair production are simulated using ALPGEN and PYTHIA. Backgrounds from other sources, such as $Z \rightarrow b\bar{b}$ and single-top-quark production are negligible.

The efficiency for selecting signal is estimated for both analyses using events generated with PYTHIA. Associated production of a ϕ boson and a b quark, $gb \rightarrow \phi b$, with subsequent decay $\phi \rightarrow b\bar{b}$, is used to model the signal. The signal cross section, experimental acceptance and the kinematics are corrected to next-to-leading order (NLO) using MCFM [42, 43], weighting events as a function of the kinematics of the leading b -quark jet not associated with the Higgs decay.

Approximately 11 500 events are selected by CDF with at least three b tags and an estimated product of signal efficiency and acceptance varying as a function of M_ϕ between 0.18% at $M_\phi = 90$ GeV and 0.8% at $M_\phi = 200$ GeV. Approximately 15 000 and 11 000 events with at least three b -tagged jets are accepted in the D0 three-jet and four-jet channels, respectively. The corresponding signal efficiencies for a Higgs of mass $M_\phi = 200$ GeV are 1.2% and 0.6%.

The channels are combined and exclusion limits set, using the modified frequentist technique [44, 45], with a log likelihood ratio (LLR) as test statistic:

$$\text{LLR} = -2 \ln \frac{p(X|H_1)}{p(X|H_0)}, \quad (1)$$

where p represents the probabilities that the data, X , are drawn from the background-only (H_0), and signal-plus-background (H_1) hypotheses, respectively. The likelihoods are constructed from the binned two-dimensional distribution of the dijet invariant mass versus the x_{tags} discriminant for CDF and the binned one-dimensional dijet invariant mass distribution for D0. Theoretical predictions of the absolute rates for the multijet backgrounds have large uncertainties. Therefore, additional scale factors are applied to the background yield in the D0 analysis and the individual templates in the CDF analysis and introduced into the likelihood as parameters with no external constraints. Systematic uncertainties are introduced as either shape or normalization variations to the model probability density functions. These systematics are governed by nuisance parameters together with Gaussian constraint terms where appropriate [46]. Sources of uncertainty related to a common component of the luminosity determination and the theoretical modeling of the signal production are considered fully correlated. All other sources of uncertainty are assumed to be uncorrelated. The modeling of the b -jet-tagging efficiency and the contamination from light-quark and gluon jets

fake are the dominant sources of uncertainty on the background model. These are implemented as uncertainties that affect the shape of the distributions entering the likelihood. Additional sub-dominant uncertainties are considered in the D0 analysis arising in the modeling of the trigger, jet efficiency, jet energy scale and jet resolution. While most of those effects have a negligible impact on the background model, effects that are dependent on the differences between b , light, and gluon jets can be significant. Dominant experimental systematic uncertainties on the signal model can be attributed to luminosity (6%), b -tagging efficiency (11-18%) and jet energy scale (2-10% depending on the ϕ boson mass hypothesis).

Limits on the product of the cross section and the branching ratio using the LLR test statistic are extracted. The limits are model-independent, apart from assuming a single narrow Higgs boson mass peak, dominated by experimental resolution effects. These are summarized in Table I, and presented in Fig. 1. The combination gives a sensitivity that is better than the D0 expected limit alone by $\approx 25\%$ at $M_\phi = 100$ GeV, steadily falling to $< 1\%$ by $M_\phi = 300$ GeV. Excesses of events above the SM background expectation are observed for $M_\phi = 120$ and 140 GeV with significances of 2.5 standard deviations and 2.6 standard deviations, respectively. These are driven by the excesses observed in the individual contributing analyses of 2.8 standard deviations at $M_\phi = 150$ GeV at CDF and 2.5 standard deviations at $M_\phi = 120$ GeV at D0. A standard convention [47] is used to account for the effect that it is more likely to find a deviation (under the background-only) hypothesis when several mass regions are probed compared with only a single hypothesis. The significance of the excesses in the combined analysis is reduced to ≈ 2 standard deviations.

Though these limits are the key results of this search, it is interesting to interpret them in terms of constraints on benchmark models within the MSSM. As a consequence of the enhanced couplings to b quarks at large $\tan\beta$, the total width of the Higgs boson increases with $\tan\beta$. When this width becomes comparable to the experimental resolution of 15-20% there is an impact on the sensitivity of the search. When interpreting the exclusion within the MSSM, the width of the Higgs boson and the enhancement of the product of cross section and branching ratio above that of the SM are calculated using FEYNHIGGS [48–53]. The width is included in the simulation of the signal as a function of mass and $\tan\beta$ by convoluting a relativistic Breit-Wigner function with the NLO cross section [42]. Additional uncertainties for these model-specific limits are considered that otherwise cancel in the model-independent limit. For comparison these are derived as in previous results [18, 22, 25]: uncertainties on the SM signal cross section are derived from varying the factorization and renormalization scales by a factor of two and from uncertainties on the parton distribution functions. The uncertainties assessed from scale

M_ϕ (GeV)	Obs. (pb)	Expected (pb)				
		-2 s.d.	-1 s.d.	median	+1 s.d.	+2 s.d.
90	38	30	41	57	81	110
100	43	22	30	42	62	86
110	43	15	20	27	39	53
120	44	12	15	20	29	39
130	25	8.1	9.6	14	19	25
140	26	5.1	7.3	10	14	20
150	18	4.1	5.5	7.4	11	15
160	12	3.3	4.5	6.0	8.5	12
170	9.4	2.5	3.6	5.0	7.0	9.5
180	7.1	2.4	3.2	4.2	6.0	8.2
190	5.7	2.2	2.5	4.0	5.0	7.0
200	5.0	2.0	2.4	3.0	4.4	6.0
210	3.8	1.6	2.2	2.6	3.7	5.0
220	3.3	1.2	1.7	2.2	3.2	4.7
230	2.5	1.0	1.5	2.0	3.0	4.1
240	2.0	0.9	1.2	1.8	2.5	3.5
250	2.0	0.9	1.1	1.6	2.3	3.2
260	1.7	0.7	1.0	1.4	2.2	2.8
270	1.3	0.7	0.9	1.2	2.0	2.4
280	1.1	0.6	0.8	1.1	1.7	2.4
290	0.82	0.6	0.8	1.1	1.9	2.4
300	0.71	0.5	0.7	1.0	1.6	2.3

TABLE I: Observed and expected upper limits at the 95% C.L. on the product of cross section and branching ratio $\sigma(gb \rightarrow \phi b) \times BR(\phi \rightarrow bb)$, within the acceptance for the highest p_T b quark not arising from the Higgs decay [43]. Expected limits are given for the median and for ± 1 and ± 2 standard deviation (s.d.) variations of the background expectation.




FIG. 1: Model independent 95% C.L. upper limits on the product of cross section and branching ratio for the combined analyses, assuming a mass degeneracy between two of the three neutral bosons and a Higgs boson width significantly smaller than the experimental resolution. The dark and light shaded regions (color online) correspond to the one and two standard deviation bands around the median expected limit.

variation are taken to be 10% and the parton distribution functions contribute an additional 3.5-13%, depending on the mass hypothesis.

The masses and couplings of the Higgs bosons in the MSSM depend, in addition to $\tan\beta$ and M_A , on other parameters through radiative corrections. Limits on $\tan\beta$ as a function of M_A are derived for the m_h^{\max} scenario [54] that favors the $b\bar{b}$ final state, assuming a CP-conserving Higgs sector [55] and a negative value of the Higgs sector bilinear coupling μ . Figure 2 shows exclusion limits in the $(\tan\beta, M_A)$ plane for this scenario. Adding a further potential theoretical uncertainty on the signal cross section of 20%, independent of M_{ϕ} , would lead to an increase of 5% in the $\tan\beta$ limit.




FIG. 2: 95% C.L. lower limit in the $(M_A, \tan\beta)$ plane for the m_h^{\max} , $\mu = -200$ GeV, including Higgs boson width effects. The exclusion limit obtained from the LEP experiments is also shown.

In summary, the combination of results on neutral Higgs boson searches in multi- b -jet events from CDF and D0 has been presented. Upper limits are set on the product of the cross section and branching ratio and constraints are placed in the $(\tan\beta, M_A)$ plane for a particular MSSM scenario. This combination, with more than half the integrated luminosity from the Tevatron still to be analyzed, provides the most stringent limits on neutral Higgs boson production and decay in the multi- b -jet mode.

We thank the Fermilab staff and technical staffs of the participating institutions for their vital contributions and acknowledge support from the DOE and NSF (USA), ARC (Australia), CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil), NSERC (Canada), NSC, CAS and CNSF (China), Colciencias (Colombia), MSMT and GACR (Czech Republic), the Academy of Finland, CEA and CNRS/IN2P3 (France), BMBF and DFG (Germany), DAE and DST (India), SFI (Ireland), INFN

(Italy), MEXT (Japan), the Korean World Class University Program and NRF (Korea), CONACyT (Mexico), FOM (Netherlands), MON, NRC KI and RFBR (Russia), the Slovak R&D Agency, the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010 (Spain), The Swedish Research Council (Sweden), SNSF (Switzerland), STFC and the Royal Society (United Kingdom), and the A.P Sloan Foundation (USA).

-
- [1] F. Englert and R. Brout, Phys. Rev. Lett. **13**, 321 (1964).
 - [2] P.W. Higgs, Phys. Lett. **12**, 132 (1964).
 - [3] P.W. Higgs, Phys. Rev. Lett. **13**, 508 (1964).
 - [4] G.S. Guralnik, C.R. Hagen, and T.W.B. Kibble, Phys. Rev. Lett. **13**, 585 (1964).
 - [5] P.W. Higgs, Phys. Rev. **145**, 1156 (1966).
 - [6] T.W.B. Kibble, Phys. Rev. **155**, 1554 (1967).
 - [7] V. Barger, J.L. Hewett, and R.J.N. Phillips, Phys. Rev. D **41**, 3421 (1990).
 - [8] H.P. Nilles, Phys. Rep. **110**, 1 (1984).
 - [9] H.E. Haber and G.L. Kane, Phys. Rep. **117**, 75 (1985).
 - [10] B. Ananthanarayan, G. Lazarides, and Q. Shafi, Phys. Rev. D **44**, 1613 (1991).
 - [11] V. Barger and C. Kao, Phys. Lett. B **518**, 117 (2001).
 - [12] B.A. Dobrescu, K. Kong, and R. Mahbubani, Phys. Lett. B **670**, 119 (2008).
 - [13] M. Gerbush, T.J. Khoo, D.J. Phalen, A. Pierce, and D. Tucker-Smith, Phys. Rev. D **77**, 095003 (2008).
 - [14] Y. Bai and B.A. Dobrescu, J. High Energy Phys. **07**, 100 (2011).
 - [15] The ALEPH Collaboration, The DELPHI Collaboration, The L3 Collaboration, and The OPAL Collaboration, Eur. Phys. J. C **47**, 547 (2006).
 - [16] T. Affolder *et al.* (CDF Collaboration), Phys. Rev. Lett. **86**, 4472 (2001).
 - [17] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **96**, 011802 (2006).
 - [18] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **85**, 032005 (2012).
 - [19] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **95**, 151801 (2005).
 - [20] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **97**, 121802 (2006).
 - [21] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **101**, 071804 (2008).
 - [22] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **101**, 221802 (2008).
 - [23] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **102**, 051804 (2009).
 - [24] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **104**, 151801 (2010).
 - [25] V.M. Abazov *et al.* (D0 Collaboration), Phys. Lett. B **698**, 97 (2011).
 - [26] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **107**, 121801 (2011).
 - [27] V.M. Abazov *et al.* (D0 Collaboration), Phys. Lett. B **707**, 323 (2012).
 - [28] V.M. Abazov *et al.* (D0 Collaboration), Phys. Lett. B **710**, 569 (2012).
 - [29] CMS Collaboration, Phys. Rev. Lett. **106**, 231801

- (2011).
- [30] ATLAS Collaboration, Phys. Lett. B **705**, 174 (2011).
- [31] CMS Collaboration, Phys. Lett. B **713**, 68 (2012).
- [32] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71** 032001 (2005).
- [33] V.M. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods Phys. Res. A **565**, 463 (2006).
- [34] M. Abolins *et al.* Nucl. Instrum. Methods Phys. Res. A **584**, 75 (2008).
- [35] R. Angstadt *et al.* Nucl. Instrum. Methods Phys. Res. A **622**, 298 (2010).
- [36] G. Blazey *et al.*, arXiv:hep-ex/0005012 (2000).
- [37] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 052003 (2005).
- [38] V.M. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods in Phys. Res. Sect. A **620**, 490 (2010).
- [39] M.L. Mangano *et al.*, J. High Energy Phys. 07, 001 (2003). ALPGEN version 2.11 is used.
- [40] T. Sjöstrand, S. Mrenna, and P. Skands, J. High Energy Phys. 05, 026 (2006). Version 6.409 is used.
- [41] R. Brun and F. Carminati, CERN program library long writeup W5013 (1993).
- [42] J. Campbell, R.K. Ellis, F. Maltoni, and S. Willenbrock, Phys. Rev. D **67**, 095002 (2003).
- [43] Cross sections are quoted with respect to the requirements that the highest- p_T b quark not from the Higgs decay has $p_T > 12$ GeV and $|\eta| < 5.0$.
- [44] T. Junk, Nucl. Instrum. Methods Phys. Res. A **434**, 435 (1999).
- [45] A. Read, Nucl. Instrum. Methods Phys. Res. A **425**, 357 (1999).
- [46] W. Fisher, FERMILAB-TM-2386-E (2007).
- [47] The ALEPH Collaboration, The DELPHI Collaboration, The L3 Collaboration, and The OPAL Collaboration, Phys. Lett. B **565**, 61 (2003).
- [48] S. Heinemeyer, W. Hollik, and G. Weiglein, Eur. Phys. J. C **9**, 343 (1999). FEYNHIGGS version 2.6.8 is used.
- [49] S. Heinemeyer, W. Hollik, and G. Weiglein, Comput. Phys. Commun. **124**, 76 (2000).
- [50] G. Degrossi, S. Heinemeyer, W. Hollik, P. Slavich, and G. Weiglein, Eur. Phys. J. C **28**, 133 (2003).
- [51] M. Frank, T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, J. High Energy Phys. 02, 047 (2007).
- [52] L. Hofer, U. Nierste, and D. Shere, J. High Energy Phys. 10, 081 (2009).
- [53] D. Noth and M. Spira, Phys. Rev. Lett. **101**, 181801 (2008).
- [54] $M_{\text{SUSY}} = 1$ TeV, $X_t = 2$ TeV, $M_2 = 0.2$ TeV $|\mu| = 0.2$ TeV, and $m_g = 0.8$ TeV.
- [55] M. Carena, S. Heinemeyer, C. E. M. Wagner, and G. Weiglein, Eur. Phys. J. C **45**, 797 (2006).