



Citation for published version:

Cybulski, MK, Riddlestone, IM, Mahon, MF, Woodman, TJ & Whittlesey, MK 2015, 'Stoichiometric and catalytic C-F bond activation by the trans-dihydride complex [Ru(IEt₂Me₂)₂(PPh₃)₂H₂] (IEt₂Me₂ = 1,3-diethyl-4,5-dimethylimidazol-2-ylidene)', Dalton Transactions , vol. 44, no. 45, pp. 19597-19605.
<https://doi.org/10.1039/C5DT01996F>

DOI:

[10.1039/C5DT01996F](https://doi.org/10.1039/C5DT01996F)

Publication date:

2015

Document Version

Early version, also known as pre-print

[Link to publication](#)

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Stoichiometric and Catalytic C-F Bond Activation by the *Trans*-
Dihydride NHC Complex [Ru(IEt₂Me₂)₂(PPh₃)₂H₂] (IEt₂Me₂ = 1,3-
diethyl-4,5-dimethylimidazol-2-ylidene)**

Mateusz K. Cybulski,^a Ian M. Riddlestone,^a Mary F. Mahon,^a Timothy J. Woodman^b and
Michael K. Whittlesey^{a*}

*Departments of^aChemistry and ^bPharmacy and Pharmacology, University of Bath,
Claverton Down, Bath BA2 7AY, UK*

Submitted for the themed issue of Dalton Trans. on 'Fluorine'

Abstract

The room temperature reaction of C_6F_6 or C_6F_5H with $[Ru(IEt_2Me_2)_2(PPh_3)_2H_2]$ (**1**; $IEt_2Me_2 = 1,3$ -diethyl-4,5-dimethylimidazol-2-ylidene) generated a mixture of the *trans*-hydride fluoride complex $[Ru(IEt_2Me_2)_2(PPh_3)_2HF]$ (**2**) and the bis-carbene pentafluorophenyl species $[Ru(IEt_2Me_2)_2(PPh_3)(C_6F_5)H]$ (**3**). The formation of **3** resulted from C-H activation of C_6F_5H (formed from C_6F_6 via stoichiometric hydrodefluorination), a process which could be reversed by working under 4 atm H_2 . Upon heating **1** with C_6F_5H , the bis-phosphine derivative $[Ru(IEt_2Me_2)(PPh_3)_2(C_6F_5)H]$ (**4**) was isolated. A more efficient route to **2** involved treatment of **1** with 0.33 eq of TREAT-HF ($Et_3N \cdot 3HF$); excess reagent gave instead the $[H_2F_3]^-$ salt (**5**) of the known cation $[Ru(IEt_2Me_2)_2(PPh_3)_2H]^+$. Under catalytic conditions, **1** proved to be an active precursor for hydrodefluorination, converting C_6F_6 to a mixture of tri, di and monofluorobenzenes (TON = 37) at 363 K with 10 mol% **1** and Et_3SiH as the reductant.

increasingly more nucleophilic Ru-H ligands. Very recently, we reported an example of such a species in the form of the mixed carbene-phosphine complex

$[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)_2\text{H}_2]$ (**1**; $\text{IEt}_2\text{Me}_2 = 1,3\text{-diethyl-4,5-dimethylimidazol-2-ylidene}$).¹⁴

The unusual *trans*-arrangement of the two hydride ligands imparts highly nucleophilic character to Ru-H, as evidenced by the formation of methane and


$[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)_2\text{HI}]$ upon addition of the electrophile MeI. We now report our initial findings on both the stoichiometric and catalytic reactivity of **1** towards aromatic fluorocarbons. As hoped for, the complex displays high activity for the catalytic HDF of C_6F_6 , undergoing up to five HDF steps in generating fluorobenzene.

Results and Discussion

Stoichiometric C-F and C-H activation of C_6F_6 and $\text{C}_6\text{F}_5\text{H}$ by **1**

Monitoring by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy the room temperature reaction of $[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)_2\text{H}_2]$ (**1**) with 10 eq of either C_6F_6 or $\text{C}_6\text{F}_5\text{H}$ in C_6H_6 solution¹⁵ showed, over the course of *ca.* 5 h, complete loss of starting material and the appearance of two new product peaks at δ 45 and 59. These were assigned to the hydride fluoride complex $[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)_2\text{HF}]$ (**2**) and the pentafluorophenyl complex $[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)(\text{C}_6\text{F}_5)\text{H}]$ (**3**) respectively (Scheme 2). The formation of the two products, which were present after 5 h in an approximate ratio (by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy) of 1:0.2 from C_6F_6 and 1:0.5 from $\text{C}_6\text{F}_5\text{H}$, arise from competing C-F and C-H activation respectively. C-H activation proved to be reversible. Thus, addition of H_2 (4 atm) to an in-situ generated mixture of **2**, **3** and PPh_3 led to the complete disappearance

of **3** over 4 h at room temperature. In a more controlled experiment, addition of 4 atm H₂ to a solution containing an isolated, crystalline sample of **3** (vide infra) and an equivalent of PPh₃ led to the complete conversion of the former to a mixture of **1** and **2** within 4 h at 298 K. Generation of the latter could be rationalised following analysis of the ¹H and ¹⁹F NMR spectra of the volatile materials from the C₆F₆ reaction. This revealed the presence of the hydrodefluorination products C₆F₅H (major species) and both 1,2,3,4- and 1,2,4,5-C₆F₄H₂, indicating that **1** must initially activate the C-F bond in C₆F₆ to give **2** and C₆F₅H, which then proved to be at least as reactive a substrate as C₆F₆, undergoing C-F activation to give the tetrafluorobenzene isomers (and additional **2**), as well as C-H activation to produce **3**. The competitive nature of C-H activation is clearly shown by the higher ratio of **3**:**2** formed in the reaction of **1** with C₆F₅H.¹⁶



Scheme 2

Both **2** and **3** could be isolated from the reaction mixture following removal of the volatile components and recrystallization of the residue. The X-ray structure of **2** (Figure 1) revealed retention of the *cis* arrangement of the two NHC ligands and two PPh₃ groups

from **1** and, as a result, very little change in either Ru-C/Ru-P distances and angles. Of particular interest was the long Ru-F distance of 2.264(2) Å. This is comparable to the value (2.284(5) Å) in [Ru(dmpe)₂H(FHF)] (dmpe = 1,2-bis(dimethylphosphino)ethane), which is the only other *trans* H-Ru-F complex we could find that has been structurally verified.¹⁷ Surprisingly, crystallographically characterised examples of Ru(L)₄H(halide) (L = PR₃ or NHC) species with *trans* H-Ru-halide geometries in general are not that common,¹⁸ despite complexes of this type being known for over 50 years.¹⁹ Elongation in the Ru-F distance in both **2** and [Ru(dmpe)₂H(FHF)] compared to those in *cis*-[Ru(dppp)₂F₂] (dppp = 1,4-bis(diphenylphosphino)ethane) and *trans*-[Ru(dppe)₂F₂] (dppe = 1,2-bis(diphenylphosphino)ethane) of 2.056(3)/2.069(3) and 2.1729(18) Å respectively²⁰ presumably results from the presence of a *trans*-labilising hydride ligand.




Fig. 1 Molecular structure of **2**. Solvent, minor disordered component and hydrogen atoms (with the exception of the hydride ligand) have been omitted for clarity. Ellipsoids

are shown at the 30% probability level. Selected bond lengths (Å) and angles (°): Ru(1)-C(1) 2.115(2), Ru(1)-C(10) 2.109(2), Ru(1)-P(1) 2.3343(6), Ru(1)-P(2) 2.3493(6), Ru(1)-F(1) 2.264(2), P(1)-Ru(1)-P(2) 98.66(2), C(1)-Ru(1)-P(1) 170.54(6), C(10)-Ru(1)-F(1) 91.64(8).

A very clear low frequency doublet of triplets Ru-H signal was apparent for **2** in the room temperature ^1H NMR spectrum in toluene- d_8 at δ -21.7 (with diagnostic J_{HF} and J_{HP} values of 52.0 and 19.7 Hz respectively). The IEt_2Me_2 signals were broad and overlapping, but resolved upon cooling to 228 K into eight sets of N- CH_2 and four sets N- CH_2CH_3 signals respectively. The hydride signal at 228 K now appeared as a doublet of doublets ($J_{\text{HF}} = 51.6$ Hz, $J_{\text{HP}} = 25.6$ Hz, $J_{\text{HP}} = 14.1$ Hz), indicating that the two PPh_3 ligands became inequivalent at low temperature. In line with this, the $^{31}\text{P}\{^1\text{H}\}$ spectrum changed from a broad singlet at room temperature to what is best described as two very broad, overlapping multiplets spread over *ca.* 1 ppm at 228 K. We were unable to resolve J_{PP} or J_{PF} splittings even at this low temperature. The ^{19}F NMR spectrum showed a broad fluoride resonance at δ -354 in both THF- d_8 and toluene- d_8 at room temperature, although the doublet hydride splitting of *ca.* 52 Hz was partially resolved in the THF case. Altering the temperature over the range 248-318 K failed to resolve any further couplings, while the addition of CsF also made no effect.²¹

The X-ray structure of the second product, the bis- IEt_2Me_2 pentafluorophenyl complex $[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)(\text{C}_6\text{F}_5)\text{H}]$ (**3**), revealed the anticipated square based pyramidal geometry, with the hydride *trans* to the vacant site (Figure 2). The two carbenes were now oriented *trans* to one another, forcing the PPh_3 and fluoroaryl ring

doublet splitting of 30.6 Hz, and a triplet splitting of 7.2 Hz arising from interaction with the two *ortho*-fluorine atoms of the C₆F₅ ring.

Efforts to accelerate the reaction of **1** with C₆F₅H using higher temperatures resulted instead in the isolation of the bis-phosphine pentafluorophenyl complex [Ru(IEt₂Me₂)(PPh₃)₂(C₆F₅)H] (**4**, Scheme 2) as the major ruthenium containing product of the reaction following overnight heating at 343 K. It was found that **4** could also be formed at room temperature, although very much as the minor partner alongside **2** and **3** if a sample of **1** and C₆F₆ (10 eq) was left at room temperature for *ca.* 100 h (ratio **2**: **3**: **4** = 1:0.4:0.1). Heating an isolated sample of **3** with PPh₃ (2 eq) at 343-363 K in C₆H₆ for 5 h failed to give **4**, implying (unsurprisingly)²² that simple substitution of NHC by phosphine does not account for the formation of **4**.

Crystals of the red compound **4** suitable for X-ray crystallography were isolated from benzene/hexane and displayed the structure shown in Figure 3. Most noticeable was the distorted octahedral geometry now present that resulted from an agostic interaction involving one of the NHC-Et groups occupying the site opposite the Ru-H. The need for the agostic stabilisation must reflect the instability of the five-coordinate 16e Ru(II) species upon replacing the strongly donating IEt₂Me₂ ligand in **3** for PPh₃ in forming **4**. The agostic distances (Ru...C(5), 2.752 Å; Ru...H(5A), 2.052 Å) lie in between those in the related NHC complexes [Ru(IⁱPr₂Me₂)₂(IⁱPr₂Me₂)'Cl] and [Ru(IEt₂Me₂)(PPh₃)₂HCl] previously described by our group²³ and are within the range considered to be strong interactions.²⁴ The Ru-C distance to the pentafluorophenyl ligand was 2.160(2) Å.

Evidence for the agostic interaction being retained in solution was apparent from small, but very clear, doublet ¹⁹F splittings on low frequency resonances for Ru...H-C at δ

0.5 and δ 6.4 in the ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra respectively. Use of ^1H - ^{19}F HMBC spectroscopy established that the coupling resulted from the *ortho*-F signal at δ -112 (see ESI). The hydride resonance in **4** (δ -24.7) resonated to higher frequency of that in five-coordinate (non-agostic) **3**, and appeared as triplet of doublets, the doublet splitting now arising from coupling to the other *ortho*-F signal at δ -106 (see ESI). These couplings help to emphasise the restricted rotation of the C_6F_5 ring suggested by the steric crowding in the crystal structure and proven by the presence of five different ^{19}F NMR resonances.





Fig. 3 Molecular structure of **4**. All hydrogen atoms, except Ru-H and those in the agostic methyl group, are omitted. Ellipsoids are shown at the 30% probability level. Selected bond lengths (Å) and angles (°): Ru(1)-C(1) 2.060(2), Ru(1)-C(10) 2.160(2), Ru(1)-P(1) 2.3452(6), Ru(1)-P(2) 2.3188(6), P(1)-Ru(1)-P(2) 168.093(19), C(10)-Ru(1)-P(1) 91.56(6), C(1)-Ru(1)-P(2) 89.41(6).

Formation of **2** via reaction of **1** with 'HF'

In an effort to find a higher yielding route to the hydride fluoride complex **2**, the reaction of **1** with Et₃N·3HF (TREAT-HF) was investigated. This reagent has become quite commonplace for the formation of transition metal bifluoride ([FHF]⁻) complexes,²⁵ but has, on occasion, also produced metal fluoride species.²⁶ As shown in Scheme 3, **2** was formed as the sole Ru containing product upon reaction of **1** with Et₃N·3HF in a precise 1:1 ratio (i.e. 0.33 eq TREAT-HF). However, increasing the stoichiometry to 1:3 Ru:HF gave instead the previously reported cation [Ru(IEt₂Me₂)₂(PPh₃)₂H]⁺,¹⁴ which was shown crystallographically to be formed as the relatively unusual [H₂F₃]⁻ salt, **5** (see ESI).²⁷




Scheme 3

Catalytic HDF using **1**

Initial catalytic studies have shown that **1** is far more active for HDF than our previously reported [Ru(NHC)(PPh₃)₂(CO)H₂] system^{10,11} bearing unsaturated or

saturated N-aryl substituted carbenes. As shown in Scheme 4, this reacted via initial phosphine loss to give 16-electron $[\text{Ru}(\text{NHC})(\text{PPh}_3)(\text{CO})\text{H}_2]$, which was converted to the hydride fluoride complex $[\text{Ru}(\text{NHC})(\text{PPh}_3)(\text{CO})\text{HF}]$ following hydrodefluorination. Back reaction with the alkylsilane reductant regenerated the dihydride complex, forming a strong Si-F bond in R_3SiF in the process which provides the driving force for the reaction.



Scheme 4

The mixture of C-F and C-H activation products formed in the stoichiometric reaction of **1** and C_6F_6 suggests that the first step of a comparable catalytic cycle with **1** might be more complex, and so the individual stoichiometric reactions of **2**, **3** and **4** with Et_3SiH were investigated to establish the viability of the return reduction steps necessary to complete the catalytic cycle. It was found that: (i) Treatment of the hydride fluoride complex **2** with 1 eq Et_3SiH led to the instantaneous reformation of **1**, along with Et_3SiF ; (ii) There was no reaction between **3** and Et_3SiH (1.5 eq) at room temperature over 6 h, or even upon heating at 343 K for 4 h; (iii) No reaction occurred between the bis-phosphine fluoroaryl complex **4** and silane (1.5 eq) at room temperature overnight, although following addition of $\text{I}(\text{Et}_2\text{Me})_2$ (5 eq), both $\text{C}_6\text{F}_4\text{H}_2$ and Et_3SiF appeared very quickly in

the ^{19}F NMR spectrum. Over the course of *ca.* 2 h, however, the sample began to decompose, shown by the deposition of black solid material




Fig. 4 Product distribution from the catalytic HDF of C_6F_6 using 10 mol% **1** with 80 eq Et_3SiH in C_6H_6 at 363 K. Reactions run under (top line) Ar (1 atm) and (bottom line) H_2 (4 atm) with percentage of products (average of 3 runs) shown after 72 h and (in parentheses) 144 h. HDF products were assigned by ^{19}F NMR spectroscopy.

Figure 4 shows the product distribution from the HDF of C_6F_6 with 10 mol% **1** carried out with Et_3SiH as reductant (80 eq) in C_6H_6 at 363 K. The elevated temperature was adopted in an effort to both push catalysis through at a reasonable rate and also to try to drive HDF through to lower fluorine containing products, which are typically more

difficult to obtain.²⁸ Remarkably, **1** proved capable of bringing about three and four HDF steps to a significant extent, affording 96% of the reaction mixture as isomers of tri- and difluorobenzenes over 72 h.²⁹ Doubling the reaction time increased the amount of 1,2- and 1,4-C₆F₂H₄ and even generated a small amount of fluorobenzene through completion of five HDF steps, giving an overall turnover number of 37. While an in-depth study of the regioselectivity of HDF remains to be carried out, the presence of both 1,2,4,5- and 1,2,3,4-isomers of C₆F₄H₂ after 72 h suggests that the very high *ortho*-regioselectivity found with [Ru(NHC)(PPh₃)₂(CO)H₂] (which converted C₆F₅H overwhelmingly to 1,2,3,4-C₆F₄H₂) is less apparent with **1**. When HDF of C₆F₆ was repeated but now under 4 atm H₂, the amount of difluorobenzene products increased (TON = 38), while the relative ratio of the 1,2:1,3:1,4 difluorobenzene isomers also altered. Interestingly, no turnover of the reaction between C₆F₆ and 10 mol% **1** took place under 4 atm H₂/ excess NEt₃ (80 equivalents)³⁰ in the absence of the silane.

Conclusions

In conclusion, we have shown that [Ru(IEt₂Me₂)₂(PPh₃)₂H₂] (**1**) is a far more active catalyst for the hydrodefluorination of C₆F₆ than the previously reported [Ru(NHC)(PPh₃)(CO)H₂] systems, as reflected in the reduction of C₆F₆ down as far as fluorobenzene. Given the previous mechanistic studies on Ru-H catalysed HDF, this enhanced activity most likely arises from the greater nucleophilicity of the hydride ligands in **1**, arising as a result of their *trans* H-Ru-H geometry. A mechanistic study of **1** is ongoing to confirm the role of the Ru-H bond, and also to help rationalise the lower

regioselectivity compared to $[\text{Ru}(\text{NHC})(\text{PPh}_3)(\text{CO})\text{H}_2]$. Moreover, we hope to be able to explain why **1** is so catalytically competent in spite of appearing, at least on the basis of stoichiometric experiments, to be far more prone than $[\text{Ru}(\text{NHC})(\text{PPh}_3)(\text{CO})\text{H}_2]$ to unfavourable C-H activation reactions with partially fluorinated substrates like $\text{C}_6\text{F}_5\text{H}$. Given that the catalysis was run under high temperature conditions where C-H activated products like $[\text{Ru}(\text{IEt}_2\text{Me}_2)(\text{PPh}_3)_2(\text{C}_6\text{F}_5)\text{H}]$ (**4**) are observed, one possibility is that such species are not dead-ends, but can be recycled into the catalytic cycle, allowing propagation of HDF to continue. While this may help to explain the bias towards more of the lower fluorine containing products with a moderate pressure of H_2 shown in Figure 4, it fails to explain the change in isomer distribution. We hope to present answers to these questions in the near future.

Experimental

General considerations

All manipulations were carried out using standard Schlenk, high vacuum and glovebox techniques. Solvents were purified using an MBraun SPS solvent system (hexane, Et_2O) or under a nitrogen atmosphere from sodium benzophenone ketyl (benzene). C_6D_6 and $\text{C}_6\text{D}_5\text{CD}_3$ were vacuum transferred from potassium. NMR spectra were recorded on Bruker Avance 400/500 and Avance III 500 MHz NMR spectrometers and referenced as follows: ^1H , δ 7.15 ($\text{C}_6\text{D}_5\text{H}$), δ 2.09 ($\text{C}_6\text{D}_5\text{CD}_2\text{H}$) and δ 3.58 ($\text{THF-}d_7$); $^{13}\text{C}\{^1\text{H}\}$, δ 128.0 (C_6D_6) and δ 21.3 ($\text{C}_6\text{D}_5\text{CD}_3$); $^{31}\text{P}\{^1\text{H}\}$, externally to 85% H_3PO_4 (δ 0.0); ^{19}F , externally to CFCl_3 (δ 0.0). PPh_3 resonances are excluded unless they could be assigned unequivocally. Elemental analyses were performed by Elemental Microanalysis Ltd,

Okehampton, Devon. $[\text{Ru}(\text{PPh}_3)_4\text{H}_2]$ ³¹ and IEt_2Me_2 ³² were prepared according to literature methods.

$[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)_2\text{H}_2]$ (1). An alternative synthesis of **1** carried out in benzene rather than as previously described¹⁴ in THF is reported here. This new approach afforded **1** in shorter time and in higher yield. $[\text{Ru}(\text{PPh}_3)_4\text{H}_2]$ (500 mg, 0.43 mmol) and IEt_2Me_2 (130 mg, 0.86 mmol) were dissolved in benzene (5 mL) and stirred in an ampoule sealed with a J. Youngs PTFE tap for 5 min at 298 K. The solution was filtered by cannula into a fresh ampoule and the volatiles were removed *in vacuo* to leave a sticky orange residue. This was washed with hexane (2 x 2 mL) to afford **1** as pale yellow solid. Yield: 279 mg, 70 %. Spectroscopic data matched those in the original report.¹⁴

$[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)_2\text{HF}]$ (2). C_6F_6 (50 μL , 0.45 mmol) was added to a benzene (5 mL) solution of **1** (140 mg, 0.15 mmol) in an ampoule fitted with J. Youngs PTFE tap. The reaction mixture was stirred vigorously for 24 h, filtered by cannula and evaporated to dryness to afford an oily red residue. Addition of hexane (1 mL) under the action of vigorous stirring resulted in a formation of a deep orange suspension (of **3**), which was filtered by cannula. Leaving the hexane filtrate at room temperature for few days afforded yellow crystals of **2**, which were manually separated from red needles of residual **3**. Yield of **2**: 43 mg, 30 %. A more efficient route to **2** involved treatment of **1** with $\text{Et}_3\text{N}\cdot 3\text{HF}$ (TREAT-HF). Thus, TREAT-HF (6.1 μL , 0.037 mmol) was added by syringe to a benzene solution (5 mL) of **1** (100 mg, 0.11 mmol) in an ampoule fitted with a J. Youngs PTFE tap. The reaction mixture was stirred for 30 min, the volatiles then removed under vacuum and the sticky yellow solid washed with hexane (2 mL) to afford **2** as a pale yellow solid. Yield: 75 mg, 72 %. ¹H NMR: δ_{H} ($\text{C}_6\text{D}_5\text{CD}_3$, 400 MHz, 228 K) -21.58

(ddd, 1H, $J_{HF} = 51.6$ Hz, $J_{HP} = 25.0$ Hz, $J_{HP} = 14.1$ Hz, Ru-*H*), 0.26 (t, 3H, $J_{HH} = 6.8$ Hz, NCH₂CH₃), 0.34 (t, 3H, $J_{HH} = 6.8$ Hz, NCH₂CH₃), 1.10 (t, 3H, $J_{HH} = 6.8$ Hz, NCH₂CH₃), 1.16 (s, 3H, NCCH₃), 1.21 (s, 3H, NCCH₃), 1.39 (t, 3H, $J_{HH} = 6.8$ Hz, NCH₂CH₃), 1.49 (s, 3H, NCCH₃), 1.56 (s, 3H, NCCH₃), 2.32 (m, 1H, $J_{HH} = 6.8$ Hz, NCHHCH₃), 2.61 (m, 1H, $J_{HH} = 6.8$ Hz, NCHHCH₃), 3.13 (m, 1H, $J_{HH} = 6.8$ Hz, NCHHCH₃), 3.36 (m, 1H, $J_{HH} = 6.8$ Hz, NCHHCH₃), 5.60 (br m, 1H, $J_{HH} = 6.8$ Hz, NCHHCH₃), 5.83 (br m, 1H, $J_{HH} = 6.8$ Hz, NCHHCH₃), 6.45 (br s, 1H, NCHHCH₃), 6.80 (br s, 1H, NCHHCH₃)*. * = chemical shift established by ¹H COSY. ³¹P {¹H} NMR: δ_P (C₆D₅CD₃, 121.5 MHz, 298 K): 43.1 (br s). ¹³C {¹H} NMR: δ_C (C₆D₅CD₃, 100 MHz, 228 K) 8.7 (s, NCCH₃), 8.8 (s, NCCH₃), 9.1 (s, NCCH₃), 9.4 (s, NCCH₃), 13.6 (s, NCH₂CH₃), 14.2 (s, NCH₂CH₃), 15.0 (s, NCH₂CH₃), 16.2 (s, NCH₂CH₃), 40.5 (d, J_{CP} or $J_{CF} = 32.2$ Hz, NCH₂CH₃), 42.0 (d, J_{CP} or $J_{CF} = 16.4$ Hz, NCH₂CH₃), 43.2 (s, NCH₂CH₃), 122.3 (s, NCCH₃), 122.9 (s, NCCH₃), 123.5 (s, NCCH₃), 124.4 (s, NCCH₃), 191.4 (m, Ru-C_{NHC}). ¹⁹F NMR (THF-*d*₈, 470 MHz, 298 K): δ -354.4 (br d, $J_{FH} = 51.6$ Hz). Analysis found: C, 68.99; H, 7.15; N, 5.62%. C₅₇H₆₃N₄FP₂Ru·0.5C₆H₁₄ requires: C, 68.93; H, 7.10; N, 5.64%.

[Ru(IEt₂Me₂)₂(PPh₃)(C₆F₅)H] (3). C₆F₅H (120 μ L, 1.1 mmol) was syringed into a J. Youngs resealable ampoule containing a hexane suspension (5 mL) of **1** (100 mg, 0.11 mmol). The reaction mixture was stirred vigorously at room temperature for 24 h to give a dark orange solid, which was isolated by cannula filtration, washed with hexane (2 x 5 mL) and dried in *vacuo*. Yield 53 mg, 58 %. ¹H NMR: δ_H (C₆D₆, 500 MHz, 298 K) - 32.95 (dt, 1H, $J_{HP} = 30.6$ Hz, $J_{HF} = 7.2$ Hz, Ru-*H*), 0.98 (t, 6H, $J_{HH} = 7.3$ Hz, NCH₂CH₃), 1.02 (t, 6H, $J_{HH} = 7.3$ Hz, NCH₂CH₃), 1.45 (s, 6H, NCCH₃), 1.48 (s, 6H, NCCH₃), 3.05 (m, 2H, NCH₂CH₃), 3.60 (m, 4H, NCH₂CH₃), 4.77 (m, 2H, NCH₂CH₃), 6.90-7.05 (br m,

9H, PC₆H₅), 7.43-7.49 (m, 6H, PC₆H₅). ³¹P{¹H} NMR: δ_P (C₆D₆, 121.5 MHz, 298 K) 59.5 (tt, *J*_{PF} = 20.7 Hz, *J*_{PF} = 9.7 Hz). ¹³C{¹H} NMR: δ_C (C₆D₆, 126 MHz, 298 K) 9.0 (s, NCH₂CH₃), 9.2 (s, NCH₂CH₃), 15.3 (s, NCCH₃), 15.4 (s, NCCH₃), 42.4 (s, NCH₂CH₃), 43.2 (s, NCH₂CH₃), 123.2 (s, NCCH₃), 123.6 (s, NCCH₃), 127.1 (d, *J*_{CP} = 7.3 Hz, PC₆H₅), 127.4 (s, PC₆H₅), 133.6 (d, *J*_{CP} = 11.0 Hz, PC₆H₅), 142.8 (d, *J*_{CP} = 26.7 Hz, PC₆H₅), 195.9 (d, *J*_{CP} = 12.1 Hz, Ru-C_{NHC}). ¹⁹F NMR: δ_F (C₆D₆, 470 MHz, 298 K) -166.4 (1F, t, *J*_{FF} = 20.3 Hz, *p*-C₆F₅), -165.6 (2F, m, *m*-C₆F₅), -111.5 (2F, br s, *o*-C₆F₅). Analysis found: C, 60.36; H, 5.74; N, 6.72. C₄₂H₄₈N₄F₅PRu requires: C, 60.34; H, 5.79; N, 6.70.

[Ru(IEt₂Me₂)(PPh₃)₂(C₆F₅)H] (4). A J. Young NMR tube containing **1** (45 mg, 48 μmol) and C₆F₅H (16 μL, 145 μmol) was heated in C₆H₆ (0.5 mL) at 343 K overnight to afford a deep red solution. This was filtered by cannula and the filtrate evaporated to dryness. After washing with hexane (3 x 0.5 mL), the residue was redissolved in a minimal amount of THF and layered with hexane to afford deep red crystals of **4**. Yield: 13 mg, 28%. ¹H NMR: δ_H (THF-*d*₈, 500 MHz, 298 K) -24.66 (1H, td, *J*_{PH} = 23.5 Hz, *J*_{HF} = 6.9 Hz, Ru-H), 0.34 (3H, t, *J*_{HH} = 7.3 Hz, NCH₂CH₃), 0.48 (3H, td, *J*_{HH} = 7.3 Hz, *J*_{HF} = 1.5 Hz, NCH₂CH₃), 1.92 (3H, s, NCCH₃), 1.96 (3H, s, NCCH₃), 2.90 (2H, q, *J*_{HH} = 7.3 Hz, NCH₂CH₃), 3.38 (2H, q, *J*_{HH} = 7.3 Hz, NCH₂CH₃), 7.02-7.24 (30H, br m, PC₆H₅). ³¹P{¹H} NMR: δ_P (THF-*d*₈, 202 MHz, 298 K) 52.3 (s). ¹³C{¹H} NMR: δ_C (THF-*d*₈, 126 MHz, 298 K) 6.4 (d, *J*_{CF} = 7.5 Hz, NCH₂CH₃), 9.4 (s, NCCH₃), 9.8 (s, NCCH₃), 14.5 (s, NCH₂CH₃), 42.5 (s, NCH₂CH₃), 44.0 (s, NCH₂CH₃), 124.7 (s, NCCH₃), 126.2 (s, NCCH₃), 127.9 (virtual triplet ('vt'), *J* = 4 Hz, PC₆H₅), 129.0 (s, PC₆H₅), 134.6 ('vt', *J* = 6 Hz, PC₆H₅), 139.0 ('vt', *J* = 17 Hz, PC₆H₅), 194.0 (m, Ru-C_{NHC}). ¹⁹F NMR: δ_F (THF-*d*₈, 470 MHz, 298 K) -171.5 (1F, t, *J*_{FF} = 20.2 Hz, *p*-C₆F₅), -170.1 (1F, m, *p*-C₆F₅), -168.9

(1F, m, *m*-C₆F₅), -111.8 (1F, m, *o*-C₆F₅), -105.5 (1F, m, *o*-C₆F₅). Analysis found: C, 64.89; H, 4.98; N, 3.01. C₅₁H₄₇N₂F₅P₂Ru requires: C, 64.75; H, 5.01; N, 2.96.

[Ru(IEt₂Me₂)₂(PPh₃)₂H][H₂F₃] (5). TREAT-HF (17.5 μL, 0.11 mmol) was added to a benzene (5 mL) solution of **1** (100 mg, 0.11 mmol) in a J. Youngs resealable ampoule. The reaction mixture was stirred at room temperature for 30 min at 298 K, before the sample was reduced to dryness. The sticky orange/red residue was washed with hexane (2 x 2 mL) and Et₂O (2 x 2 mL) and then redissolved in THF (5 mL). Addition of Et₂O resulted in the precipitation of **5** as an orange solid, which was washed further with Et₂O (2 x 5 mL) and then dried *in vacuo*. Yield: 76 mg, 69%. Crystals suitable for X-ray diffraction were obtained upon layering a concentrated THF-*d*₈ solution with hexane. ¹H NMR: δ_H (THF-*d*₈, 500 MHz, 298 K) -29.65 (1H, t, *J*_{HP} = 24.0 Hz, Ru-*H*), 0.44 (6H, t, *J*_{HH} = 7.3 Hz, NCH₂CH₃), 0.88 (6H, t, *J*_{HH} = 7.3 Hz, NCH₂CH₃), 1.81 (6H, s, NCCH₃), 2.01 (6H, s, NCCH₃), 2.75 (4H, q, *J*_{HH} = 7.3 Hz, NCH₂CH₃), 3.36 (4H, q, *J*_{HH} = 7.3 Hz, NCH₂CH₃), 7.16-7.34 (30H, m, P(C₆H₅)₃), 13.68 (2H, br s, [H₂F₃]⁻). ³¹P {¹H} NMR: δ_P (THF-*d*₈, 202 MHz, 298 K) 46.1 (s). ¹⁹F NMR: δ_F (THF-*d*₈, 470 MHz, 298 K) -115.2 (br s). Analysis found: C, 64.38; H, 5.69; N, 4.84.

C₅₄H₆₅N₄F₃P₂Ru·2C₄D₈O requires C, 64.73; H, 5.70; N, 4.87.

Procedures for Catalytic HDF. A stock solution of **1** was prepared by dissolving 0.0184 g (0.02 mmol) of the complex in 2 mL C₆H₆ in the glovebox. 0.5 mL aliquots of this solution was syringed into three J. Young's resealable NMR tubes, and C₆F₆ (5.8 μL, 0.05 mmol) and Et₃SiH (63 μL, 0.4 mmol) added to each tube. These were then placed in a pre-heated oil bath at 363 K and monitored by ¹⁹F NMR spectroscopy after 72 and 144 h. For reactions performed under H₂, a C₆H₆ (0.3 mL) sample of **1** (0.0046 g, 0.005

mmol) was placed into a medium-walled NMR tube fitted with a resealable valve and freeze-pump-thaw degassed (3 cycles). A mixture of C₆F₆ (5.8 μL, 0.05 mmol), Et₃SiH (63 μL, 0.4 mmol) and C₆H₆ (0.1 mL) was vacuum transferred into the pressure tube, which was then put under 4 atm H₂ and placed in a pre-heated oil bath at 363 K. The reaction was monitored by ¹⁹F NMR spectroscopy after 72 and 144 h.

X-ray crystallography. Data for **2** and **4** were obtained using a Nonius Kappa CCD diffractometer, while those for **3** and **5** (see ESI) were collected using Agilent SuperNova and Agilent Excalibur diffractometers, respectively. Details of the data collections, solutions and refinements are given in Table 1. All diffraction measurements were conducted at 150 K using Mo(Kα) radiation and hydride ligands were uniformly refined subject to being a distance of 1.6 Å from the relevant metal centre. Convergence was straightforward in all cases, and only exceptional details merit note. In particular, the asymmetric unit in **2** was seen to comprise one bis-carbene complex and half of a hexane molecule. The latter is proximate to an inversion centre which serves to generate the remainder. The fluoride and hydride ligands were modelled subject to being disordered with each other in a 53:47 ratio. Fractional occupancy hydride atoms were refined with a common isotropic displacement parameter. In **3**, H5A, H5B and H5C were readily located and refined with the single restraint of being at a distance of 0.98 Å from C5. The structures were solved using SHELXS-97³³ and refined using full-matrix least squares in SHELXL-97.³³ Crystallographic data for compounds have been deposited with the Cambridge Crystallographic Data Centre as supplementary publications CCDC 1400863-1400866. Copies of the data can be obtained free of charge on application to CCDC, 12

Union Road, Cambridge CB2 1EZ, UK [fax(+44) 1223 336033, e-mail:
deposit@ccdc.cam.ac.uk].

Acknowledgements. We thank the University of Bath and EPSRC for financial support.

References

1. V. P. Reddy, A. Vadapalli, E. Sinn and N. Hosmane, *J. Organomet. Chem.*, 2013, **747**, 43.
2. Y.-M. Zhao, M. S. Cheung, Z. Lin and J. Sun, *Angew. Chem. Int. Ed.*, 2012, **51**, 10359; J. Mo, X. Chen and Y. R. Chi, *J. Am. Chem. Soc.*, 2012, **134**, 8810; X. Dong, Y.-M. Zhao and J. Sun, *Synlett*, 2013, **24**, 1221; F. Li, Z. Wu and J. Wang, *Angew. Chem. Int. Ed.*, 2015, **54**, 656; X. Dong, W. Yang, W. Hu and J. Sun, *Angew. Chem. Int. Ed.*, 2015, **54**, 660.
3. J. J. Song, Z. Tan, J. T. Reeves, F. Gallou, N. K. Yee and C. H. Senanayake, *Org. Lett.*, 2005, **7**, 2193; L.-T. Shen, P.-L. Shao and S. Ye, *Adv. Synth. Catal.*, 2011, **354**, 1943; X. Dong and J. Sun, *Org. Lett.*, 2014, **16**, 2450. See also, reports on the use of PhenoFluor and PhenFluorMix for C-F bond formation. P. Tang, W. Wang and T. Ritter, *J. Am. Chem. Soc.*, 2011, **133**, 11482; T. Fujimoto and T. Ritter, *Org. Lett.*, 2015, **17**, 544.
4. H. U. Vora and T. Rovis, *J. Am. Chem. Soc.*, 2010, **132**, 2860; P. Wheeler, H. U. Vora and T. Rovis, *Chem. Sci.*, 2013, **4**, 1674; M. Kajjout, M. Smietana, J. Leroy and C. Rolando, *Tetrahedron Lett.*, 2013, **54**, 1658.
5. J. A. Akana, K. X. Bhattacharyya, P. Müller and J. P. Sadighi, *J. Am. Chem. Soc.*, 2007, **129**, 7736; B. C. Gorske, C. T. Mbofana and S. J. Miller, *Org. Lett.*, 2009,

- 11**, 4318; H. Lv, J.-H. Zhan, Y.-B. Cai, Y. Yu, B. Wang and J.-L. Zhang, *J. Am. Chem. Soc.*, 2012, **134**, 16216; C. M. Wys, B. K. Tate, J. Bacsá, M. Wieliczko and J. P. Sadighi, *Polyhedron*, 2014, **84**, 87.
6. F. Nagra, S. R. Patrick, D. Bello, M. Brill, A. Obled, D. B. Cordes, A. M. Z. Slawin, D. O'Hagan and S. P. Nolan, *ChemCatChem*, 2015, **7**, 240.
7. Trifluoromethylation reactions are also possible, but are stoichiometric: G. G. Dubinina, H. Furutachi and D. A. Vicic, *J. Am. Chem. Soc.*, 2008, **130**, 8600; G. G. Dubinina, J. Ogikubo and D. A. Vicic, *Organometallics*, 2008, **27**, 6233.
8. T. Schaub, M. Backes and U. Radius, *J. Am. Chem. Soc.*, 2006, **128**, 15964; Z. Mo, Q. Zhang and L. Deng, *Organometallics*, 2012, **31**, 6518. For reviews of the field, see: A. D. Sun and J. A. Love, *Dalton Trans.*, 2010, **39**, 10362; T. Ahrens, J. Kohlmann, M. Ahrens and T. Braun, *Chem. Rev.*, 2015, **115**, 931.
9. S. Kuhl, R. Schneider and Y. Fort, *Adv. Synth. Catal.*, 2003, **345**, 341; P. Fischer, K. Götz, A. Eichhorn and U. Radius, *Organometallics*, 2012, **31**, 1374; S. Sabater, J. A. Mata and E. Peris, *Nat. Commun.*, 2013, **4**, 2553; L. Schwartsburd, M. F. Mahon, R. C. Poulten, M. R. Warren and M. K. Whittlesey, *Organometallics*, 2014, **33**, 6165; S. Sabater, J. A. Mata and E. Peris, *Organometallics*, 2015, **34**, 1186.
10. S. P. Reade, M. F. Mahon and M. K. Whittlesey, *J. Am. Chem. Soc.*, 2009, **131**, 1847; J. A. Panetier, S. A. Macgregor and M. K. Whittlesey, *Angew. Chem. Int. Ed.*, 2011, **50**, 2783; S. A. Macgregor, D. McKay, J. A. Panetier and M. K. Whittlesey, *Dalton Trans.*, 2013, **42**, 7386.

11. D. McKay, I. M. Riddlestone, S. A. Macgregor, M. F. Mahon and M. K. Whittlesey, *ACS Catal.*, 2015, **5**, 776.
12. S. P. Reade, A. L. Acton, M. F. Mahon, T. A. Martin and M. K. Whittlesey, *Eur. J. Inorg. Chem.*, 2009, 1774.
13. L. M. Guard, A. E. W. Ledger, S. P. Reade, C. E. Ellul, M. F. Mahon and M. K. Whittlesey, *J. Organomet. Chem.*, 2011, **696**, 780; M. K. Whittlesey and E. Peris, *ACS Catal.*, 2014, **4**, 3152.
14. C. J. E. Davies, J. P. Lowe, M. F. Mahon, R. C. Poulten and M. K. Whittlesey, *Organometallics*, 2013, **32**, 4927.
15. C₆D₆ needs to be avoided because of facile H/D exchange with **1**. See ref 14.
16. For a discussion of C-F versus C-H bond activation, see: S. A. Johnson, J. A. Hatnean and M. E. Doster, *Prog. Inorg. Chem.*, 2012, **57**, 255.
17. M. K. Whittlesey, R. N. Perutz, B. Greener and M. H. Moore, *Chem. Commun.*, 1997, 187.
18. R. G. Ball, B. R. James, J. Trotter, D. K. W. Wang and K. R. Dixon, *J. Chem. Soc., Chem. Commun.*, 1979, 460; J. Figueira, M. G. Jardim, J. Rodrigues, A. Valkonen and K. Rissanen, *Inorg. Chem. Commun.*, 2013, **29**, 123; M. V. Câmpain, R. N. Perutz, B. Procacci, R. J. Thatcher, O. Torres and A. C. Whitwood, *J. Am. Chem. Soc.*, 2012, **134**, 3480.
19. J. Chatt and R. G. Hayter, *J. Chem. Soc.*, 1961, 2605.
20. D. A. J. Harding, E. G. Hope, G. A. Solan and J. Fawcett, *Acta Crystallogr., Sect. C*, 2007, **63**, m383; P. Barthazy, R. M. Stoop, M. Wörle, A. Togni and A. Mezzetti, *Organometallics*, 2000, **19**, 2844.

21. S. A. Macgregor, D. C. Roe, W. J. Marshall, K. M. Bloch, V. I. Bakhmutov and V. V. Grushin, *J. Am. Chem. Soc.*, 2005, **127**, 15304; S. P. Reade, D. Nama, M. F. Mahon, P. S. Pregosin and M. K. Whittlesey, *Organometallics*, 2007, **26**, 3484.
22. J. Huang, E. D. Stevens, S. P. Nolan and J. L. Petersen, *J. Am. Chem. Soc.*, 1999, **121**, 2674.
23. S. Burling, E. Mas-Marzá, J. E. V. Valpuesta, M. F. Mahon and M. K. Whittlesey, *Organometallics*, 2009, **28**, 6676.
24. W. Baratta, E. Herdtweck and P. Rigo, *Angew. Chem. Int. Ed.*, 1999, **38**, 1629; W. Baratta, C. Mealli, E. Herdtweck, A. Ienco, S. A. Mason and P. Rigo, *J. Am. Chem. Soc.*, 2004, **126**, 5549.
25. N. A. Jasim and R. N. Perutz, *J. Am. Chem. Soc.*, 2000, **122**, 8685; D. C. Roe, W. J. Marshall, F. Davidson, P. D. Soper and V. V. Grushin, *Organometallics*, 2000, **19**, 4575; N. A. Jasim, R. N. Perutz, S. P. Foxon and P. H. Walton, *J. Chem. Soc., Dalton Trans.*, 2001, 1676; J. Vicente, J. Gil-Rubio, D. Bautista, A. Sironi and N. Masciocchi, *Inorg. Chem.*, 2004, **43**, 5665; T. Vergote, F. Nahra, A. Welle, M. Luhmer, J. Wouters, N. Mager, O. Riant and T. Leyssens, *Chem. Eur. J.*, 2012, **18**, 793; C. Segarra, E. Mas-Marzá, J. P. Lowe, M. F. Mahon, R. C. Poulten and M. K. Whittlesey, *Organometallics*, 2012, **31**, 8584; N. Bramanathan, M. Carmona, J. P. Lowe, M. F. Mahon, R. C. Poulten and M. K. Whittlesey, *Organometallics*, 2014, **33**, 1986; B. J. Truscott, F. Nahra, A. M. Z. Slawin, D. B. Cordes and S. P. Nolan, *Chem. Commun.*, 2015, **51**, 62.
26. M. C. Pilon and V. V. Grushin, *Organometallics*, 1998, **17**, 1774; D. Noveski, T. Braun and S. Krückemeier, *J. Fluorine Chem.*, 2004, **125**, 959; L. M. Martínez-

- Prieto, C. Melero, D. del Río, P. Palma, J. Cámpora and E. Álvarez, *Organometallics*, 2012, **31**, 1425.
27. T. Mathieson, A. Schier and H. Schmidbaur, *Z. Naturforsch.*, 2000, **55b**, 1000; F. J. Fernández, M. Alfonso, H. W. Schmalle and H. Berke, *Organometallics*, 2001, **20**, 3122; B. G. Harvey, A. M. Arif, A. Glöckner and R. D. Ernst, *Organometallics*, 2007, **26**, 2872; I. Bernal and S. F. Watkins, *C. R. Chimie*, 2014, **17**, 586.
28. For a review of catalytic HDF, see: M. Kuehnel, D. Lentz and T. Braun, *Angew. Chem. Int. Ed.*, 2013, **52**, 3328.
29. For examples of catalytic HDF in which highly fluorinated substrates undergo multiple defluorination steps, see: J. Wu and S. Cao, *ChemCatChem* 2011, **3**, 1582; W. Zhao, J. Wu and S. Cao, *Adv. Synth. Catal.*, 2012, **354**, 574; J. Xiao, J. Wu, W. Zhao and S. Cao, *J. Fluorine Chem.*, 2013, **146**, 76; G. Podolan, D. Lentz and H.-U. Reissig, *Angew. Chem. Int. Ed.*, 2013, **52**, 9491; L. Zámostná, M. Ahrens and T. Braun, *J. Fluorine Chem.* 2013, **155**, 132; O. Ekkert, S. D. A. Strudley, A. Rozenfeld, A. J. P. White and M. R. Crimmin, *Organometallics*, 2014, **33**, 7027.; S. M. Senaweera, A. Singh and J. D. Weaver, *J. Am. Chem. Soc.*, 2014, **136**, 3002.
30. H. Nakai, K. Jeong, T. Matsumoto and S. Ogo, *Organometallics*, **2014**, **33**, 4349.
31. R. Young and G. Wilkinson, *Inorg. Synth.*, 1990, **28**, 337.
32. N. Kühn and T. Kratz, *Synthesis*, 1993, 561.

33. G. M. Sheldrick, *Acta. Cryst.* 1990, **467-473**, A46. Sheldrick, G. M, SHELXL-97, a computer program for crystal structure refinement, University of Göttingen, 1997.

Compound reference	2	3	4
Chemical formula	C ₅₇ H ₇₀ N ₄ FP ₂ Ru	C ₄₂ H ₄₈ N ₄ F ₅ PRu	C ₅₁ H ₄₇ N ₂ F ₅ P ₂ Ru
Formula Mass	993.18	835.88	945.92
Crystal system	Monoclinic	Orthorhombic	Triclinic
<i>a</i> /Å	16.7400(1)	31.0451(7)	12.6690(2)
<i>b</i> /Å	16.5550(1)	8.9768(2)	13.1530(2)
<i>c</i> /Å	19.4760(1)	14.4306(3)	15.5900(3)
α /°	90.00	90.0	96.300(1)
β /°	109.727(1)	90.0	105.626(1)
γ /°	90.00	90.0	116.893(1)
Unit cell volume/Å ³	5080.63(5)	4021.62(17)	2148.63(6)
Space group	<i>P</i> 2 ₁ / <i>a</i>	<i>Pna</i> 2 ₁	<i>P</i> $\bar{1}$
No. of formula units per unit cell, <i>Z</i>	4	4	2
No. of reflections measured	94717	22071	40103
No. of independent reflections	11623	7401	9725
<i>R</i> _{int}	0.0678	0.0684	0.0478
Final <i>R</i> 1 values (<i>I</i> > 2σ(<i>I</i>))	0.0370	0.0402	0.0338
Final <i>wR</i> (<i>F</i> ²) values (<i>I</i> > 2σ(<i>I</i>))	0.0767	0.0877	0.0685
Final <i>R</i> 1 values (all data)	0.0582	0.0502	0.0516
Final <i>wR</i> (<i>F</i> ²) values (all data)	0.0851	0.0942	0.0746


Table 1. Crystal data and structure refinement details for compounds **2**, **3** and **4**.

Graphical Abstract

Stoichiometric and Catalytic C-F Bond Activation by the *Trans*-Dihydride NHC

Complex $[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)_2\text{H}_2]$ ($\text{IEt}_2\text{Me}_2 = 1,3\text{-diethyl-4,5-dimethylimidazol-2-ylidene}$)

Mateusz K. Cybulski, Ian M. Riddlestone, Mary F. Mahon, Timothy J. Woodman and Michael K. Whittlesey



Multiple catalytic hydrodefluorination steps take place with the *trans*-dihydride complex $[\text{Ru}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_3)_2\text{H}_2]$ (**1**), taking C_6F_6 to tri-, di- and mono-fluorobenzenes.

