



Citation for published version:

Grammatopoulos, G, Wales, J, Kothari, A, Gill, HS, Wainwright, A & Theologis, T 2015, 'What is the early/mid-term survivorship and functional outcome after Bernese Periacetabular Osteotomy in a Pediatric Surgeon Practice?', *Clinical Orthopaedics and Related Research*. <https://doi.org/10.1007/s11999-015-4386-7>

DOI:

[10.1007/s11999-015-4386-7](https://doi.org/10.1007/s11999-015-4386-7)

Publication date:

2015

Document Version

Early version, also known as pre-print

[Link to publication](#)

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.




Figure 1: Graph illustrating Kaplan-Meier survivorship curve with conversion to hip arthroplasty as the end point for whole cohort.




Figure 2: Graph illustrating Kaplan-Meier survivorship curve as worse case scenario with conversion to hip arthroplasty, poor WOMAC score (<40) and lost to follow-ups treated as end points for whole cohort.





Figure 3: Scatter-plot illustrating the effect of AI (A) and CEA (B) on WOMAC at follow-up.