



Citation for published version:

McCullen, N, Wolfrum, M & Wagenknecht, T 2015, 'Snaking on Networks: From Local Solutions to Turing Patterns' SIAM Conference on Applications of Dynamical Systems, Snowbird, USA United States, 17/05/05 - 21/08/15, .

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link to publication](#)

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Background

- Patterns are found throughout nature and identifying them and understanding their origin forms the basis of much of our scientific understanding.
- Alan Turing laid down the basic mechanisms for spatial pattern formation, based the loss of stability of an equilibrium state to a patterned state [1].
- Recent attention has been focussed on non-regular *complex network* topologies, which are a model for many phenomena in a variety of fields including biology and sociology [2].

The Mimura-Murray model

- Reaction-diffusion system:

$$\begin{aligned} \dot{u} &= f(u, v) - DLu \\ \dot{v} &= g(u, v) - \sigma DLv \end{aligned}$$

with

$$f(u, v) = \frac{au + bu^2 - u^3}{c} - uv, \quad g(u, v) = uv - v - dv^2$$

at the parameter values $a = 35$, $b = 16$, $c = 9$, $d = 2/5$.

- u : activator, v : inhibitor; $L = (L_{(i,j)})$ network Laplacian;
- the model has a stable equilibrium at $(\bar{u}, \bar{v}) = (5, 10)$,
- Turing instability: equilibrium destabilized on increase of σ ,
– undergoes a *supercritical* bifurcation at $\sigma = \sigma_T \approx 15.5$.

Scale-Free “Complex” Networks

- Preferential-attachment networks such as the Barabási–Albert model are found throughout nature and have a *scale-free* power-law degree distribution [5].




Figure 1: A “Complex” Network

Mimura-Murray on Complex Networks

- Coexistence and multi-stability of a huge variety of patterns was previously found [4].




Figure 2: (a) from Nakao & Mikhailov (2010) [4]; (b) from Wolfrum [3].

- Explained analytically by considering single differentiated nodes (SDN) with others fixed at equilibrium [3], multiple bifurcation curves found – one for each node.

Outline of this Work

- The emergence of patterns of activity on complex networks with reaction-diffusion dynamics on the nodes has been numerically studied.
- The transition between previously analysed single-node solutions [3] and fully developed global activation states (so called *Turing states*) has been investigated.
- We reveal *snaking bifurcations* connecting different localised solutions, as found on regular lattices, shedding light on the origin of previously reported multistable “*Turing patterns*” [4].

Localised Patterns and Snaking Bifurcations

- Theory exists for development of patterns on reaction-diffusion systems on regular lattices:




Figure 3: (a) from Burke & Knobloch [6]; (b) from Beck et al. [7]

- localised patterns develop via homoclinic “snaking bifurcations” from a ground-state;
- each level of the “snake” is associated with increasingly more *activated nodes*.

Snaking Bifurcations on Complex Mimura-Murray




Figure 4: We find that snaking is also the route to multiply differentiated patterns on Mimura-Murray systems on complex scale-free networks.

The Bigger Picture




Figure 5: (a) The “zoo” of solutions is immense; (b) Attenuating the influence of neighbouring nodes reveals the deviation of the minimum (σ^*) of the region of existence of the differentiated states from the single-node (SDN) theory.

Conclusions

- Reaction-diffusion systems on complex scale-free networks display a rich set of coexisting solutions, from single differentiated nodes to full-scale *Turing patterns*.
- The patterns of activity, originating in singly-differentiated-node solutions, develop into full-scale *patterns* of activity via *snaking bifurcations*, as found in other related systems.
- Simple analysis can begin to reveal the origins of the full complexity of these systems.

References

- [1] Alan Mathison Turing, *The chemical basis of morphogenesis*, Philosophical Transactions of the Royal Society of London 237 (641): 37–72.
- [2] Gergely Palla et al. *Uncovering the overlapping community structure of complex networks in nature and society* Nature 435, (2005): 814–818.
- [3] Matthias Wolfrum, *The Turing bifurcation in network systems: Collective patterns and single differentiated nodes*, Physica D: Nonlinear Phenomena, (2012).
- [4] Hiroya Nakao and Alexander S Mikhailov, *Turing patterns in network-organized activator-inhibitor systems*, Nature Physics, 6 (2010): pp. 544–550.
- [5] Albert-László Barabási and Réka Albert, *Emergence of scaling in random networks*, Science, 286 (1999): pp. 509–512.
- [6] John Burke and Edgar Knobloch, *Snakes and ladders: localized states in the Swift-Hohenberg equation*, Physics Letters A 360.6 (2007): 681–688.
- [7] Margaret Beck, Jürgen Knobloch, David JB Lloyd, Björn Sandstede and Thomas Wagenknecht. *Snakes, ladders, and isolas of localized patterns*, SIAM Journal on Mathematical Analysis 41, (2009): no. 3 936–972.