



*Citation for published version:*

Knobloch, R & Kyprianou, A 2014, 'Survival of homogenous fragmentation processes with killing', *Annales de l'Institut Henri Poincaré B Probabilités et Statistiques*, vol. 50, no. 2, pp. 467 - 491. <https://doi.org/10.1214/12-AIHP520>

*DOI:*

[10.1214/12-AIHP520](https://doi.org/10.1214/12-AIHP520)

*Publication date:*

2014

*Document Version*

Peer reviewed version

[Link to publication](#)

The published version is available via: <http://dx.doi.org/10.1214/12-AIHP520>

## University of Bath

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Survival of homogeneous fragmentation processes with killing

Robert Knobloch\* and Andreas E. Kyprianou†

August 20, 2012

## Abstract

We consider a homogeneous fragmentation process with killing at an exponential barrier. With the help of two families of martingales we analyse the decay of the largest fragment for parameter values that allow for survival. In this respect the present paper is also concerned with the probability of extinction of the killed process.

**AMS 2000 Mathematics Subject Classification:** 60J25, 60G09.

**Keywords and phrases:** homogeneous fragmentation, scale functions, additive martingales, multiplicative martingales, largest fragment.

## 1 Introduction and main results

This paper is concerned with a homogeneous fragmentation process in which there is an additional killing upon crossing a certain space-time barrier (this killing mechanism is defined rigorously in Section 1.2). In particular, we consider the decay of the largest fragment in this process with killing.

The motivation for the killing procedure that we introduce in the present paper, partly stems from its relation to the Fisher-Kolmogorov-Petrovskii-Piskounov (FKPP) equation. In the context of fragmentation processes this connection is studied in [18]. The role an analogous killing plays with regard to solutions of the FKPP equation in the setting of branching Brownian motions (BBM) was investigated in [15]. Furthermore, this kind of killing for random multi-particle systems was considered also in various other contexts and in the literature there is some interesting recent activity in this regard. The killing of BBM at a linear space-time barrier was also studied in [14] and recently in [2], where a relation of the killed BBM and its genealogy to continuous-state branching processes and the Bolthausen-Sznitman coalescent was revealed. Regarding similar killing schemes for branching random walks we refer e.g. to [1], [10], [11] and [13]. In the context of fragmentation processes such a killing mechanism has not been considered so far. However, the above-mentioned papers which are concerned with related types of spatial branching processes suggest that this kind of killing has interesting applications.

---

\*Institut für Mathematik, Goethe-Universität Frankfurt am Main, 60054 Frankfurt am Main, Germany  
e-mail: knobloch@math.uni-frankfurt.de

†Department of Mathematical Sciences, University of Bath, Claverton Down, Bath, BA2 7AY, U.K.  
e-mail: a.kyprianou@bath.ac.uk

We begin our exposition by briefly reviewing what is meant by a homogeneous fragmentation process, thereby introducing some notation.

## 1.1 homogeneous fragmentation processes

Below we give a brief overview of the definition and structure of a homogeneous fragmentation process. The reader is referred to Bertoin [6] for a more detailed overview. Let  $\mathcal{P}$  be the space of partitions of the natural numbers. Here a partition of  $\mathbb{N}$  is a sequence  $\pi = (\pi_1, \pi_2, \dots)$  of disjoint sets, called blocks, such that  $\bigcup_{i \in \mathbb{N}} \pi_i = \mathbb{N}$ . The blocks of a partition are enumerated in the increasing order of their least element, that is to say  $\min \pi_i \leq \min \pi_j$  when  $i \leq j$  (with the convention that  $\min \emptyset = \infty$ ). Now consider the measure  $\mu$  on  $\mathcal{P}$ , given by

$$\mu(d\pi) = \int_{\mathcal{S}} \varrho_{\mathbf{s}}(d\pi) \nu(d\mathbf{s}),$$

where  $\varrho_{\mathbf{s}}$  is the law of Kingman's paint-box based on  $\mathbf{s} \in \mathcal{S}$  (cf. page 98 of Bertoin [6]) with

$$\mathcal{S} := \left\{ \mathbf{s} = (s_1, s_2, \dots) : s_1 \geq s_2 \geq \dots \geq 0, \sum_{i=1}^{\infty} s_i \leq 1 \right\},$$

and the so-called *dislocation measure*  $\nu \neq 0$  is a measure on  $\mathcal{S}$  such that

$$\nu(\mathbf{s} \in \mathcal{S} : s_2 = 0) = 0 \tag{1}$$

as well as

$$\int_{\mathcal{S}} (1 - s_1) \nu(d\mathbf{s}) < \infty. \tag{2}$$

It is known that  $\mu$  is an exchangeable partition measure, meaning that it is invariant under the action of finite permutations on  $\mathcal{P}$ . It is also known (cf. Chapter 3 of Bertoin [6]) that it is possible to construct a fragmentation process on the space of partitions  $\mathcal{P}$  with the help of a Poisson point process  $\{(\pi(t), k(t)) : t \geq 0\}$  on  $\mathcal{P} \times \mathbb{N}$  which has intensity measure  $\mu \otimes \sharp$ , where  $\sharp$  is the counting measure. The aforementioned  $\mathcal{P}$ -valued fragmentation process is a Markov process which we denote by  $\Pi = \{\Pi(t) : t \geq 0\}$ , where  $\Pi(t) = (\Pi_1(t), \Pi_2(t), \dots) \in \mathcal{P}$  is such that at all times  $t \geq 0$  for which an atom  $(\pi(t), k(t))$  occurs in  $(\mathcal{P} \setminus (\mathbb{N}, \emptyset, \dots)) \times \mathbb{N}$ ,  $\Pi(t)$  is obtained from  $\Pi(t-)$  by partitioning the  $k(t)$ -th block into the sub-blocks  $(\Pi_{k(t)}(t-) \cap \pi_j(t) : j = 1, 2, \dots)$ . When  $\nu$  is a finite measure each block experiences an exponential holding time before it fragments.

Thanks to the properties of the exchangeable partition measure  $\mu$  it can be shown that for each  $t \geq 0$  the distribution of  $\Pi(t)$  is exchangeable and that the blocks of  $\Pi(t)$  have asymptotic frequencies in the sense that for each  $i \in \mathbb{N}$  the limit

$$|\Pi_i(t)| := \lim_{n \rightarrow \infty} \frac{1}{n} \sharp \{ \Pi_i(t) \cap \{1, \dots, n\} \}$$

exists almost surely. Moreover, Bertoin showed that  $|\Pi_i(t)|$  exists  $\mathbb{P}$ -a.s. simultaneously for all  $t \geq 0$  and  $i \in \mathbb{N}$ .

We denote the countable random jump times of  $\Pi$  by  $\mathcal{I} \subseteq \mathbb{R}_0^+$ . Further, let  $\mathcal{F} := (\mathcal{F}_t)_{t \in \mathbb{R}_0^+}$  denote the filtration generated by  $\Pi$ . In addition, let  $\mathcal{G} := (\mathcal{G}_t)_{t \in \mathbb{R}_0^+}$  be the sub-filtration generated by the asymptotic frequencies of  $\Pi$  and let  $\mathcal{F}^1 := (\mathcal{F}_t^1)_{t \in \mathbb{R}_0^+}$  denote the filtration generated by  $(\Pi_1(t))_{t \in \mathbb{R}_0^+}$ .

Let us define  $\xi(t) := -\log |\Pi_1(t)|$  for every  $t \geq 0$ , with the convention  $-\log 0 := \infty$ . Resorting to the Poissonian construction of the fragmentation process, Bertoin proved that  $\xi = \{\xi(t) : t \geq 0\}$  is a killed subordinator with cemetery state  $\infty$  and killing rate

$$\kappa := \int_{\mathcal{S}} \left( 1 - \sum_{k \in \mathbb{N}} s_k \right) \nu(ds). \quad (3)$$

Moreover, it is well known that its Laplace exponent  $\Phi$ , given by

$$e^{-\Phi(p)} := \mathbb{E}(e^{-p\xi(1)}),$$

can be characterised over an appropriate domain of  $p$  through the dislocation measure  $\nu$  as follows. Define the constant

$$\underline{p} = \inf \left\{ p \in \mathbb{R} : \int_{\mathcal{S}} \left| 1 - \sum_{i=1}^{\infty} s_i^{1+p} \right| \nu(ds) < \infty \right\}$$

which is necessarily in  $[-1, 0]$ . Then

$$\Phi(p) = \int_{\mathcal{S}} \left( 1 - \sum_{i=1}^{\infty} s_i^{1+p} \right) \nu(ds)$$

for all  $p > \underline{p}$  (and we understand  $\Phi(\underline{p}) = \Phi(\underline{p}+)$ ). The tagged fragment  $\Pi_1$ , and in particular its Laplace exponent  $\Phi$ , can be used to extract information about the decay and spatial distribution of blocks in the fragmentation process. A case in point concerns the asymptotic rate of decay of the largest block

$$\lambda_1(t) := \sup_{n \in \mathbb{N}} |\Pi_n(t)|, \quad t \geq 0.$$

To this end, note that  $\Phi$  is strictly increasing, concave and differentiable. We shall assume that

$$(p+1)\Phi'(p) > \Phi(p) \text{ for some } p \in (\underline{p}, \infty). \quad (4)$$

This assumption is automatically satisfied if there exists some  $p^* \geq \underline{p}$  with  $\Phi(p^*) = 0$ , hence in particular in the conservative case where  $\nu(\mathbf{s} \in \mathcal{S} : \sum_{k \in \mathbb{N}} s_k < 1) = 0$  and thus  $p^* = 0$ . Following the reasoning in the proof of Lemma 1 in [5] one may proceed with (4) in hand to show that there exists a unique maximal value of the function

$$p \mapsto c_p := \frac{\Phi(p)}{p+1}$$

in  $(\underline{p}, \infty)$ , which is achieved at some  $\bar{p} > \underline{p}$  and which is also equal to  $\Phi'(\bar{p})$ . This maximal value turns out to characterise the asymptotic rate of decay of the largest block, as shown in the following proposition that is lifted from Bertoin [6].

**Proposition 1** (cf. Corollary 1.4 of [6]). *We have*

$$\lim_{t \rightarrow \infty} \frac{-\log \lambda_1(t)}{t} = c_{\bar{p}}$$

*$\mathbb{P}$ -almost surely.*

In Corollary 1.4 of [6] Bertoin proves this result for fragmentation chains, but in view of Lemma 1.35 of [17] the same line of argument works for fragmentation processes.

## 1.2 Killed homogeneous fragmentation processes

Now let  $c > 0$  and  $x \in \mathbb{R}_0^+$ . We want to introduce killing of  $\Pi$  upon hitting the space–time barrier

$$\left\{ (y, t) \in \mathbb{R}_0^+ \times \mathbb{R}_0^+ : y < e^{-(x+ct)} \right\}$$

as follows. A block  $\Pi_n(t)$  is killed at the moment of its creation  $t \in \mathcal{I}$  if  $|\Pi_n(t)| < e^{-(x+ct)}$ , see Figure 1. Here, killing a block means that it is sent to a *cemetery state*, which we shall identify by  $\emptyset$ .




Figure 1: Realisation of a fragmentation process with finite dislocation measure without killing, in (a), and with killing, in (b).

Suppose that for  $t \geq 0$  we define  $\mathcal{N}_t^x$  to be the index set of the blocks in  $(\Pi_n(t))_{n \in \mathbb{N}}$  that are not yet killed by time  $t$ . It is important to note that  $N_t^x := \text{card}(\mathcal{N}_t^x)$  is finite for each  $t$ . Indeed, as  $\sum_{n \in \mathbb{N}} |\Pi_n(t)| \leq 1$  we infer that  $|\Pi_n(t)| \geq e^{-(x+ct)}$  for at most  $e^{x+ct}$ -many  $n \in \mathbb{N}$ . That is

$$N_t^x \leq e^{x+ct}$$

for all  $t \in \mathbb{R}_0^+$ . Denote by  $\Pi^x := (\Pi^x(t) : t \geq 0)$ , where  $\Pi^x(t) = (\Pi_n(t))_{n \in \mathcal{N}_t^x}$ , the resulting killed fragmentation process and note that  $\Pi^x$  is not  $\mathcal{P}$ -valued.

For each  $n \in \mathbb{N}$  the block of  $\Pi^x$  containing  $n$  has a killing time that may be finite or infinite. Note that the killed fragmentation process  $\Pi^x$  also depends on the constant  $c > 0$ . However, in order to keep the notation as simple as possible we do not include the parameter  $c$  in the notation as this constant does not change within the results or proofs of this paper.

In this paper we shall answer the question whether it is possible that the supremum over all the aforementioned respective individual killing times, which is henceforth denoted by  $\zeta^x$ , is finite. We

say that  $\Pi^x$  becomes *extinct* if  $\{\zeta^x < \infty\}$ . Note that  $\zeta^x > 0$   $\mathbb{P}$ -a.s., that is to say almost surely instantaneous extinction is not possible, on account of the fact that the spectrally negative Lévy process  $\{ct + \log |\Pi_1(t)| : t \geq 0\}$  is irregular for  $(-\infty, 0)$  when issued from the origin. Our first main result in this respect is the following.

**Theorem 2.** *For all  $c \leq c_{\bar{p}}$  we have  $\mathbb{P}(\zeta^x < \infty) = 1$  for every  $x \in \mathbb{R}_0^+$ . If  $c > c_{\bar{p}}$ , then  $x \mapsto \mathbb{P}(\zeta^x < \infty)$  is a nonincreasing,  $(0, 1)$ -valued function on  $\mathbb{R}_0^+$ .*

In the case that extinction does not occur with probability 1, we shall give two qualitative results concerning the evolution of the process on survival. The first result shows that the total number of fragments in the surviving process explodes.

**Theorem 3.** *Let  $c > c_{\bar{p}}$ . Then we have that*

$$\limsup_{t \rightarrow \infty} N_t^x = \infty$$

*holds  $\mathbb{P}(\cdot | \zeta^x = \infty)$ -a.s. for any  $x \in \mathbb{R}_0^+$ .*

The second result shows that the asymptotic exponential rate of decay of the largest fragment,

$$\lambda_1^x(t) := \max_{n \in \mathbb{N}} |\Pi_n^x(t)|, \quad t \geq 0,$$

is the same as when the killing scheme is not in effect, cf. Proposition 1.

**Theorem 4.** *Let  $c > c_{\bar{p}}$  and  $x \in \mathbb{R}_0^+$ . Then we have*

$$\lim_{t \rightarrow \infty} \frac{-\log \lambda_1^x(t)}{t} = c_{\bar{p}}$$

*$\mathbb{P}(\cdot | \zeta^x = \infty)$ -almost surely.*

What lies fundamentally behind the proofs of our main results is a detailed study of the interaction between two classes of martingales.

The outline of this paper is as follows. In the next section we provide some general notions that are used in the subsequent parts of the present paper and in particular we employ the connection between fragmentations and Lévy processes. Section 3 is concerned with the proof of Theorem 2 and in Section 4 we provide the proof of Theorem 3. Then, in Section 5, we introduce a multiplicative process and examine when this process is a martingale. The object under consideration in Section 6 is an additive process which also turns out to be a martingale and whose limit we study with regard to strict positivity. In the final section of this paper we prove Theorem 4.

## 2 Preliminaries

Let  $B_n(t)$ ,  $t \in \mathbb{R}_0^+$ , denote the block in  $\Pi(t)$  that contains the element  $n \in \mathbb{N}$  and recall from (3) that under  $\mathbb{P}$  the process  $\xi_n = (-\log |B_n(t)|)_{t \in \mathbb{R}_0^+}$  is a killed subordinator (with cemetery state  $+\infty$  and killing rate  $\kappa$ ).

**Definition 5.** For every  $n \in \mathbb{N}$  let the process  $X_n := (X_n(t))_{t \in \mathbb{R}_0^+}$  be defined by

$$X_n(t) := ct - \xi_n(t)$$

for all  $t \in \mathbb{R}_0^+$ .

Notice that under  $\mathbb{P}$  the dynamics of the process  $X_n$  are those of a killed spectrally negative Lévy process of bounded variation (with cemetery state  $-\infty$  and killing rate  $\kappa$ ). Moreover, the jump times of  $X_n$ , henceforth denoted by the countably infinite set,  $\mathcal{I}_n \subseteq \mathbb{R}^+$ , are the set of dislocation times of  $(B_n(t))_{t \in \mathbb{R}_0^+}$ . That is,  $X_n$  jumps exactly when the subordinator  $\xi_n$  jumps. For any  $n \in \mathbb{N}$  and  $x \in \mathbb{R}_0^+$  consider the following  $\mathcal{F}$ -stopping times:

$$\tau_{n,x}^+ := \inf\{t \in \mathbb{R}_0^+ : X_n(t) > x\} \quad \text{as well as} \quad \tau_{n,x}^- := \inf\{t \in \mathbb{R}_0^+ : X_n(t) < -x\}.$$

For any  $p \in (\underline{p}, \infty)$  consider the change of measure given by

$$\left. \frac{d\mathbb{P}^{(p)}}{d\mathbb{P}} \right|_{\mathcal{F}_t} = e^{\Phi(p)t - p\xi(t)} = e^{pX_1(t) - \psi(p)t}, \quad (5)$$

where

$$\psi(p) = \frac{1}{t} \log \mathbb{E}(e^{pX_1(t)}) = cp - \Phi(p)$$

is the Laplace exponent of  $X_1$ . Moreover, considering the projection of (5) onto the sub-filtration  $\mathcal{G}$  results in

$$\left. \frac{d\mathbb{P}^{(p)}}{d\mathbb{P}} \right|_{\mathcal{G}_t} = M_t(p) := \sum_{n \in \mathbb{N}} |\Pi_n(t)|^{1+p} e^{\Phi(p)t},$$

for all  $p \in (\underline{p}, \infty)$  and  $t \in \mathbb{R}_0^+$ .

**Remark 6.** Let  $p \in (\underline{p}, \bar{p})$  and denote by  $M_\infty(p)$  the  $\mathbb{P}$ -a.s. limit of the nonnegative martingale  $M(p) := (M_t(p))_{t \in \mathbb{R}_0^+}$ . According to Theorem 1 of [7] (cf. also Theorem 4 of [8] for the conservative case) the unit-mean martingale  $M(p)$  is uniformly integrable. Hence,  $\mathbb{E}(M_\infty(p)) = 1$  and thus  $\mathbb{P}^{(p)}$  is a probability measure on  $\mathcal{G}_\infty := \bigcup_{t \in \mathbb{R}_0^+} \mathcal{G}_t$ . Moreover, using that  $\mathbb{E}(M_\infty(p)) > 0$  one obtains that  $M_\infty(p) > 0$   $\mathbb{P}$ -a.s., see Lemma 1.35 of [17] (or Theorem 2 of [5] for the conservative case). Consequently, restricted to the  $\sigma$ -algebra  $\mathcal{G}_\infty$ , the measures  $\mathbb{P}^{(p)}$  and  $\mathbb{P}$  are equivalent.  $\diamond$

Corollary 3.10 in [19] shows that under the measure  $\mathbb{P}^{(p)}$  the process  $X_1$  is again a spectrally negative Lévy process such that

$$\psi_p(\lambda) := \frac{1}{t} \log \mathbb{E}^{(p)}(e^{\lambda X_1(t)}) = \psi(\lambda + p) - \psi(p) = c\lambda - \Phi(\lambda + p) + \Phi(p) \quad (6)$$

for all  $\lambda > \underline{p} - p$ . Let  $W_p$  be the scale function of the spectrally negative Lévy process  $X_1$  under  $\mathbb{P}^{(p)}$ . That is to say,  $W_p$  is the unique increasing and continuous function on  $(0, \infty)$  that is defined through the Laplace transform

$$\int_0^\infty e^{-\lambda x} W_p(x) dx = \frac{1}{\psi_p(\lambda)},$$

for all  $\lambda > p - \underline{p}$ .

A fundamental identity involving the scale function  $W_p$  that we shall appeal to later is the following result taken from Theorem 8.1, equation (8.7), in [19]:

$$\mathbb{P}^{(p)}(\tau_{1,x}^- = \infty) = (\psi_p'(0+) \vee 0)W_p(x) \quad (7)$$

for all  $x > 0$ . Another important fact that we shall also use concerns the value of  $W_p$  at zero. Indeed, thanks to the fact that  $X_1$  has paths of bounded variation, it turns out that for all  $p \geq 0$ ,  $W_p(0+) = 1/c$ . See for example Lemma 8.6 in [19].

An important role in what follows will be played by  $X_n$  killed upon hitting  $(-\infty, -x)$  for  $n \in \mathbb{N}$  and  $x \in \mathbb{R}_0^+$ . For  $t \in \mathbb{R}_0^+$  set

$$X_n^x(t) := (X_n(t) + x)\mathbb{1}_{\{\tau_{n,x}^- > t\}} = (x + ct + \log |B_n(t)|)\mathbb{1}_{\{\tau_{n,x}^- > t\}}.$$

### 3 Properties of the extinction probability

In this section we prove Theorem 2 by dealing with the cases  $c \in (0, c_{\bar{p}}]$  and  $c > c_{\bar{p}}$  as two separate lemmas. The first lemma below deals with the easier, but less interesting, case that  $c \in (0, c_{\bar{p}}]$ .

**Lemma 7.** *Let  $c \in (0, c_{\bar{p}}]$ . Then  $\mathbb{P}(\zeta^x < \infty) = 1$  for all  $x \in \mathbb{R}_0^+$ .*

*Proof.* Using stochastic monotonicity it suffices to consider the case that  $c = c_{\bar{p}} = \Phi(\bar{p})/(1 + \bar{p})$ . It was shown in Theorem 4 in [8] (cf. also Theorem 1 in [7]) that  $M_t(\bar{p}) \rightarrow 0$   $\mathbb{P}$ -a.s. as  $t \rightarrow \infty$ . Since  $M_t(\bar{p}) \geq e^{\Phi(\bar{p})t} \lambda_1^{1+\bar{p}}(t)$  for all  $t \in \mathbb{R}_0^+$ , we thus deduce that

$$(c_{\bar{p}}t + \log(\lambda_1(t))) \rightarrow -\infty$$

as  $t \rightarrow \infty$  and hence  $\mathbb{P}(\zeta^x < \infty) = 1$  for all  $x \in \mathbb{R}_0^+$ .  $\square$

Notice that the statement of the previous lemma is obvious for  $c \in (0, c_{\bar{p}})$  as the asymptotic decay of the largest fragment in the non-killed setting is given by  $c_{\bar{p}}$ , see Proposition 1, and thus the fragmentation process eventually crosses the killing line almost surely. However, for the critical value  $c = c_{\bar{p}}$  this argument does not work as one needs to rule out the possibility that the largest fragment could approach the killing line without intersecting it.

The following result deals with the more interesting case that  $c > c_{\bar{p}}$ .

**Lemma 8.** *Let  $c > c_{\bar{p}}$ . Then*

$$\mathbb{P}(\zeta^x < \infty) \in (0, 1)$$

for all  $x \in \mathbb{R}_0^+$ .

*Proof.* The proof is divided into two parts. The first part shows that  $\mathbb{P}(\zeta^x < \infty) < 1$  and the second part proves that  $\mathbb{P}(\zeta^x < \infty) > 0$  for all  $x \in \mathbb{R}_0^+$ .



Part I. Note that  $c > c_{\bar{p}} = \Phi'(\bar{p})$  and hence, since  $\Phi'$  is continuous, we may always choose  $p \in (\underline{p}, \bar{p})$  such that  $c > \Phi'(p)$ . In that case  $\psi'_p(0+) = \psi'(p) = c - \Phi'(p) > 0$ . Hence, by means of the nondecreasingness of  $\mathbb{P}(\zeta^{(\cdot)} = \infty)$ , we deduce from (7) that

$$\mathbb{P}^{(p)}(\zeta^x = \infty) \geq \mathbb{P}^{(p)}(\tau_{1,0}^- = \infty) = \psi'_p(0+)W_p(0+) = \frac{\psi'_p(0+)}{c} \in (0, 1)$$

for all  $x \in \mathbb{R}_0^+$ . According to Remark 6 this results in

$$\mathbb{P}(\zeta^x = \infty) > 0, \quad \text{i.e.} \quad \mathbb{P}(\zeta^x < \infty) < 1.$$

Part II. Let  $x \in \mathbb{R}_0^+$ . In order to show that  $\mathbb{P}(\zeta^x < \infty) > 0$  we fix some  $a > x$  and some  $y_0 \in (1/2 \vee (1 - e^{-a}), 1)$  such that

$$q := \mu(\pi \in \mathcal{P} : |\pi_1| \in (0, y_0]) \in (0, \infty).$$

The last inclusion is possible since, on the one hand,  $\mu(\pi \in \mathcal{P} : |\pi_1| \in (0, y_0]) = \mu(\pi \in \mathcal{P} : -\log |\pi_1| \geq -\log y_0)$  is the tail of the jump measure of the subordinator  $\xi$  which is necessarily finite. On the other hand, there exists some  $x \in (0, 1)$  such that

$$\mu(\pi \in \mathcal{P} : |\pi_1| \in (0, x]) > 0,$$

as otherwise the Lévy measure of  $\xi$  has no mass in  $(0, \infty)$  which contradicts the fact that  $\xi$  is a subordinator.

Recall that  $\{\pi(t) : t \in \mathcal{I}_1\}$  are the atoms of the Poisson point process on  $\mathcal{P}$  that determines  $\xi$ . Further, denote the (possibly infinite) killing time of  $\xi$  by  $\tau_\xi$  and recall from (3) that  $\tau_\xi$  is independent of the dynamics of the process  $\xi$ , up to its moment of killing, and exponentially distributed with parameter  $\kappa$ . Moreover, by means of Proposition 2 in Section 0.5 of [4] we have that  $\tau(y_0) := \inf\{t \in \mathcal{I}_1 : |\pi_1(t)| \in (0, y_0]\}$  is exponentially distributed with parameter  $q$ . It is straightforward to check that every block which does not contain 1 and which is produced at some dislocation before the time  $\tau(y_0) \wedge \tau_\xi$  of the block containing 1 will be no larger than a proportion  $e^{-a}$  of its parent. This follows directly from the inequality that for all  $t \in \mathcal{I}_1$  with  $t < \tau(y_0)$  and all  $j \in \mathbb{N} \setminus \{1\}$ ,

$$|\pi_j(t)| \leq \sum_{n \in \mathbb{N} \setminus \{1\}} |\pi_n(t)| \leq 1 - |\pi_1(t)| \leq 1 - y_0 \leq e^{-a}.$$

The classical Thinning Theorem for Poisson point processes (e.g. Proposition 2 in Section 0.5 of [4]) allows us to conclude that  $(X_1^x(u))_{u \in [0, \tau(y_0) \wedge \tau_\xi]}$  has the law of a Lévy process, say  $\tilde{X}_1^x$ , which is the difference of a linear drift with constant rate  $c$  and a driftless subordinator with Lévy measure  $\mu(\pi \in \mathcal{P} : -\log |\pi_1| \in dx) |_{(0, -\log y_0]}$ , sampled up to a time which is the minimum of two independent and exponentially distributed random times, say  $\mathbf{e}_q$  and  $\mathbf{e}_\kappa$ , with respective rates  $q$  and  $\kappa$ .

Now define

$$R^{(q+\kappa)}(a, x, dy) = \int_0^\infty e^{-(q+\kappa)t} dt \cdot \mathbb{P} \left( \tilde{X}_1^x(t) \in dy, \sup_{s \leq t} \tilde{X}_1^x(s) \leq a, \inf_{s \leq t} \tilde{X}_1^x(s) \geq 0 \right), \quad y \in (0, a).$$

Theorem 8.7 in [19] shows that  $R^{(q+\kappa)}(a, x, dy)$  is absolutely continuous with strictly positive Lebesgue density in the neighbourhood of the origin (this is at least immediately obvious for

$y \in (0, x)$  by inspecting the expression for the resolvent in the aforementioned theorem). A little thought in light of the above remarks reveals that, on the event  $\{\sup_{s < \tau(y_0) \wedge \tau_\xi} X_1^x(s) \leq a, \inf_{s < \tau(y_0) \wedge \tau_\xi} X_1^x(s) \geq 0\}$ , the process  $(X_1^x(u))_{u \in [0, \tau(y_0) \wedge \tau_\xi]}$  describes (on the negative-logarithmic scale and relative to the killing barrier) the **only** surviving block in the process  $\Pi^x$  over the time horizon  $[0, \tau(y_0) \wedge \tau_\xi]$ .

Using these facts, as well as the observation that  $\tau(y_0)$  is almost surely not a jump time for  $\tilde{X}_1^x$ , we now have the estimate

$$\begin{aligned} \mathbb{P}(\zeta^x < \infty) &\geq \mathbb{P}\left(X_1^x(\tau(y_0)-) \in [0, -\log(y_0)], \sup_{s < \tau(y_0)} X_1^x(s) \leq a, \inf_{s < \tau(y_0)} X_1^x(s) \geq 0, \tau(y_0) < \tau_\xi\right) \\ &\geq \mathbb{P}\left(\tilde{X}_1^x(\mathbf{e}_q-) \in [0, -\log(y_0)], \sup_{s < \mathbf{e}_q} \tilde{X}_1^x(s) \leq a, \inf_{s < \mathbf{e}_q} \tilde{X}_1^x(s) \geq 0, \mathbf{e}_q < \mathbf{e}_\kappa\right) \\ &\geq \mathbb{E}\left(e^{-\kappa \mathbf{e}_q}; \tilde{X}_1^x(\mathbf{e}_q-) \in [0, -\log(y_0)], \sup_{s < \mathbf{e}_q} \tilde{X}_1^x(s) \leq a, \inf_{s < \mathbf{e}_q} \tilde{X}_1^x(s) \geq 0\right) \\ &= qR^{(q+\kappa)}(a, x, [0, -\log(y_0)]) > 0 \end{aligned}$$

as required.  $\square$

## 4 Explosion of the number of blocks on survival

In this section we provide the proof of Theorem 3. To this end, we shall use the following auxiliary lemma which states that for any  $n \in \mathbb{N}$  there exists a time such that with positive probability the fragmentation process has at least  $n$  blocks. More precisely, we have the following result.

**Lemma 9.** *Let  $c > c_{\bar{p}}$ . Then for any  $n \in \mathbb{N}$  there exists a  $t > 0$  such that*

$$\mathbb{P}(N_t^0 \geq n) > 0. \quad (8)$$

*Proof.* In the first part of the proof we show that the probability of the event  $\{N_t^0 \geq 2\}$  is positive for some  $t \in \mathbb{R}_0^+$  and in the second part we use this in conjunction with an induction argument to prove the assertion.

Part I. Let us first show that there exists some  $z_0 \in (1/2, 1)$  such that

$$\mu(\pi \in \mathcal{P} : |\pi|_2^\downarrow \geq 1 - z_0, |\pi_1| > 0) > 0, \quad (9)$$

where  $\{|\pi|_i^\downarrow : i \geq 1\}$  represents the asymptotic frequencies of  $\pi \in \mathcal{P}$  when ranked in descending order. To this end, assume  $\mu(\pi \in \mathcal{P} : |\pi|_2^\downarrow \geq a, |\pi_1| > 0) = 0$  for all  $a \in (0, 1)$ . This assumption implies that  $\mu(\pi \in \mathcal{P} : |\pi|_2^\downarrow \neq 0, |\pi_1| > 0) = 0$ , which in view of (1) results in  $\mu(\pi \in \mathcal{P} : |\pi_1| > 0) = 0$  and thus contradicts  $\nu \neq 0$ . Consequently, there exists some  $z_0 \in (1/2, 1)$  such that (9) holds. Next note that, on account of the inequality  $|\pi|_1^\downarrow + |\pi|_2^\downarrow \leq 1$ ,

$$p := \mu(\pi \in \mathcal{P} : |\pi|_1^\downarrow \leq z_0, |\pi_1| > 0) \geq \mu(\pi \in \mathcal{P} : |\pi|_2^\downarrow \geq 1 - z_0, |\pi_1| > 0) > 0.$$

Observe that  $\nu(\mathbf{s} \in \mathcal{S} : s_1 \in (0, z_0]) < \infty$ , as otherwise

$$\int_{\mathcal{S}} (1 - s_1) \nu(d\mathbf{s}) \geq \int_{\{\mathbf{s} \in \mathcal{S} : s_1 \in (0, z_0]\}} (1 - s_1) \nu(d\mathbf{s}) \geq (1 - z_0) \nu(\mathbf{s} \in \mathcal{S} : s_1 \in (0, z_0]) = \infty,$$

which contradicts (2). Therefore, we infer from formula (3) in [16] that

$$p \leq \mu(\pi \in \mathcal{P} : |\pi|_1^\downarrow \leq z_0) \leq \nu(\mathbf{s} \in \mathcal{S} : s_1 \in (0, z_0]) < \infty.$$

Now let  $\eta(z_0) := \inf\{t \in \mathcal{I}_1 : |\pi(t)|_1^\downarrow \leq z_0, |\pi_1(t)| > 0\}$ . The classical Thinning Theorem for Poisson point processes shows that  $|\pi(\eta(z_0))|_2^\downarrow$  and  $\eta(z_0)$  are independent and that

$$\mathbb{P}\left(|\pi(\eta(z_0))|_2^\downarrow \geq 1 - z_0\right) = \frac{\mu(\pi \in \mathcal{P} : |\pi|_2^\downarrow \geq 1 - z_0, |\pi_1| > 0)}{\mu(\pi \in \mathcal{P} : |\pi|_1^\downarrow \leq z_0, |\pi_1| > 0)} > 0. \quad (10)$$

Moreover  $|\pi(\eta(z_0))|_2^\downarrow$  and  $(X_1^x(u))_{u \in [0, \eta(z_0))}$  are independent and  $\eta(z_0)$  is exponentially distributed with parameter  $p$ . Now let  $\hat{X}_1^x$  be a spectrally negative Lévy process, shifted by  $x \in \mathbb{R}_0^+$ , which is written as the difference of a linear drift with rate  $c$  and a driftless subordinator with Lévy measure  $\mu(\pi \in \mathcal{P} : |\pi|_1^\downarrow > z_0; -\log |\pi_1| \in dy), y > 0$ , which is independent of all other previously mentioned random objects.

We want to work with its resolvent on the half line

$$R^{(p+\kappa)}(x, dy) := \int_0^\infty e^{-(p+\kappa)t} dt \cdot \mathbb{P}(\hat{X}_1^x(t) \in dy, \inf_{s \leq t} \hat{X}_1^x(s) \geq 0), \quad y > 0$$

which is known to have a strictly positive density for all  $x \geq 0$ , cf. Corollary 8.8 of [19]. Note that in the case  $x = 0$  the process  $\hat{X}_1^x$  will take a strictly positive amount of time to exit the domain  $[0, \infty)$  on account of path irregularity, see the introduction of Chapter 8 of [19]. Let  $\mathbf{e}_p$  and  $\mathbf{e}_\kappa$  be two independent (of everything) exponentially distributed random variables with respective rates  $p$  and  $\kappa$ . Since almost surely neither  $\mathbf{e}_p$  nor  $\mathbf{e}_\kappa$  is a jump time for  $\hat{X}_1^0$ , it follows from (10) that

$$\begin{aligned} & \mathbb{P}\left(N_{\eta(z_0)}^0 \geq 2\right) \\ & \geq \mathbb{P}\left(X_1^0(\eta(z_0)-) > -\log(1 - z_0), |\pi(\eta(z_0))|_2^\downarrow \geq 1 - z_0, \eta(z_0) < \tau_\xi\right) \\ & = \mathbb{E}\left(e^{-\kappa \mathbf{e}_p}; \hat{X}_1^0(\mathbf{e}_p-) > -\log(1 - z_0), \inf_{s < \mathbf{e}_p} \hat{X}_1^0(s) \geq 0\right) \mathbb{P}\left(|\pi(\eta(z_0))|_2^\downarrow \geq 1 - z_0\right) \\ & = pR^{(p+\kappa)}(0, (-\log(1 - z_0), \infty)) \frac{\mu(\pi \in \mathcal{P} : |\pi|_2^\downarrow \geq 1 - z_0, |\pi_1| > 0)}{\mu(\pi \in \mathcal{P} : |\pi|_1^\downarrow \leq z_0, |\pi_1| > 0)} > 0. \end{aligned}$$

Given that  $\eta(z_0)$  is exponentially distributed, it is now a standard argument to deduce that there must exist a  $t > 0$  such that

$$\mathbb{P}(N_t^0 \geq 2) > 0. \quad (11)$$

Part II. We prove (8) by resorting to the principle of mathematical induction. To this end, let  $n \in \mathbb{N}$ , fix some  $u_0 > 0$  such that (11) holds and, as the induction hypothesis, assume that

$$\mathbb{P}(N_{nu_0}^0 \geq n + 1) > 0.$$

To provide an estimate for  $\mathbb{P}(N_{(n+1)u_0}^0 \geq n+2)$  note that the event  $\{N_{(n+1)u_0}^0 \geq n+2\}$  contains the event that  $N_{nu_0}^0 \geq n+1$  and subsequently  $n$  of the blocks alive at time  $nu_0$  survive for a further  $u_0$  units of time, whilst one of the blocks at time  $nu_0$  succeeds in fragmenting further to produce at least two further particles  $u_0$  units of time later. A lower bound on the probability of the latter event that makes use of the fragmentation property and the monotonicity in  $x$  of  $\mathbb{P}(N_{nu_0}^x \geq n+1)$  and  $P(\zeta^x > u_0)$ , produces the estimate,

$$\mathbb{P}\left(N_{(n+1)u_0}^0 \geq n+2\right) \geq \mathbb{P}\left(N_{nu_0}^0 \geq n+1\right) \mathbb{P}\left(N_{u_0}^0 \geq 2\right) \mathbb{P}\left(\zeta^0 > u_0\right)^n > 0.$$

Coupled with (11), which closes the argument by induction, the proof of the lemma is complete.  $\square$

Having established the previous lemma we are now in a position to tackle the proof of Theorem 3.

*Proof of Theorem 3.* By Lemma 9, fix some  $k \in \mathbb{N}$  as well as  $t_0 > 0$  such that  $\mathbb{P}(N_{t_0}^0 \geq k) > 0$  and for every  $n \in \mathbb{N}$  and  $x \in \mathbb{R}_0^+$  define

$$E_n^x := \{\omega \in \Omega : N_{nt_0}^x(\omega) \geq k\}.$$

By means of the fragmentation property and the monotonicity in  $x$  of  $\mathbb{P}(N_{t_0}^x \geq k)$

$$\mathbb{P}(E_n^x | \mathcal{F}_{(n-1)t_0}) \geq \mathbb{P}(N_{t_0}^0 \geq k) > 0 \tag{12}$$

on  $\{\zeta^x = \infty\}$ . As a consequence of (12) we obtain that

$$\sum_{n \in \mathbb{N}} \mathbb{P}(E_n^x | \mathcal{F}_{(n-1)t_0}) = \infty \tag{13}$$

$\mathbb{P}$ -a.s. on  $\{\zeta^x = \infty\}$  for any  $x \in \mathbb{R}_0^+$ .

Since  $E_n^x$  is  $\mathcal{F}_{nt_0}$ -measurable, we can apply the extended Borel–Cantelli lemma (see e.g. Corollary (3.2) in Chapter 4 of [12] or Corollary 5.29 in [9]) to deduce that

$$\{E_n^x \text{ happens infinitely often}\} = \left\{ \sum_{n \in \mathbb{N}} \mathbb{P}(E_n^x | \mathcal{F}_{(n-1)t_0}) = \infty \right\},$$

and thus (13) shows that on the event  $\{\zeta^x = \infty\}$ ,  $x \in \mathbb{R}_0^+$ , the event  $E_n^x$  happens for infinitely many  $n \in \mathbb{N}$ . Consequently, we infer by monotonicity in  $x$  of  $N_t^x$  that

$$\mathbb{P}\left(\limsup_{t \rightarrow \infty} N_t^x \geq k \mid \zeta^x = \infty\right) = 1,$$

which proves the assertion on account of the fact that  $k$  may be taken arbitrarily large.  $\square$

## 5 Multiplicative martingales

Like many different types of spatial branching processes, the probability of extinction of our killed fragmentation process turns out to be intimately related to certain product martingales which we now introduce.

More specifically, the object under consideration in the present section is the stochastic process defined as follows. For any function  $f : \mathbb{R} \rightarrow [0, 1]$  and  $x \in \mathbb{R}_0^+$  let  $Z^{x,f} := \{Z_t^{x,f} : t \geq 0\}$  be given by

$$Z_t^{x,f} = \prod_{n \in \mathcal{N}_t^x} f(x + ct + \log |\Pi_n^x(t)|), \quad t \geq 0.$$

We are interested in understanding which functions  $f$  make the above process a martingale. In that case we refer to  $Z^{x,f}$  as a *multiplicative martingale*. The following theorem shows that within the class of nonincreasing functions which are valued zero at  $\infty$ , there is a unique choice of  $f$ .

**Theorem 10.** *Let  $c > c_{\bar{p}}$  and let  $f : \mathbb{R} \rightarrow [0, 1]$  be a monotone function. Then the following two statements are equivalent.*

(i) *For any  $x \in \mathbb{R}_0^+$  the process  $Z^{x,f}$  is a martingale with respect to the filtration  $\mathcal{F}$  and*

$$\lim_{x \rightarrow \infty} f(x) = 0.$$

(ii) *For all  $x \in \mathbb{R}_0^+$  :*

$$f(x) = \mathbb{P}(\zeta^x < \infty).$$

For any  $c > c_{\bar{p}}$  and  $t, x \in \mathbb{R}_0^+$  define

$$R_1^x(t) = x + ct + \log \lambda_1(t).$$

In order to prove Theorem 10 we shall use the following lemma which states that on survival of the killed fragmentation process the process  $(R_1^x(t))_{t \in \mathbb{R}_0^+}$  is unbounded.

**Lemma 11.** *Let  $c > c_{\bar{p}}$  and  $x \in \mathbb{R}_0^+$ . Then we have*

$$\limsup_{t \rightarrow \infty} R_1^x(t) = \infty$$

$\mathbb{P}(\cdot | \zeta^x = \infty)$ -almost surely.

*Proof.* Let  $z > x$  and set

$$\Gamma_z^x := \left\{ \omega \in \Omega : \inf\{t \in \mathbb{R}_0^+ : X_n^x(t)(\omega) \notin [0, z]\} = \infty \forall n \in \mathbb{N} \right\}.$$

Theorem 12 in Section VI.3 of [4] shows that the probability that a spectrally negative Lévy process never leaves the interval  $(0, z)$  when started in its interior is zero. Consequently, we have that

$$\tau_{n,x}^- < \tau_{n,z-x}^+ = \infty \quad \text{on} \quad \Gamma_z^x.$$

For each  $n \in \mathbb{N}$  set

$$\sigma_n := \inf\{t \in \mathbb{R}_0^+ : N_t^x \geq n\}$$

and note that Theorem 3 implies that  $\sigma_n$  is a  $\mathbb{P}$ -a.s. finite stopping time on  $\{\zeta^x = \infty\}$ . Let  $\tilde{\mathcal{N}}_t^x = \{n \in \mathbb{N} : X_n^x(t) \geq 0\}$  and introduce the equivalence relation  $\sim$  on  $\tilde{\mathcal{N}}_t^x$  such that  $n \sim m$  when  $n \in B_m(t)$ . The cardinality of  $\tilde{\mathcal{N}}_t^x / \sim$  is equal to  $N_t^x$ . Further, let  $p \in (\underline{p}, \bar{p})$ . By means of Lemma 8.6 of [19], we then infer from the strong fragmentation property and equation (8.8) of Theorem 8.1 of [19] that

$$\begin{aligned} \mathbb{P}^{(p)}(\Gamma_z^x | \mathcal{F}_{\sigma_n}) &\leq \prod_{k \in \tilde{\mathcal{N}}_{\sigma_n}^x / \sim} \mathbb{P}^{(p)}(\Gamma_z^y) \Big|_{y=X_k^x(\sigma_n)} \\ &= \prod_{k \in \tilde{\mathcal{N}}_{\sigma_n}^x / \sim} \mathbb{P}^{(p)}(\tau_{k,y}^- < \tau_{k,z-y}^+) \Big|_{y=X_k^x(\sigma_n)} \\ &= \prod_{k \in \tilde{\mathcal{N}}_{\sigma_n}^x / \sim} \left(1 - \frac{W_p(X_k^x(\sigma_k))}{W_p(z)}\right) \\ &\leq \left(1 - \frac{1}{cW_p(z)}\right)^{N_{\sigma_n}^x} \\ &\leq \left(1 - \frac{1}{cW_p(z)}\right)^n \end{aligned}$$

$\mathbb{P}^{(p)}$ -a.s. on  $\{\zeta^x = \infty\}$  for any  $n \in \mathbb{N}$ . Therefore, since  $\{R_1^x(s) < z \forall s \in \mathbb{R}_0^+\} = \Gamma_z^x$ , we have

$$\begin{aligned} \mathbb{P}^{(p)}\left(\left\{\sup_{s \in \mathbb{R}_0^+} R_1^x(s) < z\right\} \cap \{\zeta^x = \infty\}\right) &= \mathbb{P}^{(p)}(\Gamma_z^x \cap \{\zeta^x = \infty\}) \\ &= \lim_{n \rightarrow \infty} \mathbb{E}^{(p)}\left(\mathbb{P}^{(p)}(\Gamma_z^x \cap \{\zeta^x = \infty\} | \mathcal{F}_{\sigma_n})\right) \\ &= \mathbb{E}^{(p)}\left(\lim_{n \rightarrow \infty} \mathbb{P}^{(p)}(\Gamma_z^x \cap \{\zeta^x = \infty\} | \mathcal{F}_{\sigma_n})\right) \\ &= 0. \end{aligned}$$

From this last equality and the fact that  $z > x$  is arbitrary, one readily deduces that

$$\mathbb{P}^{(p)}\left(\left\{\limsup_{s \rightarrow \infty} R_1^x(s) < \infty\right\} \cap \{\zeta^x = \infty\}\right) = 0.$$

Since both events  $\{\limsup_{s \rightarrow \infty} R_1^x(s) < \infty\}$  and  $\{\zeta^x = \infty\}$  are  $\mathcal{G}_\infty$ -measurable, we therefore infer from Remark 6 that

$$\mathbb{P}\left(\left\{\limsup_{s \rightarrow \infty} R_1^x(s) < \infty\right\} \cap \{\zeta^x = \infty\}\right) = 0,$$

which proves the assertion.  $\square$

Let us now tackle the proof of Theorem 10.

*Proof of Theorem 10.* The proof is guided by a similar result in Harris et al. [15] for branching Brownian motion. We divide the proof into two parts. The first part proves the uniqueness of monotone functions  $f$  satisfying  $\lim_{y \rightarrow \infty} f(y) = 0$  for which  $Z^{x,f}$  is a martingale. Part II of the proof shows that the probability of extinction constitutes a function that makes  $Z^{x,f}$  a martingale.

Part I. By the martingale convergence theorem we have that  $Z^{x,f}$  being a nonnegative martingale implies that  $Z_\infty^{x,f} := \lim_{t \rightarrow \infty} Z_t^{x,f}$  exists  $\mathbb{P}$ -almost surely. Since the empty product equals 1 it is immediately clear that

$$Z_\infty^{x,f} = 1 \tag{14}$$

holds  $\mathbb{P}$ -a.s. on  $\{\zeta^x < \infty\}$ . Moreover, according to Lemma 11 we have that  $\limsup_{t \rightarrow \infty} R_1^x(t) = \infty$   $\mathbb{P}$ -a.s. on  $\{\zeta^x = \infty\}$ . Since  $\lim_{y \rightarrow \infty} f(y) = 0$ , we thus deduce that

$$0 \leq Z_\infty^{x,f} \leq \liminf_{t \rightarrow \infty} f(R_1^x(t)) = 0 \tag{15}$$

$\mathbb{P}$ -a.s. on  $\{\zeta^x = \infty\}$ . Hence, in view of (14) and (15) we infer that

$$Z_\infty^{x,f} = \mathbb{1}_{\{\zeta^x < \infty\}} \tag{16}$$

holds true  $\mathbb{P}$ -almost surely. As a consequence of  $Z^{x,f}$  being a bounded, and hence uniformly integrable, martingale we conclude from (16) that

$$f(x) = \mathbb{E}(Z_0^{x,f}) = \mathbb{E}(Z_\infty^{x,f}) = \mathbb{P}(\zeta^x < \infty).$$

Part II. Recalling Lemma 8, let  $g : \mathbb{R} \rightarrow (0, 1)$  be given by  $g(x) = \mathbb{P}(\zeta^x < \infty)$ . Since  $g$  is monotone and bounded, the limit  $g(+\infty) := \lim_{x \rightarrow \infty} g(x)$  exists in  $[0, 1)$ . Furthermore, for any  $t \in \mathbb{R}_0^+$  we have  $\lim_{x \rightarrow \infty} \mathbb{1}_{\mathcal{N}_t^x}(n) = 1$   $\mathbb{P}$ -a.s. for every  $n \in \mathbb{N}$ . In addition, we have that  $X_n^x(t) \rightarrow \infty$   $\mathbb{P}$ -a.s. for any  $n \in \mathbb{N}$  and  $t \in \mathbb{R}_0^+$  as  $x \rightarrow \infty$ . Resorting to the fragmentation property of  $\Pi$  we deduce that

$$g(x) = \mathbb{E}(\mathbb{P}(\zeta^x < \infty | \mathcal{F}_t)) = \mathbb{E}\left(\prod_{n \in \mathcal{N}_t^x} g(x + ct + \log |\Pi_n^x(t)|)\right) = \mathbb{E}(Z_t^{x,g}) \tag{17}$$

holds for all  $t \in \mathbb{R}_0^+$ . By means of the fragmentation property we thus have that

$$\mathbb{E}(Z_{t+s}^{x,g} | \mathcal{F}_t) = \prod_{n \in \mathcal{N}_t^x} g(x + ct + \log |\Pi_n^x(t)|) = Z_t^{x,g}$$

$\mathbb{P}$ -almost surely. Hence,  $Z^{x,g}$  is a  $\mathbb{P}$ -martingale. Moreover, by the Dominated Convergence Theorem, we deduce from (17) that

$$\begin{aligned} g(+\infty) &= \lim_{x \rightarrow \infty} \mathbb{E}\left(\prod_{n \in \mathcal{N}_t^x} g(x + ct + \log |\Pi_n^x(t)|)\right) \\ &= \mathbb{E}\left(\lim_{y \rightarrow \infty} \prod_{n \in \mathcal{N}_t^y} \lim_{x \rightarrow \infty} g(x)\right) \end{aligned}$$

$$= \mathbb{E} \left( \lim_{y \rightarrow \infty} \lim_{x \rightarrow \infty} g(x)^{N_t^y} \right).$$

Consequently,  $g(+\infty) \in \{0, 1\}$ . Since  $g$  is decreasing and  $g(x) \in (0, 1)$  for all  $x \in \mathbb{R}_0^+$ , this forces us to choose  $g(+\infty) = 0$ .  $\square$

## 6 Additive martingales

In this section we deal with an additive stochastic process  $M^x(p) := (M_t^x(p))_{t \in \mathbb{R}_0^+}$ ,  $p \in (\underline{p}, \bar{p})$ , that for  $c > c_{\bar{p}}$  and  $x \in \mathbb{R}_0^+$ , is given by

$$M_t^x(p) = \sum_{n \in \mathcal{N}_t^x} W_p(x + ct + \log |\Pi_n^x(t)|) e^{\Phi(p)t} |\Pi_n^x(t)|^{1+p}.$$

The main result of this section is the following theorem.

**Theorem 12.** *Let  $c > c_{\bar{p}}$  and let  $p \in (\underline{p}, \bar{p})$  be such that  $c > \Phi'(p)$ . Then the process  $M^x(p)$  is a nonnegative  $\mathcal{F}$ -martingale with  $\mathbb{P}$ -a.s. limit  $M_\infty^x(p)$ . Moreover, this martingale limit satisfies*

$$\mathbb{P}(\{M_\infty^x(p) = 0\} \Delta \{\zeta^x < \infty\}) = 0,$$

where  $\Delta$  denotes the symmetric difference.

The following lemma is a version of the so-called *many-to-one identity*. To state it, let us introduce, for each  $n \in \mathbb{N}$  and  $t \geq 0$ , the notation  $\{\overleftarrow{\Pi}_n(s) : s \leq t\}$  to mean ancestral evolution of the block  $\Pi_i(t)$ . That is to say, if  $\Pi_n(t)$  is the block containing  $k \in \mathbb{N}$ , then  $\{\overleftarrow{\Pi}_n(s) : s \leq t\} = \{B_k(s) : s \leq t\}$ .

**Lemma 13.** *We have*

$$\mathbb{E} \left( \sum_{n \in \mathbb{N}} |\Pi_n(t)| f(\{\overleftarrow{\Pi}_n(s) : s \leq t\}) \right) = \mathbb{E} (f(\{|\Pi_1(s)| : s \leq t\}))$$

for every  $t \in \mathbb{R}_0^+$  and  $f : \text{RCLL}([0, t], [0, 1]) \rightarrow \mathbb{R}_0^+$ , where RCLL denotes the space of càdlàg functions.

*Proof.* The proof follows directly as a consequence of the fact that  $\Pi_1(t)$  has the law of a size-biased pick from  $\Pi(t)$ . See for example Lemma 2 of Berestycki et al. [3].  $\square$

The next lemma establishes the first assertion of Theorem 12 in that it shows that under  $\mathbb{P}$  the process  $M^x(p)$  is a martingale for suitable  $c$  and  $p$ .

**Lemma 14.** *Let  $c > c_{\bar{p}}$  and let  $p \in (\underline{p}, \bar{p})$  be such that  $c > \Phi'(p)$ . Further, let  $x \in \mathbb{R}_0^+$ . Then the process  $M^x(p)$  is a  $\mathbb{P}$ -martingale with respect to the filtration  $\mathcal{F}$ .*



*Proof.* Let us first show that for any  $t \in \mathbb{R}_0^+$  the process  $(W_p(X_1^x(s))\mathbb{1}_{\{s < \tau_{1,x}^-\}})_{s \in \mathbb{R}_0^+}$  is a  $\mathbb{P}^{(p)}$ -martingale with respect to  $\mathcal{F}$ . It is a straightforward exercise using (6) to show that  $\psi'_p(0+) = c - \Phi'(p) > 0$ . By the Markov property of  $X_1$  under  $\mathbb{P}^{(p)}$  we thus infer from (7) that

$$\begin{aligned} \mathbb{E}^{(p)} \left( \mathbb{1}_{\{\tau_{1,x}^- = \infty\}} \middle| \mathcal{F}_s \right) &= \mathbb{P}^{(p)} \left( \tau_{1,y}^- = \infty \right) \Big|_{y=x+X_1(s)} \mathbb{1}_{\{s < \tau_{1,x}^-\}} \\ &= \psi'_p(0+) W_p(x + X_1(s)) \mathbb{1}_{\{s < \tau_{1,x}^-\}} \end{aligned} \quad (18)$$

holds  $\mathbb{P}^{(p)}$ -a.s. for any  $s \in \mathbb{R}_0^+$ . Note that the left-hand side of (18) defines a closed  $\mathbb{P}^{(p)}$ -martingale. Further, observe that  $x + X_1(s) = X_1^x(s)$  on the event  $\{s < \tau_{1,x}^-\}$ .

By means of Lemma 13 we deduce that

$$\begin{aligned} \mathbb{E}(M_t^x(p)) &= e^{\Phi(p)s} \mathbb{E} \left( \sum_{n \in \mathcal{N}_t^x} W_p(x + ct + \log |\Pi_n^x(t)|) e^{\Phi(p)t} |\Pi_n^x(t)|^{1+p} \right) \\ &= \mathbb{E} \left( W_p(X_1^x(t)) \mathbb{1}_{\{t < \tau_{1,x}^-\}} e^{\Phi(p)t - p\xi(t)} \right) \\ &= \mathbb{E}^{(p)} \left( W_p(X_1^x(t)) \mathbb{1}_{\{t < \tau_{1,x}^-\}} \right) \\ &= W_p(x) \end{aligned} \quad (19)$$

for all  $t \in \mathbb{R}_0^+$ , where the final equality is a consequence of the above-mentioned martingale property of  $(W_p(X_1^x(s))\mathbb{1}_{\{s < \tau_{1,x}^-\}})_{s \in \mathbb{R}_0^+}$ . In view of (19) we infer from the fragmentation property of  $\Pi$  that

$$\begin{aligned} \mathbb{E}(M_{t+s}^x(p) | \mathcal{F}_t) &= \sum_{n \in \mathcal{N}_t^x} e^{\Phi(p)t} |\Pi_n^x(t)|^{1+p} \mathbb{E}(M^{(n)} | \mathcal{F}_t) \\ &= \sum_{n \in \mathcal{N}_t^x} e^{\Phi(p)t} |\Pi_n^x(t)|^{1+p} W_p(x + ct + \log |\Pi_n^x(t)|) \\ &= M_t^x(p) \end{aligned}$$

$\mathbb{P}$ -a.s. for all  $s, t \in \mathbb{R}_0^+$ , where conditional on  $\mathcal{F}_t$  the  $M^{(n)}$  are independent and satisfy

$$\mathbb{P} \left( M^{(n)} \in \cdot \middle| \mathcal{F}_t \right) = \mathbb{P} \left( M_s^y(p) \in \cdot \right) \Big|_{y=x+ct+\log |\Pi_n^x(t)|}$$

$\mathbb{P}$ -almost surely. □

Let us now turn to the proof of Theorem 12. The main ingredient in the proof of Theorem 12 turns out to be Theorem 10, which deals with the product martingale  $Z^{x,f}$ .

*Proof of Theorem 12.* According to Lemma 14 we have that  $M^x(p)$  is a nonnegative martingale and by the Martingale Convergence Theorem it follows that  $M_\infty^x(p) := \lim_{t \rightarrow \infty} M_t^x(p)$  exists  $\mathbb{P}$ -almost surely. It remains to show that the symmetric difference  $\{M_\infty^x(p) = 0\} \Delta \{\zeta^x < \infty\}$  is a  $\mathbb{P}$ -null set.

Define the function  $g_p : \mathbb{R}_0^+ \rightarrow [0, 1]$  given by

$$g_p(x) = \mathbb{P}(M_\infty^x(p) = 0)$$

for any  $x \in \mathbb{R}_0^+$ . Resorting to the fragmentation property we deduce that

$$\mathbb{P}(M_\infty^x(p) = 0 | \mathcal{F}_t) = \prod_{n \in \mathcal{N}_t^x} g_p(x + ct + \log |\Pi_n^x(t)|) = Z_t^{x, g_p}$$

holds  $\mathbb{P}$ -almost surely for all  $t \in \mathbb{R}_0^+$ . Therefore,  $Z^{x, g_p}$  is a  $\mathbb{P}$ -martingale. Note also that, thanks to the fact that both  $\mathcal{N}_t^x$  and  $W_p(x)$  are monotone increasing in  $x$ , for all  $\epsilon > 0$ ,  $M_\infty^{x+\epsilon}(p) \geq M_\infty^x(p)$  and hence  $g_p(\cdot)$  is a monotone function. It follows that  $g_p(+\infty)$  exists in  $[0, 1]$  and moreover, by taking the expectation and then the limit as  $x \rightarrow \infty$  in (23), we infer that

$$g_p(+\infty) \in \{0, 1\} \tag{20}$$

as otherwise we are led to the contradictory statement that  $g_p(+\infty) < g_p(+\infty)$ . Recall the martingale  $M(p)$  that we defined in Remark 6. Taking account of (7) we have that  $M_\infty^x(p) \leq M_\infty(p)/\psi'_p(0+)$ . Hence, since  $M(p)$  is an  $L^q$ -convergent martingale (cf. Theorem 2 of [5] and Proposition 3.5 of [17]) for some  $q > 1$ , it follows that  $M^x(p)$  is too. Coupled with the stochastic monotonicity of  $M_\infty^x(p)$  in  $x$ , this implies in view of (20) that necessarily  $g_p(+\infty) = 0$ .

We may now apply Theorem 10 and infer that  $g_p(x) = \mathbb{P}(\zeta^x < \infty)$ . Since  $\{\zeta^x < \infty\} \subseteq \{M_\infty^x(p) = 0\}$  for each  $x > 0$  this implies that

$$\mathbb{P}(\{\zeta^x < \infty\} \triangle \{M_\infty^x(p) = 0\}) = 0$$

for every  $x > 0$  as required.  $\square$

## 7 Exponential decay rate of the largest fragment

The final section of this paper is devoted to the proof of Theorem 4. That is, in this section we deal with the asymptotic behaviour of the largest fragment in the killed fragmentation process.

*Proof of Theorem 4.* Our approach is based on the method of proof for Corollary 1.4 in [6] and makes use of the martingale  $M^x(p)$  that we considered in the previous section.

First note that since  $\lambda_1^x(t) \leq \lambda_1(t)$ , it follows from Proposition 1 that  $\mathbb{P}$ -a.s.,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log(\lambda_1^x(t)) \leq c_{\bar{p}}. \tag{21}$$

Recall that we are assuming  $c > c_{\bar{p}}$ . In order to deal with the liminf set

$$\hat{p} := \inf \{p \in (\underline{p}, \bar{p}) : \Phi'(p) < c\},$$

and let  $p \in (\hat{p}, \bar{p})$  as well as  $\epsilon \in (0, p - \hat{p})$ . Since

$$\psi'_p(0+) = c - \Phi'(p) > \psi'_{p-\epsilon}(0+) = c - \Phi'(p - \epsilon) > c - \Phi'(\hat{p}) \geq 0,$$

where  $\Phi'(-\infty) := 0$ , we infer from (7) that the scale functions  $W_p$  and  $W_{p-\epsilon}$  are uniformly bounded from above by  $1/\psi'_p(0+)$  and  $1/\psi'_{p-\epsilon}(0+)$ , respectively. Moreover, according to Lemma 8.6 in [19]

we have  $W_p(0+) = W_{p-\epsilon}(0+) = c^{-1}$ . Hence, there exists a constant  $K > 0$  such that  $W_p(y) \leq KW_{p-\epsilon}(y)$  for all  $y \geq 0$ . Observe that

$$\begin{aligned} M_t^x(p) &= \sum_{n \in \mathcal{N}_t^x} W_p(x + ct + \log |\Pi_n^x(t)|) e^{\Phi(p)t} |\Pi_n^x(t)|^{1+p} \\ &\leq K e^{(\Phi(p) - \Phi(p-\epsilon))t} [\lambda_1^x(t)]^\epsilon e^{\Phi(p-\epsilon)t} \sum_{n \in \mathcal{N}_t^x} W_{p-\epsilon}(x + ct + \log |\Pi_n^x(t)|) |\Pi_n^x(t)|^{1+p-\epsilon} \\ &= K e^{(\Phi(p) - \Phi(p-\epsilon))t} [\lambda_1^x(t)]^\epsilon M_t^x(p-\epsilon). \end{aligned} \tag{22}$$

According to Theorem 12 we have that both  $M_\infty^x(p-\epsilon)$  and  $M_\infty^x(p)$  are  $(0, \infty)$ -valued  $\mathbb{P}(\cdot | \zeta^x = \infty)$ -almost surely. Consequently, taking the logarithm, dividing by  $t$  and taking the limit inferior as  $t \rightarrow \infty$  we thus deduce from (22) that

$$\liminf_{t \rightarrow \infty} \frac{1}{t} \log(\lambda_1^x(t)) \geq -\frac{\Phi(p) - \Phi(p-\epsilon)}{\epsilon}$$

$\mathbb{P}(\cdot | \zeta^x = \infty)$ -almost surely. Therefore, we have

$$\liminf_{t \rightarrow \infty} \frac{1}{t} \log(\lambda_1^x(t)) \geq -\lim_{\epsilon \rightarrow 0} \frac{\Phi(p) - \Phi(p-\epsilon)}{\epsilon} = -\Phi'(p) \tag{23}$$

$\mathbb{P}(\cdot | \zeta^x = \infty)$ -almost surely. Letting  $p \rightarrow \bar{p}$  and resorting to the fact that  $\Phi$  is the Laplace exponent of  $\xi$ , which ensures the continuity of  $\Phi'$ , (23) results in

$$\liminf_{t \rightarrow \infty} \frac{1}{t} \log(\lambda_1^x(t)) \geq -\Phi'(\bar{p}) \tag{24}$$

$\mathbb{P}(\cdot | \zeta^x = \infty)$ -almost surely.

Recalling that  $c_{\bar{p}} = \Phi'(\bar{p})$ , (21) and (24) imply the assertion of the theorem.  $\square$

## Acknowledgements

Both authors would like to thank Julien Berestycki and an anonymous referee for their valuable comments on this work.

## References

- [1] Bérard, J., Gouéré, J.-B. (2010) Brunet-Derrida behavior of branching-selection particle systems on the line, *Comm. Math. Phys.* **298**, 323-342.
- [2] Berestycki, J., Berestycki N. and Schweinsberg, J. (2011) The genealogy of branching Brownian motion with absorption, *To appear in Ann. Probab.*
- [3] Berestycki, J., Harris, S.C. and Kyprianou, A.E. (2011) Travelling waves and homogeneous fragmentation, *Ann. Appl. Probab.* **21**, 1749-1794.

- [4] Bertoin, J. (1996) *Lévy Processes*, Cambridge University Press.
- [5] Bertoin, J. (2003) Asymptotic behaviour of fragmentation processes, *J. Europ. Math. Soc.* **5**, 395 -416.
- [6] Bertoin, J. (2006) *Random Fragmentation and Coagulation Processes*, Cambridge University Press.
- [7] Bertoin, J., Rouault, A. (2003) Additive martingales and probability tilting for homogeneous fragmentations, Preprint.
- [8] Bertoin, J., Rouault, A. (2005) Discretization methods for homogeneous fragmentations, *J. London Math. Soc.* **72**, 91-109.
- [9] Breiman, L. (1992) *Probability*, second edition, SIAM.
- [10] Derrida, B., Simon, D. (2007) The survival probability of a branching random walk in presence of an absorbing wall. *Europhys. Lett. EPL* **78**, Art. 60006.
- [11] Derrida, B. and Simon, D. (2008) Quasi-stationary regime of a branching random walk in presence of an absorbing wall. *J. Stat. Phys.* **131**, 203-233.
- [12] Durrett, R. (1991) *Probability: theory and examples*, Duxbury Press.
- [13] Gantert N., Hu Y., and Shi Z. (2011) Asymptotics for the survival probability in a supercritical branching random walk, *Ann. Inst. H. Poincaré Probab. Statist.* **47**, 111-129.
- [14] Harris, J.W. and Harris, S.C. (2007), Survival probabilities for branching Brownian motion with absorption, *Elect. Comm. Probab.* **12**, 81-92.
- [15] Harris, J., Harris, S.C. and Kyprianou, A.E. (2006) Further probabilistic analysis of the Fisher-Kolmogorov-Petrovskii-Piscounov equation: one sided travelling waves, *Ann. Inst. H. Poincaré Probab. Statist.* **42**, 125-145.
- [16] Harris, S.C., Knobloch, R. and Kyprianou, A.E. (2010) Strong law of large numbers for fragmentation processes, *Ann. Inst. H. Poincaré Probab. Statist.* **46**, 119-134.
- [17] Knobloch, R. (2011) *Asymptotic properties of fragmentation processes*, PhD Thesis, University of Bath.
- [18] Knobloch, R. (2012) One-sided FKPP travelling waves in the context of homogeneous fragmentation processes, Preprint, arXiv: 1204.0758.
- [19] Kyprianou, A.E. (2006) *Introductory Lectures on Fluctuations of Lévy Processes with Applications*, Springer.