



Citation for published version:

Tabor, A, Keogh, E & Eccleston, C 2017, 'Embodied pain—negotiating the boundaries of possible action', Pain, vol. 158, no. 6, pp. 1007-1011. https://doi.org/10.1097/j.pain.0000000000000875

DOI:

10.1097/j.pain.0000000000000875

Publication date: 2017

Document Version Peer reviewed version

Link to publication

The final published version of this article is available at: https://doi.org/10.1097/j.pain.0000000000000875

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policyIf you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 13. May. 2019

1	Embodied pain—negotiating the boundaries of possible action
2	
3	Abby Tabor*, Edmund Keogh, Christopher Eccleston
4	Centre for Pain Research, University of Bath, UK.
5	
6	
7	* Corresponding author
8	Abby Tabor, PhD
9	Centre for Pain Research
10	Department for Health (1 West)
11	University of Bath
12	BA2 7AY, UK
13	+441225 384225
L 4	
15	
16	
17	
18	Number of words: 2001
19	Number of pages: 18
20	
21	
22	Keywords: embodied pain, action, inference, liminality, defence

1. Introduction.

Pain is a protective strategy, which emerges from on-going interaction between body and world. Pain is, however, often thought of as a unitary output—an end product experienced as an intrusion upon an often unsuspecting perceiver [56]. We know a lot about how nociception relates to pain, informed by both biological and psychological influences [30,70,98], about how pain intrudes into awareness [5,26,29,34], and how it relates to clinical variables such as suffering and disability [35]. However, despite significant advances, the mechanisms of pain intrusion remain elusive [63]. In this paper we stress a functional view of pain as more than experience; as defensive action operating in the context of uncertain threat.

Although traditional characterisations of perception as a product of sensory information have been critiqued [19,41,53], including in pain [89,96], there is now a well advanced contemporary view that all perception is embodied and embedded [41,67,79,88]. Here, *embodied* is defined by action, the premise that cognition extends beyond the brain so that an ever-changing body is at the core of how our experiences are shaped; this may be the unconscious workings of our immune system or the collaborative efforts made to avoid movement. *Embedded* refers to the situated interaction between the embodied being and the external environment, in both place (current context) and time (evolutionary context).

From this view, all experience is inferential [78], dynamic [22,55], and related to action in the world [2,21,24]. Thus, to describe the experience of pain we must understand it within its evolved, learned, and ultimately threat-defined context [33,101]. Theories of embodied experience are well advanced elsewhere, most notably in cybernetics [4,23,81], evolutionary biology [39,75,82] and consciousness [83,84]. Its provenance can be traced to structural psychology [93], phenomenology [47,53,62], and perception

[41,77]. However, embodied domains have avoided pain, considering it either too simple [32] or paradoxically too difficult [6].

Our embodied view, in many ways complements existing literature [18,27,36,42,97] supporting the growing understanding of pain as an experience inferred from uncertain information [3,17,85,100]. However, it critically looks to extend this work beyond a passive, information processing model that has come to dominate [49]. Here, we emphasise the body, not separate from the brain nor the world, but part of the facility that actively shapes our experience of pain. This perspective defines pain in terms of action: an experience which, as part of a protective strategy, attempts to defend one's self in the presence of inferred threat.

We start with a consideration of the core features of embodied pain. Next, we review the few studies that have been attempted on embodied perception and pain. Finally, we discuss how this approach can be applied usefully to pain, exploring both the research and clinical implications of embodied pain.

2. Inferring experience in an uncertain world

In proposing a view of pain as embodied and embedded, we draw upon three principles from the broader literature on embodied experience: inference, liminality, and defence. First, all experience is inferred, and inference functions principally to maintain coherence in complex and inherently uncertain environments—inference. Second, all experience is fundamentally defined by the boundaries of possible action—liminality. Third, all experience can be disturbed by bodily threat: pain is an action that functions to reduce threat; promoting defence and maintaining the integrity of coherent

2.1. Inference

behaviour—defence.

We know now that our experiences are inferred [47, 89]: we fill in the gaps [44], selectively attend [1,31], unconsciously prime [10,50], and in essence prioritise efficiency over accuracy [52,94]. Perception results from attempts to accommodate information that has deviated from our predictions [20]. It is only through the actions of our body and our predictions of the consequences of these actions that we are able to disambiguate the world [39]. Thus, the reciprocal relationship between action and prediction continually reshapes our experience of pain.

Perception as inference can be characterised computationally [103], and has been explored in pain [3,17,61]. Critically, however, the role of the body is often relegated in these more reductionist models, overshadowed by the dominant view of pain as a phenomenon of the brain [99]. In contrast, experience from an embodied perspective is borne out of the hierarchical, sensorimotor interactions we have with the world [40,73,74]. Importantly, this accounts for the changing ability of the individual to act in their environment, as well as what the environment affords. When pain is included within this sensorimotor interaction, it can be considered an action that deliberately alters the way in which we are able to interact with our environment and so in turn, changes what the environment affords.

2.2. Liminality

Experience can be thought of as a strategy generated from the need to continually adjust our actions when our predictions emerge as inadequate, i.e., a mismatch that does not provide a coherent basis for action [23,51]. The need for homeostatic coherence above all else drives experience [9,25,81]. Pain, along with other bodily experiences (e.g. fatigue, itch, temperature, pressure and disequilibrium) that intrude upon awareness indicate that boundaries have been reached and action must be taken—they are liminal experiences.

2.3 Defence

Much of the active inference we describe occurs outside of awareness. Like a stream following a well-worn channel defined by natural banks that guide and constrain, so felt experience flows largely uninterrupted, embodied by physical constraints and embedded within social constraints. To stray outside of these bounds produces specific alerts that function to modify our actions or alter our predictions. Each physical sense has a specific threat tied to specific defensive actions, which attempt to return the individual to within viable constraints [28].

In some circumstances those defensive actions are insufficient and the result is experienced as disturbing, e.g., das unheimliche phenomena in which we experience incoherent perceptions of familiarity; an illusion of relationship, in which objects are uncannily personal [38]. When all defensive actions fail there emerge whole system delusional experiences, including repression, de-realization, and—as the final defence—dissociation [12,13,58].

3. Embodied pain motivating action

First we review research on how pain influences non-pain perceptual judgement, and the obverse- *inference*. Second, we consider studies of action constrained when it meets the boundaries imposed by the body in pain, studied as illusions that alter the experience of pain- *liminal*. Third, we consider examples of whole body disturbances for their accounting of pain, studied as specific experiences of pain related dissociation, or global experiences of delusion, in a final defence by departure
defence.

There is a small body of experimental work on how the experience of pain can alter non-pain perception. For example, we have shown that pain affects judgements of distance when the object-distance being judged is threat-related [91], an observation previously made in patients with clinical pain [102]. Similarly, pain can affect judgements of the weight of external objects [90], and the weight, size, and shape of one's own body [67,69]. Clinically, reports of pain, temperature, stiffness, and imbalance are hard to disentangle, so often appear together [68], and have yet to be experimentally separated. Without such finesse, attempts to capture embodied experience rightly faces scrutiny and challenge [37]; although studies have replicated the effects of higher order cognition and mood on pain [11,92]. There are also studies of counter-stimulation offered in competition to pain as distraction [59]. Evidence from direct experimental studies conducted shows pain to be dynamic, flexible, and connected; a reflection of inference in an uncertain world.

Illusionary experience goes beyond altered sensory judgements. 'Illusionary' is normally judged as impossible or improbable perception based on a common agreement on the world; for example, if I perceive a limb that every external observer knows me to have lost. Painful missing body parts are a common experience for amputees [72], although they are rarely reported in isolation from temperature, pressure, weight, size and itch phenomena. Visual counter-stimulation using mirrors or virtual reality can alter aspects of size, position, and ownership, but also pain [15,60,76]. Some illusions may be harder to identify than others. For example, patients with osteoarthritis demonstrate an altered sensorimotor relationship with the affected limb in addition to the experience of pain [43,87,88]. Evidence from studies of illusionary physical experience can be seen usefully as examples of pain operating as a liminal phenomenon, unstable and malleable.

Embodied pain involves an elision between perception and action, such that pain without action should be considered unusual, abnormal, or extreme. From this perspective, chronic pain involves persistent action that attempts to reduce threat over time. Inescapable pain, where action is inadequate, may be a signal feature of severe

distress eg., total pain, or locked in syndrome) [7]. At risk in inescapable pain is the coherence of all behaviour. There are studies of altered bodily coherence in individuals with CRPS I [67] and observations of dissociation from ownership of a limb [57]. But there are few experimental studies of what can be considered a final defence by departure, in repression, de-realization, or dissociation. In anthropology there are qualitative accounts of specific rites of passage [65], and in social psychology of deviant social practice [8]. In the history of medicine we find rich description of inescapable surgical pain without anaesthesia [14] and in contemporary medicine there are similar accounts, such as in emergency care, or burns care [66]. There is no meta-synthesis of this literature, however, accounts of inescapable pain—of pain denied action—all feature what we call a final defence in a dissociative departure from our body. Although these departures are well studied in clinical neurology, and so have a structure [54] they have not been studied in pain. Evidence from studies of final defence show that only in extreme circumstances does perception cleave from action.

4. Discussion

Pain as embodied and embedded—inferred, liminal, and functioning for defence—has far reaching research and clinical implications (Fig. 1.). Our focus should shift from pain as a passive, sensory experience to pain as a dynamic, motor experience. Pain is always about action [96].

For research, our focus should be on the critical gaps. First, there is a need to explore the changing interactions between experience of the body and associated action (conscious and non-conscious). Studies of proprioception [45], peri-personal space [79], and bodily size [68] have offered the best entry points, but a programme of research into other liminal bodily experiences, such as itch, fatigue, disequilibrium, and respiration are also needed.




Fig. 1. Embodied Pain: proposed research and clinical agendas.

192

193

194

195

196

197

198

199

The clinical study of treatments aimed at altering experience should consider actions associated with threat. In part, this approach is concerned with gaining detailed accounts of real-life interactions. In acute pain, there are unexplored opportunities in going beyond simple distraction, making use of the inherent uncertainty associated with our bodily experiences; recognising that we act continually to reduce uncertainty. This line of work is already being pursued with the use of bodily illusions [45,71,76]. In chronic pain, interesting are e-health and m-health innovations that now allow for

moment-by-moment measurement of functional, physiological and experiential parameters in the real word. Clinically, treatments framed within a motivational context of how pain interferes with purposeful goal-orientated behaviour (e.g, completing a work task) may be improved by studying how threat to bodily coherence is managed [16,80]. In particular, accounting for how action and prediction influence individually defined boundaries. We are beginning to think of therapy as the attempt to redefine a stable coherence of one's identity in line with the context of a persistent urge for defence [66].

5. Conclusion

We propose that pain is inescapably embodied and embedded; an action that reflects the uncertainty of body and world. '*Embodied pain*' provides a theoretical platform from which novel investigations can aim to understand coherent action in complex, goal-rich environments.

- 214 References
- 215 [1] Allport DA. Attention and control: have we been asking the wrong questions?
- A critical review of twenty-five years. In: Meyer E, Kornblurn S, editors.
- 217 Attention and Performance XVI: Synergies in Experimental Psychology,
- 218 Artificial Intelligence, and Cognitive Neuroscience. Cambridge, MA: MIT Press,
- 219 1993. pp. 182–218.
- 220 [2] Allport DA. Selection for action: some behavioral and neurophysiological
- considerations of attention and action. In: Heuer H, Sanders HF, editors.
- Perspectives on Perception and Action. Hillsdale, NJ: Lawrence Erlbaum
- 223 Associates, 1987. pp. 395–419.
- 224 [3] Anchisi D, Zanon M. A Bayesian perspective on sensory and cognitive
- integration in pain perception and placebo analgesia. PLoS One 2015;10:1–
- 226 20.
- 227 [4] Ashby WR. An introduction to cybernetics. London: Chapman & Hail Ltd.,
- 228 1956.
- 229 [5] Attridge N, Crombez G, Van Ryckeghem D, Keogh E, Eccleston C. The
- experience of cognitive intrusion of pain. Pain 2015;156:1978–90.
- 231 [6] Aydede M. Is feeling pain the perception of something? J. Philos.
- 232 2009;106:531–567.
- 233 [7] Bauby J-D. The diving bell and the butterfly. London: Fourth Estate, 1997.
- 234 [8] Baumeister RF. The enigmatic appeal of sexual masochism: why people
- desire pain, bondage, and humiliation sex. J. Soc. Clin. Psychol. 1997;16:133–
- 236 150.
- 237 [9] Bechara A, Damasio AR. The somatic marker hypothesis: a neural theory of
- economic decision. Games Econ. Behav. 2005;52:336–372.
- 239 [10] Beierholm UR, Quartz SR, Shams L. Bayesian priors are encoded
- independently from likelihoods in human multisensory perception. J. Vis.
- 241 2009;9:1–9.

242 [11] Benedetti F, Pollo A, Lopiano L, Lanotte M, Vighetti S, Rainero I. Conscious 243 expectation and unconscious conditioning in analgesic, motor, and hormonal 244 placebo/nocebo responses. J. Neurosci. 2003;23:4315-4323. 245 Blackmore SJ. Beyond the body: an investigation of the out-of-the-body [12] 246 experiences. London: Heinemann, 1982. 247 Blanke O. Out of body experiences and their neural basis. BMJ [13] 248 2004;329:1414–1415. 249 [14] Bourke J. The story of pain: from prayer to painkillers. Oxford: Oxford 250 University Press. 2014. 251 [15] Bowering KJ, O'Connell NE, Tabor A, Catley MJ, Leake HB, Moseley GL, 252 Stanton TR. The effects of graded motor imagery and its components on 253 chronic pain: a systematic review and meta-analysis. J. Pain 2013;14:3–13. 254 [16] Brandtstadter J, Rothermund K. The life-course dynamics of goal pursuit and 255 goal adjustment: a two-process framework. Dev. Rev. 2002;22:117–150. 256 [17] Buchel C, Geuter S, Sprenger C, Eippert F. Placebo analgesia: a predictive 257 coding perspective. Neuron 2014;81:1223-1239. 258 [18] Butler DS, Moseley GL. Explain Pain: revised and updated. 2nd ed. Adelaide, 259 SA: Noigroup Publications, 2013. 260 [19] Chemero A. An outline of a theory of affordances. Ecol. Psychol. 261 2003;15:181–195. 262 Clark. Busting out: predictive brains, embodied minds, and the puzzle of the [20] 263 evidentiary veil. Nous 2016:1-27. 264 [21] Clark A. An embodied cognitive science? Trends Cogn. Sci. 1999;3:345–351. 265 doi:10.1016/S1364-6613(99)01361-3. 266 Clark A. Embodied Prediction. In: Metzinger T, Windt JM, editors. Open MIND. [22] 267 Frankfurt am Main: MIND Group, 2015. p. 7. doi:10.15502/9783958570115.

Clark A. Whatever next? Predictive brains, situated agents, and the future of

cognitive science. Behav. Brain Sci. 2013;36:181-204.

268

269

[23]

- 270 [24] Clark A, Chalmers D. The extended mind. Analysis 1998;58:7–19.
- 271 [25] Craig AD. A new view of pain as a homeostatic emotion. Trends Neurosci.
- 272 2003;26:303–307.
- 273 [26] Crombez G, Eccleston C, Baeyens F, Eelen P. Disruptive nature of pain: an
- experimental investigation. Behav. Res. Ther. 1996;34:911–918.
- 275 [27] Crombez G, Eccleston C, Van Damme S, Vlaeyen JWS, Karoly P. Fear-
- avoidance model of chronic pain: the next generation. Clin. J. Pain
- 277 2012;28:475–83.
- 278 [28] Damasio A, Carvalho GB. The nature of feelings: evolutionary and
- 279 neurobiological origins. Nat. Rev. Neurosci. 2013;14:143–52.
- 280 [29] Van Damme S, Legrain V, Vogt J, Crombez G. Keeping pain in mind: a
- motivational account of attention to pain. Neurosci. Biobehav. Rev.
- 282 2010;34:204–213.
- 283 [30] Davis KD. Neuroimaging of pain: what does it tell us? Curr. Opin. Support.
- 284 Palliat. Care 2011;5:116–121.
- 285 [31] Dayan P, Kakade S, Montague PR. Learning and selective attention. Nat.
- 286 Neurosci. 2000;3:1218–1223.
- 287 [32] Dennett DC. Quining qualia. In: Marcel A, Bisiach E, editors. Consciousness in
- 288 Modern Science. Oxford: Oxford University Press, 1988.
- 289 [33] Eccleston C, Crombez G. Worry and chronic pain: a misdirected problem
- 290 solving model. Pain, 132; 233-236.
- 291 [34] Eccleston C, Crombez G. Pain demands attention: a cognitive-affective model
- of the interruptive function of pain. Psychol Bull 1999;125:356–366.
- 293 [35] Eccleston C, Crombez G, Aldrich S, Stannard C. Attention and somatic
- 294 awareness in chronic pain. Pain 1997;72:209–215.
- 295 [36] Engel GL. The need for a new medical model: A challenge for biomedicine.
- 296 Science. 1977;196:129–136.
- 297 [37] Firestone C, Scholl BJ. Cognition does not affect perception: Evaluating the

298 evidence for "top-down" effects. Behav. Brain Sci. 2015:1–72. Available: 299 doi.org/10.1017/s0140525x15000965. 300 [38] Freud S. The Uncanny. In: Strachey J, editor. The standard edition of the 301 complete psychological works of Sigmund Freud (Vol 17). London: The 302 Hogarth Press, 1919. pp. 218-256. 303 Friston K. The free-energy principle: a unified brain theory? Nat. Rev. [39] 304 Neurosci. 2010;11:127–138. 305 Gallagher S, Bower M. Making enactivism even more embodied. Avant Trends [40] 306 Interdiscip. Stud. 2014:2:232–247. Gibson JJ. The Theory of Affordances. In: Shaw R, Bransford J, editors. 307 [41] 308 Perceiving, Acting, and Knowing. Towards an Ecological Psychology. 309 Hoboken, NJ: John Wiley & Sons Inc., 1977. pp. 127-142. 310 [42] Gifford L. Pain, the Tissues and the Nervous System: A conceptual model. 311 Physiotherapy 1998;84:27–36. 312 [43] Gilpin HR, Moseley GL, Stanton TR, Newport R. Evidence for distorted mental 313 representation of the hand in osteoarthritis. Rheumatology 2014;54:678–682. 314 [44] Gregory RL. Perceptions as hypotheses. Philos. Trans. R. Soc. B Biol. Sci. 315 1980;290:181-197. 316 [45] Harvie DS, Broecker M, Smith RT, Meulders A, Madden VJ, Moseley GL. 317 Bogus visual feedback alters onset of movement-evoked pain in people with 318 neck pain. Psychol. Sci. 2015;26:385-92. 319 Haugeland J. Mind embodied and embedded. In: Haugeland J, editor. Having [46] 320 Thought: Essays in the Metaphysics of Mind. Cambridge, MA: Harvard 321 University Press, 1998. 322 Heidegger M. Being and Time. trans. J. Macquarrie and E Robinson, editor [47] 323 Tubingen: Max Niemeyer Verlag, 1962. 324 [48] Helmholtz H von. Handbuch der physiologischen optik, vol 3. (Trans.). 325 Southall JPC, editor New York, NY: Dover Publications, 1962.

326 [49] Helmholtz H von. Treatise on physiological optics, vol. II. 1924. 327 [50] Hohwy J. Attention and conscious perception in the hypothesis testing brain. 328 Front. Psychol. 2012;2:96. 329 Hohwy J. The predictive mind. Oxford: Oxford University Press, 2013. [51] 330 [52] Humphrey N. The placebo effect. In: Gregory R, editor. Oxford companion to 331 the mind. Oxford: Oxford University Press, 2005. 332 [53] Husserl E. Ideas: a general introduction to pure phenomenology. Trans. W. R 333 Boyce Gibson, editor New York: Collier Books, 1931. 334 Kihlstrom JF. Dissociative Disorders. Annu. Rev. Clin. Psychol. 2005;1:227-[54] 335 253. 336 [55] Körding KP, Wolpert DM. Bayesian integration in sensorimotor learning. 337 Nature 2004;427:244-247. 338 Legrain V, Damme S Van, Eccleston C, Davis KD, Seminowicz DA, Crombez [56] 339 G. A neurocognitive model of attention to pain: behavioral and neuroimaging 340 evidence. Pain 2009;144:230-232. 341 Lewis JS, Schweinhardt P. Perceptions of the painful body: the relationship [57] 342 between body perception disturbance, pain and tactile discrimination in 343 complex regional pain syndrome. Eur. J. Pain 2012;16:1320-1330. 344 [58] Lopez C, Halje P, Blanke O. Body ownership and embodiment: vestibular and 345 multisensory mechanisms. Neurophysiol. Clin. 2008;38:149–161. 346 Malloy KM, Milling LS. The effectiveness of virtual reality distraction for pain [59] 347 reduction: a systematic review. Clin. Psychol. Rev. 2010;30:1011–1018. 348 [60] Mancini F, Longo MR, Kammers MPM, Haggard P. Visual distortion of body 349 size modulates pain perception. Psychol. Sci. 2011;22:325-330. 350 Mano H, Seymour B. Pain: a distributed brain information network? PLoS Biol. [61] 351 2015;13:e1002037. Merleau-Ponty M. Phenomenology of Perception (Trans.). Smith C, editor 352 [62]

London: Routledge & Kegan Paul, 1962.

353

- 354 [63] Moayedi M, Davis KD. Theories of pain: from Specificity to Gate Control. J.
- 355 Neurophysiol. 2013;109:5–12.
- 356 [64] Moore A, Derry S, Eccleston C, Kalso E. Expect analgesic failure; pursue
- 357 analgesic success. Br. Med. J. 2013;346:f2690–f2690.
- 358 [65] Morinis A. The ritual experience: pain and the transformation of consciousness
- in ordeals of initiation. Ethos 1985;13:150–174.
- 360 [66] Morse JM, Mitcham C. The experience of agonizing pain and signals of
- 361 disembodiment. J. Psychosom. Res. 1998;44:667–680.
- 362 [67] Moseley GL. Distorted body image in complex regional pain syndrome.
- 363 Neurology 2005;65:773–778.
- 364 [68] Moseley GL, Gallace A, Spence C. Bodily illusions in health and disease:
- physiological and clinical perspectives and the concept of a cortical "body
- 366 matrix." Neurosci. Biobehav. Rev. 2012;36:34–46.
- 367 [69] Moseley GL, Parsons TJ, Spence C. Visual distortion of a limb modulates the
- pain and swelling evoked by movement. Curr. Biol. 2008;18:R1047-8.
- 369 [70] Moseley GL, Vlaeyen JWS. Beyond nociception: the imprecision hypothesis of
- 370 chronic pain. Pain 2015;156:35–38.
- 371 [71] Murray CD, Pettifer S, Howard T, Patchick EL, Caillette F, Kulkarni J, Bamford
- 372 C. The treatment of phantom limb pain using immersive virtual reality: three
- 373 case studies. Disabil. Rehabil. 2007;29:1465–1469.
- 374 [72] Nikolajsen L, Jensen ST. Phantom limb pain. Br. J. Anaesth. 2001;87:107-
- 375 116.
- 376 [73] Noe A. Action in perception. Cambridge, MA: MIT Press, 2004 p.
- 377 [74] O'Regan JK, Dagenaar J. Consciousness without inner models: a
- sensorimotor account of what is going on in our heads. Proc. AISB 2014.
- 379 [75] Prescott TJ, Bryson JJ, Seth AK. Introduction. Modelling natural action
- 380 selection. Philos. Trans. R. Soc. B Biol. Sci. 2007;362:1521–1529.
- 381 [76] Preston C, Newport R. Analgesic effects of multisensory illusions in

382 osteoarthritis. Rheumatology 2011;50:2314–2315. 383 [77] Proffitt DR. An embodied approach to perception: by what units are visual 384 perceptions scaled? Perspect. Psychol. Sci. 2013;8:474-483. 385 [78] Rao RPN, Ballard DH. Predictive coding in the visual cortex: a functional 386 interpretation of some extra-classical receptive-field effects. Nat. Neurosci. 387 1999;2:79–87. 388 Sambo CF, Iannetti GD. Better safe than sorry? The safety margin [79] 389 surrounding the body is increased by anxiety. J. Neurosci. 2013;33:14225-30. 390 [80] Schmitz U, Saile H, Nilges P. Coping with chronic pain: flexible goal 391 adjustment as an interactive buffer against pain-related distress. Pain 392 1996;67:41-51. 393 Seth AK. The Cybernetic Bayesian Brain. In: Metzinger T, Windt JM, editors. [81] 394 Open MIND. Frankfurt am Main: MIND Group, 2015;35. 395 doi:10.15502/9783958570108. 396 [82] Seth AK. The ecology of action selection: insights from artificial life. Philos. 397 Trans. R. Soc. Lond. B. Biol. Sci. 2007;362:1545-1558. 398 [83] Seth AK. Why fish pain cannot and should not be ruled out. Anim. Sentience 399 2016;3:1-5. 400 [84] Seth AK, Suzuki K, Critchley HD. An interoceptive predictive coding model of 401 conscious presence. Front. Psychol. 2012;3:1–16. 402 Seymour B, Dolan RJ. Emotion, Motivation, and Pain. In: McMahon S, [85] 403 Koltzenburg M, Tracey I, Turk DC, editors. Wall and Melzack's Textbook of 404 Pain. Philadelphia, PA: Saunders, Elsevier Ltd, 2013. pp. 248–255. 405 Shapiro LA. The mind incarnate. Cambridge, MA: MIT Press, 2004. [86] 406 Stanton TR, Lin CWC, Bray H, Smeets RJEM, Taylor D, Law RYW, Moseley [87] 407 GL. Tactile acuity is disrupted in osteoarthritis but is unrelated to disruptions in 408 motor imagery performance. Rheumatology 2013;52:1509–1519.

Stanton TR, Lin CWC, Smeets RJEM, Taylor D, Law R, Lorimer Moseley G.

409

[88]

- Spatially defined disruption of motor imagery performance in people with
- osteoarthritis. Rheumatology 2012;51:1455–1464.
- 412 [89] Sullivan MD. Pain in language. From sentience to Sapience. J. Pain 1995;4:3-
- 413 14.
- 414 [90] Sullivan MJ, Thibault P, Savard A, Catchlove R, Kozey J, Stanish WD. The
- influence of communication goals and physical demands on different
- dimensions of pain behavior. Pain 2006;125:270–277.
- 417 [91] Tabor A, Catley MJ, Gandevia SC, Thacker M a., Spence C, Moseley GL. The
- close proximity of threat: altered distance perception in the anticipation of pain.
- 419 Front. Psychol. 2015;6:1–6.
- 420 [92] Tang NKY, Salkovskis PM, Hodges A, Wright KJ, Hanna M, Hester J. Effects
- of mood on pain responses and pain tolerance: an experimental study in
- 422 chronic back pain patients. Pain 2008;138:392–401.
- 423 [93] Titchener EB. Structural and functional psychology. Philos. Rev. 1899;8:290-
- 424 299.
- 425 [94] Trimmer PC, Marshall JAR, Fromhage L, McNamara JM, Houston AI.
- 426 Understanding the placebo effect from an evolutionary perspective. Evol.
- 427 Hum. Behav. 2013;34:8–15.
- 428 [95] Varela F, Rosch E, Thompson E. The embodied mind: cognitive science and
- human experience. Cambridge, MA: MIT Press, 1991.
- 430 [96] Wall PD. On the relation of injury to pain. The John J. Bonica Lecture. Pain
- 431 1979;6:253–264.
- 432 [97] Wall PD. Pain: the science of suffering. New York, NY: Columbia University
- 433 Press, 2000.
- 434 [98] Wiech K, Ploner M, Tracey I. Neurocognitive aspects of pain perception.
- 435 Trends Cogn. Sci. 2008;12:306–313.
- Wiech K, Tracey I. Pain, decisions, and actions: a motivational perspective.
- 437 Front. Neurosci. 2013;7:1–12.

438	[100]	Wiech K, Vandekerckhove J, Zaman J, Tuerlinckx F, Vlaeyen JWS, Tracey I.
439		Influence of prior information on pain involves biased perceptual decision-
440		making. Curr. Biol. 2014:R679–R681.
441	[101]	Williams AC de C. What can evolutionary theory tell us about chronic pain?
442		Pain 2015;157:1.
443	[102]	Witt JK, Linkenauger SA, Bakdash JZ, Augustyn JS, Cook A, Proffitt DR. The
444		long road of pain: chronic pain increases perceived distance. Exp. Brain Res.
445		2009;192:145–148.
446	[103]	Yuille AL, Bulthoff HH, Kersten D, Mamassian P. Perception as Bayesian
447		Inference. Annu. Rev. Psychol. 1996;55:271–304.
448		