



Citation for published version:

Guo, S & Zimmer, J 2018, 'Travelling wavefronts in nonlocal diffusion equations with nonlocal delay effects', *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 41, no. 2, pp. 919-943.
<https://doi.org/10.1007/s40840-017-0481-0>

DOI:

[10.1007/s40840-017-0481-0](https://doi.org/10.1007/s40840-017-0481-0)

Publication date:

2018

Document Version

Peer reviewed version

[Link to publication](#)

This is a post-peer-review, pre-copyedit version of an article published in *Bulletin of the Malaysian Mathematical Sciences Society*. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s40840-017-0481-0>.

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Travelling wavefronts in nonlocal diffusion equations with nonlocal delay effects

Shangjiang Guo*

College of Mathematics and Econometrics, Hunan University
Changsha, Hunan 410082, People's Republic of China

Johannes Zimmer

Department of Mathematical Sciences, University of Bath,
Bath BA2 7AY, United Kingdom

Abstract

This paper deals with the existence, monotonicity, uniqueness and asymptotic behaviour of travelling wavefronts for a class of temporally delayed, spatially nonlocal diffusion equations.

Keywords. Travelling waves, time delay, Fisher-KPP equation

AMS subject classifications. 35K57, 34K05, 92D25.

1 Introduction

Travelling wavefront solutions play an important role in the description of the long-term behaviour of solutions to initial value problems in reaction-diffusion equations, both in the spatially continuous case and in spatially discrete situations. Such solutions are also of interest in their own right, for example to understand transitions between different states of a physical system, propagation of patterns, and domain invasion of species in population biology (see, e.g., [3, 4, 8, 13, 37]). In this paper, we study the existence, uniqueness and asymptotic stability of travelling wavefronts of the equation:

$$u_t(x, t) = pu_{xx}(x, t) + d(J * u - u)(x, t) + f(u(x, t), (h ** u)(x, t)), \quad (1)$$

where $x \in \mathbb{R}$, $d \geq 0$, $p \geq 0$, and

$$(J * u)(x, t) := \int_{\mathbb{R}} J(x - y)u(y, t)dy,$$
$$(h ** u)(x, t) := \int_{-\infty}^t \int_{\mathbb{R}} h(x - y, t - s)u(y, s)dyds.$$

Equation (1) mixes a continuous Laplacian with a nonlocal diffusion $d(J * u - u)(x, t)$, which describes that the diffusion of density u at a point x and time t depends not only on $u(x, t)$ but also on all the values of u in a neighbourhood of x through the convolution term $J * u$. In population dynamics, the reaction term $f(u, h ** u)$ is usually used to describe the recruits of population, and $h ** u$ represents a weighted average of the population density both in past time and space [6, 7]. The nonlinear functions $f(u, v)$ and $h(u)$ satisfy the following hypotheses:

(F1) $f \in C([0, K] \times [0, K], \mathbb{R})$, $f(0, 0) = f(K, K) = 0$, $f(u, h ** u) > 0$ for all $u \in (0, K)$,
 $\partial_2 f(u, v) \geq 0$ for all $(u, v) \in [0, K] \times [0, K]$, where K is a positive constant.

*E-mail: shangjguo@hnu.edu.cn

(F2) There exist some $M > 0$ and $\sigma \in (0, 1]$ such that $0 \leq \partial_1 f(0, 0)u + \partial_2 f(0, 0)v - f(u, v) \leq M(u + v)^{1+\sigma}$ for all $(u, v) \in [0, K] \times [0, K]$ and $\partial_1 f(K, K) + \partial_2 f(K, K) < 0$.

(H1) Both $J(\cdot)$ and $h(\cdot, t)$ are nonnegative, even, integrable and satisfies $\int_{\mathbb{R}} J(x)dx = 1$ and $\int_0^\infty \int_{\mathbb{R}} h(x, t)dxdt = 1$.

(H2) There exists some $\lambda_0 > 0$ (possibly equal to ∞) such that $\int_0^\infty J(x) \exp\{\lambda x\}dx < \infty$ and $\int_0^\infty \int_0^\infty h(x, t) \exp\{\lambda(x - ct)\}dxdt < \infty$ for all $c \geq 0$ and $\lambda \in [0, \lambda_0)$.

Assumptions (F1) and (F2) are standard. From (F1) we can see that (1) has two equilibria 0 and K . Furthermore, condition (F2) together with (F1) and (H1) implies that $\partial_1 f(0, 0) + \partial_2 f(0, 0) \geq \frac{2}{K}f(\frac{K}{2}, \frac{K}{2}) = \frac{2}{K}f(\frac{K}{2}, h * *(\frac{K}{2})) > 0$, hence 0 is unstable and K is stable. In this article, we will not require that $\partial_2 f(0, 0) > 0$.

Since Equation (1) involves a general diffusion kernel and delayed nonlinearity, it can be reduced to some well-known equations if J , h , and f are chosen to take a a some special form (see, for example, [1, 16, 35, 39, 40, 43, 44]). In particular, special cases of (1) include a host-vector disease model, a nonlocal population model with age structure, and a nonlocal Nicholson's blowflies model with delay; these cases are discussed in a second paper investigating the stability of the system [20]. For example, choosing $d = 0$, $f(u, v) = -\tau u + \tau \beta v e^{-v}$, equation (1) can be reduced to the following Nicholson's Blowflies equation with spatio-temporal delays

$$u_t(x, t) = pu_{xx}(x, t) - \tau u(x, t) + \tau \beta (h * *u)(x, t) \exp\{-(h * *u)(x, t)\},$$

which was studied by Li, Ruan, and Wang [26], and Lin [28]. If $p = 0$ and $J(x) = \frac{1}{2}[\delta(-1) + \delta(1)]$ and $h(x, t) = k(x)\delta(t - \tau)$, where $\delta(\cdot)$ is the Dirac delta function, then (1) reduces to the discrete reaction-diffusion equation

$$u_t(x, t) = d \cdot \Delta_1 u(x, t) + f(u(x, t), (k * u)(x, t - \tau)), \quad (2)$$

where $\Delta_1 u(x, t) = \frac{1}{2}[u(x + 1, t) - 2u(x, t) + u(x - 1, t)]$. If $f(u, v) = -au + b(v)$ and $k(x) = \delta(x)$, then (2) reduces to the local equation

$$u_t(x, t) = d \cdot \Delta_1 u(x, t) - au(x, t) + b(u(x, t - \tau)), \quad x \in \mathbb{R}, t \geq 0, \quad (3)$$

where $b \in C^1([0, \infty], \mathbb{R})$. If $f(u, v) = g(u)$ and $g(u)$ denotes a Lipschitz continuous function satisfying $g(u) > 0 = g(0) = g(1)$ for all $u \in (0, 1)$, equation (2) becomes

$$u_t(x, t) = d \cdot \Delta_1 u(x, t) + g(u(x, t)). \quad (4)$$

On the other hand, when $d = 0$ and $h(x, t) = k(x)\delta(t - \tau)$, (1) reduces to the following reaction-diffusion equation with discrete time delay

$$u_t(x, t) = pu_{xx}(x, t) + f(u(x, t), (k * u)(x, t - \tau)). \quad (5)$$

Moreover, the equation (2) is a spatially discrete version of (5) with p replaced by d . In recently years, spatially non-local differential equations such as (5) have attracted significant attention (see, e.g., [15, 19, 33, 37, 38, 39]). Under some monostable assumption, Wang *et al.* [39] investigated the existence, uniqueness, and global asymptotical stability of travelling wave fronts. We also refer to So *et al.* [37] for more details and some specific forms of f , obtained from integrating along characteristics of a structured population model, an idea from the work of Smith and Thieme [36]. See also [37] for a similar model and [18] for a survey on the history and the current status of the study of reaction diffusion equations with non-local delayed interactions. In particular, when $f(u, v) = v(1 - u)$ and $k(u) = \delta(u)$, the equation (5) is delayed *Fisher's equation* [17] or *KPP equation* [25], which arises in the study of gene development or population dynamics. When $f(u, v) = -au + b(v)$ and $k(u) = \delta(u)$, the equation (5) is the local *Nicholson's blowflies equation* and has been investigated in [19, 21, 27, 32]. When $f(u, v) = -au + b(1 - u)v$, equation (5) is called the *vector disease model* as proposed by Ruan and Xiao [34]. When $f(u, v) = bv \exp\{-\gamma\tau\} - \delta u^2$

and $k(u) = \frac{1}{\sqrt{4\pi\alpha\tau}} \exp\{\frac{-y^2}{4\alpha\tau}\}$, equation (5) is the age-structured reaction diffusion model of a single species proposed by Al-Omari & Gourley [2]. Existence and stability of travelling wavefronts for the reaction-diffusion equation (5) and its special forms has been extensively studied in the literature.

We are interested in wave propagation phenomena. In particular, we are interested in monotone travelling waves $u(x, t) = \phi(x + ct)$ for (1), with ϕ saturating at 0 and K . We call c the *travelling wave speed* and ϕ the *profile* of the wavefront. In order to address these questions, we need to find an increasing function $\phi(\xi)$, where $\xi = x + ct$, which is a solution of the associated travelling wave equation

$$\begin{aligned} -c\phi'(\xi) + p\phi''(\xi) + d(J * \phi - \phi)(\xi) + f(\phi(\xi), (h * * \phi)(\xi)) &= 0, \quad \xi \in \mathbb{R}, \\ \lim_{\xi \rightarrow -\infty} \phi(\xi) &= 0, \quad \lim_{\xi \rightarrow \infty} \phi(\xi) = K, \\ 0 \leq \phi(\xi) &\leq K, \quad \xi \in \mathbb{R}, \end{aligned} \tag{6}$$

where

$$\begin{aligned} (J * \phi)(\xi) &= \int_{\mathbb{R}} J(y)\phi(\xi - y)dy, \\ (h * * \phi)(\xi) &= \int_0^\infty \int_{\mathbb{R}} h(y, s)\phi(\xi - y - cs)dyds. \end{aligned}$$

For convenience, we write $\phi(-\infty)$ and $\phi(\infty)$ as abbreviations for $\lim_{\xi \rightarrow -\infty} \phi(\xi)$ and $\lim_{\xi \rightarrow \infty} \phi(\xi)$, respectively. Travelling wavefronts of (3) have been intensively studied in recent years, see, e.g., [8, 9, 10, 11, 12, 13, 22, 23, 24, 29, 30, 31, 42, 45, 46, 47]. Zinner *et al.* [47] addressed the existence and minimal speed of travelling wavefront for (4). Recently, based on [11, 12], Chen *et al.* [10] investigated the uniqueness and asymptotic behaviour of travelling waves for (2) with $d = 2$. To the best of our knowledge, however, there are no results regarding the existence, uniqueness, monotonicity, asymptotic behaviour, and asymptotic stability of travelling waves for an equation as general as (1).

There is an enormous amount of work on related equations which is impossible even to sketch. We only mention the work of Coville and coworkers, where the nonlinearity is local, but general nonlocal expressions instead of the nonlocal diffusion equation are considered (e.g., [14]). Some methods are similar, such as the use of super- and subsolutions. For interesting work on a Fisher-KPP equation with a non-local saturation effect, where no maximum principle holds, we refer to [5].

We shall establish the existence, uniqueness, monotonicity, asymptotic behaviour of travelling waves for (1) under the assumptions (F1), (F2), (H1), and (H2).

Theorem 1.1 *Under assumptions (F1), (F2), (H1), and (H2), there exists a minimal wave speed $c^* > 0$ such that for each $c \geq c^*$, equation (1) has a travelling wavefront $\phi(x + ct)$ satisfying (6). Moreover,*

1. *the solution ϕ of (6) is unique up to a translation.*
2. *Every solution ϕ of (6) is strictly monotone, i.e., $\phi'(\xi) > 0$ for all $\xi \in \mathbb{R}$.*
3. *Every solution ϕ of (6) satisfies $0 < \phi(\cdot) < K$ on \mathbb{R} .*
4. *Any solution of (6) satisfies $\lim_{\xi \rightarrow -\infty} \phi'(\xi)/\phi(\xi) = \lambda$, with λ being the minimal positive root of*

$$c\lambda - p\lambda^2 - d[H(\lambda) - 1] - \partial_1 f(0, 0) - \partial_2 f(0, 0)G(c, \lambda) = 0, \tag{7}$$

where

$$\begin{aligned} H(\lambda) &= \int_{\mathbb{R}} J(y) \exp\{-\lambda y\} dy, \\ G(c, \lambda) &= \int_0^\infty \int_{\mathbb{R}} h(y, s) \exp\{-\lambda(y + cs)\} dyds. \end{aligned}$$

for $\lambda \in \mathbb{C}$ with $\text{Re}\lambda < \lambda_0$.

5. Any solution of (6) satisfies $\lim_{\xi \rightarrow \infty} \phi'(\xi)/[K - \phi(\xi)] = \gamma$, with γ being the unique positive root of

$$c\gamma + p\gamma^2 + d[H(-\gamma) - 1] + \partial_1 f(K, K) + \partial_2 f(K, K)G(c, -\gamma) = 0. \quad (8)$$

We remark that if $c > c^*$, equation (7) has exactly two real roots, both positive.

This paper is organised as follows. In Section 2, we provide some preliminary results; in Section 3, we establish the existence of a travelling wavefront, using the monotone iteration method developed by Wu and Zou [42] with a pair of super- and sub-solutions. In particular, Theorems 3.1 and 3.2 establish the existence part of Theorem 1.1. To derive the monotonicity and uniqueness of wave profiles (Section 5), we shall first apply Ikehara's theorem in Section 4 to study the asymptotic behaviour of wave profiles. This idea originated in Carr and Chmaj's paper [9], where the authors study the uniqueness of waves for a nonlocal monostable equation. Theorem 5.1 establishes the monotonicity part of Theorem 1.1, and uniqueness is discussed in Theorem 5.2, and nonexistence of travelling waves for $c < c^*$ is the content of Theorem 5.3.

2 Notation and auxiliary results

Throughout this paper, $C > 0$ denotes a generic constant, while C_i ($i = 1, 2, \dots$) represents a specific constant. Let I be an interval, typically $I = \mathbb{R}$. Let $T > 0$ be a real number and \mathcal{B} be a Banach space. We denote by $C([0, T], \mathcal{B})$ the space of the \mathcal{B} -valued continuous functions on $[0, T]$, while $L^2([0, T], \mathcal{B})$ is the space of the \mathcal{B} -valued L^2 -functions on $[0, T]$. The corresponding spaces of the \mathcal{B} -valued functions on $[0, \infty)$ are defined similarly.

For a given travelling wave ϕ of (1) satisfying (6), define

$$\begin{aligned} G_j(\xi) &= \partial_j f \left(\phi(\xi), \int_0^\infty \int_{\mathbb{R}} h(y, s) \phi(\xi - y - cs) dy ds \right), \quad j = 1, 2, \\ B(\xi) &= \int_0^\infty \int_{\mathbb{R}} h(y, s) G_2(\xi + y + cs) dy ds. \end{aligned} \quad (9)$$

Obviously, $B(\xi)$ and $G_j(\xi)$, $j = 1, 2$ are non-increasing and satisfy

$$G_1(\infty) = \partial_1 f(K, K), \quad B(\infty) = G_2(\infty) = \partial_2 f(K, K). \quad (10)$$

Moreover, both $G(c, \lambda)$ and $H(\lambda)$ are twice differentiable in $\lambda \in [0, \lambda_0)$. Moreover, $G(c, 0) = 1$, $H(0) = 1$, $H'(\lambda) > 0$, $G_{\lambda\lambda}(c, \lambda) > 0$, and $H''(\lambda) > 0$. Set

$$\Delta(c, \lambda) = c\lambda - p\lambda^2 - d[H(\lambda) - 1] - \partial_1 f(0, 0) - \partial_2 f(0, 0)G(c, \lambda) \quad (11)$$

and

$$\tilde{\Delta}(c, \lambda) = c\lambda + p\lambda^2 + d[H(-\lambda) - 1] + \partial_1 f(K, K) + \partial_2 f(K, K)G(c, -\lambda) \quad (12)$$

for all $c \in \mathbb{R}$ and $\lambda \in \mathbb{C}$ with $c \geq 0$ and $\text{Re}\lambda < \lambda^+$, where $\lambda^+ = \lambda_0$ if $\partial_2 f(0, 0) > 0$ and $\lambda^+ = +\infty$ if $\partial_2 f(0, 0) = 0$.

We require two simple technical statements.

Lemma 2.1 *There exist $c^* > 0$ and $\lambda^* \in (0, \lambda^+)$ such that $\Delta(c^*, \lambda^*) = 0$ and $\Delta_\lambda(c^*, \lambda^*) = 0$. Furthermore,*

- (i) *if $0 < c < c^*$, then $\Delta(c, \lambda) < 0$ for all $\lambda \geq 0$;*
- (ii) *if $c > c^*$, then the equation $\Delta(c, \cdot) = 0$ has two positive real roots $\lambda_1(c)$ and $\lambda_2(c)$ with $0 < \lambda_1(c) < \lambda^* < \lambda_2(c) < \lambda^+$ such that $\lambda_1'(c) < 0$, $\lambda_2'(c) > 0$, $\Delta(c, \lambda) > 0$ for all $\lambda \in (\lambda_1(c), \lambda_2(c))$, and $\Delta(c, \lambda) < 0$ for all $(-\infty, \lambda^+) \setminus [\lambda_1(c), \lambda_2(c)]$.*

Proof. Note that for all $\lambda \in (0, \lambda^+)$,

$$\begin{aligned}\Delta_\lambda(c, \lambda) &= c - 2p\lambda - dH'(\lambda) - \partial_2 f(0, 0)G_\lambda(c, \lambda), \\ \Delta_{\lambda\lambda}(c, \lambda) &= -2p - dH''(\lambda) - \partial_2 f(0, 0)G_{\lambda\lambda}(c, \lambda) < 0, \\ \Delta_c(c, \lambda) &= \lambda - \partial_2 f(0, 0)G_c(c, \lambda) > 0, \\ \Delta(c, 0) &= -\partial_1 f(0, 0) - \partial_2 f(0, 0) < 0, \\ \Delta(0, \lambda) &= -2p\lambda^2 - d[H(\lambda) - 1] - \partial_1 f(0, 0) - \partial_2 f(0, 0)G(0, \lambda) < 0\end{aligned}$$

and

$$\lim_{\lambda \rightarrow \lambda^+ - 0} \Delta(c, \lambda) = -\infty.$$

Then the conclusion of this lemma follows. \square

Lemma 2.2 *Under assumptions (F1) and (F2), for each fixed $c \geq 0$, $\tilde{\Delta}(c, \cdot)$ has exactly one positive zero $v(c)$.*

Proof. In view of (F1) and (F2), we have

$$\tilde{\Delta}(c, 0) = \partial_1 f(K, K) + \partial_2 f(K, K) < 0$$

and

$$\lim_{\lambda \rightarrow \lambda^+ - 0} \tilde{\Delta}(c, \lambda) = +\infty.$$

Therefore, $\tilde{\Delta}(c, \cdot)$ has at least one positive zero. Note that

$$\tilde{\Delta}_\lambda(c, \lambda) = c + 2p\lambda - dH'(-\lambda) - \partial_2 f(K, K)G_\lambda(c, -\lambda)$$

and for $\lambda > 0$

$$G_\lambda(c, -\lambda) = \int_0^\infty \int_0^\infty h(y, s) \left[(y - cs)e^{-\lambda(y+cs)} - (y + cs)e^{\lambda(y-cs)} \right] dy ds < 0.$$

Then we have $\tilde{\Delta}_\lambda(c, \lambda) > 0$ for all $\lambda \in (0, \lambda^+)$. This implies that $\tilde{\Delta}_\lambda(c, \lambda)$ is increasing in λ and so it has exactly one positive zero. \square

We now define the notion of super- and sub-solutions. For any absolutely continuous function $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ satisfying that φ' and φ'' exist almost everywhere and are essentially bounded on \mathbb{R} , we set

$$N_c[\varphi](\xi) \triangleq c\varphi'(\xi) - p\varphi''(\xi) - d(J * \phi - \phi)(\xi) - f(\varphi(\xi), (h * * \varphi)(\xi)).$$

Given a positive constant c , a non-decreasing continuous function φ^+ is called a *super-solution* of (6) if $\varphi^+(-\infty) = 0$ and φ^+ is differentiable almost everywhere in \mathbb{R} such that $N_c[\varphi^+](\xi) \geq 0$ for almost every $\xi \in \mathbb{R}$. A continuous function φ^- is called a *sub-solution* of (6) if $\varphi^-(-\infty) = 0$, $\varphi^-(\xi)$ is not identically equal to 0 and φ^- is differentiable almost everywhere in \mathbb{R} such that $N_c[\varphi^-](\xi) \leq 0$ for almost every $\xi \in \mathbb{R}$.

Next, we introduce the operator $\mathcal{H}_\mu: C(\mathbb{R}) \rightarrow C(\mathbb{R})$ by

$$\mathcal{H}_\mu(\varphi) = d(J * \varphi - \varphi) + f(\varphi, h * * \varphi) + \mu\varphi.$$

It is easy to see that φ satisfies (6) if and only if φ satisfies

$$\varphi(\xi) = T_\mu(\varphi)(\xi), \tag{13}$$

where

$$T_\mu(\varphi)(\xi) = \frac{1}{c} \int_0^\infty \exp\left\{-\frac{\mu x}{c}\right\} \mathcal{H}_\mu(\varphi)(\xi + x) dx$$

if $p = 0$, and

$$T_\mu(\varphi)(\xi) = \frac{1}{p(\varsigma_2 - \varsigma_1)} \left[\int_{-\infty}^{\xi} e^{\varsigma_1(\xi-x)} \mathcal{H}_\mu(\varphi)(x) dx + \int_{\xi}^{\infty} e^{\varsigma_2(\xi-x)} \mathcal{H}_\mu(\varphi)(x) dx \right]$$

if $p > 0$, and

$$\varsigma_1 = \frac{c - \sqrt{c^2 + 4p\mu}}{2p} < 0, \quad \varsigma_2 = \frac{c + \sqrt{c^2 + 4p\mu}}{2p} > 0. \quad (14)$$

Choose $\mu > 2d + \max\{|\partial_j f(u, v)| : (u, v) \in [0, K] \times [0, K], j = 1, 2\}$. Then the operator T_μ is well-defined for functions ϕ of a growth rate less than $e^{\mu x}$. Furthermore, since f is monotone in the second argument by (F1), we have for $\varphi \leq \psi$

$$\mathcal{H}_\mu(\varphi) - \mathcal{H}_\mu(\psi) = \mu[\varphi - \psi] + d[J * (\varphi - \psi) - (\varphi - \psi)] + \partial_1 f(\tilde{\varphi}, h * \varphi)[\varphi - \psi] \leq 0,$$

where $\tilde{\varphi}(y)$ lies between $\varphi(y)$ and $\psi(y)$. Then the choice of μ shows that $\mathcal{H}_\mu(\varphi)$ is monotone in φ ,

$$\mathcal{H}_\mu(\varphi)(\xi) \leq \mathcal{H}_\mu(\psi)(\xi) \quad \text{if } 0 \leq \varphi \leq \psi \leq K \text{ in } \mathbb{R}. \quad (15)$$

Thus, we have the following result on the monotonic travelling waves.

Lemma 2.3 *Under assumptions (F1) and (H1), assume that there exists a super-solution φ^+ and a sub-solution φ^- of (6) such that $0 \leq \varphi^- \leq \varphi^+ \leq K$ on \mathbb{R} . Then (6) has a solution φ satisfying $\varphi'(\xi) \geq 0$ for all $\xi \in \mathbb{R}$.*

Proof. Assume that there exist a super-solution φ^+ and a sub-solution φ^- of (6) such that $0 \leq \varphi^- \leq \varphi^+ \leq K$ in \mathbb{R} . Define $\varphi_1 = T_\mu(\varphi^+)$. Then φ_1 is a well-defined C^1 function. From the definition of super-solution, we have

$$\varphi^+ \geq T_\mu(\varphi^+) = \varphi_1.$$

Also, by the definition of sub-solution and the property (15) of \mathcal{H}_μ , we get

$$\varphi^- \leq T_\mu(\varphi^-) \leq T_\mu(\varphi^+) = \varphi_1.$$

Hence $\varphi^-(\xi) \leq \varphi_1(\xi) \leq \varphi^+(\xi)$ for all $\xi \in \mathbb{R}$. Moreover, using the fact that φ^+ is non-decreasing and $\mu > 2d + \max\{|\partial_j f(u, v)| : u, v \in [0, K], j = 1, 2\}$, we have $\mathcal{H}_\mu(\varphi^+)(s) \geq \mathcal{H}_\mu(\varphi^+)(\xi)$ for all $s \geq \xi$ and hence $\varphi_1'(\xi) \geq 0$. Now define $\varphi_{n+1} = T_\mu(\varphi_n)$ for all $n \in \mathbb{N}$. By induction, it is easy to see that $0 \leq \varphi^- \leq \varphi_{n+1} \leq \varphi_n \leq \varphi^+ \leq K$ and $\varphi_{n+1}' \geq 0$ on \mathbb{R} for all $n \in \mathbb{N}$. Then the limit $\varphi(\xi) \triangleq \lim_{n \rightarrow \infty} \varphi_n(\xi)$ exists for all $\xi \in \mathbb{R}$ and $\varphi(\xi)$ is non-decreasing on \mathbb{R} . By Lebesgue's dominated convergence theorem, φ satisfies (13) and hence satisfies (6).

3 Existence

In this section, we shall establish the existence of travelling waves by constructing a suitable pair of super- and subsolutions. First, we derive two properties of possible solutions of (6).

Lemma 3.1 *Under assumptions (F1) and (H1), every solution (c, φ) of (6) satisfies $0 < \varphi(\xi) < K$ for all $\xi \in \mathbb{R}$.*

Proof. Let (c, φ) be a solution of (6). Suppose that there exists $\xi_0 \in \mathbb{R}$ such that $\varphi(\xi_0) = 0$. In view of $\varphi(\infty) = K$, without loss of generality, we may assume ξ_0 is the right-most point such that $\varphi(\xi_0) = 0$. Since $\varphi(\xi) \geq 0$ for all $\xi \geq \xi_0$, we have $\varphi'(\xi_0) = \varphi''(\xi_0) = 0$. It follows from (6) and (F1) that $\varphi(\xi) \equiv 0$ for all $\xi \in \mathbb{R}$, which contradicts the definition of ξ_0 . Therefore, $\varphi > 0$ on \mathbb{R} . Similarly, $\varphi < K$ on \mathbb{R} . This completes the proof. \square

Lemma 3.2 Under assumptions (F1) and (H1), every solution (c, φ) of (6) satisfying $\varphi' \geq 0$ on \mathbb{R} satisfies $\varphi' > 0$ on \mathbb{R} .

Proof. Suppose on the contrary that there exists $\xi_0 \in \mathbb{R}$ such that $\varphi'(\xi_0) = 0$. By differentiating (13) with respect to ξ , we obtain $\mathcal{H}_\mu(\varphi)(s) = \mathcal{H}_\mu(\varphi)(\xi_0)$ for all $s \geq \xi_0$. Letting $s \rightarrow \infty$, we obtain $\mathcal{H}_\mu(\varphi)(\xi_0) = c\mu K$. This, together with $\varphi'(\xi_0) = 0$ and (6), implies that $\varphi(\xi_0) = K$, which contradicts Lemma 3.1. Hence the lemma is proved. \square

Lemma 3.3 Assume that (F1), (F2), (H1) and (H2) hold. Let c^* , $\lambda_1(c)$ and $\lambda_2(c)$ be defined as in Lemma 2.1. Let $c > c^*$ be any number. Then for every $\gamma \in (0, \min\{\sigma\lambda_1(c), \lambda_2(c) - \lambda_1(c)\})$ there exists $Q(c, \gamma) > 1$ such that for every $q > Q(c, \gamma)$, the functions ϕ^\pm defined by

$$\phi^+(\xi) = \min\{K, \exp\{\lambda_1(c)\xi\}\}, \quad \xi \in \mathbb{R} \quad (16)$$

and

$$\phi^-(\xi) = \exp\{\lambda_1(c)\xi\} \max\{0, 1 - q \exp\{\gamma\xi\}\}, \quad \xi \in \mathbb{R} \quad (17)$$

are a super-solution and a sub-solution to (6), respectively.

Proof. We begin by proving that ϕ^\pm are a pair of super- and sub-solutions of (6). We only consider the case $\partial_2 f(0, 0) > 0$ because the proof of the case $\partial_2 f(0, 0) = 0$ is similar. It follows from (16) that $\phi^+(\xi) \leq \exp\{\lambda_1(c)\xi\}$ for all $\xi \in \mathbb{R}$ and hence

$$J * \phi^+(\xi) \leq \int_{\mathbb{R}} J(y) \exp\{\lambda_1(c)(\xi - y)\} dy = \exp\{\lambda_1(c)\xi\} H(\lambda_1(c))$$

and

$$h * * \phi^+(\xi) \leq \int_0^\infty \int_{\mathbb{R}} h(y, s) \exp\{\lambda_1(c)(\xi - y - cs)\} dy ds = \exp\{\lambda_1(c)\xi\} G(c, \lambda_1(c)).$$

Moreover, there exists $\xi^* > 0$ satisfying $\exp\{\lambda_1(c)\xi^*\} = K$, $\phi^+(\xi) = K$ for $\xi > \xi^*$ and $\phi^+(\xi) = \exp\{\lambda_1(c)\xi\}$ for $\xi \leq \xi^*$. For $\xi > \xi^*$, we have

$$N_c[\phi^+](\xi) = -d[J * \phi^+(\xi) - K] - f(K, (h * * \phi^+)(\xi)) \geq -f(K, K) = 0.$$

For $\xi \leq \xi^*$, we have

$$\begin{aligned} N_c[\phi^+](\xi) &\geq \phi^+(\xi) \{c\lambda_1(c) - p\lambda_1^2(c) - d[H(\lambda_1(c)) - 1]\} - f(\phi^+(\xi), (h * * \phi^+)(\xi)) \\ &= \phi^+(\xi) \Delta(c, \lambda_1(c)) - f(\phi^+(\xi), (h * * \phi^+)(\xi)) + \partial_1 f(0, 0) \phi^+(\xi) \\ &\quad + \phi^+(\xi) \partial_2 f(0, 0) G(c, \lambda_1(c)) \\ &\geq -f(\phi^+(\xi), (h * * \phi^+)(\xi)) + \partial_1 f(0, 0) \phi^+(\xi) + \partial_2 f(0, 0) (h * * \phi^+)(\xi) \geq 0, \end{aligned}$$

where we have used the condition (F2) in the last inequality. Therefore, ϕ^+ is a supersolution of (6).

It follows from (17) that $\exp\{\lambda_1(c)\xi\} \geq \phi^-(\xi) \geq \exp\{\lambda_1(c)\xi\} (1 - q \exp\{\gamma\xi\})$ for all $\xi \in \mathbb{R}$ and hence

$$\begin{aligned} J * \phi^-(\xi) &\geq \int_{\mathbb{R}} J(y) \exp\{\lambda_1(c)(\xi - y)\} (1 - q \exp\{\gamma(\xi - y)\}) dy \\ &= \exp\{\lambda_1(c)\xi\} H(\lambda_1(c)) - q \exp\{[\gamma + \lambda_1(c)]\xi\} H(\gamma + \lambda_1(c)), \\ h * * \phi^-(\xi) &\geq \int_0^\infty \int_{\mathbb{R}} h(y, s) \exp\{\lambda_1(c)(\xi - y - cs)\} (1 - q \exp\{\gamma(\xi - y - cs)\}) dy ds \\ &= \exp\{\lambda_1(c)\xi\} G(c, \lambda_1(c)) - q \exp\{[\gamma + \lambda_1(c)]\xi\} G(c, \gamma + \lambda_1(c)). \end{aligned}$$

Let $\xi_0 = -\frac{1}{\gamma} \ln q$. Clearly, $\phi^-(\xi) = 0$ for $\xi > \xi_0$ and $\phi^-(\xi) = \exp\{\lambda_1(c)\xi\} (1 - q \exp\{\gamma\xi\})$ for $\xi \leq \xi_0$. For $\xi > \xi_0$, we have

$$N_c[\phi^-](\xi) = -dJ * \phi^-(\xi) - f(0, (h * * \phi^-)(\xi)) \leq -f(0, 0) = 0.$$

For $\xi \leq \xi_0$, we have

$$\begin{aligned}
N_c[\phi^-](\xi) &\leq \exp\{\lambda_1(c)\xi\}\{c\lambda_1(c) - p\lambda_1^2(c) - d[H(\lambda_1(c)) - 1]\} \\
&\quad - q \exp\{[\gamma + \lambda_1(c)]\xi\}\{c[\gamma + \lambda_1(c)] - p[\gamma + \lambda_1(c)]^2 - d[H(\gamma + \lambda_1(c)) - 1]\} \\
&\quad - f(\phi^-(\xi), (h * \phi^-)(\xi)) \\
&= \exp\{\lambda_1(c)\xi\}\Delta(c, \lambda_1(c)) - q \exp\{[\gamma + \lambda_1(c)]\xi\}\Delta(c, \gamma + \lambda_1(c)) \\
&\quad - f(\phi^-(\xi), (h * \phi^-)(\xi)) + \partial_1 f(0, 0)\phi^-(\xi) + \partial_2 f(0, 0) \exp\{\lambda_1(c)\xi\}G(c, \lambda_1(c)) \\
&\quad - q\partial_2(0, 0) \exp\{[\gamma + \lambda_1(c)]\xi\}G(c, \gamma + \lambda_1(c)) \\
&\leq -q \exp\{[\gamma + \lambda_1(c)]\xi\}\Delta(c, \gamma + \lambda_1(c)) - f(\phi^-(\xi), (h * \phi^-)(\xi)) \\
&\quad + \partial_1 f(0, 0)\phi^-(\xi) + \partial_2 f(0, 0)(h * \phi^-)(\xi).
\end{aligned}$$

In view of (F2), we have

$$\begin{aligned}
N_c[\phi^-](\xi) &\leq -q \exp\{[\gamma + \lambda_1(c)]\xi\}\Delta(c, \gamma + \lambda_1(c)) + M[\phi^-(\xi) + (h * \phi^-)(\xi)]^{1+\sigma} \\
&\leq -q \exp\{[\gamma + \lambda_1(c)]\xi\}\Delta(c, \gamma + \lambda_1(c)) + M[1 + G(c, \gamma + \lambda_1(c))]^{1+\sigma} \exp\{(1 + \sigma)\lambda_1(c)\xi\} \\
&\leq \exp\{[\gamma + \lambda_1(c)]\xi\}\Delta(c, \gamma + \lambda_1(c)) \left\{ \frac{M[1 + G(c, \gamma + \lambda_1(c))]^{1+\sigma}}{\Delta(c, \gamma + \lambda_1(c))} - q \right\} \leq 0,
\end{aligned}$$

provided that

$$q \geq Q(c, \eta) \triangleq \max \left\{ 1, \frac{M[1 + G(c, \gamma + \lambda_1(c))]^{1+\sigma}}{\Delta(c, \gamma + \lambda_1(c))} \right\}.$$

Therefore, ϕ^- is a subsolution of (6). The proof is complete. \square

As a consequence of Lemmas 3.1, 3.2, and 3.3, we have the following result on the existence of increasing travelling waves.

Theorem 3.1 *Under the conditions (F1), (F2), (H1) and (H2), let c^* , $\lambda_1(c)$ and $\lambda_2(c)$ be defined as in Lemma 2.1. Then for each $c > c^*$, (6) admits a solution (c, ϕ) satisfying $0 < \phi(\xi) < K$, $\phi'(\xi) > 0$ for all $\xi \in \mathbb{R}$, and*

$$\lim_{\xi \rightarrow -\infty} \phi(\xi) \exp\{-\lambda\xi\} = 1, \quad \lim_{\xi \rightarrow -\infty} \phi'(\xi) \exp\{-\lambda\xi\} = \lambda, \quad (18)$$

where $\lambda = \lambda_1(c)$ is the smallest positive zero of $\Delta(c, \cdot)$.

Proof. It follows from Lemmas 2.3, 3.1, 3.2, and 3.3 that there exists a strictly increasing solution $\phi(\xi)$ to (6), which will be denoted by (c, ϕ) and satisfies

$$\exp\{\lambda_1(c)\xi\} (1 - q \exp\{\gamma\xi\}) \leq \phi(\xi) \leq \exp\{\lambda_1(c)\xi\}, \quad \xi \in \mathbb{R}. \quad (19)$$

It then follows from (19) that

$$\lim_{\xi \rightarrow -\infty} |\phi(\xi) \exp\{-\lambda_1(c)\xi\} - 1| \leq \lim_{\xi \rightarrow -\infty} q \exp\{\gamma\xi\} = 0.$$

In view of condition (F2), we have

$$\begin{aligned}
&\lim_{\xi \rightarrow -\infty} |f(\phi(\xi), (h * \phi)(\xi)) - \partial_1 f(0, 0)\phi(\xi) - \partial_2 f(0, 0)(h * \phi)(\xi)| \exp\{-\lambda_1(c)\xi\} \\
&\leq M \lim_{\xi \rightarrow -\infty} [\phi(\xi) + (h * \phi)(\xi)]^{1+\sigma} \exp\{-\lambda_1(c)\xi\} \\
&= M \left[\lim_{\xi \rightarrow -\infty} \phi(\xi) e^{-\lambda_1(c)\xi/(1+\sigma)} + \lim_{\xi \rightarrow -\infty} (h * \phi)(\xi) \exp\{-\lambda_1(c)\xi/(1 + \sigma)\} \right]^{1+\sigma} \\
&= M \left[\lim_{\xi \rightarrow -\infty} (h * \phi)(\xi) e^{-\lambda_1(c)\xi/(1+\sigma)} \right]^{1+\sigma} \\
&\leq M \left[g'(0) \lim_{\xi \rightarrow -\infty} (h * \phi)(\xi) e^{-\lambda_1(c)\xi/(1+\sigma)} \right]^{1+\sigma}
\end{aligned}$$

and

$$\begin{aligned}
& \lim_{\xi \rightarrow -\infty} (h * \phi)(\xi) \exp\{-\lambda_1(c)\xi/(1+\sigma)\} \\
&= \lim_{\xi \rightarrow -\infty} \int_0^\infty \int_{-\infty}^\infty h(y,t)\phi(\xi-y-ct) \exp\{-\lambda_1(c)\xi/(1+\sigma)\} dy dt \\
&= \int_0^\infty \int_{-\infty}^\infty h(y,t) \left[\lim_{\xi \rightarrow -\infty} \phi(\xi-y-ct) \exp\{-\lambda_1(c)\xi/(1+\sigma)\} \right] dy dt = 0.
\end{aligned}$$

Hence, if $p = 0$ then for $c > c^*$, we have

$$\begin{aligned}
& c \lim_{\xi \rightarrow -\infty} \phi'(\xi) \exp\{-\lambda_1(c)\xi\} \\
&= \lim_{\xi \rightarrow -\infty} [d(J * \phi - \phi)(\xi) + f(\phi(\xi), (h * \phi)(\xi))] \exp\{-\lambda_1(c)\xi\} \\
&= d[H(\lambda_1(c)) - 1] + \lim_{\xi \rightarrow -\infty} f(\phi(\xi), (h * \phi)(\xi)) \exp\{-\lambda_1(c)\xi\} \\
&= d[H(\lambda_1(c)) - 1] + \lim_{\xi \rightarrow -\infty} [\partial_1 f(0,0)\phi(\xi) + \partial_2 f(0,0)(h * \phi)(\xi)] \exp\{-\lambda_1(c)\xi\} \\
&= d[H(\lambda_1(c)) - 1] + \partial_1 f(0,0) + \partial_2 f(0,0)G(c, \lambda_1(c)) = c\lambda_1(c).
\end{aligned}$$

If $p \neq 0$ then for $c > c^*$, using $\phi'(-\infty) = 0$ and integrating both sides of (6) from $-\infty$ to ξ , we have

$$\begin{aligned}
& \lim_{\xi \rightarrow -\infty} \phi'(\xi) \exp\{-\lambda_1(c)\xi\} \\
&= \frac{c}{p} - \frac{1}{p} \lim_{\xi \rightarrow -\infty} \exp\{-\lambda_1(c)\xi\} \int_{-\infty}^\xi [d(J * \phi - \phi)(s) + f(\phi(s), (h * \phi)(s))] ds \\
&= \frac{c}{p} - \lim_{\xi \rightarrow -\infty} \frac{\exp\{-\lambda_1(c)\xi\} [d(J * \phi - \phi)(\xi) + f(\phi(\xi), (h * \phi)(\xi))]}{p\lambda_1(c)} \\
&= \frac{c}{p} - \frac{d[H(\lambda_1(c)) - 1] + \partial_1 f(0,0) + \partial_2 f(0,0)G(c, \lambda_1(c))}{p\lambda_1(c)} = \lambda_1(c).
\end{aligned}$$

This completes the proof. \square

Remark 3.1 In Theorem 3.1, by Lebesgue's dominated convergence theorem, we also have

$$\lim_{\xi \rightarrow -\infty} (h * \phi)(\xi) \exp\{-\lambda_1(c)\xi\} = G(c, \lambda_1(c)).$$

Remark 3.2 Theorem 3.1 implies that the asymptotic behaviours of wave profiles of the travelling waves obtained by super- and sub-solutions satisfy (18) for $c > c^*$. Furthermore, in the subsequent section, we shall show that the wave profile φ of every travelling wave of (1) satisfying (6) has similar asymptotic behaviours.

Next, we prove that (6) has a solution (c, ϕ) with $0 < \phi < K$ and $\phi' > 0$ on \mathbb{R} for $c = c^*$.

Theorem 3.2 Under the conditions (F1), (F2), (H1) and (H2), (6) has a solution (c, ϕ) with $0 < \phi < K$ and $\phi' > 0$ on \mathbb{R} for $c = c^*$.

Proof. We choose a sequence $\{c_j\} \subseteq (c^*, \infty)$ such that $\lim_{j \rightarrow \infty} c_j = c^*$. Then for each j there exists a strictly increasing travelling wave (c_j, ϕ_j) of (1) such that $\phi_j(-\infty) = 0$ and $\phi_j(+\infty) = K$. Since $\phi_j(\cdot + \zeta)$, $\zeta \in \mathbb{R}$, is also a travelling wave, we can assume that $\phi_j(0) = \alpha$ and $\phi_j(x) \leq K$ for a fixed $\alpha \in (0, K)$ and all $x \in \mathbb{R}$ and $j \geq 1$. Note that ϕ_j is a fixed point of operator T_μ in E with $c = c_j$ and $T_\mu(\phi_j)(\xi)$ can be differentiated with respect to ξ , where E is the Banach space of bounded and uniformly continuous functions on \mathbb{R} equipped with the maximum norm. Moreover, we differentiate both sides of (6) with respect to ξ to obtain

$$\begin{aligned}
0 &= -c\phi_j''(\xi) + p\phi_j'''(\xi) + d[J * \phi_j' - \phi_j'](\xi) + \partial_1 f(\phi_j(\xi), (h * \phi_j)(\xi))\phi_j'(\xi) \\
&\quad + \partial_2 f(\phi_j(\xi), (h * \phi_j)(\xi))h * \phi_j'(\xi).
\end{aligned}$$

By the definition of \mathcal{H}_μ , it follows that there exist three positive numbers N_1 , N_2 , and N_3 (if $p \neq 0$) such that

$$|\phi'_j(\xi)| \leq N_1, \quad |\phi''_j(\xi)| \leq N_2, \quad |\phi'''_j(\xi)| \leq N_3$$

for all n and ξ . Therefore, ϕ'_j , ϕ''_j and ϕ'''_j (if $p \neq 0$) are uniformly bounded and equi-continuous sequences of functions on \mathbb{R} . Then the Arzelà-Ascoli theorem implies that there exists a subsequence of $\{c_j\}$ (for simplicity, denoted again by $\{c_j\}$), such that $\lim_{j \rightarrow \infty} c_j = c^*$, and ϕ'_j , ϕ''_j and ϕ'''_j (if $p \neq 0$) converge uniformly on every bounded and closed subset of \mathbb{R} . Thus, ϕ'_j , ϕ''_j and ϕ'''_j (if $p \neq 0$) converge pointwise on \mathbb{R} to ϕ'_* , ϕ''_* and ϕ'''_* (if $p \neq 0$), respectively. By Lebesgue's dominated convergence theorem, letting $j \rightarrow \infty$ in the equation $\phi_j = T_\mu(\phi_j)$, we then get $\phi_* = T_\mu(\phi_*)$. Thus, ϕ_* is a solution of (6) in the case where $c = c_*$. Clearly, ϕ_* is monotonically increasing on \mathbb{R} , $\phi_*(0) = \alpha$ and $\phi_*(x) \leq K$ for all $x \in \mathbb{R}$. One can easily verify that $\phi_*(-\infty) = 0$ and $\phi_*(+\infty) = K$. Thus, (1) has a monotone travelling wave solution connecting 0 and K with the wave speed $c = c_*$. This completes the proof. \square

4 Asymptotic behaviour

In this section, we always assume that (F1), (F2), (H1) and (H2) hold, and that c^* , λ^* , and $\lambda_1(c)$ are defined as in Lemma 2.1. We shall follow a method of Carr and Chmaj [9] and Wang, Li, and Ruan [39] to establish the exact asymptotic behaviour of the profile $\phi(\xi)$ as $\xi \rightarrow -\infty$. For this purpose, we need Ikehara's theorem on the asymptotic behaviour of a positive decreasing function whose Laplace is of a certain given shape. The proof of Ikehara's theorem can be found, e.g., in [9, 41].

Theorem 4.1 (Ikehara's theorem) *Let $\mathcal{L}[u](\mu) = \int_0^\infty \exp\{-\mu\xi\}u(\xi)d\xi$ be the Laplace transform of u , with u being a positive non-decreasing function. Assume that $\mathcal{L}[u]$ has the representation*

$$\mathcal{L}[u](\mu) = \frac{E(\mu)}{(\mu + \alpha)^{k+1}},$$

where $k > -1$ and E is analytic in the strip $-\alpha \leq \operatorname{Re}\mu < 0$. Then

$$\lim_{\xi \rightarrow \infty} \frac{u(\xi)}{\xi^k e^{-\alpha\xi}} = \frac{E(-\alpha)}{\Gamma(\alpha + 1)},$$

where Γ is the Gamma function.

Lemma 4.1 *Assume that (F1), (F2), (H1) and (H2) hold. Let (c, φ) be a solution of (6). Then there exists $\gamma > 0$ such that*

$$\sup_{\xi \in \mathbb{R}} \varphi(\xi) \exp\{-\gamma\xi\} < \infty \quad \text{and} \quad \sup_{\xi \in \mathbb{R}} \Phi(\xi) \exp\{-\gamma\xi\} < \infty, \quad (20)$$

where $\Phi(\xi) \triangleq \int_{-\infty}^\xi \varphi(s)ds$.

Proof. For each $\xi \in \mathbb{R}$, define

$$\psi(\xi) \triangleq (h * \varphi)(\xi) = \int_0^\infty \int_{\mathbb{R}} h(y, \tau) \varphi(\xi - y - c\tau) dy d\tau.$$

In view of $\partial_2 f(0, 0) + \partial_1 f(0, 0) > 0$, there exist $\delta_0 \in (0, K)$ such that

$$\varepsilon_1 \triangleq [\partial_2 f(0, 0) + \partial_1 f(0, 0)]/4 \geq M(u + v)^\sigma \quad (21)$$

for all $u, v \in [0, \delta_0]$. In view of (22) and $\varphi(-\infty) = 0$, there exists $\xi_0 < 0$ such that for all $\xi < \xi_0$, both $\varphi(\xi)$ and $\psi(\xi)$ lie in the interval $(0, \delta_0)$, where δ_0 is defined as (21). Thus, for every $\xi < \xi_0$,

$$\begin{aligned} f(\varphi(\xi), \psi(x)) &\geq \partial_1 f(0, 0)\varphi(\xi) + \partial_2 f(0, 0)\psi(\xi) - M[\varphi(\xi) + \psi(\xi)]^{1+\sigma} \\ &\geq \partial_1 f(0, 0)\varphi(\xi) + \partial_2 f(0, 0)\psi(\xi) - \varepsilon_1[\varphi(\xi) + \psi(\xi)] \\ &= (\varepsilon_1 - 2\varepsilon_2)\varphi(\xi) + (\varepsilon_1 + 2\varepsilon_2)\psi(\xi) \end{aligned} \quad (22)$$

for all $u, v \in [0, \delta_0]$, where $\varepsilon_2 = [\partial_2 f(0, 0) - \partial_1 f(0, 0)]/4$. Therefore,

$$c\varphi'(\xi) - p\varphi''(\xi) - d(J * \varphi - \varphi)(\xi) \geq (\varepsilon_1 - 2\varepsilon_2)\varphi(\xi) + (\varepsilon_1 + 2\varepsilon_2)\psi(\xi) \quad (23)$$

for all $\xi < \xi_0$. By using a similar argument as in the proof of Theorem 3.2, we can prove that $\varphi(\xi)$ and $\psi(\xi)$ are both integrable on $(-\infty, 0]$.

By Fubini's theorem and Lebesgue's dominated convergence theorem

$$\begin{aligned} \int_{-\infty}^{\xi} \psi(s) ds &= \int_{-\infty}^{\xi} \left[\int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \varphi(s - y - c\tau) dy d\tau \right] ds \\ &= \lim_{z \rightarrow -\infty} \int_z^{\xi} \left[\int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \varphi(s - y - c\tau) dy d\tau \right] ds \\ &= \lim_{z \rightarrow -\infty} \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \left[\int_z^{\xi} \varphi(s - y - c\tau) ds \right] dy d\tau \\ &= \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \left[\int_{-\infty}^{\xi} \varphi(s - y - c\tau) ds \right] dy d\tau \\ &= \int_0^{\infty} \int_{\mathbb{R}} h(y) \Phi(\xi - y - c\tau) dy d\tau. \end{aligned}$$

Integrating (23) from $-\infty$ to ξ with $\xi < \xi_0$, we have (using again Fubini's theorem)

$$\begin{aligned} c\varphi(\xi) - p\varphi'(\xi) - d(J * \Phi - \Phi)(\xi) &\geq (\varepsilon_1 - 2\varepsilon_2)\Phi(\xi) + (\varepsilon_1 + 2\varepsilon_2) \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \Phi(\xi - y - c\tau) dy d\tau \\ &= \varepsilon_1 \Phi(\xi) + \varepsilon_1 \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \Phi(\xi - y - c\tau) dy d\tau \\ &\quad + 2\varepsilon_2 \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) [\Phi(\xi - y - c\tau) - \Phi(\xi)] dy d\tau. \end{aligned} \quad (24)$$

Note that

$$\begin{aligned} &\int_z^{\xi} \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) [\Phi(s - y - c\tau) - \Phi(s)] dy d\tau ds \\ &= - \int_0^{\infty} \int_{\mathbb{R}} (y + c\tau) h(y, \tau) \int_0^1 [\Phi(\xi - t(y + c\tau)) - \Phi(z - t(y + c\tau))] dt dy d\tau \\ &\rightarrow - \int_0^{\infty} \int_{\mathbb{R}} (y + c\tau) h(y, \tau) \int_0^1 \Phi(\xi - t(y + c\tau)) dt dy d\tau \end{aligned}$$

as $z \rightarrow -\infty$. Thus, (24) means that $\Phi(\xi)$ and $\int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \Phi(s - y - c\tau) dy d\tau$ are integrable on $(-\infty, \xi]$. Moreover,

$$\begin{aligned} c\Phi(\xi) - p\varphi(\xi) - d \int_{-\infty}^{\xi} (J * \Phi - \Phi)(s) ds + 2\varepsilon_2 \int_0^{\infty} \int_{\mathbb{R}} (y + c\tau) h(y, \tau) \int_0^1 \Phi(\xi - t(y + c\tau)) dt dy d\tau \\ \geq \varepsilon_1 \int_{-\infty}^{\xi} \Phi(s) ds + \varepsilon_1 \int_{-\infty}^{\xi} \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \Phi(s - y - c\tau) dy d\tau ds. \end{aligned} \quad (25)$$

Since $\Phi(\xi)$ is increasing, for any $y \in \mathbb{R}$ we have

$$(y + c\tau) h(y, \tau) \Phi(\xi) \geq (y + c\tau) h(y, \tau) \int_0^1 \Phi(\xi - t(y + c\tau)) dt.$$

If $\varepsilon_2 \geq 0$, then it follows from (25) that

$$c\Phi(\xi) - p\varphi(\xi) - d \int_{-\infty}^{\xi} (J * \Phi - \Phi)(s)ds + 2\varepsilon_2\Phi(\xi) \int_0^{\infty} \int_{\mathbb{R}} (y + c\tau)h(y, \tau)dyd\tau \geq \varepsilon_1 \int_{-\infty}^{\xi} \Phi(s)ds. \quad (26)$$

The Mean Value Theorem for Integrals implies that for each $y > 0$, there exist $\xi_1(y) \in (\xi, \xi + y)$ and $\xi_2(y) \in (\xi - y, \xi)$ such that $\int_{\xi}^{\xi+y} \Phi(s)ds = y\Phi(\xi_1(y))$ and $\int_{\xi-y}^{\xi} \Phi(s)ds = y\Phi(\xi_2(y))$. It follows from the monotonicity of Φ that

$$\begin{aligned} \int_{-\infty}^{\xi} (J * \Phi - \Phi)(s)ds &= \int_{-\infty}^{\xi} \int_{\mathbb{R}} J(y)[\Phi(x - y) - \Phi(x)]dydx \\ &= \int_{\mathbb{R}} J(y) \int_{\xi}^{\xi-y} \Phi(x)dx dy \\ &= \int_0^{\infty} J(y) \int_{\xi}^{\xi-y} \Phi(x)dx dy + \int_{-\infty}^0 J(y) \int_{\xi}^{\xi-y} \Phi(x)dx dy \\ &= \int_0^{\infty} J(y) \int_{\xi}^{\xi-y} \Phi(x)dx dy + \int_0^{\infty} J(-y) \int_{\xi}^{\xi+y} \Phi(x)dx dy \\ &= \int_0^{\infty} J(y) \left[\int_{\xi}^{\xi+y} \Phi(x)dx - \int_{\xi-y}^{\xi} \Phi(x)dx \right] dy \\ &= \int_0^{\infty} yJ(y)[\Phi(\xi_1(y)) - \Phi(\xi_2(y))]dy > 0. \end{aligned}$$

This, together with (26), implies that

$$\left[c + 2\varepsilon_2 \int_0^{\infty} \int_{\mathbb{R}} (y + c\tau)h(y, \tau)dyd\tau \right] \Phi(\xi) \geq \varepsilon_1 \int_{-\infty}^{\xi} \Phi(s)ds + p\varphi(\xi). \quad (27)$$

If $\varepsilon_2 < 0$, that is, $\partial_1 f(0, 0) > \partial_2 f(0, 0) \geq 0$, then there exists $\xi'_0 < \xi_0$ such that

$$c\varphi'(\xi) - p\varphi''(\xi) - d(J * \varphi - \varphi)(\xi) = f(\varphi(\xi), \psi(\xi)) \geq f(\varphi(\xi), 0) \geq \frac{1}{2}\partial_1 f(0, 0)\varphi(\xi) > \varepsilon_1\varphi(\xi)$$

for all $\xi < \xi'_0$. Thus

$$c\Phi(\xi) \geq \varepsilon_1 \int_{-\infty}^{\xi} \Phi(s)ds + p\varphi(\xi). \quad (28)$$

Combing (27) and (28), we have

$$\left[c + 2|\varepsilon_2| \int_0^{\infty} \int_{\mathbb{R}} |y + c\tau|h(y, \tau)dyd\tau \right] \Phi(\xi) \geq \varepsilon_1 \int_{-\infty}^0 \Phi(s+\xi)ds \geq \varepsilon_1 \int_{-r}^0 \Phi(s+\xi)ds \geq \varepsilon_1 r\Phi(\xi-r)$$

for all $r > 0$ and $\xi < \xi'_0$. Thus there exists $r_0 > 0$ and some $\theta \in (0, 1)$ such that

$$\gamma \triangleq \frac{1}{r_0} \ln \frac{1}{\theta} \in (\lambda_1(c), \lambda_0) \quad (29)$$

and

$$\Phi(\xi - r_0) \leq \theta\Phi(\xi)$$

uniformly in ξ . Thus

$$\Phi(\xi - r_0) \exp\{-\gamma(\xi - r_0)\} \leq \Phi(\xi) \exp\{-\gamma\xi\}.$$

This, together with $\lim_{\xi \rightarrow \infty} \Phi(\xi)e^{-\gamma\xi} = 0$, implies that

$$\sup_{\xi \in \mathbb{R}} \{\Phi(\xi) \exp\{-\gamma\xi\}\} < \infty. \quad (30)$$

Moreover,

$$\begin{aligned}\exp\{-\gamma\xi\} \int_{-\infty}^{\xi} \Phi(s)ds &= \exp\{-\gamma\xi\} \int_{-\infty}^0 \Phi(\xi+s)ds \\ &= \int_{-\infty}^0 \Phi(\xi+s) \exp\{-\gamma(\xi+s)\} \exp\{\gamma s\}ds \leq \frac{1}{\gamma} \sup_{\xi \in \mathbb{R}} \{\Phi(\xi) \exp\{-\gamma\xi\}\}.\end{aligned}$$

Thus, it follows from (27), (28), and (30) that

$$\sup_{\xi \in \mathbb{R}} \{\varphi(\xi) \exp\{-\gamma\xi\}\} < \infty. \quad (31)$$

This completes the proof. \square

Lemma 4.2 *Assume that (F1), (F2), (H1) and (H2) hold. Let (c, φ) be a solution of (6). Then $\lim_{\xi \rightarrow -\infty} \varphi(\xi) \exp\{-\lambda_1(c)\xi\}$ exists for each $c > c^*$.*

Proof. Define a bilateral Laplace transform of $\varphi(\xi)$ by

$$L[\varphi](\lambda) = \int_{\mathbb{R}} \exp\{-\lambda\xi\} \varphi(\xi) d\xi.$$

By Lemma 4.1 and Fubini's theorem, we have

$$\begin{aligned}\int_{\mathbb{R}} e^{-\lambda\xi} \psi(\xi) d\xi &= \int_{\mathbb{R}} \left[e^{-\lambda\xi} \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \varphi(\xi - y - c\tau) dy \right] d\xi d\tau \\ &= \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) \left[\int_{\mathbb{R}} e^{-\lambda\xi} \varphi(\xi - y - c\tau) d\xi \right] dy d\tau \\ &= L(\lambda) \int_0^{\infty} \int_{\mathbb{R}} h(y, \tau) e^{-\lambda(y+c\tau)} dy d\tau = L(\lambda) G(c, \lambda).\end{aligned}$$

Take the bilateral Laplace transform of (6) with respect to ξ , we have (with $\Delta(c, \lambda)$ defined in (11))

$$\Delta(c, \lambda) L[\varphi](\lambda) = \mathcal{R}(\lambda), \quad (32)$$

where $\mathcal{R}(\lambda)$ is the Laplace transform of the function $f(\varphi(\xi), \psi(\xi)) - \partial_1 f(0, 0) \varphi(\xi) - \partial_2 f(0, 0) \psi(\xi)$. It is not difficult to see that $\mathcal{R}(\lambda)$ is defined for λ with $0 < \operatorname{Re}\lambda < \gamma$. In addition, it is easy to see that $\Delta(c, \cdot)$ has no zero λ with $\operatorname{Re}\lambda = \lambda_1(c)$ other than $\lambda = \lambda_1(c)$. This implies that $(\lambda - \lambda_1(c))/\Delta(c, \lambda)$ is analytic in the strip $0 < \operatorname{Re}\lambda \leq \lambda_1(c)$. If there exists some $\xi_0 > 0$ such that $\varphi(\xi)$ is increasing for all $\xi \in (-\infty, -\xi_0)$, then $u(\xi) = \varphi(-\xi)$ is a positive decreasing function on (ξ_0, ∞) . Moreover, it follows from (32) that

$$\int_{\xi_0}^{\infty} e^{\lambda\xi} u(\xi) d\xi = \int_{-\infty}^{-\xi_0} e^{-\lambda\xi} \varphi(\xi) d\xi = \frac{\mathcal{E}(\lambda)}{\lambda - \lambda_1(c)} \quad (33)$$

with

$$\mathcal{E}(\lambda) = \frac{(\lambda - \lambda_1(c))\mathcal{R}(\lambda)}{\Delta(c, \lambda)} - (\lambda - \lambda_1(c)) \int_{-\xi_0}^{\infty} e^{-\lambda\xi} \varphi(\xi) d\xi,$$

which is analytic in the strip $0 < \operatorname{Re}\lambda \leq \lambda_1(c)$ because $\int_{-\xi_0}^{\infty} \exp\{-\lambda\xi\} \varphi(\xi) d\xi$ is analytic for all $\operatorname{Re}\lambda > 0$. By means of Theorem 4.1, $\lim_{\xi \rightarrow \infty} u(\xi) \exp\{\lambda_1(c)\xi\}$, which is equal to $\lim_{\xi \rightarrow -\infty} \varphi(\xi) \exp\{-\lambda_1(c)\xi\}$, exists.

If $\varphi(\xi)$ is not monotone on any interval $(-\infty, \xi_0)$ with $|\xi_0|$ sufficiently large, let $\chi(\xi) = \exp\{q\xi\} \varphi(\xi)$, where $q = d/c$ if $p = 0$ and

$$q = \frac{-c + \sqrt{c^2 + 4pd}}{2p}$$

if $p > 0$. Then

$$(c + 2pq)\chi'(\xi) - p\chi''(\xi) = \exp\{q\xi\} [dJ * \varphi(\xi) + f(\varphi(\xi), \psi(\xi))] \geq 0.$$

Suppose that there exists $\xi_1 < \xi_2$ such that $\chi(\xi_1) > \chi(\xi_2)$. Note that $\lim_{\xi \rightarrow \infty} \chi(\xi) = +\infty$, thus there exists $\xi_3 > \xi_1$ such that $\chi'(\xi_3) = 0$ and $\chi''(\xi_2) \geq 0$, which contradicts the equation above. Thus, we have $\chi'(\xi) \geq 0$. Then for the bilateral Laplace transform of $\chi(\xi)$, $L[\chi](\lambda) = L(\lambda - q)$. It follows from (32) that

$$\Delta(c, \lambda - q)L_1(\lambda) = \mathcal{R}(\lambda - q).$$

Using a similar argument as above, we see that

$$\lim_{\xi \rightarrow -\infty} \varphi(\xi) \exp\{-\lambda_1(c)\xi\} = \lim_{\xi \rightarrow -\infty} \chi(\xi) \exp\{-(q + \lambda_1(c))\xi\}$$

exists. This completes the proof. \square

Using a similar argument as in the proof of the previous lemma, we can verify the following result.

Lemma 4.3 *Assume that (F1), (F2), (H1) and (H2) hold. Let (c^*, φ) be a solution of (6). Then $\lim_{\xi \rightarrow -\infty} \varphi(\xi)\xi^{-1} \exp\{-\lambda^*\xi\}$ exists.*

Now we are ready to summarise the asymptotic behaviour of wave profile φ as follows.

Theorem 4.2 *Under assumptions (F1), (F2), (H1) and (H2), for each solution of (c, φ) of (6) there exists $\eta = \eta(\varphi)$ such that*

$$\lim_{\xi \rightarrow -\infty} \frac{\varphi(\xi + \eta)}{\exp\{\lambda_1(c)\xi\}} = 1 \quad \text{for } c > c^* \quad (34)$$

and

$$\lim_{\xi \rightarrow -\infty} \frac{\varphi(\xi + \eta)}{\xi \exp\{\lambda_1(c)\xi\}} = 1 \quad \text{for } c = c^*. \quad (35)$$

Moreover,

$$\lim_{\xi \rightarrow -\infty} \frac{\varphi'(\xi)}{\varphi(\xi)} = \lambda_1(c) \quad \text{for } c \geq c^*. \quad (36)$$

Proof. Both (34) and (35) follow easily from Lemmas 4.2 and 4.3. It follows from (6) that

$$\begin{aligned} \lim_{\xi \rightarrow -\infty} \frac{c\varphi'(\xi)}{\varphi(\xi)} &= p \lim_{\xi \rightarrow -\infty} \frac{\varphi''(\xi)}{\varphi(\xi)} + d \lim_{\xi \rightarrow -\infty} \left[\frac{J * \varphi(\xi)}{\varphi(\xi)} - 1 \right] + \lim_{\xi \rightarrow -\infty} \frac{f(\varphi(\xi), \psi(\xi))}{\varphi(\xi)} \\ &= p\lambda_1^2(c) + d[H(\lambda_1(c)) - 1] + \partial_1 f(0, 0) + \partial_2 f(0, 0) \lim_{\xi \rightarrow -\infty} \frac{\psi(\xi)}{\varphi(\xi)} \\ &= p\lambda_1^2(c) + d[H(\lambda_1(c)) - 1] + \partial_1 f(0, 0) \\ &\quad + \partial_2 f(0, 0) \lim_{\xi \rightarrow -\infty} \int_0^\infty \int_{\mathbb{R}} h(y, \tau) \frac{\varphi(\xi - y - c\tau)}{\varphi(\xi)} dy d\tau \\ &= p\lambda_1^2(c) + d[H(\lambda_1(c)) - 1] + \partial_1 f(0, 0) + \partial_2 f(0, 0)G(c, \lambda_1(c)) \\ &= c\lambda_1(c). \end{aligned}$$

This completes the proof. \square

Theorem 4.3 *Under assumptions (F1), (F2), (H1) and (H2), for each solution of (c, φ) of (6) there exists $\eta = \eta(\varphi)$ such that*

$$\lim_{\xi \rightarrow \infty} \frac{K - \varphi(\xi + \eta)}{\exp\{-v(c)\xi\}} = 1, \quad (37)$$

where $v(c)$ is the unique positive zero of $\tilde{\Delta}(c, \cdot)$, according to Lemma 2.2. Moreover,

$$\lim_{\xi \rightarrow \infty} \frac{\varphi'(\xi)}{K - \varphi(\xi)} = v(c). \quad (38)$$

Proof. Define $\Phi(\xi) \triangleq K - \varphi(-\xi)$ and $\Psi(\xi) \triangleq K - (h * * \varphi)(-\xi)$. Obviously, $\Phi(\xi)$ satisfies that $\Phi(-\infty) = 0$, $\Phi(\infty) = K$, $0 < \Phi(\xi) < K$, and

$$c\Phi'(\xi) = -p\Phi''(\xi) + f(K - \Phi(\xi), K - \Psi(\xi)) - d(J * \Phi - \Phi)(\xi). \quad (39)$$

Then for any $\mu > 2d + \max\{|\partial_1 f(u, v)| : u, v \in [0, K]\}$, we have $[\Phi(\xi)e^{-\mu\xi}]' < 0$ for all ξ . Then, using the bilateral Laplace transform $L[\Phi]$ of $\Phi(\xi)$, we have, using Fubini's theorem again,

$$\begin{aligned} \int_{\mathbb{R}} e^{-\lambda\xi} \Psi(\xi) d\xi &= \int_{\mathbb{R}} e^{-\lambda\xi} \left[K - \int_0^\infty \int_{\mathbb{R}} h(y, \tau) \varphi(-\xi - y - c\tau) dy d\tau \right] d\xi \\ &= \int_{\mathbb{R}} \left[e^{-\lambda\xi} \int_0^\infty \int_{\mathbb{R}} h(y, \tau) \Phi(\xi + y + c\tau) dy d\tau \right] d\xi \\ &= \int_0^\infty \int_{\mathbb{R}} h(y, \tau) \left[\int_{\mathbb{R}} e^{-\lambda\xi} \Phi(\xi + y + c\tau) d\xi \right] dy d\tau \\ &= L[\Phi](\lambda) \int_0^\infty \int_{\mathbb{R}} h(y, \tau) e^{\lambda(y+c\tau)} dy d\tau = L[\Phi](\lambda) G(c, -\lambda). \end{aligned}$$

Take Laplace transform of (39) with respect to ξ , we have

$$\tilde{\Delta}(c, \lambda) \tilde{L}(\lambda) = \tilde{\mathcal{R}}(\lambda), \quad (40)$$

where $\tilde{\mathcal{R}}(\lambda)$ is the Laplace transform of the function $f(K - \Phi(\xi), K - \Psi(\xi)) - \partial_1 f(K, K)\Phi(\xi) - \partial_2 f(K, K)\Psi(\xi)$. Using a similar arguments as that in the proof of Lemma 4.2, we see that there exists $\eta = \eta(\varphi)$ such that

$$\lim_{\xi \rightarrow -\infty} \frac{\Phi(\xi + \eta)}{\exp\{v(c)\xi\}} = 1 \quad \text{and} \quad \lim_{\xi \rightarrow -\infty} \frac{\Phi'(\xi)}{\Phi(\xi)} = \lambda_1(c), \quad (41)$$

from which (37) and (38) follows. The proof is completed. \square

5 Monotonicity and Uniqueness

In this section, we investigate the monotonicity and uniqueness (up to a translation) of the travelling wavefront of (6) by using the sliding method developed in Chen and Guo [12].

Theorem 5.1 *Under assumptions (F1), (F2), (H1), and (H2), every solution (c, φ) of (6) satisfies $\varphi'(\xi) > 0$ for all $\xi \in \mathbb{R}$.*

Proof. It follows from Lemma 3.1 and Theorems 4.2 and 4.3 that there exists $M > 0$ such that $\varphi'(\xi) > 0$ for all $|\xi| \geq M$. It thus suffices to show that $\varphi'(\xi) > 0$ for all $\xi \in [-M, M]$. Suppose on the contrary that $\varphi'(\xi) \leq 0$ for some $\xi_0 \in [-M, M]$. By continuity of $\varphi'(\xi)$, there exists $\xi_1 \in [-M, M]$ such that $\varphi'(\xi_1) = 0$. Then, using the similar arguments as in the proof of Lemma 3.2, we obtain a contradiction. This completes the proof. \square

In order to prove the uniqueness up to a translation, we shall need the following strong comparison principle.

Lemma 5.1 *Let (c, ϕ_1) and (c, ϕ_2) be solutions of (6) with $\phi_1 \geq \phi_2$ on \mathbb{R} . Then either $\phi_1 \equiv \phi_2$ or $\phi_1 > \phi_2$ on \mathbb{R} .*

Proof. Suppose that there exists some $\xi_0 \in \mathbb{R}$ such that $\phi_1(\xi_0) = \phi_2(\xi_0)$. In view of $\phi_1 \geq \phi_2$ on \mathbb{R} , it follows that $\mathcal{H}_\mu(\phi_1)(x) = \mathcal{H}_\mu(\phi_2)(x)$ for all $x \geq \xi_0$. It follows from the monotonicity of \mathcal{H}_μ that $\phi_1 \equiv \phi_2$ on \mathbb{R} . \square

Lemma 5.2 *Under assumptions (F1) and (F2), there exists $\varepsilon_0 \in (0, K)$ such that*

$$f((1+s)u, (1+s)v) - (1+s)f(u, v) < 0 \quad (42)$$

for all $s \in (0, \varepsilon_0)$ and $(u, v) \in \mathbb{R}^2$ satisfying $|K - u| < \varepsilon_0$ and $|K - v| < \varepsilon_0$.

Proof. For each $s \geq 0$, define

$$F(s, u, v) = f((1+s)u, (1+s)v) - (1+s)f(u, v).$$

Then $F(0, u, v) = 0$ and $F_s(0, u, v) = u\partial_1 f(u, v) + v\partial_2 f(u, v) - f(u, v)$ for all $(u, v) \in \mathbb{R}^2$. In view of assumption (F2), we have $F_s(0, K, K) = K\partial_1 f(K, K) + K\partial_2 f(K, K) < 0$. Therefore, there exists $\varepsilon_0 > 0$ such that $F(s, u, v) < 0$ for all $s \in (0, \varepsilon_0)$ and $(u, v) \in \mathbb{R}^2$ satisfying $|K - u| < \varepsilon_0$ and $|K - v| < \varepsilon_0$. \square

Lemma 5.3 *Assume that (F1), (F2), (H1) and (H2) hold. Let (c, ϕ_1) and (c, ϕ_2) be solutions of (6). Suppose there exists a constant $\varepsilon \in (0, \varepsilon_0]$ such that $(1 + \varepsilon)\phi_1(x - \kappa\varepsilon) \geq \phi_2(x)$ on \mathbb{R} , where*

$$\kappa = \sup \left\{ \frac{\phi_1(x)}{\phi_1'(x)} : \phi_1(x) \leq K - \varepsilon_0 \right\}.$$

Then $\phi_1 \geq \phi_2$ on \mathbb{R} .

Proof. Define $W(\varepsilon, x) = (1 + \varepsilon)\phi_1(x - \kappa\varepsilon) - \phi_2(x)$ and $\varepsilon^* = \inf\{\varepsilon \geq 0 : W(\varepsilon, x) \geq 0 \text{ for all } x \in \mathbb{R}\}$. By continuity of W , $W(\varepsilon^*, x) \geq 0$ for all $x \in \mathbb{R}$. We claim $\varepsilon^* = 0$. Suppose on the contrary that $\varepsilon \in (0, \varepsilon_0]$. Then, by the definition of κ ,

$$W_\varepsilon(\varepsilon, x) = \phi_1(x - \kappa\varepsilon) - \kappa(1 + \varepsilon)\phi_1'(x - \kappa\varepsilon) < 0$$

on $\{x \in \mathbb{R} : \phi_1(x - \kappa\varepsilon) \leq K - \varepsilon_0\}$. Noting that $W(\varepsilon^*, \infty) = \varepsilon^*K > 0$, we can find x_0 with $\phi_1(x_0 - \kappa\varepsilon^*) > K - \varepsilon_0$ such that

$$0 = W(\varepsilon^*, x_0) = W_x(\varepsilon^*, x_0) = W_{xx}(\varepsilon^*, x_0).$$

Thus, $(1 + \varepsilon^*)\phi_1(\xi_0) = \phi_2(x_0)$, $(1 + \varepsilon^*)\phi_1'(\xi_0) = \phi_2'(x_0)$, and $(1 + \varepsilon^*)\phi_1''(\xi_0) = \phi_2''(x_0)$, where $\xi_0 = x_0 - \kappa\varepsilon^*$. This, together with (42), implies

$$\begin{aligned} 0 &= -c\phi_2'(x_0) + p\phi_2''(x_0) + d(J * \phi_2 - \phi_2)(x_0) + f(\phi_2(x_0), (h * \phi_2)(x_0)) \\ &\leq -c(1 + \varepsilon^*)\phi_1'(\xi_0) + p(1 + \varepsilon^*)\phi_1''(\xi_0) + d(1 + \varepsilon^*)(J * \phi_1 - \phi_1)(\xi_0) \\ &\quad + f((1 + \varepsilon^*)\phi_1(\xi_0), \{h * [(1 + \varepsilon^*)\phi_1]\}(\xi_0)) \\ &= f((1 + \varepsilon^*)\phi_1(\xi_0), (1 + \varepsilon^*)(h * \phi_1)(\xi_0)) - (1 + \varepsilon^*)f(\phi_1(\xi_0), (h * \phi_1)(\xi_0)) < 0, \end{aligned}$$

a contradiction. Hence $\varepsilon^* = 0$ and so $\phi_1 \geq \phi_2$ on \mathbb{R} . \square

Theorem 5.2 *Assume that (F1), (F2), (H1) and (H2) hold. For each $c \geq c^*$, let (c, ϕ_1) and (c, ϕ_2) be two solutions to (6). Then there exists $\gamma \in \mathbb{R}$ such that $\phi_1(\cdot) = \phi_2(\cdot + \gamma)$, i.e., travelling waves are unique up to a translation.*

Proof. By translating ϕ_2 if necessary, we can assume that $0 < \phi_1(0) = \phi_2(0) < K$. By Theorem 4.2, we have

$$\lim_{x \rightarrow -\infty} \frac{\phi_2(x)}{\phi_1(x)} = e^{\lambda_1(c)\theta}$$

for some $\theta \in \mathbb{R}$. Without loss of generality, we assume that $e^{\lambda_1(c)\theta} \leq 1$, for otherwise we can exchange ϕ_1 and ϕ_2 . Then

$$W(\xi) = \lim_{x \rightarrow -\infty} \frac{\phi_2(x)}{\phi_1(x + \xi)} < 1$$

for all $\xi > 0$. Fix $\xi = 1$; then there exists $x_1 > 0$ such that

$$\phi_1(x + 1) > \phi_2(x) \text{ for all } x \in (-\infty, -x_1). \quad (43)$$

Since $\phi_1(\infty) = K$, there exists $x_2 \gg 1$ such that $\phi_1(x) \geq K/(1 + \varepsilon_0)$ for all $x > x_2$. It follows that

$$(1 + \varepsilon_0)\phi_1(x) \geq K \geq \phi_2(x) \text{ for all } x > x_2. \quad (44)$$

Let $\eta = \max\{\phi_2(x) : x \in [-x_1, x_2]\} \in (0, K)$. In view of $\phi_1(\infty) = K$, there exists $x_3 \gg 1$ such that $\phi_1(x) \geq \eta$ for all $x > x_3$. Thus, for $x \in [-x_1, x_2]$, we have $x + x_1 + x_3 \in [x_3, x_1 + x_2 + x_3]$ and hence

$$\phi_1(x + x_1 + x_3) \geq \eta \geq \phi_2(x). \quad (45)$$

Set $z = 1 + x_1 + x_3 + \kappa\varepsilon_0$. It follows from (43)–(45) that

$$(1 + \varepsilon_0)\phi_1(x + z - \kappa\varepsilon_0) \geq \phi_2(x) \text{ for all } x \in \mathbb{R}. \quad (46)$$

By monotonicity of φ_1 and Lemma 5.3, $\phi_1(x + z) \geq \phi_2(x)$ for all $x \in \mathbb{R}$. Set

$$\xi^* = \inf\{z > 0 : \phi_1(x + z) \geq \phi_2(x) \text{ for all } x \in \mathbb{R}\}.$$

We claim that $\xi^* = 0$. If not, then $\xi^* > 0$ and so we have $\phi_1(x + \xi^*) \geq \phi_2(x)$ for all $x \in \mathbb{R}$. It follows from $W(\xi^*/2) < 1$ that there exists $x_4 > 0$ such that

$$\phi_1(x + \xi^*/2) \geq \phi_2(x) \text{ for } x \leq -x_4.$$

Consider the function $(1 + \varepsilon)\phi_1(x + \xi^* - 2\kappa\varepsilon)$. Since $\phi_1(\infty) = K$ and $\phi_1'(\infty) = 0$, there exists $x_5 \gg 1$ such that

$$\frac{d}{d\varepsilon} \{(1 + \varepsilon)\phi_1(x + \xi^* - 2\kappa\varepsilon)\} = \phi_1(x + \xi^* - 2\kappa\varepsilon) - 2\kappa(1 + \varepsilon)\phi_1'(x + \xi^* - 2\kappa\varepsilon) > 0$$

for all $x \geq x_5$ and $\varepsilon \in [0, 1]$. That is, for all $x \geq x_5$ and $\varepsilon \in [0, 1]$,

$$(1 + \varepsilon)\phi_1(x + \xi^* - 2\kappa\varepsilon) \geq \phi_1(x + \xi^*) \geq \phi_2(x).$$

Now we consider the interval $[-x_4, x_5]$, since $\phi_1(\cdot + \xi^*) \geq \phi_2(\cdot)$, by Lemma 5.1, $\phi_1(\cdot + \xi^*) > \phi_2(\cdot)$ in $[-x_4, x_5]$. Thus, there exists $\varepsilon \in (0, \min\{\varepsilon_0, \xi^*/(4\kappa)\})$ such that $\phi_1(\cdot + \xi^* - 2\kappa\varepsilon) \geq \phi_2(\cdot)$ on $[-x_4, x_5]$. Therefore, combining the estimates on $(-\infty, -x_4]$, $[-x_4, x_5]$, and $[x_5, \infty)$, we conclude that $(1 + \varepsilon)\phi_1(\cdot + \xi^* - 2\kappa\varepsilon) \geq \phi_2(\cdot)$ on \mathbb{R} . It follows from Lemma 5.3 that $\phi_1(\cdot + \xi^* - \kappa\varepsilon) \geq \phi_2(\cdot)$ on \mathbb{R} . This contradicts the definition of ξ^* . Hence, $\xi^* = 0$, i.e., $\phi_1(\cdot) \geq \phi_2(\cdot)$. Since $\phi_1(0) = \phi_2(0)$, we have $\phi_1 \equiv \phi_2$ on \mathbb{R} . \square

Finally, we show that no travelling wave solution of speed $c < c^*$ exist. The usual approach is to combine the comparison method and the finite time-delay approximation to establish the existence of the spreading speed c^* for the solutions with initial functions having compact supports. In fact, c^* coincides with the minimal wave speed for monotone travelling waves of (1). Thus, the nonexistence of travelling waves with the wave speed $c < c^*$ is a straightforward consequence of the spreading speed. In what follows, however, we shall employ a different method to investigate the nonexistence of travelling waves with the wave speed $c < c^*$.

Theorem 5.3 *Assume that (F1), (F2), (H1) and (H2) hold. Let c^* be defined as in Lemma 2.1. Then for every $c \in (0, c^*)$, (1) has no travelling wave front with (c, φ) satisfying (6).*

Proof. In view of Theorem 5.1, every solution (c, φ) of (6) satisfies $\varphi'(\xi) > 0$ for all $\xi \in \mathbb{R}$. Take a sequence $\xi_n \rightarrow -\infty$ such that $\varphi(\xi_n) \rightarrow 0$ and set $v_n(\xi) = \varphi(\xi + \xi_n)/\varphi(\xi_n)$. As φ is bounded and satisfies (6), the Harnack's inequality implies that the sequence v_n is locally uniformly bounded. This function v_n satisfies

$$-cv_n'(\xi) + pv_n''(\xi) + d(J * v_n - v_n)(\xi) + \partial_1 f(0, 0)v_n(\xi) + \partial_2 f(0, 0)(h * v_n)(\xi) + R_n(\xi) = 0 \quad (47)$$

for $\xi \in \mathbb{R}$, where

$$R_n(\xi) = \frac{f(\varphi(\xi + \xi_n), (h * \varphi)(\xi + \xi_n)) - \partial_1 f(0, 0)\varphi(\xi + \xi_n) - \partial_2 f(0, 0)(h * \varphi)(\xi + \xi_n)}{\varphi(\xi_n)}.$$

The Harnack's inequality implies that the shifted functions $R_n(\xi)$ converge to zero locally uniformly in ξ . Thus one may assume, up to extraction of a subsequence, that the sequence v_n converges to a function v that satisfies:

$$-cv'(\xi) + pv''(\xi) + d(J * v - v)(\xi) + \partial_1 f(0, 0)v(\xi) + \partial_2 f(0, 0)(h * v)(\xi) = 0, \quad \xi \in \mathbb{R}, \quad (48)$$

Moreover, v is positive since it is nonnegative and $v(0) = 1$. Equation (48) admits such a solution if and only if $c \geq c^*$. Therefore, for every $c \in (0, c^*)$, (1) has no travelling wave front with (c, φ) satisfying (6). This completes the proof. \square

6 Acknowledgements

This work was partially supported by the Natural Science Foundation of People's Republic of China (Grant No 11671123 & 11271115), the UK's EPSRC (EP/K027743/1), the Leverhulme Trust (RPG-2013-261) and a Royal Society Wolfson Research Merit Award.

References

- [1] J. F. M. Al-Omari & S. A. Gourley, A nonlocal reaction-diffusion model for a single species with stage structure and distributed maturation delay. *Eur. J. Appl. Math.* 16 (2005) 37–51.
- [2] J. Al-Omari and S.A. Gourley, Monotone traveling fronts in age-structured reaction-diffusion model of a single species, *J. Math. Biol.* 45 (2002) 294–312.
- [3] P.W. Bates, P.C. Fife, X.F. Ren and X.F. Wang, Traveling waves in a convolution model for phase transitions, *Arch. Rational Mech. Anal.* 138 (1997) 105–136.
- [4] P.W. Bates and A. Chmaj, A discrete convolution model for phase transitions, *Arch. Rational Mech. Anal.* 150 (1999) 281–305.
- [5] H. Berestycki, G. Nadin, B. Perthame, L. Ryzhik, The non-local Fisher-KPP equation: travelling waves and steady states. *Nonlinearity* 22 (2009) 2813–2844.
- [6] N. Britton, Aggregation and the competitive exclusion principle, *J. Theor. Biol.* 136 (1989) 57–66.
- [7] N. Britton, Spatial structures and periodic travelling waves in an integro-differential reaction-diffusion population model, *SIAM J. Appl. Math.* 50 (1990), 1663–1688.
- [8] J.W. Cahn, J. Mallet-Paret and E.S. van Vleck, Traveling wave solutions for systems of ODEs on a two-dimensional spatial lattice, *SIAM J. Appl. Math.* 59 (1999) 455–493.
- [9] J. Carr and A. Chmaj, Uniqueness of travelling waves for nonlocal monostable equations, *Proc. Amer. Math. Soc.* 132 (2004) 2433–2439.
- [10] X. Chen, S. Fu and J. Wu, Uniqueness and asymptotics of traveling waves of monostable dynamics on lattices, *SIAM J. Math. Anal.* 38 (2006) 233–258.
- [11] X. Chen and J.S. Guo, Existence and asymptotic stability of travelling waves of discrete quasilinear monostable equations, *J. Differential Equations* 184 (2002) 549–569.
- [12] X. Chen and J.S. Guo, Uniqueness and existence of travelling waves for discrete quasilinear monostable dynamics, *Math. Ann.* 326 (2003) 123–146.
- [13] S.-N. Chow, J. Mallet-Paret and W. Shen, Traveling waves in lattice dynamical systems, *J. Differential Equations* 149 (1998) 248–291.

- [14] J. Coville, L. Dupaigne, On a nonlocal reaction diffusion equation arising in population dynamics, *Proceedings of the Royal Society of Edinburgh* 137 (2007) A 1–29.
- [15] J. Fang, J. Wei and X.-Q. Zhao, Spatial dynamics of a nonlocal and time-delayed reaction-diffusion system, *J. Differential Equations*, 245 (2008) 2749–2770.
- [16] T. Faria, W. Huang, J.H. Wu, Traveling waves for delayed reaction-diffusion equations with global response. *Proc. R. Soc. A.* 462(2006) 229–261.
- [17] R.A. Fisher, The advance of advantageous genes, *Ann. Eugenics* 7 (1937) 355–369.
- [18] S.A. Gourley, J.W.-H. So and J.H. Wu, Non-locality of reaction-diffusion equations induced by delay: biological modelling and nonlinear dynamics, in: D.V. Anosov, A. Skubachevskii (Guest Eds.), *Contemporary Mathematics. Thematic Surveys*, Kluwer Plenum, Dordrecht, New York, 2003, pp. 84–120.
- [19] S.A. Gourley and J. Wu, Delayed non-local diffusive systems in biological invasion and disease spread, in *Nonlinear Dynamics and Evolution Equations*, Fields Inst. Commun. 48, H. Brunner, X.-Q. Zhao, and X. Zou, eds., AMS, Providence, RI, 2006, pp. 137–200.
- [20] S. Guo and J. Zimmer, Stability of travelling wavefronts in discrete reaction-diffusion equations with nonlocal delay effects, *Nonlinearity* 28 (2015) 463–492.
- [21] W.S.C. Gurney, S.P. Blythe and R. M. Nisbet, Nicholson’s blowflies revisited, *Nature* 287 (1980) 17–21.
- [22] C.-H. Hsu and S.-S. Lin, Existence and multiplicity of traveling waves in a lattice dynamical system, *J. Differential Equations* 164 (2000) 431–450.
- [23] W. Hudson and B. Zinner, Existence of traveling waves for a generalized discrete Fisher’s equation, *Comm. Appl. Nonlinear Anal.* 1 (1994) 23–46.
- [24] J.P. Keener, Propagation and its failure in coupled systems of discrete excitable cells, *SIAM J. Appl. Math.* 22 (1987) 556–572.
- [25] A.N. Kolmogorov, I.G. Petrovsky and N.S. Piskunov, Étude de l’équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique, *Bull. Univ. Moskov. Ser. Internat., Sect. A* 1 (1937) 1–25.
- [26] W.T. Li, S.G. Ruan, Z.C. Wang, On the diffusive Nicholson’s blowflies equation with nonlocal delay, *J. Nonlinear Sci.* 17 (2007) 505–525.
- [27] C.-K. Lin and M. Mei, On travelling wavefronts of the Nicholson’s blowflies equations with diffusion, *Proc. Roy. Soc. Edinburgh Sect. A*, 140 (2010) 135–152.
- [28] G.J. Lin, Traveling wave solutions in the Nicholsons blowflies equation with spatio-temporal delay, *Appl. Math. Comput.* 209 (2009) 314–326.
- [29] S. Ma, X. Liao and J. Wu, Traveling wave solutions for planar lattice differential systems with applications to neural networks, *J. Differential Equations* 182 (2002) 269–297.
- [30] S. Ma and X. Zou, Existence, uniqueness and stability of travelling waves in a discrete reaction-diffusion monostable equation with delay, *J. Differential Equations* 217 (2005) 54–87.
- [31] J. Mallet-Paret, The global structure of traveling waves in spatially discrete dynamical systems, *J. Dynam. Differential Equations* 11 (1999) 49–127.
- [32] M. Mei, C.-K. Lin, C.-T. Lin and J. W.-H. So, Traveling wavefronts for time-delayed reaction-diffusion equation: (I) local nonlinearity, *J. Differential Equations*, 247 (2009) 495–510.

- [33] C. Ou and J. Wu, Persistence of wavefronts in delayed non-local reaction-diffusion equations, *J. Differential Equations*, 235 (2007) 219–261.
- [34] S. Ruan and D. Xiao, Stability of steady states and existence of traveling waves in a vector disease model. *Proc. Roy. Soc. Edinburgh* 134 (2004) 991–1011 .
- [35] K.W. Schaaf, Asymptotic behavior and traveling wave solutions for parabolic functional differential equations. *Trans. Amer. Math. Soc.*, 302 (1987), 587–615.
- [36] H.L. Smith and H. Thieme, Strongly order preserving semiflows generated by functional differential equations, *J. Differential Equations* 93 (1991) 332–363.
- [37] J. W.-H. So, J. Wu and X. Zou, A reaction-diffusion model for a single species with age structure: (I) Traveling wavefronts on unbounded domains, *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 457 (2001) 1841–1853.
- [38] H. Thieme and X.-Q. Zhao, Asymptotic speeds of spread and traveling waves for integral equation and delayed reaction-diffusion models, *J. Differential Equations*, 195 (2003) 430–370.
- [39] Z. Wang, W. Li, S. Ruan, Traveling fronts in monostable equations with nonlocal delayed effects, *J Dyn Diff Equat* 20 (2008) 573–607.
- [40] P.X. Weng J.H. WU, Wavefronts for a nonlocal reaction-diffusion population model with general distributive maturity. *IMA J. Appl. Math.* 73 (2008) 477–495.
- [41] D.V. Widder, *The Laplace Transform*, Princeton Univ. Press, Princeton, NJ., 1941.
- [42] J. Wu and X. Zou, Asymptotic and periodic boundary value problems of mixed FDEs and wave solutions of lattice differential equations, *J. Differential Equations* 135 (1997) 315–357.
- [43] J.H. Wu, X.F. Zou, Traveling wave fronts of reaction-diffusion systems with delay. *J. Dyn. Differ. Equ.*, 13(2001), 651–687
- [44] X.Q. Zhao, D.M. Xiao, The asymptotic speed of spread and traveling waves for a vector disease model. *J. Dyn. Differ. Equ.*, 18(2006), 1001–1019.
- [45] B. Zinner, Stability of traveling wavefronts for the discrete Nagumo equation, *SIAM J. Math. Anal.* 22 (1991) 1016–1020.
- [46] B. Zinner, Existence of traveling wavefront solution for the discrete Nagumo equation, *J. Differential Equations* 96 (1992) 1–27.
- [47] B. Zinner, G. Harris and W. Hudson, Travelling wavefronts for the discrete Fisher’s equation, *J. Differential Equations* 105 (1993) 46–62.