



Citation for published version:

Silva, DAS, Chaput, J-P, Katzmarzyk, PT, Fogelholm, M, Hu, G, Maher, C, Olds, T, Onywera, V, Sarmiento, OL, Standage, M, Tudor-Locke, C & Tremblay, MS 2018, 'Physical Education Classes, Physical Activity, and Sedentary Behavior in Children', *Medicine & Science in Sports & Exercise*, vol. 50, no. 5, pp. 995-1004. <https://doi.org/10.1249/MSS.0000000000001524>

DOI:

[10.1249/MSS.0000000000001524](https://doi.org/10.1249/MSS.0000000000001524)

Publication date:

2018

Document Version

Peer reviewed version

[Link to publication](#)

This is not the final published version of the article. The version of record is available to subscribers via <https://doi.org/10.1249/MSS.0000000000001524>.

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

26 **ABSTRACT**

27 **Purpose:** To examine the associations between participation frequency in Physical Education
28 (PE) classes and objective measures of physical activity (PA) and sedentary behavior (SB) in
29 children from 12 countries at different levels of development.

30 **Methods:** This multinational, cross-sectional study included 5,874 children aged 9-11 years
31 from sites in Australia, Brazil, Canada, China, Colombia, Finland, India, Kenya, Portugal,
32 South Africa, the United Kingdom and the United States. PA and SB were monitored over 7
33 consecutive days using a waist-worn accelerometer. PA and SB data were presented for
34 weekdays (times in- and out-of-school) and weekend days. Participation frequency in PE
35 classes was determined by questionnaire. Multilevel modeling analyses stratified by sex were
36 used.

37 **Results:** Overall, 24.8% of children self-reported participation in PE classes ≥ 3 times/week
38 (25.3% in high-income countries [HIC], and 24.3% in low- and middle-income countries
39 [LMIC]). After adjusting for age, sex, parental education and body mass index z-score,
40 results showed that children from LMIC who took PE classes 1-2 times/week were more
41 likely to present better indicators of PA and shorter time in SB in- and out-of-school. In HIC,
42 boys that participated in PE classes were more likely to meet the moderate-to-vigorous PA
43 (MVPA) recommendations and to present better indicators of PA (in school) and shorter time
44 in SB in- and out-of-school. For girls in HIC, attending PE classes increased the likelihood of
45 spending more time in MVPA, especially if they attended ≥ 3 times/week.

46 **Conclusion:** Attending PE classes is associated with a higher level of PA and lower level of
47 SB in- and out-of-school during weekdays in children from countries at various levels of
48 development.

49 **Keywords:** CROSS-SECTIONAL STUDY, EPIDEMIOLOGY, EXERCISE, GLOBAL
50 HEALTH, SCHOOL HEALTH.

51 INTRODUCTION

52 Physical activity (PA) and sedentary behavior (SB) are distinct behaviors that affect the
53 health of children (1-3). The two movement behaviors have different determinants and
54 because levels of PA and SB around of the world are concerning, these behaviors represent
55 major challenges for public health authorities (4). According to a study coordinated by the
56 Active Healthy Kids Global Alliance (4), which compiled data from 38 countries, more than
57 60% of children do not meet the recommended 60 minutes of daily moderate-to-vigorous PA
58 (MVPA). Likewise, more than 60% of children spend more than two hours a day on
59 recreational screen time (4), exceeding public health recommendations of some countries (5).

60 The school environment is a context where young people spend much of the day
61 assimilating knowledge about different topics, including health. In this way, the school
62 environment is conducive to informing children about the benefits of regular PA and the
63 importance of preventing excessive SB throughout the day (4). In addition to guiding and
64 educating young people about these lifestyle behaviors, schools play a key role in providing
65 structured and unstructured PA (4). One opportunity for this to happen is through Physical
66 Education (PE) classes (6,7). The importance of PE classes for the PA of children has been
67 highlighted in guidelines from various countries where it is recommended that young people
68 engage in MVPA for at least 50% of PE class time (6,7).

69 The discussion of PE classes in the guidelines comes from a series of studies that
70 showed that participation frequency in PE was associated with higher levels of PA and lower
71 levels of SB on days with PE classes and on days without PE classes (8,9). However,
72 equivocal findings are also reported in the literature (18). Specifically, the association
73 between participation frequency in PE classes and PA and SB may differ depending on age,
74 sex, or weight status (10). Two explanations have been proposed to explain these equivocal
75 observations. First, Dishman et al. (11) suggested that the practice of PA, at any intensity,

76 causes changes at the level of the brain that stimulate vigor and more movement throughout
77 the day. Thus, it can be inferred that children who engage in PE classes will tend to be more
78 physically active on different days (week days and weekend days) or vice versa. Second, the
79 "activitystat" hypothesis suggests that there is an energy expenditure threshold for children
80 and, once reached, the rest of the daily time may be compensated with little or no PA on the
81 same day or on other days of the week (12,13). From this theory, it can be inferred that
82 participation frequency in PE classes may help to reach this threshold of energy expenditure
83 and, on PE days, the PA time for the rest of the day will be decreased. Both explanations (11-
84 13) do not give details about the differences that may exist between boys and girls that have
85 been identified as one of the factors that can modify the association between participation
86 frequency in PE classes and PA and SB time (10).

87 Previous studies that have investigated the association between participation frequency
88 in PE classes and PA and SB levels throughout the day have typically been developed with
89 samples of students from a single city or country, and most of these studies have been
90 conducted in high-income countries (HIC) (9,14,15). The few studies in low- and middle-
91 income countries (LMIC) have used subjective measures of PA and SB, which likely impacts
92 the accuracy of estimates reported (16,17). In order to overcome limitations of previous
93 studies, the present analysis aims to examine the association between participation frequency
94 in PE classes and objective measures of PA and SB throughout the day, both on weekdays
95 (in- and out-of-school) and weekend days in children from 12 countries.

96 **METHODS**

97 **Study Design and Setting**

98 The International Study of Childhood Obesity, Lifestyle and the Environment
99 (ISCOLE) is a cross-sectional, multinational study designed to examine the relationships
100 between lifestyle behaviors and obesity in children in 12 study sites located in Australia,

101 Brazil, Canada, China, Colombia, Finland, India, Kenya, Portugal, South Africa, the United
102 Kingdom, and the United States. These countries represent a wide range of economic
103 development (low to high income), Human Development Index (0.509 in Kenya to 0.929 in
104 Australia), and income inequality (Gini index of 26.9 in Finland to 63.1 in South Africa) (18).
105 The rationale, design, and methods of ISCOLE have previously been published in detail
106 elsewhere (18). The primary sampling frame was schools, which were typically stratified by
107 an indicator of socioeconomic status (SES) to maximize variability within sites. Each study
108 site identified one or more school districts (within reasonably close proximity to the local
109 study center), to provide a sufficient population to draw a sample of 500 students. A
110 standardized protocol was used to collect data across all sites, and all study personnel
111 underwent rigorous training and certification to ensure data quality. The Pennington
112 Biomedical Research Center Institutional Review Board as well as Institutional/Ethical
113 Review Boards at each site approved the study. Written informed consent was obtained from
114 parents/legal guardians, and child assent was also obtained as required by local ethics review
115 boards. Data were collected during the school year at each study site and occurred between
116 September 2011 and December 2013.

117 **Participants**

118 ISCOLE targeted grade levels likely to ensure minimal variability around a mean age of
119 10 years. All children within the targeted grade level in a sampled school were eligible to
120 participate; hence, the sample included 9-11 year-old children. Based on a priori sample size
121 and power calculations (18), each site aimed to recruit a sex-balanced sample of at least 500
122 children. The response rate in ISCOLE was 60.0% (children with consent to participate
123 divided by consents distributed). Of the 7,372 children who participated in ISCOLE, a total
124 of 5,874 remained in the final analytical sample. We excluded participants without valid
125 accelerometry (n=1,214), information on participation frequency in PE classes (n=24),

126 reported level of parental education (n=255), or body mass index (BMI) z-score (n=5).
127 Children who were excluded for missing data did not significantly differ in their descriptive
128 characteristics from those who were included in the present analysis (see Table,
129 Supplemental Digital Content 1).

130 **Measurements**

131 **Physical Activity and Sedentary Time**

132 Vigorous PA (VPA), moderate PA (MPA), MVPA, light PA (LPA), total PA (TPA =
133 MVPA + LPA), and total sedentary time were objectively-assessed using 24-h, waist-worn
134 accelerometry (19). An Actigraph GT3X+ accelerometer (ActiGraph LLC, Pensacola, FL,
135 USA) was worn at the waist on an elasticized belt at the right mid-axillary line. Participants
136 were encouraged to wear the accelerometer 24 h per day (removing only for water-based
137 activities) for at least 7 days, including 2 weekend days. Overall, mean 24-h wear time and
138 wake wear time were 22.6 h/day and 14.9 h/day in ISCOLE, respectively. The minimum
139 amount of non-sleep data that was considered acceptable for inclusion was at least 4 days
140 with at least 10 h of wake wear time per day, including at least 1 weekend day (19). Data
141 were collected at a sampling rate of 80 Hz, downloaded in 1-s epochs with the low-frequency
142 extension filter using the ActiLife software version 5.6 or higher (ActiGraph LLC, Pensacola,
143 FL, USA), and reintegrated to 15-s epochs for analysis. After exclusion of sleep period time
144 (20,21) and awake non-wear time (any sequence of ≥ 20 consecutive minutes of zero activity
145 counts), VPA was defined as activity ≥ 1003 counts/15 s, MPA was defined as all activity
146 ≥ 574 counts/15 s and < 1003 counts/15 s, MVPA was defined as all activity ≥ 574 counts/15 s,
147 LPA was defined as all activity > 25 counts/15 s and < 574 counts/15 s, and total sedentary
148 time as all movement ≤ 25 counts/15 s, consistent with the widely used Evenson cutoffs (22).
149 MVPA and SB were analyzed considering the average of weekdays and weekend days. The
150 weekdays were divided as “in-school” and “out-of-school”, with out-of-school comprising

151 the time before and after school. The before school time period was considered from sleep
152 onset wake (established using a validated algorithm (20)) until school start time, “in-school”
153 was defined as the time between school start and end times (determined individually for each
154 participating school), and the after school period was from school end time through the
155 child’s bedtime (also determined by our validated algorithm (20) determined from
156 accelerometry). Full details on how wake time was determined are reported in previous
157 studies by the ISCOLE group (19-21).

158 Children were classified as meeting the MVPA recommendation of an average of ≥ 60
159 min/day, or not (5,23). In addition, each PA intensity (min/day) and the overall SB time
160 (min/day) were categorized according to the distribution tertile for boys and girls. We chose
161 this classification because there are no specific recommendations for each PA intensity and
162 the overall SB time. Sensitivity analyses using other cut-points for PA and SB revealed
163 similar patterns of associations; therefore, only tertiles are presented in the present paper.

164 **Participation frequency in PE classes**

165 Participation frequency in PE classes was assessed using a self-report questionnaire
166 item from the U.S. Youth Risk Behavior Surveillance System (24). The students were asked:
167 “In the last week you were in school, on how many days did you go to physical education
168 (PE) classes?” Response options ranged from zero days to five days. In the present study,
169 responses were classified into three categories, "0 days/week", "1-2 days/week", and " ≥ 3
170 days/week" to maximize power and because the results of the associations between
171 participation frequency in PE classes and PA/SB for 3, 4, and 5 days/week were similar.

172 **Covariates**

173 Age, sex, level of parental education, and BMI z-scores were included as covariates.
174 These variables were chosen because of their association with the dependent and independent
175 variables in previous studies (9,15,25,26). Age was computed from birthdates and

176 measurement dates, and sex was self-reported on a questionnaire. The level of parental
177 education was reported by the parents themselves in a questionnaire sent to them (18).
178 Highest level of parental education was used as a proxy measure for family SES and coded
179 into three categories based on the highest level of education attained by either parent: “did
180 not complete high school”, “completed high school or some college”, or “completed
181 bachelor’s or postgraduate degree”. Body mass was measured with a Tanita SC-240 scale
182 (Arlington Heights, IL, USA), after all outer clothing, heavy pocket items and shoes were
183 removed. Body height was measured without shoes using a Seca 213 portable stadiometer
184 (Hamburg, Germany), with the participant’s head in the Frankfort horizontal plane and after a
185 deep inspiration. Each measurement was repeated, and the average was used for analysis (a
186 third measurement was obtained if the first two measurements were greater than 0.5 kg or 0.5
187 cm apart for body mass and body height, respectively, and the average of the two closest
188 measurements was used for analysis). BMI (kg/m^2) was calculated, and BMI z-scores were
189 computed using age- and sex-specific reference data from the World Health Organization
190 (27).

191 **Statistical Analysis**

192 The analyses were performed for the whole sample, stratified by sex and country-level
193 World Bank classification of economic development. Descriptive statistics (absolute
194 frequency, relative frequency, mean values and standard deviations) on the characteristics of
195 the sample were presented for each site and for the set of sites from countries classified as
196 HICs (Australia, Canada, Finland, Portugal, United Kingdom, United States) and LMICs
197 (Brazil, China, Colombia, India, Kenya, South Africa), in accordance with the
198 recommendations of the World Bank (28). A multi-level logistic mixed regression model was
199 used where the dependent variable was meeting the MVPA recommendations (5,23). Multi-
200 level polytomous logistic mixed regression models were used (29,30), where the dependent

201 variables were the various PA indices, with a model for each (VPA, MPA, MVPA, LPA,
202 TPA), and SB (sex-specific reference categories = 1st (lowest) tertile for PA indices and 3rd
203 (highest) tertile for SB). Odds ratios and 95% confidence intervals were estimated. Study
204 sites were considered to have fixed effects and schools nested within study sites were viewed
205 as having random effects. Age, sex (with models containing the whole sample), parental
206 education, and BMI z-scores were included as covariates. Statistical analyses were performed
207 using Stata 13.0 software (STATA Corp. College Station, Texas, USA). The level of
208 statistical significance was set at $p < 0.05$.

209 **RESULTS**

210 This study analyzed data from 5,874 children (45.6% male) from 12 countries, with an
211 average age of 10.4 ± 0.6 years old (Table 1). Average MVPA was 60.3 ± 24.8 min/day,
212 average VPA was 17.9 ± 11.1 min/day, average MPA was 42.4 ± 5.6 min/day, and average
213 LPA was 314.9 ± 52.7 min/day. Total sedentary time was 513.4 ± 69.2 min/day. Table 1 also
214 presents this information for each country and according to the country-level World Bank
215 classification of economic development (HIC and LMIC). Supplemental Digital Content 2
216 and 3 present these results for boys (see Table, Supplemental Digital Content 2) and girls (see
217 Table, Supplemental Digital Content 3).

218 In the full sample ($n = 5,874$) (see Figure 1), 24.8% of children self-reported
219 participation frequency in PE classes three or more times per week (25.3% in HIC and 24.3%
220 LMIC, $p > 0.05$). The country that had most students reporting attending PE classes for three
221 days or more in the week was Canada (64.8%). Portugal was the country with the greatest
222 proportion of children reporting attending PE classes once or twice a week (99.0%). South
223 Africa was the country with the greatest proportion of students not attending PE classes
224 (32.1%). These results were similar when analyzing boys (see Figure, Supplemental Digital
225 Content 4) and girls (see Figure, Supplemental Digital Content 5) separately.

226 In general (Table 2), 44.4% of the sample met the recommendations for MVPA, with a
227 greater adherence in Finland (63.8%) and a lower adherence in China (15.9%). The highest
228 proportion of children who spent more time in VPA was found in Finland (53.0%) and the
229 lowest in India (12.3%). The highest proportion of children who spent more time in MPA
230 was found in Colombia (55.7%) and the lowest in China (8.1%). The highest proportion of
231 children who spent more time in LPA was found in Brazil (49.0%) and the lowest in Finland
232 (14.0%) and the UK (14.0%). The highest proportion of children who spent less time on SB
233 was found in Australia (52.5%) and the lowest in China (9.5%). Full details for all PA indices
234 and SB in- and out-of-school are provided in Table 2 (full sample) and in Supplemental
235 Digital Content (boys: see Table, Supplemental Digital Content 6; girls, see Table,
236 Supplemental Digital Content 7).

237 As shown in Table 3, children from LMIC who participated in PE classes at least 1-2
238 times/week were more likely to meet the MVPA recommendations (male – OR: 1.80; 95%
239 CI: 1.17; 2.77; female – OR: 2.17; 95% CI: 1.44; 3.27), to spend more time at different PA
240 intensities, and to have shorter SB time than those who did not attend PE classes. In HIC,
241 there were differences between boys and girls, where boys participating in PE classes were
242 more likely to meet the recommendations for time spent in MVPA and VPA, and shorter SB
243 time. For girls from HIC, attending PE classes increased the likelihood of spending more time
244 in MVPA (OR: 2.42; 95% CI: 1.22; 4.80) and MPA (OR: 2.44; 95% CI: 1.25; 4.75),
245 especially if they attended three or more classes/week.

246 As reported in Table 4, children from LMIC who participated in PE classes 1-2
247 times/week were more likely to spend more time in MVPA out of school on weekdays (male
248 - OR: 2.39; 95% CI: 1.43; 3.99; female: 2.79, 95% CI: 1.71, 4.58). Children from LMIC
249 (male - OR: 3.23; 95% CI: 1.82; 5.71; female - OR: 7.27; 95% CI: 4.39; 12.05) and HIC
250 (male - OR: 5.87; 95% CI: 2.78; 12.35; female - OR: 3.47; 95% CI: 1.77; 6.82) who

251 participated in PE classes three or more times/week were more likely to spend more time on
252 MVPA at school. Girls from LMIC that had at least 1-2 PE classes/week were more likely to
253 spend more time in MVPA on weekends (OR: 1.86; 95% CI: 1.17; 2.95).

254 As shown in Table 4, children of both sexes from LMIC (male - OR: 2.23; 95% CI:
255 1.23; 4.05; female - OR: 2.76; 95% CI: 1.56; 4.88) and the boys from HIC (OR: 2.45; 95% CI
256 : 1.09; 5.51) who participated in 3 or more PE classes/week were more likely to spend less
257 time in SB out of school on weekdays. For in-school SB time, the children who presented the
258 highest odds of spending shorter time in SB were boys (OR: 3.33, 95% CI: 1.97; 6.63) and
259 girls (OR: 4.87; 95% CI: 2.94; 8.06) from LMIC, and boys (OR: 2.40; 95% CI: 1.01; 5.81)
260 from HIC who took 3 or more PE classes/week. For the time in SB on weekends, no
261 significant associations were found.

262 **DISCUSSION**

263 The main finding of the present study was that participation frequency in PE classes
264 was related to greater odds of children demonstrating desirable indicators of PA and SB daily
265 behaviors. In addition, participation frequency in PE classes was associated with longer time
266 spent in MVPA and shorter time in SB in- and out-of-school on weekdays, especially in
267 children from LMIC. These results were more evident in LMIC, where at least one PE
268 class/week represented improvements in PA and SB indicators. For HIC, the results were
269 more evident in boys and with participation frequency in ≥ 3 PE classes/week.

270 The health benefits of participation frequency in PE classes for children have been
271 reported in previous studies (31-34). A shared explanation provided by these authors for the
272 improvement in health indicators of children who participated in PE classes was that PE class
273 attendance was associated with greater involvement in PA in general. The present study
274 supports these findings by showing better indicators of PA and SB in- and out-of-school for
275 children who report attending PE classes.

276 Similar to the present study, other studies have also found an association between
277 participation frequency in PE classes and higher PA levels and lower SB levels in children
278 (8,9,25). A cross-sectional study with children from Estonia measured PA and SB by
279 accelerometry during PE classes and during the remainder of the day. The authors found that
280 only one third of PE classes was spent on MVPA and the rest was spent on LPA and SB.
281 However, Estonian children who participated in PE classes had shorter SB time and greater
282 MVPA involvement throughout the day than peers who did not take PE classes (9). A study
283 conducted with school-aged children from the US measured PA using SenseWear armband
284 monitor (BodyMedia®, Pittsburg, US) during weekdays and weekends. The authors found
285 that the students were less sedentary and more physically active for the rest of the day
286 following the classes on PE days (8). In Brazilian children, it was reported that at least one
287 PE class/week was associated with less time in SB throughout the day (25). Although the
288 present study did not discriminate the measurement of PA and SB during the PE class, we
289 can hypothesize that involvement in PE classes is a stimulus to keep young people more
290 physically active and less sedentary throughout the day. This hypothesis was supported
291 because the children in this study who participated in PE classes had more time in MVPA and
292 less time in SB in- and out-of-school on weekdays. The rationale for this hypothesis is that
293 the practice of PA, regardless of intensity, causes changes in the cerebral cortex and
294 neurophysiological stimulation and can reduce the sensation of fatigue throughout the day,
295 while improving mood and the willingness to keep moving (11). Another possible
296 explanation would be that PE classes make children familiar and more confident with
297 physical activity, which increases the possibility of engaging in other activities out of school
298 (35).

299 Some studies examining the moderators of the relationship between participation
300 frequency in PE classes and PA/SB throughout the day have found no associations

301 (10,14,15). According to these authors, sex, weight status and lesson content are key
302 moderators of this association. For example, no association was found between participation
303 frequency in PE classes and PA/SB levels throughout the day in overweight and obese girls.
304 On the other hand, participation frequency in PE classes contributed significantly to higher
305 LPA in the normal-weight girls and boys during the school day (15). Nettlefold et al. (10)
306 also reported no association between participation frequency in PE classes and PA levels
307 throughout the day for children in Vancouver, Canada. One justification found in the
308 literature for the null findings observed between involvement in PE classes and PA/SB levels
309 throughout the day is the “activitystat” hypothesis, which suggests that children have an
310 energy expenditure set point and, therefore, compensate for increased PA in one part of the
311 day by decreasing PA later (12,13). The present study did not confirm this hypothesis
312 because the participation in PE classes was associated with greater involvement in MVPA
313 and less time spent in SB in- and out-of-school on weekdays.

314 The present study investigated whether the country-level World Bank classification of
315 economic development modified the associations found between participation frequency in
316 PE classes and PA/SB. We observed that the relationship between participation frequency in
317 PE classes and PA/SB levels of children was related to the development status/income level
318 of the country. The literature has already reported that levels of PA and SB vary according to
319 the development status/income level of the country and family SES (15,17,28). A recent
320 systematic review and meta-analysis of studies with school-aged children revealed that in
321 HIC, family SES was inversely associated with SB, whereas in low/middle-income countries,
322 there was a positive association between family SES and SB (36). A systematic review
323 revealed that school-aged children in Sub-Saharan Africa (low income region) with higher
324 family SES were less physically active and spent more time on SB than those with lower
325 family SES (37). The present study added the component of PE classes to this discussion

326 between different income levels with the understanding that PE classes are opportune
327 moments in the school day for young people to practice PA. However, we did not stratify
328 analyses by income level of the families in each group of countries to protect against type 2
329 error (i.e., we sought to retain power in the statistical analyses). Thus, it is suggested that
330 future studies with larger samples make this stratification.

331 This study found that in LMIC, participation frequency in at least one PE class was
332 associated with higher levels of PA and lower SB levels. In HIC, this result was most evident
333 in male children only or those attending three or more PE classes/week. A possible
334 explanation for these differences is that countries implement different strategies and policies
335 to promote PA and healthy habits for children (4). The Active Healthy Kids Global Alliance
336 organizes and prepares Report Cards on the PA of children and youth from different countries
337 around the world. This program revealed that government actions and policies to encourage
338 and promote PA are more evident in HIC, and some LMIC do not have government policies
339 to encourage PA, including in the school environment (4). Based on this evidence and the
340 results of the present study, it is deduced that participation frequency in PE classes in LMIC
341 represents an important avenue for young people to engage in PA throughout the day in- and
342 out-of-school. In this study, young people who did at least one PE class/week were more
343 likely to be classified as having higher levels of PA of different intensities and with lower SB
344 than those who did not take PE classes.

345 For HIC, the literature reports greater incentives and government actions to promote
346 PA, both in- and out-of-school (4,38,39). For activities at school, the justification given is
347 that schools in HIC provide built environments more suitable for the practice of PA when
348 compared to LMIC (4,38,39). For out-of-school activities, the justification is not centered
349 only on the built environment but also on the opportunities for young people to engage in
350 targeted PA practice in periods outside of school (4,39). It is possible that these other

351 structured and unstructured activities (compared to PE classes), inside and outside of the
352 school, represent the largest portion of the students' time spent on PA (38).

353 This study found that for some PA and SB indicators, at least 1-2 PE classes/week
354 represented higher odds to spend more time in PA and shorter time in SB, and that to do three
355 or more PE classes/week did not represent greater odds. Thus, there was no dose-response
356 relationship between participation frequency in PE classes and PA and SB throughout the
357 day. We can speculate that the quality of PE classes is as important as the frequency of these
358 classes. Fröberg et al. (14) reported that the contents that are discussed in PE classes can be
359 facilitators or barriers to PA in children. Perhaps even a small dose (e.g., 1 or 2 PE classes per
360 week) can achieve desirable outcomes. In this regard, it is recommended that the contents of
361 the PE classes be promoters of an active lifestyle of the children.

362 There are several limitations in the present study. First, the cross-sectional design
363 precludes us from making cause-and-effect inferences. Second, we cannot exclude the
364 possibility that unmeasured confounding variables may explain some of the observed
365 relationships. Third, we did not evaluate the PA level and the SB level during PE classes and,
366 therefore, we do not know the behavior pattern of the students during PE classes. Fourth, we
367 objectively assessed PA and SB using accelerometry; however, a limitation of this approach
368 is the inability to quantify some physical activities such as cycling and swimming. Five, the
369 sample of 12 countries involved in the survey was not nationally representative and not
370 represent the behavior pattern of any particular country, though the samples were generally
371 quite comparable to other nationally representative surveys (40). Finally, the participation
372 frequency in PE classes was self-reported and information on the PE curriculum was not
373 collected. The strength of this study was in the recruitment of a large multinational sample of
374 children from LMIC across several regions of the world, the highly standardized

375 measurement protocol, the use of objective measurements whenever possible, and the
376 rigorous quality control program.

377 In conclusion, participation frequency in PE classes was associated with healthy
378 indicators of PA and SB throughout the day in children from around the world. Participation
379 frequency in PE classes also resulted in longer time spent in MVPA and shorter time in SB
380 in- and out-of-school on weekdays. These results were more evident in LMIC, where at least
381 one PE class/week was associated with the desirable indicators of PA and SB. For HIC, these
382 results were more evident in boys and those attending three or more PE classes/week.
383 Collectively, PE classes seem to represent an opportunity to positively influence PA and SB
384 levels of children, especially those from LMIC.

385 **Funding**

386 The International Study of Childhood Obesity, Lifestyle, and the Environment (ISCOLE) was
387 funded by The Coca-Cola Company. GH was supported by a grant from the National Institute
388 of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health under
389 Award Number R01DK100790. The funder had no role in the design and conduct of the
390 study; collection, management, analysis and interpretation of the data; and preparation,
391 review, or approval of the manuscript.

392 **Acknowledgments**

393 We wish to thank the ISCOLE External Advisory Board, the ISCOLE participants and their
394 families who made this study possible, and the ISCOLE Research Group that was quoted in
395 Katzmarzyk et al. (11).

396 **Conflict of Interest**

397 The authors declare no conflicts of interest.

398 The results of the present study do not constitute endorsement by the American College of
399 Sports Medicine.

400 The authors declare that the results of the study are presented clearly, honestly, and without
401 fabrication, falsification, or inappropriate data manipulation.

402

403 REFERENCES

- 404 1. Carson V, Hunter S, Kuzik N, et al. Systematic review of sedentary behaviour and health
405 indicators in school-aged children and youth: an update. *Appl Physiol Nutr Metab.*
406 2016;41(6 Suppl 3):S240-65.
- 407 2. Poitras VJ, Gray CE, Borghese MM, et al. Systematic review of the relationships between
408 objectively measured physical activity and health indicators in school-aged children and
409 youth. *Appl Physiol Nutr Metab.* 2016;41(6 Suppl 3):S197-239.
- 410 3. Tremblay MS, Aubert S, Barnes JD, et al. Sedentary Behavior Research Network (SBRN)
411 - Terminology Consensus Project process and outcome. *Int J Behav Nutr Phys Act.*
412 2017;14(1):75.
- 413 4. Tremblay MS, Barnes JD, González SA, et al. Global Matrix 2.0: Report Card Grades on
414 the Physical Activity of Children and Youth Comparing 38 Countries. *J Phys Act Health.*
415 2016;13(11 Suppl 2):S343-S366.
- 416 5. Tremblay MS, Carson V, Chaput JP, et al. Canadian 24-hour Movement Guidelines for
417 Children and Youth: An Integration of Physical Activity, Sedentary Behaviour, and
418 Sleep. *Appl Physiol Nutr Metab.* 2016;41(6 Suppl 3):S311-S27.
- 419 6. Association for Physical Education. Health position paper. *Phys Educ Matters.* 2008;
420 3(1):8-12.
- 421 7. U.S. Department of Health and Human Services, Centers for Disease Control and
422 Prevention, National Center for Chronic Disease Prevention and Health Promotion,
423 Division of Adolescent and School Health. *Strategies to improve the quality of Physical*
424 *Education.* [Internet]. 2010; [cited 2016 Dec 30]. Available from:
425 https://www.cdc.gov/healthyschools/pecat/quality_pe.pdf.
- 426 8. Chen S, Kim Y, Gao Z. The contributing role of physical education in youth's daily
427 physical activity and sedentary behavior. *BMC Public Health.* 2014;14:110.

- 428 9. Mooses K, Pihu M, Riso EM, Hannus A, Kaasik P, Kull M. Physical Education Increases
429 Daily Moderate to Vigorous Physical Activity and Reduces Sedentary Time. *J Sch*
430 *Health*. 2017;87(8):602-7.
- 431 10. Nettlefold L, McKay HA, Warburton DE, McGuire KA, Bredin SS, Naylor PJ. The
432 challenge of low physical activity during the school day: at recess, lunch and in physical
433 education. *Br J Sports Med*. 2011;45(10):813-9.
- 434 11. Dishman RK, Thom NJ, Puetz TW, O'Connor PJ, Clementz BA. Effects of cycling
435 exercise on vigor, fatigue, and electroencephalographic activity among young adults who
436 report persistent fatigue. *Psychophysiology*. 2010;47(6):1066-74.
- 437 12. Rowland TW. The biological basis of physical activity. *Med Sci Sports Exerc*.
438 1998;30(3):392-9.
- 439 13. Wilkin TJ. Can we modulate physical activity in children? No. *Int J Obes*.
440 2011;35(10):1270-6.
- 441 14. Fröberg A, Raustorp A, Pagels P, Larsson C, Boldemann C. Levels of physical activity
442 during physical education lessons in Sweden. *Acta Paediatr*. 2017;106(1):135-41.
- 443 15. Sigmund E, Sigmundová D, Hamrik Z, Madarásová Gecková A. Does participation in
444 physical education reduce sedentary behavior in school and throughout the day among
445 normal-weight and overweight-to-obese Czech children aged 9-11 years? *Int J Environ*
446 *Res Public Health*. 2014;11(1):1076-93.
- 447 16. Santos SJ, Hardman CM, Barros SS, Franca CS, Barros MV. Association between
448 physical activity, participation in Physical Education classes, and social isolation in
449 adolescents. *J Pediatr (Rio J)*. 2015;91(6):543-50.
- 450 17. Tassitano RM, Barros MV, Tenório MC, Bezerra J, Florindo AA, Reis RS. Enrollment in
451 physical education is associated with health-related behavior among high school students.
452 *J Sch Health*. 2010;80(3):126-33.
- 453 18. Katzmarzyk PT, Barreira TV, Broyles ST, et al. The International Study of Childhood
454 Obesity, Lifestyle and the Environment (ISCOLE): design and methods. *BMC Public*
455 *Health*. 2013;13:900.

- 456 19. Tudor-Locke C, Barreira TV, Schuna JM Jr, et al. Improving wear time compliance with
457 a 24-hour waist-worn accelerometer protocol in the International Study of Childhood
458 Obesity, Lifestyle and the Environment (ISCOLE). *Int J Behav Nutr Phys Act.*
459 2015;12:11.
- 460 20. Barreira TV, Schuna Jr JM, Mire EF, et al. Identifying children's nocturnal sleep using a
461 24-h waist accelerometry. *Med Sci Sports Exerc.* 2015; 47(5):937-43.
- 462 21. Tudor-Locke C, Barreira TV, Schuna Jr JM, Mire EF, Katzmarzyk PT. Fully automated
463 waist-worn accelerometer algorithm for detecting children's sleep-period time separate
464 from 24-h physical activity or sedentary behaviors. *Appl Physiol Nutr Metab.*
465 2014;39(1):53-7.
- 466 22. Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG. Calibration of two
467 objective measures of physical activity for children. *J Sports Sci* 2008;26(14):1557-65.
- 468 23. World Health Organization. *Global Recommendations on Physical Activity for Health.*
469 [Internet]. 2010; [cited 2013 Nov 30]. Available from:
470 http://www.who.int/dietphysicalactivity/factsheet_recommendations/en/.
- 471 24. U.S. Centers for Disease Control and Prevention. *Youth Risk Behavior Surveillance*
472 *System (YRBSS)* [Internet]. 2012; [cited 2016 Dec 30]. Available from:
473 <https://www.cdc.gov/healthyyouth/data/yrbs/index.htm>.
- 474 25. Costa BG, Silva KS, George AM, Assis MA. Sedentary behavior during school-time:
475 Sociodemographic, weight status, physical education class, and school performance
476 correlates in Brazilian schoolchildren. *J Sci Med Sport.* 2017;20(1):70-4.
- 477 26. Sutherland R, Campbell E, Lubans DR, et al. Physical education in secondary schools
478 located in low-income communities: Physical activity levels, lesson context and teacher
479 interaction. *J Sci Med Sport.* 2016;19(2):135-41.
- 480 27. de Onis M, Onyanga AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a
481 WHO growth reference for school-aged children and adolescents. *Bull World Health*
482 *Organ.* 2007;85(9):660-7.

- 483 28. World Bank. *World Development Indicators*. [Internet]. 2012; [cited 2013 Nov 30].
484 Available from: <https://openknowledge.worldbank.org/handle/10986/6014>.
- 485 29. Skrondal A, Rabe-Hesketh S. Multilevel logistic regression for polytomous data and
486 rankings. *Psychometrika*. 2003;68(2):267-87.
- 487 30. Skrondal A, Rabe-Hesketh S. (2004). *Generalized Latent Variable Modeling: Multilevel,*
488 *Longitudinal and Structural Equation Modeling*. Boca Raton (FL): Chapman &
489 Hall/CRC; 2004. 528p.
- 490 31. Kim SY, So WY. The relationship between school performance and the number of
491 physical education classes attended by Korean adolescent students. *J Sports Sci Med*.
492 2012;11(2):226-30.
- 493 32. Klakk H, Andersen LB, Heidemann M, Møller NC, Wedderkopp N. Six physical
494 education lessons a week can reduce cardiovascular risk in school children aged 6-13
495 years: a longitudinal study. *Scand J Public Health*. 2014;42(2):128-36.
- 496 33. Park JW, Park SH, Koo CM, et al. Regular physical education class enhances sociality
497 and physical fitness while reducing psychological problems in children of multicultural
498 families. *J Exerc Rehabil*. 2017;13(2):168-78.
- 499 34. Telford RD, Cunningham RB, Fitzgerald R, et al. Physical education, obesity, and
500 academic achievement: a 2-year longitudinal investigation of Australian elementary
501 school children. *Am J Public Health*. 2012;102(2):368-74.
- 502 35. Palmer SE, Bycura DK, Warren M. A Physical Education Intervention Effects on
503 Correlates of Physical Activity and Motivation. *Health Promot Pract*. [Epub ahead of
504 print]
- 505 36. Mielke GI, Brown WJ, Nunes BP, Silva IC, Hallal PC. Socioeconomic Correlates of
506 Sedentary Behavior in Adolescents: Systematic Review and Meta-Analysis. *Sports Med*.
507 2017;47(1):61-75.
- 508 37. Muthuri SK, Wachira LJ, Leblanc AG, et al. Temporal trends and correlates of physical
509 activity, sedentary behaviour, and physical fitness among school-aged children in Sub-

510 Saharan Africa: a systematic review. *Int J Environ Res Public Health*. 2014;11(3):3327-
511 59.

512 38. García Bengoechea E, Sabiston CM, Ahmed R, Farnoush M. Exploring links to
513 unorganized and organized physical activity during adolescence: the role of gender,
514 socioeconomic status, weight status, and enjoyment of physical education. *Res Q Exerc*
515 *Sport*. 2010;81(1):7-16.

516 39. Sanchez-Vaznaugh EV, Goldman Rosas L, Fernández-Peña JR, Baek J, Egerter S,
517 Sánchez BN. Physical education policy compliance and Latino children's fitness: Does
518 the association vary by school neighborhood socioeconomic advantage? *Plos One*.
519 2017;12(6):e0178980.

520 40. LeBlanc AG, Katzmarzyk PT, Barreira TV, et al. Are participant characteristics from
521 ISCOLE study sites comparable to the rest of their country? *Int J Obes Suppl*.
522 2015;5(Suppl 2):S9-S16.

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537 **Table 1.** Descriptive characteristics of the sample.

538

Country (site)	Participants	Age (years)	MVPA	VPA	MPA	LPA	TPA	SB	MVPA	MVPA	MVPA	SB out	SB in	SB on
									out of school*	in school*	on weekends	of school*	school* weekends	
	(n, % males)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)	Mean (S.D.)
All sites	5874 (45.6)	10.4 (0.6)	60.3 (24.8)	17.9 (11.1)	42.4 (15.6)	314.9 (52.7)	375.2 (64.6)	513.4 (69.2)	36.3 (19.4)	25.8 (13.6)	55.4 (31.9)	292.9 (67.5)	233.8 (61.3)	488.7 (92.3)
HIC	2779 (43.5)	10.5 (0.5)	60.4 (23.0)	19.0 (11.4)	41.4 (13.5)	301.9 (48.2)	362.3 (58.6)	516.3 (66.8)	35.5 (17.1)	27.6 (12.6)	53.4 (29.5)	301.6 (61.0)	224.1 (47.5)	502.9 (91.6)
Australia (Adelaide)	439 (46.0)	10.7 (0.4)	65.2 (23.1)	22.5 (11.8)	42.7 (13.0)	310.1 (47.8)	375.3 (59.3)	476.8 (59.9)	35.0 (14.7)	34.6 (13.4)	55.7 (30.4)	269.9 (45.9)	210.8 (32.1)	475.7 (88.9)
Canada (Ottawa)	502 (41.2)	10.5 (0.4)	58.6 (19.4)	16.8 (9.0)	41.8 (12.1)	304.7 (44.5)	363.3 (54.6)	511.6 (62.9)	32.4 (14.2)	29.6 (9.9)	49.7 (24.2)	295.9 (49.2)	215.3 (29.9)	518.2 (86.6)
Finland (Helsinki, Espoo and Vantaa)	434 (46.4)	10.5 (0.4)	70.8 (26.2)	23.2 (13.9)	47.6 (15.1)	293.4 (43.7)	364.2 (58.4)	528.9 (67.1)	48.7 (21.8)	26.3 (12.0)	60.9 (34.2)	372.7 (66.3)	164.2 (30.0)	519.5 (89.2)
Portugal (Porto)	578 (43.4)	10.5 (0.3)	56.0 (21.8)	16.8 (10.1)	39.2 (13.4)	301.8 (50.0)	357.8 (59.7)	552.6 (61.3)	30.6 (14.7)	29.4 (12.9)	43.9 (25.4)	312.7 (48.7)	248.1 (34.7)	543.6 (81.3)
UK (Bath and North East Somerset)	377 (43.5)	10.9 (0.5)	64.2 (22.4)	20.9 (11.5)	43.3 (12.9)	286.1 (45.5)	350.3 (55.9)	495.2 (58.8)	39.3 (16.7)	27.8 (11.1)	58.3 (29.8)	283.7 (44.1)	223.2 (31.1)	483.1 (85.4)
USA (Baton Rouge)	449 (41.1)	9.9 (0.6)	49.9 (18.8)	15.1 (9.2)	34.8 (11.1)	313.7 (50.9)	363.6 (60.5)	520.4 (61.3)	30.2 (12.9)	17.3 (8.7)	56.1 (30.3)	271.2 (46.4)	273.8 (34.1)	461.9 (90.7)
LMIC	3095 (47.4)	10.3 (0.6)	60.1 (26.3)	16.9 (10.7)	43.2 (17.4)	326.9 (53.8)	387.0 (67.6)	510.6 (71.2)	36.9 (21.1)	24.2 (14.3)	57.3 (33.8)	285.1 (72.0)	242.6 (70.3)	475.9 (90.9)

Brazil (Sao Paulo)	435 (49.0)	10.5 (0.5)	59.4 (26.2)	17.7 (11.3)	41.7 (16.6)	337.9 (53.1)	397.3 (67.9)	500.6 (68.4)	36.4 (21.4)	24.8 (18.8)	54.1 (34.9)	315.1 (84.7)	194.6 (68.2)	483.0 (97.7)
China (Tianjin)	463 (52.0)	9.9 (0.5)	45.2 (15.9)	12.6 (6.7)	32.6 (10.5)	293.4 (53.6)	338.6 (62.3)	564.7 (67.7)	21.9 (10.3)	24.1 (11.3)	41.6 (21.5)	279.8 (42.4)	308.4 (43.1)	521.8 (82.8)
Colombia (Bogotá)	821 (49.3)	10.5 (0.6)	68.1 (24.8)	17.9 (10.1)	50.2 (16.8)	333.0 (49.4)	401.1 (63.2)	500.1 (67.1)	46.8 (21.9)	24.0 (13.0)	62.0 (32.3)	329.1 (63.6)	187.4 (48.5)	466.0 (85.4)
India (Bangalore)	526 (45.6)	10.5 (0.5)	48.7 (20.8)	12.8 (7.8)	35.9 (14.1)	340.1 (50.5)	388.8 (60.8)	516.7 (66.2)	27.6 (13.5)	21.5 (10.7)	47.0 (27.5)	286.9 (47.8)	244.1 (37.9)	488.1 (83.2)
Kenya (Nairobi)	459 (46.5)	10.3 (0.7)	71.6 (31.3)	22.5 (13.8)	49.1 (19.6)	329.9 (51.7)	401.5 (65.3)	494.7 (65.9)	37.0 (20.8)	33.7 (16.9)	74.8 (40.7)	201.5 (49.9)	314.6 (55.6)	454.3 (90.9)
South Africa (Cape Town)	391 (39.7)	10.3 (0.7)	63.5 (25.4)	17.9 (10.4)	45.6 (16.8)	321.5 (53.1)	385.0 (66.7)	489.1 (66.3)	47.2 (21.4)	16.8 (8.2)	62.8 (33.6)	260.9 (50.7)	247.8 (33.5)	443.6 (88.6)

539 HIC: High income countries; LMIC: Low- and middle-income countries; MVPA: moderate to vigorous physical activity; VPA: vigorous physical activity; MPA: moderate
540 physical activity; LPA: light physical activity; TPA: total physical activity; SB: sedentary behavior; *week days (from Monday to Friday).

541

542

543

544 **Figure 1.** Distribution of participation frequency in Physical Education (PE) classes by
545 study site.

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572 **List of Supplemental Digital Content**

573 Supplemental Digital Content 1. doc(x)

574 Descriptive characteristics of children who were excluded for missing data and of children
575 who participated in the present analysis.

576

577 Supplemental Digital Content 2. doc(x)

578 Descriptive characteristics of boys in the present study (n = 2,678).

579

580 Supplemental Digital Content 3. doc(x)

581 Descriptive characteristics of girls in the present study (n = 3,196).

582

583 Supplemental Digital Content 4. JPEG

584 Distribution of participation frequency in Physical Education (PE) classes of boys by study
585 site (n = 2,678).

586

587 Supplemental Digital Content 5. JPEG

588 Distribution of participation frequency in Physical Education (PE) classes of girls by study
589 site (n = 3,196).

590

591 Supplemental Digital Content 6. doc(x)

592 Distribution of the sample of boys in relation to meeting the recommendations for physical
593 activity, and highest tertile of time in physical activity (most active) and lowest tertile of time
594 in sedentary behavior (least sedentary) (n = 2,678).

595

596 Supplemental Digital Content 7. doc(x)

597 Distribution of the sample of girls in relation to meeting the recommendations for physical
598 activity, and highest tertile of time in physical activity (most active) and lowest tertile of time
599 in sedentary behavior (least sedentary) (n = 3,196).

600 **Supplemental Digital Content 1.** Descriptive characteristics of children who were excluded for
 601 missing data and of children who participated in the present analysis.

	Male (n = 3,422)		
	Children who were excluded	Children who were included in	p-value
	for missing data (n = 744)	the present analysis (n = 2,678)	
Age – years (Mean, S.D.)	10.4 (0.6)	10.5 (0.5)	0.95*
Level of parental education (%)			
Less high school	9.8	11.4	0.523†
Some high school	10.5	8.1	
Completed high school	22.6	23.3	
Some college	21.2	19.7	
Bachelor’s degree	17.7	18.2	
Post-graduate degree	18.2	19.3	
Countries (%)			
High-income	49.9	45.0	0.06†
Low- and middle-income	50.1	55.0	
Participation frequency in Physical Education classes/week (%)			
0	6.3	6.5	0.80†
1-2	68.5	69.7	
≥ 3	25.2	23.8	
BMI z-score (WHO)			
Severe thinness	0.4	0.1	0.12†
Thinness	1.5	1.5	
Normal	60.1	64.2	
Overweight	23.4	19.1	
Obesity	14.5	15.1	
	Female (n = 3,950)		
	Children who were excluded	Children who were included in	
	for missing data	the present analysis	

	(n = 754)	(n = 3,196)	
Age – years (Mean, S.D.)	10.4 (0.6)	10.4 (0.5)	0.54*
Level of parental education (%)			
Less high school	11.7	10.7	0.30†
Some high school	10.1	9.2	
Completed high school	26.4	23.1	
Some college	18.5	18.8	
Bachelor's degree	16.9	16.8	
Post-graduate degree	16.4	21.4	
Countries (%)			
High-income	52.2	49.2	0.24†
Low- and middle-income	47.8	50.8	
Participation frequency in Physical Education classes/week (%)			
0	8.0	6.9	0.71†
1-2	66.6	67.5	
≥ 3	25.4	25.6	
BMI z-score (WHO)			
Severe thinness	0.0	0.2	0.88†
Thinness	1.5	1.8	
Normal	66.1	66.8	
Overweight	21.5	21.4	
Obesity	10.9	9.9	

602 S.D.: standard deviation; BMI: body mass index; WHO: World Health Organization; * Student's t-
603 test; † Chi-squared test.

604

Country (site)	Age	MVPA	VPA	MPA	LPA	TPA	SB	MVPA	MVPA	MVPA	SB out of	SB in	SB on
	(years)							out of	in	on	school*	school*	weekends
	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean
	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)
All sites	10.5	69.8	21.7	48.1	319.6	389.4	503.9	40.7	31.2	64.1	290.8	223.4	488.7
	(0.5)	(25.8)	(12.2)	(15.9)	(51.5)	(63.6)	(69.9)	(20.8)	(14.9)	(34.9)	(69.1)	(60.8)	(96.6)
HIC	10.5	70.7	23.4	47.3	305.6	376.3	509.1	40.3	33.2	61.9	303.1	211.6	507.6
	(0.5)	(24.2)	(12.7)	(13.9)	(47.5)	(57.8)	(68.6)	(18.5)	(13.6)	(33.2)	(63.5)	(46.5)	(97.3)
Australia (Adelaide)	10.8	74.9	26.7	48.2	313.2	388.1	470.9	38.6	41.2	62.0	270.7	197.6	487.9
	(0.4)	(23.7)	(12.6)	(13.1)	(48.5)	(58.8)	(63.9)	(15.1)	(13.6)	(32.6)	(48.1)	(28.9)	(96.2)
Canada (Ottawa)	10.5	67.1	19.5	47.6	310.0	377.1	507.0	37.5	34.5	54.7	297.7	203.8	524.3
	(0.3)	(19.4)	(10.0)	(11.7)	(43.6)	(51.9)	(65.9)	(14.8)	(9.4)	(25.8)	(53.6)	(26.8)	(89.3)
Finland (Helsinki, Espoo and Vantaa)	10.5	81.6	27.8	53.8	298.6	380.2	524.1	54.0	31.6	72.0	374.1	154.2	519.5
	(0.4)	(27.9)	(15.5)	(15.6)	(42.8)	(58.0)	(70.6)	(24.1)	(13.2)	(38.8)	(70.2)	(37.5)	(97.0)
Portugal (Porto)	10.5	68.1	21.8	46.3	306.2	374.3	538.3	36.5	36.3	53.3	308.5	233.6	543.0
	(0.2)	(23.0)	(11.5)	(13.7)	(48.6)	(58.3)	(65.1)	(16.0)	(13.5)	(29.5)	(51.6)	(33.8)	(90.0)
UK (Bath and North East Somerset)	10.9	73.9	26.0	47.9	285.5	359.4	493.4	42.3	33.9	65.0	291.7	213.1	500.3
	(0.4)	(23.9)	(12.5)	(13.7)	(43.7)	(55.8)	(61.2)	(18.5)	(10.9)	(33.7)	(45.2)	(31.4)	(88.9)
USA (Baton Rouge)	10.0	58.1	18.4	39.7	318.3	376.4	512.3	34.1	19.8	67.9	270.0	265.6	454.1
	(0.6)	(19.9)	(10.1)	(11.4)	(50.7)	(60.5)	(62.1)	(13.5)	(9.8)	(34.2)	(47.1)	(34.0)	(97.5)
LMIC	10.3	69.0	20.2	48.8	331.4	400.4	499.5	40.9	29.5	65.9	280.2	233.2	472.7
	(0.6)	(26.9)	(11.5)	(17.5)	(51.7)	(66.2)	(70.8)	(22.4)	(15.7)	(36.2)	(71.7)	(69.1)	(92.9)
Brazil (Sao Paulo)	10.5	71.4	22.6	48.8	342.3	413.7	491.4	42.2	31.3	65.7	303.9	194.9	482.7
	(0.5)	(27.9)	(13.0)	(16.9)	(52.4)	(67.8)	(70.1)	(24.2)	(21.4)	(40.2)	(88.2)	(66.1)	(101.3)

China (Tianjin)	9.9	49.5	14.1	35.4	304.6	354.1	551.6	24.2	26.0	45.2	277.0	297.4	514.2
	(0.5)	(16.2)	(7.2)	(10.7)	(51.3)	(59.2)	(64.9)	(10.8)	(11.2)	(22.2)	(42.2)	(43.3)	(80.4)
Colombia (Bogotá)	10.5	76.3	21.4	54.9	336.7	413.0	491.4	50.0	29.3	69.1	323.6	180.8	467.4
	(0.6)	(25.8)	(11.1)	(17.3)	(48.0)	(63.3)	(66.9)	(23.6)	(14.7)	(34.9)	(63.9)	(50.6)	(88.0)
India (Bangalore)	10.5	61.5	16.8	44.7	348.4	409.9	490.3	33.0	28.9	61.4	272.8	225.1	473.8
	(0.5)	(10.7)	(8.4)	(13.9)	(49.0)	(58.2)	(61.0)	(14.8)	(10.4)	(31.7)	(45.1)	(32.5)	(83.5)
Kenya (Nairobi)	10.2	80.5	25.8	54.7	330.0	410.5	486.4	38.6	40.5	83.8	202.3	306.6	446.8
	(0.7)	(30.5)	(13.9)	(19.1)	(48.5)	(62.9)	(69.3)	(18.9)	(17.6)	(40.9)	(53.7)	(55.1)	(95.3)
South Africa (Cape Town)	10.4	72.8	21.2	51.6	321.6	394.4	481.2	54.9	20.2	71.5	256.5	239.9	444.3
	(0.7)	(25.9)	(10.5)	(17.0)	(51.5)	(66.2)	(70.4)	(22.6)	(8.8)	(34.7)	(52.4)	(34.6)	(95.8)

606 HIC: High income countries; LMIC: Low- and middle-income countries; MVPA: moderate to vigorous physical activity; VPA: vigorous physical activity; MPA: moderate
607 physical activity; LPA: light physical activity; TPA: total physical activity; SB: sedentary behavior; *week days (from Monday to Friday).

608


609

Country (site)	Age	MVPA	VPA	MPA	LPA	TPA	SB	MVPA	MVPA	MVPA	SB out	SB in	SB on
	(years)							out of	in	on	of	school*	weekends
	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean	Mean
	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)	(S.D.)
All sites	10.4	52.3	14.8	37.5	311.1	363.4	521.3	32.9	21.2	48.5	294.8	241.7	489.3
	(0.5)	(20.8)	(8.9)	(13.6)	(53.2)	(62.9)	(67.4)	(17.4)	(10.4)	(27.5)	(66.3)	(59.7)	(89.6)
HIC	10.5	52.6	15.7	36.9	299.1	351.7	521.8	31.7	23.2	46.3	301.2	233.6	502.1
	(0.5)	(18.4)	(8.8)	(11.3)	(48.5)	(56.8)	(64.7)	(14.7)	(9.6)	(24.4)	(59.4)	(45.8)	(88.0)
Australia (Adelaide)	10.7	56.9	18.9	38.0	307.5	364.4	481.7	31.4	28.6	49.8	269.4	221.7	466.2
	(0.4)	(19.1)	(9.7)	(10.8)	(47.1)	(57.5)	(56.0)	(13.3)	(9.9)	(26.8)	(43.4)	(30.3)	(80.8)
Canada (Ottawa)	10.5	52.7	14.9	37.8	300.9	353.6	514.8	28.6	26.1	46.3	294.6	223.0	513.2
	(0.4)	(16.9)	(7.6)	(10.7)	(45.4)	(54.3)	(60.6)	(12.3)	(8.8)	(22.3)	(46.2)	(29.3)	(84.2)
Finland (Helsinki, Espoo and Vantaa)	10.5	61.5	19.2	42.3	288.8	350.4	533.1	43.9	21.9	50.5	373.4	172.2	523.5
	(0.4)	(20.5)	(10.7)	(12.5)	(44.0)	(55.1)	(63.6)	(18.5)	(8.7)	(26.8)	(64.0)	(37.8)	(83.9)
Portugal (Porto)	10.4	46.7	13.0	33.7	298.4	345.1	563.4	26.4	24.1	37.6	313.5	259.3	543.7
	(0.3)	(15.1)	(6.5)	(10.0)	(50.8)	(57.6)	(56.1)	(11.3)	(9.0)	(20.0)	(46.3)	(31.0)	(78.7)
UK (Bath and North East Somerset)	10.9	56.7	16.9	39.7	286.6	343.3	496.7	35.8	22.9	50.4	279.1	232.0	477.9
	(0.4)	(17.8)	(8.7)	(11.0)	(46.9)	(54.9)	(56.9)	(14.0)	(8.7)	(25.1)	(43.3)	(28.0)	(83.4)
USA (Baton Rouge)	9.9	44.1	12.8	31.3	310.5	354.6	526.1	27.4	15.5	47.8	272.4	279.4	467.2
	(0.5)	(15.7)	(7.7)	(9.4)	(50.9)	(58.8)	(60.2)	(11.6)	(7.3)	(23.7)	(45.7)	(32.8)	(85.5)
LMIC	10.3	52.1	13.9	38.2	322.8	374.9	520.6	34.0	19.3	50.5	288.5	249.6	476.7
	(0.6)	(22.9)	(8.9)	(15.6)	(55.2)	(66.4)	(70.0)	(19.5)	(10.8)	(30.0)	(71.8)	(69.8)	(89.5)
Brazil (Sao Paulo)	10.4	47.8	12.9	34.9	333.7	381.5	509.3	31.1	18.6	43.7	324.4	194.0	482.3
	(0.5)	(18.3)	(6.4)	(13.1)	(53.6)	(64.2)	(65.7)	(16.7)	(13.1)	(24.2)	(80.7)	(70.2)	(93.4)


China (Tianjin)	9.9	40.5	11.0	29.5	281.2	321.7	578.8	19.2	22.0	37.5	283.0	320.0	530.5
	(0.5)	(14.1)	(5.6)	(9.3)	(53.3)	(61.2)	(68.0)	(8.8)	(10.8)	(19.8)	(42.4)	(39.6)	(84.5)
Colombia (Bogotá)	10.4	60.1	14.5	45.6	329.4	389.5	508.6	43.6	18.8	55.0	334.4	193.6	464.7
	(0.6)	(20.8)	(7.8)	(14.9)	(50.5)	(61.0)	(66.3)	(19.6)	(8.3)	(27.8)	(62.9)	(45.5)	(82.8)
India (Bangalore)	10.4	37.9	9.3	28.6	333.1	371.0	538.7	23.6	15.5	35.9	297.7	259.0	497.8
	(0.5)	(13.5)	(5.1)	(9.2)	(50.6)	(57.3)	(62.2)	(11.0)	(6.2)	(17.4)	(47.3)	(35.5)	(82.8)
Kenya (Nairobi)	10.2	63.8	19.6	44.2	329.8	393.6	501.9	35.6	28.0	67.3	200.7	321.3	460.4
	(0.6)	(29.9)	(13.0)	(18.7)	(54.4)	(66.3)	(61.9)	(22.0)	(13.8)	(39.0)	(46.6)	(55.1)	(86.6)
South Africa (Cape Town)	10.2	57.3	15.7	41.6	321.3	378.6	494.3	43.8	14.9	60.5	261.6	251.3	438.7
	(0.6)	(23.1)	(9.7)	(15.4)	(54.3)	(66.5)	(62.9)	(19.1)	(7.0)	(32.7)	(46.7)	(31.9)	(82.6)

611 HIC: High income countries; LMIC: Low- and middle-income countries; MVPA: moderate to vigorous physical activity; VPA: vigorous physical activity; MPA: moderate
612 physical activity; LPA: light physical activity; TPA: total physical activity; SB: sedentary behavior; *week days (from Monday to Friday).

613



Female



615

616

617 **Supplemental Digital Content 6.** Distribution of the sample of boys in relation to meeting the recommendations for physical activity, and highest tertile of time in physical
 618 activity (most active) and lowest tertile of time in sedentary behavior (least sedentary) (n = 2,678).

Country (site)	Meeting the MVPA recommendations	MVPA	VPA	MPA	LPA	TPA	SB	MVPA	MVPA	MVPA	SB	SB in	SB on	
								out of	in	on	out of	school	weekends	
									school	school	weekends	school		
			3 rd	3 rd	3 rd	3 rd	3 rd	1 st	3 rd	3 rd	3 rd	1 st	1 st	1 st
	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	
	%	%	%	%	%	%	%	%	%	%	%	%	%	
All sites	61.2	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	
HIC	64.7	33.0	38.4	30.4	22.5	24.5	29.3	31.3	40.5	31.3	23.6	35.4	25.1	
Australia (Adelaide)	73.2	39.1	51.4	34.5	26.8	32.3	48.6	32.5	62.7	33.0	41.1	46.9	34.4	
Canada (Ottawa)	60.1	27.2	24.4	31.5	24.9	23.9	30.5	25.4	46.9	20.1	23.0	33.0	18.7	
Finland (Helsinki, Espoo and Vantaa)	78.4	50.5	52.3	46.8	15.6	26.6	24.3	58.4	33.5	42.1	4.3	85.6	21.1	
Portugal (Porto)	61.0	29.0	32.7	26.4	24.5	23.0	14.5	24.5	48.9	23.7	13.5	15.3	10.6	
UK (Bath and North East Somerset)	71.1	37.4	47.1	29.4	10.7	16.0	39.0	30.9	37.6	37.6	22.7	32.0	26.0	
USA (Baton Rouge)	43.6	14.4	23.1	12.8	31.3	24.1	22.6	16.8	7.9	35.1	41.4	2.6	46.6	
LMIC	58.2	33.5	29.4	35.9	42.9	40.8	36.6	35.2	27.3	35.1	38.6	31.2	37.3	
Brazil (Sao Paulo)	64.8	36.5	39.7	36.5	48.4	42.5	39.3	39.7	36.3	34.2	34.2	59.0	35.9	
China (Tianjin)	23.3	6.2	8.9	6.2	21.7	13.6	10.5	3.7	17.8	10.0	32.8	0.8	16.2	
Colombia (Bogotá)	72.0	46.2	34.4	52.1	48.1	51.2	41.9	53.3	24.3	41.4	14.9	66.3	38.0	
India (Bangalore)	47.8	19.7	16.1	24.1	57.0	45.8	35.7	20.3	22.0	28.6	36.1	16.2	34.0	
Kenya (Nairobi)	71.2	47.2	44.2	44.6	41.2	47.2	45.1	24.9	58.9	52.6	87.1	3.3	53.1	
South Africa (Cape Town)	66.7	41.5	34.6	45.9	35.8	37.7	49.7	62.9	5.1	44.4	52.8	9.6	51.7	

619 HIC: High income countries; LMIC: Low- and middle-income countries; PE: physical education; MVPA: moderate-to-vigorous physical activity; VPA: vigorous physical
620 activity; MPA: moderate physical activity; LPA: light physical activity; TPA: total physical activity; SB: sedentary behavior.
621

622 **Supplemental Digital Content 7.** Distribution of the sample of girls in relation to meeting the recommendations for physical activity, and highest tertile of time in physical
 623 activity (most active) and lowest tertile of time in sedentary behavior (least sedentary) (n = 3,196).

Country (site)	Meeting the MVPA recommendations	MVPA	VPA	MPA	LPA	TPA	SB	MVPA	MVPA	MVPA	SB	SB in	SB on	
								out of	in	on	out of	school	weekends	
									school	school	weekends	school		
		3 rd	3 rd	3 rd	3 rd	3 rd	1 st	3 rd	3 rd	3 rd	1 st	1 st	1 st	
	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	tertile	
	%	%	%	%	%	%	%	%	%	%	%	%	%	
All sites	30.4	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	33.3	
HIC	30.7	33.6	37.8	30.5	23.6	25.0	32.9	29.9	42.2	30.8	27.8	32.7	27.2	
Australia (Adelaide)	39.9	42.2	52.3	32.9	28.3	30.6	55.8	28.4	63.8	32.5	46.5	39.9	42.4	
Canada (Ottawa)	31.6	35.5	34.2	34.5	25.0	25.3	34.9	21.1	53.5	30.8	26.4	37.5	19.4	
Finland (Helsinki, Espoo and Vantaa)	51.2	53.2	53.6	46.8	12.7	23.0	24.2	60.5	37.9	41.1	1.6	87.4	19.4	
Portugal (Porto)	16.5	19.1	25.1	18.2	23.9	23.1	10.8	15.1	43.8	16.4	14.0	10.4	11.5	
UK (Bath and North East Somerset)	38.3	42.8	43.2	42.4	16.5	21.0	47.7	45.7	41.9	35.5	38.5	29.5	35.5	
USA (Baton Rouge)	14.3	16.1	25.4	13.9	33.6	27.1	27.5	18.7	12.5	34.8	46.5	2.9	43.6	
LMIC	30.0	32.6	28.7	35.6	42.9	41.2	34.3	36.6	24.7	35.7	36.0	34.0	37.2	
Brazil (Sao Paulo)	25.0	28.1	27.2	30.7	49.6	44.3	36.4	30.5	24.0	29.7	23.6	72.8	35.4	
China (Tianjin)	8.0	10.1	17.2	10.1	16.4	14.3	8.4	3.1	30.4	17.2	28.2	0.9	15.0	
Colombia (Bogotá)	47.2	50.5	33.4	59.2	48.4	49.5	40.6	61.6	21.5	46.1	12.6	76.1	41.5	
India (Bangalore)	5.7	6.4	9.1	8.1	47.8	36.7	21.5	10.7	8.6	13.1	22.0	11.0	27.8	
Kenya (Nairobi)	45.1	47.4	48.1	47.0	47.4	54.1	44.0	36.7	57.4	54.2	91.2	3.2	46.6	
South Africa (Cape Town)	38.4	43.0	35.5	44.2	41.7	41.3	46.7	59.0	12.9	47.8	53.6	15.1	51.8	

624 HIC: High income countries; LMIC: Low- and middle-income countries; PE: physical education; MVPA: moderate-to-vigorous physical activity; VPA: vigorous physical
625 activity; MPA: moderate physical activity; LPA: light physical activity; TPA: total physical activity; SB: sedentary behavior.

626

627