

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/108198>

Please be advised that this information was generated on 2017-12-06 and may be subject to change.

Identification of a novel *FBN1* gene mutation in a large Pakistani family with Marfan syndrome

Shazia Micheal,^{1,2} Muhammad Imran Khan,^{1,3} Farah Akhtar,⁴ Marjan M. Weiss,⁵ Farah Islam,⁴ Mehmood Ali,⁴ Raheel Qamar,^{1,6} Alessandra Maugeri,⁵ Anneke I. den Hollander^{2,3}

¹Department of Biosciences, COMSATS Institute of Information Technology, Islamabad, Pakistan; ²Department of Ophthalmology, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands; ³Department of Human Genetics, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands; ⁴Al-Shifa Eye Trust Hospital, Rawalpindi, Pakistan; ⁵Department of Clinical Genetics, VU University Medical Center, Amsterdam, the Netherlands; ⁶Shifa College of Medicine, Islamabad, Pakistan

Purpose: To describe a novel mutation in the fibrillin-1 (*FBN1*) gene in a large Pakistani family with autosomal dominant Marfan syndrome (MFS).

Methods: Blood samples were collected of 11 family members affected with Marfan syndrome, and DNA was isolated by phenol-extraction. The coding exons of *FBN1* were analyzed by polymerase chain reaction (PCR) and direct sequencing. One hundred-thirty controls were screened for a mutation in the *FBN1* gene that was identified in this family by restriction fragment length polymorphism (RFLP) analysis.

Results: A novel heterozygous missense mutation c.2368T>A; p.Cys790Ser was observed in exon 19. This mutation substitutes a highly conserved cysteine residue by serine in a calcium binding epidermal growth factor-like domain (cbEGF) of *FBN1*. This mutation was present in all affected members and absent from unaffected individuals of the family in addition to 130 healthy Pakistani controls. Interestingly all affected family members presented with ectopia lentis, myopia and glaucoma, but lacked the cardinal cardiovascular features of MFS.

Conclusions: This is a first report of a mutation in *FBN1* in MFS patients of Pakistani origin. The identification of a *FBN1* mutation in this family confirms the diagnosis of MFS patients and expands the worldwide spectrum of *FBN1* mutations.

Marfan syndrome (MFS) is an autosomal dominantly inherited syndrome with a prevalence of 1 in 5,000–10,000 individuals. The major clinical manifestations of the syndrome include three major systems according to the Ghent criteria i.e., the ocular, skeletal and cardiovascular systems [1,2].

Ocular features mainly involve ectopia lentis, which is observed in around 80% of MFS patients. Ectopia lentis is characterized by the dislocation of the lens, which typically occurs in patients between birth and 20 years of age after that lens is stabilized. Other features include high myopic eyes and retinal detachment in individuals aged 50–59 years [3,4]. According to the new Ghent criteria another cardinal feature of MFS is aortic root aneurysm/dissection. The most common physical features were craniofacial characteristics, high-arched palate, positive thumb and wrist signs. If the family history of the patient is not positive, the involvement of at least two organ systems is required to establish the diagnosis of MFS. Genetic screening of MFS can aid the diagnosis, as the presence of a mutation in the fibrillin-1 gene (*FBN1*) in the

presence of a major manifestation of one organ system is sufficient to make the diagnosis [5].

There are three types of fibrillins in humans: *FBN1*, *FBN2* and *FBN3*. Fibrillins are extracellular matrix fibrillar components which are essential for the correct function of elastic and nonelastic tissues including blood vessels, bone and eye [6]. Fibrillin-1 is a 350-kDa protein responsible for head-to-tail assembly of 10–12-nm fibrillin monomers in presence of calcium-constituting microfibrils. *FBN1* forms a large multimeric protein complex by interacting with transforming growth factor beta ($TGF\beta$), latent $TGF\beta$ binding protein (LTBPs), and microfibrils that interact with bone morphogenetic protein (BMP) complexes. Depending on the requirements of a cell or tissue the *FBN1* complex can activate extracellular matrix (ECM) sequestered growth factors or inhibit activated growth factors. Thereby defects in fibrillin and its associated structure e.g., due to mutation, potentially could activate the growth factor signaling pathway, that can lead to MFS and related disorders of the connective tissue [7].

FBN1 is composed of three types of repeated modules. The epidermal growth factor (EGF)-like modules contain six highly conserved cysteine residues, which form disulphide bonds with each other and are critical for the stabilized folding of the domain. *FBN1* has 47 such modules, and 43 of them contain a calcium binding (cb) consensus sequence and are

Correspondence to: Anneke I. den Hollander, Department of Ophthalmology, Radboud University Nijmegen Medical Centre, P.O. Box 9101, 6500 HB, Nijmegen, The Netherlands; Phone: +31-24-3610402; FAX: +31-24-3540522; email: a.denhollander@ohk.umcn.nl

known as cbEGF-like modules [8]. The calcium ion bound in the cbEGF-like domain performs a crucial structural role in restricting the interdomain flexibility, which might have a role in protein-protein interaction [9]. The second type of module of FBN1 are transforming growth factor β 1-binding (or TB) protein-like module (TGF β 1-BP-like module, or 8- Cys/TB), which is found seven times in FBN1. This module contains eight cysteine residues that form four disulfide bonds. The third type of module is a hybrid module, which occurs twice [5].

Currently more than 1,200 mutations are known in the *FBN1* gene and missense mutations account for a major proportion (60%) of these mutations [10]. The majority of these mutations affect one of the cbEGF domains; often involving one of the six highly conserved cysteine residues within the cbEGF domains. Most mutations leading to a severe disease are found to be clustered in exons 24–32, which encodes a central stretch of 12 cbEGF repeats. This stretch is important in the formation of a rigid rod-like structure, which might be involved in the formation of microfibril assembly [11].

In this study we analyzed a five-generation family from Pakistan. All eleven affected family members lacked the cardinal cardiovascular features, but the diagnosis of MFS was confirmed by the identification of a novel mutation in the *FBN1* gene.

METHODS

Patients and clinical data: In this study we recruited a five-generation consanguineous family with two loops, one from central Punjab and the other from the Azad Jamu and Kashmir area of Pakistan. Eleven out of 14 affected individuals and 5 of 6 normal healthy individuals participated in the study. 130 additional normal healthy controls were recruited for the study. After obtaining informed consent, thorough physical, ocular and cardiovascular examinations were performed for all participating family members.

Molecular genetic analysis: Genomic DNA was extracted from whole blood using a conventional phenol-chloroform method [12]. PCR amplification of the 65 coding exons and flanking regions of the *FBN1* gene was performed in the proband (IV:13) using a PE 9700 thermocycler (Applied Biosystems, Foster City, CA). Primers used for PCR amplification are presented in Table 1. Briefly, for all amplicons the following cycling conditions were applied: initial denaturation at 94 °C for 5 min followed by 35 cycles of 94 °C for 30 s, 64 °C for 30 s, and 72 °C for 30 s. Sequencing reactions were performed using an ABI 3730 DNA analyzer from Applied Biosystems. For the detection of deletions and duplications, the SALSA MLPA kits P065 and P066 from MRC Holland (Amsterdam, the Netherlands) were used.

Segregation of a novel missense change in exon 19 was performed by direct sequencing of this exon in other family

members using standard conditions. The forward primer 5'-CAG GAG TTT TGC CTT TTT GC-3' and reverse primer 5'-TGC CAT GTA GAA CCA CAG AA-3' were used to amplify a 394 base pair (bp) product containing exon 19. PCR products were visualized on 2% agarose gel and purified by using PCR clean-up purification plates (NucleoFast® 96 PCR; MACHEREY-NAGEL, Düren, Germany), according to the manufacturer's protocol. Purified PCR products were analyzed by Sanger sequencing in an automated DNA sequencer (Big Dye Terminator, version 3 on a 3730 DNA analyzer; Applied Biosystems). Sequencing results were assembled and analyzed by using Vector NTI Advance™ 2011 software from Life technologies/Invitrogen (Bleiswijk, The Netherlands).

Unrelated control individuals were analyzed for the novel mutation by restriction fragment length polymorphism (RFLP) analysis using the restriction enzyme AlwNI (New England Biolabs, Ipswich, MA). PCR products were digested by using 2 U of enzyme, 1× PCR buffer 4 (20 mM Tris-acetate, 50 mM potassium acetate, 10 mM Magnesium Acetate, 1mM Dithiothreitol pH 7.9) and 16 μ l of amplified PCR product, and incubated for 2 h at 37 °C. After heat inactivation at 65 °C the product was separated on a 3% agarose gel.

RESULTS

Clinical characteristics: Eleven affected individuals of the family including 8 males and 3 females participated in the study (Figure 1). All patients showed similar clinical symptoms. In all affected members bilateral lens dislocation occurred and lenstectomy was performed. Other ocular symptoms included high myopia and glaucoma. None of them displayed any cardiovascular system abnormalities on echocardiography. Abnormalities of the skeletal system in MFS such as tall stature, long limbs, joint hypermobility, long narrow head, arachnodactyly, flat feet and medial displacement of medial malleoli, hollow cheeks and recessed or protruding jaw were noted in all individuals (Table 2).

Identification of a novel FBN1 mutation: Direct sequencing of *FBN1* revealed a novel heterozygous mutation (c. 2368T>A) in exon 19 which results in a change from a cysteine to a serine (p.Cys790Ser; Figure 2A,B). This missense mutation was present heterozygously in all affected members of the family while it was absent from normal individuals of the family as well as 130 unrelated healthy controls.

The p.Cys790Ser mutation resides in the 12th cbEGF domain, where it affects one of the highly conserved cysteine residues. Cys790 is the 4th cysteine residue of this cbEGF domain, which is predicted to form a disulphide bond with the 2nd cysteine residue (Cys776; Figure 3).

DISCUSSION

Of all mutations that have globally been identified in the *FBN1* gene, 38.6% result in a truncated FBN1 protein, and

TABLE 1. PRIMER SEQUENCES OF *FBN1*.

Primer name	Forward primer sequence	Reverse primer sequence	T °C		Product size
			annealing		
FBN1-E01-M13-01F & 01R	CAAGAGGGGGGGGAG	CGAAGGGGTGGGGACTAAACA	64	annealing	489
FBN1-E02-M13-01F & 01R	TATTTGGCCATCTCTCTCT	CCATGCAACCAACAACA	64		240
FBN1-E03-M13-02F & 02R	TGGTCCCCTATACAAATCGT	ATTGCAGGAAAGAGAAAGC	64		288
FBN1-E04-M13-01F & 01R	AGCTGTGCAATCTATGCATTTA	ATTACTTGTCTACAAAACAGGT	64		246
FBN1-E05-M13-01F & 01R	CCACAAGTGTACTTCATTAGCA	GTGCAAAATTAGTAACAGCTTAGG	64		260
FBN1-E06A-M13-01F & 01R	GCATGATGGTTCCTGCTT	AGACAATCCCGCTGAGTT	64		160
FBN1-E06B-M13-01F & 01R	CTGTGATCAGCAACCAGATG	GGCTCCAGAGCAAAATAAG	64		262
FBN1-E07-M13-01F & 01R	CTGCAATGAATTTCAITAGATTT	TTTTGCCTGCCCCCACTA	64		267
FBN1-E08-M13-01F & 01R	ACTGACGAATGGTTTTATATTGTG	AGTTGTTGTTATGGAACGTACTT	64		279
FBN1-E09-M13-01F & 01R	GTITTTACAAGTATTAATCTCAGCG	GGCTGGATGGATATTTCT	64		292
FBN1-E10-M13-01F & 01R	GCCTACGCTCAGCTGTG	AATGTTAACTTGAACAATGCAAGA	64		310
FBN1-E11-M13-01F & 01R	CAACATCTTGTTCATTAITGTTCAG	CAAGAAACAGAAITTAACAACAGAC	64		314
FBN1-E12-M13-01F & 01R	GGAAACCAGAAAGCTTAGAATTA	GTTAGCATATATGTCCCACATCC	64		270
FBN1-E13-M13-01F & 01R	ACTCCCCTAAATAAAGCTATTTCT	GCAATGGAAAGGAGAGGACT	64		245
FBN1-E14-M13-01F & 01R	GCCTACTCTTGGTCATAAGAAA	AAAGGCACGTGAAGAACA	64		243
FBN1-E15-M13-01F & 01R	GCCTGATGCTGCATATATTTCCCTA	TGAGTGACAGAGGCTGAAC	64		329
FBN1-E16-M13-01F & 01R	GGGTTCTCATCTGTTTGAAGT	CTCAATGGTGGCAAGG	64		294
FBN1-E17-M13-01F & 01R	CTGCAAAACAAGGGAATCAT	TGATGCTGCCCTGCACATA	64		182
FBN1-E18-M13-01F & 01R	TCCTGTAGCTCCTAAGGTCT	ATTATGCAGGCAATGTTTTAGA	64		308
FBN1-E19-M13-01F & 01R	AGATACAGGCAAAAGTTTGGG	CTAATGGCAITCCAAAAGATAGC	64		271
FBN1-E20-M13-01F & 01R	GGGTCAAAGTTGAAGTACTCT	GCAGGAAAAGCTGACATTAAG	64		296
FBN1-E21-M13-01F & 01R	ATTCCAAAGGTGTATGTTGAATTT	AGACCATTGGAGTGTGTATAGG	64		278
FBN1-E22-M13-01F & 01R	ACTATGTCAGAACTGCAAAAGTC	TATGACAGCTTTATCCAGTCCGA	64		243
FBN1-E23-M13-01F & 01R	ACTTACCAGTTCAAAAATGGG	CTAAGTGCTCAGCTATATCTGT	64		348
FBN1-E24A-M13-01F & 01R	ACAGAGTGTGGCAGTTTG	CTCCTCGTACTCAGGAGTATTT	64		268
FBN1-E24B-M13-01F & 01R	TGAGGAATCCGAGGAGTG	TGGGATGATCAAGTAGAGTGC	64		237
FBN1-E25-M13-01F & 01R	GCAATGAGACTCCTGACT	GCCTTAATCTTTGCCACAATATG	64		307
FBN1-E26-M13-01F & 01R	AAGCTGTCTGAGACTC	GCCTCATGGAAATCCCTCTCTT	64		259
FBN1-E27-M13-01F & 01R	TGGTGGAGGAGATGAGGC	GCAATGATGCAITCAAAAACACTG	64		270
FBN1-E28-M13-01F & 01R	TTCACACCAATTTACTTGTGGTC	ACATAGAGTGTTTAGGGAGAGAT	64		367
FBN1-E29-M13-01F & 01R	ACGAGATTTGGAGGGGAC	TAGGAACTACTGAGAGATTC AAC	64		266
FBN1-E30-M13-01F & 01R	ACTGAAACAGTGAACCAATATCAA	TGCTTATGACTAACAAAGACAAGAT	64		263
FBN1-E31A-M13-01F & 01R	GTCAATGTTATTAITGTTCTCGAGG	GTGGCAGATAAATGAGCCCTT	64		193
FBN1-E31B-M13-01F & 01R	ATCTGCCTAAGTGGGACC	ACAAAATTCAAAGAAAGTGGAAAGC	64		291
FBN1-E32-M13-01F & 01R	GACATTTGCTGAGCCT	ATGTTAATCTATGCAGTCTCTTG	64		254
FBN1-E33-M13-01F & 01R	ATTGGTTTTAAATACCACCCCTTC	TGGCTTCTGACTAGTGTG	64		258
FBN1-E34-M13-01F & 01R	CTAAACCGAGGAAAGATAACG	TCTCATCAAGCCCAAGCAAG	64		245
FBN1-E35-M13-01F & 01R	GATTGGTTAGATACTCTGCAT	GACACCAAGGAGCTGAT	64		315
FBN1-E36-M13-01F & 01R	CTCACACTGGCTCAGGTGATAA	ACACAGTATGCTTGTCTCTC	64		245
FBN1-E37-M13-01F & 01R	AGAAAGATTCGCTGATGC	GAGAACTGGCTGGATGA	64		292
FBN1-E38-M13-02F & 02R	ACAAAAGGTTAACCTACTTCAGAC	TAAACCCAAAGGAAATCAAGTTGTG	64		311
FBN1-E39-M13-01F & 01R	TTCCTGGGTTTATTACAAATGCT	CAGGTGACTTGTGATCTGC	64		265
FBN1-E40-M13-01F & 01R	GGCCATTCCAAAAATGTGAAG	GATGAGAACCAAAACATGCATTAAC	64		256
FBN1-E41-M13-01F & 01R	GTGATTTCCACATGGCA	TTTCCCAACAATTCATGGGTAA	64		314
FBN1-E42-M13-01F & 01R	TCCAATATTGTTCTTTGCTGACC	AAATAATGCTAACCAAAAGGCAAA	64		225
FBN1-E43-M13-01F & 01R	TGTGCTGCTGTCATC	GTAGCTCATCAGTTAGCTCTT	64		253
FBN1-E44-M13-01F & 01R	GTTGACTGGACACCAGATT	GAAGACAAAACCTTTGGGTAGG	64		260
FBN1-E45-M13-01F & 01R	CTCCTGAGAAATGATAGTAGAAGT	CAAATGAAGCTTTCAACAGCATA	64		294
FBN1-E46-M13-01F & 01R	CCGTGTAACCACTTTTCTACT	CTGGAACACATGAGATGATGTAA	64		291
FBN1-E47-M13-01F & 01R	TTGATTATGCTGGGATTAATGACA	ATGATTCCTTGTGATGTTCTCT	64		288
FBN1-E48-M13-01F & 01R	CAGTGGGAACCTCTTCTCTA	CCGACACTCCCTCATTTGCT	64		231
FBN1-E49-M13-01F & 01R	CTGATGATGCTCCATCTGTTG	TGCAGATTGAAAAGCCCA	64		250
FBN1-E50-M13-01F & 01R	TGCAATACGGACTCAGTAGG	TACTTACATCATGGCCAGTCT	64		302

TABLE 1. CONTINUED.

Primer name	Forward primer sequence	Reverse primer sequence	T °C		Product size
			annealing		
FBN1-E51-M13-02F & 02R	AGCATGTAGCAATTTTCTACCT	AAGAATAACTAGAGAAGAAGCAGAT	64	64	259
FBN1-E52-M13-01F & 01R	CTTCACGTTTAAAAAATACCTTGT	AGTCCATCTTGGTACCTAT	64	64	301
FBN1-E53-M13-01F & 01R	ACAACAACAACAACAATAATACAG	TGTTCCCAGGATCAGTACAC	64	64	303
FBN1-E54-M13-01F & 01R	TTGCTGTCCATGATCCCT	TTGCTGTCCATGATCCCT	64	64	245
FBN1-E55-M13-01F & 01R	AAAGTCAGGTAATTAAGGCAGATA	CTTCTGATGCACTCAAAGTCT	64	64	274
FBN1-E56-M13-01F & 01R	AAATGGTCAGATGACTCTTCTTG	GTGTGGAGGCTGAGGTTAG	64	64	253
FBN1-E57-M13-01F & 01R	TGCTCTTAAAAATTTCTTGACATCC	ACAAATAAATAGATTCCTGCAAG	64	64	338
FBN1-E58-M13-01F & 01R	AGTATTTACACTGAAGTGACCC	AAAATTTCCACTTGAGGATAAGC	64	64	285
FBN1-E59-M13-01F & 01R	TGAGCGTGTACACATCATTT	GGAATGCAGCCAATGTGTC	64	64	268
FBN1-E60-M13-01F & 01R	CCTGTTTGTGGCTTGAC	GAATCGCTACAATCCATGTAGG	64	64	240
FBN1-E61-M13-01F & 01R	ATGATACAAAAGAGAGCTTTGGG	CCTCCACAAGGATTCACCA	64	64	269
FBN1-E62-M13-01F & 01R	CTTCAGAGAGAGATGTGAGTTG	TGTTTTGCTTCATAGGACCTGATA	64	64	263
FBN1-E63A-M13-01F & 01R	TCAATAGAAAATCTCTGGCTGCT	TCCTCCACTGAACTGTTCCATAC	64	64	202
FBN1-E63B-M13-01F & 01R	TACAAAGTGCATGTGTC	ACGAAATGAAAGAATCTCCAACC	64	64	245
FBN1-E64-M13-01F & 01R	CCTACCTTGTCTTCCCATTTTAA	TTCCACCACAGGAGACAT	64	64	305
FBN1-E65A-M13-01F & 01R	CTTTAATATGAGAGCTAAGTGCCA	AGCCATCTTCATTTCCAGATTC	64	64	278
FBN1-E65B-M13-01F & 01R	CTCTGACGAATCAACACAGATAC	ATATGATGATTTCTGATTTGGGGGA	64	64	377




Figure 1. Pedigree of a Pakistani family with MFS. Squares indicates males and circles females, black symbols represents affected and white unaffected individuals, gray symbols indicates unknown affection status. Slashed symbols represent the deceased subjects. + indicates the normal allele, and M indicates the Cys790Ser mutation.

60.3% represent missense mutations and of which most (78%) are localized in the cbEGF-like modules [13]. In the 43 cbEGF domains each have six highly conserved cysteine residues (C1-C6), which form disulphide bonds among each other (C1-C3, C2-C4, C5-C6). We identified a novel mutation in exon 19 of the *FBNI* gene in a large MFS family from Pakistan, which, is predicted to abolish the C2-C4 (Cys776-Cys790) disulphide bond of the 12th cbEGF domain, as the 4th cysteine residue is replaced with serine (Cys790Ser). Missense mutations in *FBNI* that affect the cysteines, which are essential for the correct EGF-like domain structure, act in a dominant negative manner. Since the monomers from the mutated allele are folded incorrectly, they assemble with the normal monomers from the other allele creating abnormal multimers [14,15].

So far, no clear genotype-phenotype correlations in *FBNI* have yet been established, though some correlations have been suggested in some studies, which included a large number of individuals (n=101, 93, 57, 81, and 76 patients). Interestingly, a higher frequency of cysteine substitutions was

observed in MFS patients with ectopia lentis, opposed to premature termination codon mutations [16-20]. This is in line with the clinical findings in the family described in this study, as all affected family members developed ectopia lentis.

Previously most of the *FBNI* mutations were found in other exons rather than exon 19. To date only four mutations have been identified in exon 19: a frameshift mutation observed in one Italian patient with clinical symptoms mainly involving the skeletal and cardiovascular systems [21], and three missense mutations in 3 sporadic patients from Belgium with classical MFS and involvement of the cardiovascular system [1]. All three missense mutations were present in the 8th cbEGF domain affecting the 2nd and 3rd cysteine residues. Though all mutations identified in exon 19 have so far been associated with involvement of the cardiovascular system, this is not the case in the family described in this study, as none of the affected family members showed cardiovascular system abnormalities on echocardiography.

It has been reported that mutations in exons 24-32 are found in MFS individuals with a more severe and complete

TABLE 2. CLINICAL EVALUATION OF AFFECTED FAMILY MEMBERS.

Patient ID	Manifestation	III:4	III:5	III:7	IV:4	IV:5	IV:7	IV:8	IV:9	IV:11	IV:12	IV:13
Age (years)		55	40	35	30	28	12	16	8	20	6	6
Gender		M	M	M	M	M	F	F	F	M	M	M
Ocular System												
(i)	Ectopia lentis	+	+	+	+	+	+	+	+	+	+	+
(ii)	Myopia	+	+	+	+	+	+	+	+	+	+	+
(iii)	Abnormally flat cornea	-	-	-	-	-	-	-	-	-	-	-
(iv)	Early development of nuclear cataract	+	+	+	-	-	-	+	-	+	+	+
(V)	Glaucoma (IOP)	R34 L34	15 15	R14 L14	R15 L12	R14 L16	R20 L24	R16 L16	R12 L14	R18 R16	R14 L12	R18 L20
(Vi)	Retinal detachment	+	-	+	+	-	-	-	-	-	-	-
Cardiovascular system												
(i)	Aortic root dimension (mm)*	32	31.8	23.6	35.1	27.5	25.1	28.7	20	34	35	20
(ii)	Mitral valve prolapse	-	-	-	-	-	-	-	-	-	-	-
(iii)	Dilation of pulmonary artery	-	-	-	-	-	-	-	-	-	-	-
Skeletal system												
(i)	Height (H; cm)	176.7	173.4	167.6	170.6	167.4	146.3	158.5	121	167.6	106.6	106.6
(ii)	Arm span (AS; cm)	182	177	172	175	173	153	164	125	175	110	110
(iii)	AS/H (normal <1.05)	1.03	1.02	1.02	1.02	1.03	1.04	1.03	1.03	1.04	1.03	1.03
(iv)	Pectus carinatum	-	-	-	-	-	-	-	-	-	-	-
(v)	Pectus excavatum	-	-	-	-	-	-	-	-	-	-	-
(vi)	Arachnodactyly	+	+	+	+	+	+	+	+	+	+	+
(vii)	High palate with dental crowding	+	+	+	+	+	+	+	+	+	+	+
(viii)	Flatfoot	+	+	+	+	+	+	+	+	+	+	+
(ix)	Characteristic face (long narrow head, hollow cheeks, recessed or protruding jaw)	+	+	+	+	+	+	+	+	+	+	+
Other manifestations												
(i)	Hyperextensible skin	-	-	-	-	-	-	-	-	-	-	-
(ii)	Striae	-	-	-	-	-	-	-	-	-	-	-
(iii)	Hernia	+	+	+	+	+	-	-	+	+	-	-

IOP: intraocular pressure, R: right eye, L: left eye. *normal range 20–37 mm.

phenotype, including a younger age at diagnosis and a higher probability of developing ectopia lentis, ascending aortic dilatation, aortic surgery, mitral valve abnormalities, scoliosis, and shorter survival [11]. The ocular manifestations in this family, carrying a mutation in exon 19 of *FBNI*, were severe. All patients, including 4 young children, developed ectopia lentis, myopia and glaucoma. However, no cardiovascular system defects have been developed in the affected individuals so far. Moreover, among skeletal manifestations no pectus abnormalities or scoliosis have been noted.

Loeys et al. [1] did not find distinguishing features in patients with MFS with or without a *FBNI* mutation except for the presence of ectopia lentis, which was significantly higher in individuals with mutations in *FBNI*. A significant difference was observed between the clinical phenotype of

patients carrying a missense mutation in 11th and 12th EGF-like domains compared with patients carrying a missense mutation in EGF-like domains 13–18. Patients with mutations in EGF-like domains 11 and 12 have a shorter survival, a younger age at diagnosis, a higher probability of neonatal presentation, and a higher risk of developing ascending aortic dilatation than patients with a missense mutation affecting EGF-like domains 13–17 [11]. In the Pakistani family, carrying a mutation in the 12th EGF-like domain, patients were also diagnosed at a young age and some of them have a neonatal presentation, but no cardiovascular symptoms were observed.

Marfan syndrome is characterized by a high clinical heterogeneity. The presentation of cardinal symptoms varies among families but also within families. The prevalence of cardiovascular features such as mitral valve prolapse is 43%




Figure 2. Sequence traces of the novel *FBNI* missense mutation Cys790Ser in exon 19. **A:** A heterozygous change T>A (indicated by arrow) is identified in an affected family (IV:13) member. **B:** The corresponding normal sequence in an unaffected family member (IV:10).

and ascending aortic dilation is 53% in individuals in the age group <30. It increases steadily with age to 75% and 96%, respectively, in individuals of ≤ 60 years with *FBNI* mutations [22]. The affected individuals of the 5-generation family we present here lack the cardinal cardiovascular features of MFS, but do have the typical skeletal and ocular symptoms. The novel mutation in the *FBNI* gene identified in this family supports the diagnosis of MFS.

Although cardiovascular features are absent in the family, it has been reported that in some cases these features may not manifest until adulthood. Black et al. [23] reported the development of aortic root dilatation in the fifth decade of life in patients presenting familial ectopia lentis at the age of 58 and 70 years. In the present family only one affected individual has reached age 50, whereas three are in the age group of 30–40 years, four affected individuals are in their

childhood and 3 are infants. Therefore the young age might explain the absence of cardiovascular manifestation in these individuals. The study of Black et al. [23] confirms the need for life-long screening in adult patients carrying a *FBNI* mutation in the absence of major manifestations.

In this family of Pakistani origin affected individuals experience major involvement of the ocular and skeletal systems. Interestingly, all patients in this family developed ectopia lentis, had high myopia and glaucoma. In literature it has been reported that glaucoma occurs in 64% of patients with MFS and in 10% of patients with isolated ectopia lentis [24]. Glaucoma has been observed more frequently in spontaneous late subluxation of the lens than in the congenital type. However, in the family described here glaucoma was observed in combination with early subluxation of the lens in all affected individuals, suggesting that predisposing

factors may be present in this family, adding to the risk of developing glaucoma.

To the best of our knowledge this is the first report of a family with *FBN1* mutations, in which all affected individuals developed glaucoma and myopia. The presence of these particular clinical features in all affected individuals might be due to the novel mutation p.Cys790Ser, or alternatively may be associated with additional genetic factors that contribute to the disease phenotype in this family. We believe that the p.Cys790Ser mutation affects the structure of the protein by disturbing the arrangement of the microfilaments, and acts in a dominant negative manner. Although more than 1,200 mutations in the *FBN1* gene have been identified for MFS, this is a first report from Pakistan, which expands the worldwide mutation spectrum and adds to the existing knowledge of genotype-phenotype comparisons for MFS.




Figure 3. A schematic representation of the consensus secondary structure of a prototypical cbEGF-like domain. Calcium binding in the NH₂-terminal region of the wild-type domain is mediated by the highly conserved amino acids highlighted in blue. The highly conserved cysteines of the cbEGF-like domain are marked with a C, and the lines between the cysteine residues represent disulfide bridges. The mutation Cys790Ser affects the 4th cysteine residue, which is predicted to form a disulfide bridge with the 2nd cysteine residue (Cys776). The mutation abolishes the disulfide bond formation and thus causes misfolding of the protein.

ACKNOWLEDGMENTS

We would like to thank all the family members and control individuals who participated in the study.

REFERENCES

- Loeys B, Nuytinck L, Delvaux I, De Bie S, De Paepe A. Genotype and phenotype analysis of 171 patients referred for molecular study of the fibrillin-1 gene *FBN1* because of suspected Marfan syndrome. *Arch Intern Med* 2001; 161:2447-54. [PMID: 11700157]
- Loeys BL, Dietz HC, Braverman AC, Callewaert BL, De Backer J, Devereux RB, Hilhorst-Hofstee Y, Jondeau G, Faivre L, Milewicz DM, Pyeritz RE, Sponseller PD, Wordworth P, De Paepe AM. The revised Ghent nosology for the Marfan syndrome. *J Med Genet* 2010; 47:476-85. [PMID: 20591885]
- Charteris DG. Ophthalmic abnormalities in Marfan syndrome. In: Child, AH Briggs MJ, editors. *The Marfan Syndrome: A Clinical Guide*. London: British Heart Foundation; 2002. p. 52-53.
- Hasan A, Poloniecki J, Child A. Ageing in Marfan syndrome. *Int J Clin Pract* 2007; 61:1308-20. [PMID: 17627709]
- Colod-Bérout G, Boileau C. Marfan syndrome in the third Millennium. *Eur J Hum Genet* 2002; 10:673-81. [PMID: 12404097]
- Hubmacher D, Tiedemann K, Reinhardt DP. Fibrillins: from biogenesis of microfibrils to signaling functions. *Curr Top Dev Biol* 2006; 75:93-123. [PMID: 16984811]
- Ramirez F, Sakai LY. Biogenesis and function of fibrillin assemblies. *Cell Tissue Res* 2010; 339:71-82. [PMID: 19513754]
- Handford PA, Mayhew M, Baron M, Winship PR, Campbell ID, Brownlee GG. Key residues involved in calcium-binding motifs in EGF-like domains. *Nature* 1991; 351:164-7. [PMID: 2030732]
- Werner JM, Knott V, Handford PA, Campbell ID, Downing AK. Backbone dynamics of a cbEGF domain pair in the presence of calcium. *J Mol Biol* 2000; 296:1065-78. [PMID: 10686104]
- Robinson PN, Booms P, Katzke S, Ladewig M, Neumann L, Palz M, Pregla R, Tiecke F, Rosenberg T. Mutations of *FBN1* and genotype phenotype correlations in Marfan syndrome and related fibrillinopathies. *Hum Mutat* 2002; 20:153-61. [PMID: 12203987]
- Faivre L, Colod-Beroud G, Loeys BL, Child A, Binquet C, Gautier E, Callewaert B, Arbustini E, Mayer K, Arslan-Kirchner M, Kiotsekoglou A, Comeglio P, Marziliano N, Dietz HC, Halliday D, Beroud C, Bonithon-Kopp C, Claustres M, Muti C, Plauchu H, Robinson PN, Adès LC, Biggin A, Benetts B, Brett M, Holman KJ, De Backer J, Coucke P, Francke U, De Paepe A, Jondeau G, Boileau C. Effect of mutation type and location on clinical outcome in 1,013 probands with Marfan syndrome or related phenotypes and *FBN1* mutations: an international study. *Am J Hum Genet* 2007; 81:454-66. [PMID: 17701892]
- Sambrook J, Russell DW, Sambrook J. *The condensed protocols from Molecular cloning: a laboratory manual*. Cold

- Spring Harbor (NY): Cold Spring Harbor Laboratory Press; 2006.
13. Collod-B eroud G, Boileau C. Marfan Syndrome: A Primer for Clinicians and Scientists, Eureka.com and Luwer Academic/Plenum Publishers; 2004. p. 1–11.
 14. Sakai LY, Keene DR, Glanville RW, B achinger HP. Purification and partial characterization of fibrillin, a cysteine-rich structural component of connective tissue microfibrils. *J Biol Chem* 1991; 266:14763-70. [PMID: 1860873]
 15. Dietz HC, McIntosh I, Sakai LY, Corson GM, Chalberg SC, Pyeritz RE, Francomano CA. Four novel FBN1 mutations: significance for mutant transcript level and EGF-like domain calcium binding in the pathogenesis of Marfan syndrome. *Genomics* 1993; 17:468-75. [PMID: 8406497]
 16. Schrijver I, Liu W, Odom R, Brenn T, Oefner P, Furthmayr H, Francke U. Premature termination mutations in FBN1: distinct effects on differential allelic expression and on protein and clinical phenotypes. *Am J Hum Genet* 2002; 71:223-37. [PMID: 12068374]
 17. Loeys B, De Backer J, Van Acker P, Wettinck K, Pals G, Nuytinck L, Coucke P, De Paepe A. Comprehensive molecular screening of the FBN1 gene favors locus homogeneity of classical Marfan syndrome. *Hum Mutat* 2004; 24:140-6. [PMID: 15241795]
 18. Biggin A, Holman K, Brett M, Bennetts B, Ade's L. Detection of thirty novel FBN1 mutations in patients with Marfan syndrome or a related fibrillinopathy. *Hum Mutat* 2004; 23:99. [PMID: 14695540]
 19. Arbustini E, Grasso M, Ansaldi S, Malattia C, Pilotto A, Porcu E, Disabella E, Marziliano N, Pisani A, Lanzarini L, Mosconi M, Antoniazzi E, Zoia MC, Meloni G, Magrassi L, Brega A, Bedeschi MF, Torrente I, Mari F, Tavazzi L. Identification of sixty-two novel and twelve known FBN1 mutations in eighty-one unrelated probands with Marfan syndrome and other fibrillinopathies. *Hum Mutat* 2005; 26:494. [PMID: 16222657]
 20. Rommel K, Karck M, Haverich A, von Kodolitsch Y, Rybczynski M, Muller G, Singh KK, Schmidtke J, Arslan-Kirchner M. Identification of 29 novel and nine recurrent fibrillin-1 (FBN1) mutations and genotype-phenotype correlations in 76 patients with Marfan syndrome. *Hum Mutat* 2005; 26:529-39. [PMID: 16220557]
 21. Pepe G, Giusti B, Evangelisti L, Porciani MC, Brunelli T, Giurlani L, Attanasio M, Fattori R, Bagni C, Comeglio P, Abbate R, Gensini GF. Fibrillin-1 (FBN1) gene frameshift mutations in Marfan patients: genotype-phenotype correlation. *Clin Genet* 2001; 59:444-50. [PMID: 11453977]
 22. D etaint D, Faivre L, Collod-Beroud G, Child AH, Loeys BL, Binquet C, Gautier E, Arbustini E, Mayer K, Arslan-Kirchner M, Stheneur C, Halliday D, Beroud C, Bonithon-Kopp C, Claustres M, Plauchu H, Robinson PN, Kiotseoglou A, De Backer J, Ad es L, Francke U, De Paepe A, Boileau C, Jondeau G. Cardiovascular manifestations in men and women carrying a FBN1 mutation. *Eur Heart J* 2010; 31:2223-9. [PMID: 20709720]
 23. Black C, Withers AP, Gray JR, Bridges AB, Craig A, Baty DU, Boxer M. Correlation of a recurrent FBN1 mutation (R122C) with an atypical familial Marfan syndrome phenotype. *Hum Mutat* 1998; S198-200. [PMID: 9452085]
 24. Nelson LB, Maumenee IH. Ectopia lentis. *Surv Ophthalmol* 1982; 27:143-60. [PMID: 6984233]

Articles are provided courtesy of Emory University and the Zhongshan Ophthalmic Center, Sun Yat-sen University, P.R. China. The print version of this article was created on 14 July 2012. This reflects all typographical corrections and errata to the article through that date. Details of any changes may be found in the online version of the article.