



eResearch: the open access repository of the research output of Queen Margaret University, Edinburgh

This is an author-formatted version of a presentation published as:

Roberston, J and MacDonald, Kath (2009) *Bone loss in CF: A fragmented Picture*. In: European Cystic Fibrosis Conference, Brest, France.

Accessed from:

<http://eresearch.qmu.ac.uk/961/>

Repository Use Policy

The full-text may be used and/or reproduced, and given to third parties for personal research or study, educational or not-for-profit purposes providing that:

- The full-text is not changed in any way
- A full bibliographic reference is made
- A hyperlink is given to the original metadata page in eResearch

eResearch policies on access and re-use can be viewed on our Policies page:
<http://eresearch.qmu.ac.uk/policies.html>

Copyright © and Moral Rights for this article are retained by the individual authors and/or other copyright owners.


<http://eresearch.qmu.ac.uk>

Bone loss in CF: A fragmented picture

Robertson J¹, MacDonald K², Innes JA¹.

¹Scottish Adult CF Service, Western General Hospital, Edinburgh

²Queen Margaret University Edinburgh



INTRODUCTION

Bone loss is highly prevalent in our CF population (70%) and is associated with risk factors: age, low vitamin D levels, low BMI and CFRDM.¹ The evidence that low BMD is a predictor of fracture risk is inconclusive and whilst there are some data to suggest that oral bisphosphonates^{2,3} increase BMD, what remains unclear is the clinical significance of bone loss we have documented in our centre. We hypothesise that bone loss occurs in patients with risk factors described above, and that treatment using the CF Trust Bone Mineralisation Guidelines will result in improvement in BMD. However, we present 4 case studies which show that not all patients “fit this picture” and that treatment does not always result in the predicted outcome.



Case study 1

- o Male aged 37: DF 508 Homozygote, PI, FEV₁ 17%, BMI 20.4, CFRDM (on insulin), Vitamin D*23nmol/L, frequent exacerbations, referred for lung transplant.
- o 1st DEXA: Z Scores -3.9 (LS), -2.4 (H) Treated with Alendronate 70mg orally weekly.
- o Rpt DEXA @ 2yrs Z Scores -2.7 (LS) -2.2 (H), (FEV₁ 21% BMI 25.9, Vitamin D 53nmol/L)
- o Rpt DEXA @ 4yrs Z Scores -2.6 (LS), -2.0 (H) (FEV₁ 16%, BMI 24.9, Vitamin D 98nmol/L)
- o Rpt DEXA @ 5.5yrs Z Scores -3.9 (LS), -1.9 (H), (FEV₁ 15%, BMI 24.8 Vitamin D 41nmol/L)

Summary *Despite treatment for 5 yrs, no improvement in bone loss, however overall decline in health*

Case study 2

- o Female aged 43: DF508/unknown, PI, FEV₁ 25%, BMI 17, CFRDM (no insulin) Vitamin D 68nmol/L, frequent exacerbations, NG feeding
- o 1st DEXA: Z Scores -0.4 (LS), -0.5 (H)
- o No treatment (died 6 months post scan)

Summary *No bone loss in patient with severe CF despite multiple risk factors.*

Case study 3

- o Male aged 39: DF508 Homozygote, PI, FEV₁ 31%, BMI 21.9, CRDM (on insulin), Vitamin D 82nmol/L
- o 1st DEXA: Z Scores -0.3(LS), -0.7 (H)- No treatment
- o Rpt DEXA @3yrs Z Scores -0.3 (LS), -0.6(H) FEV₁ 31%, BM1 22.2, Vitamin D 53nmols/L
- o now frequent exacerbations, referred for transplant.

Summary *DEXA scan continues to be in normal range despite worsening health and falling Vitamin D levels.*

Case study 4

- o Male aged 48:DF508/5T-9T, PS, FEV₁ 100%, BMI 31.5, Vitamin D 25nmols/L,
- o 1st DEXA: Z Scores -2.5(LS), -1.3(H) Treated with Alendronate 70mg orally weekly
- o Rpt DEXA @ 2.5yrs Z Scores -2.7 (LS), -1.0 (H), (FEV₁ 101%, BM1 32.4, Vit D 30nmols/L)

Summary *Deterioration in DEXA scan at 2.5yrs post Alendronate treatment, despite stable health in well patient with mild CF.*

*Vitamin D measured as 25-OH Vitamin D (nmols/L range:25-170) LS=Lumbar Spine H=hip

CONCLUSION

These 4 case studies illustrate the diversity of presentation and variability in outcomes with or without treatment for bone loss. Adherence to therapy and individual differences in response to treatment are hard to predict. We conclude that detection and management of bone loss in CF continues to be challenging. Individual treatment decisions should be based on individual measurements, as protocol-based decisions do not suit the circumstances of all individuals.

References

1. Cystic Fibrosis Trust (2007) Bone Mineralisation in Cystic Fibrosis, CF Trust Kent.
2. Aris RM, Lester GE, Caminetti M, et al (2004) Efficacy of alendronate in adults with cystic fibrosis with low bone density, American Journal of Respiratory Critical Care Medicine 169:77-82.
3. Cawood T.J, McKenna M.J, Gallagher C.G. et al (2005) Oral bisphosphonates improve bone mineral density in adults with cystic fibrosis, Irish Medical Journal:98:270-273.