

RESEARCH ARTICLE

Open Access



The head shaft angle is associated with hip displacement in children at GMFCS levels III-V - a population based study

L Finlayson¹, T Czuba², M S Gaston³, G Hägglund^{4*}  and J E Robb⁵

Abstract

Background: An increased Head Shaft Angle (HSA) has been reported as a risk factor for hip displacement in children with cerebral palsy (CP) but opinions differ in the literature. The purpose of this study was to re-evaluate the relationship between HSA and hip displacement in a different population of children with CP.

Methods: The Cerebral Palsy Integrated Pathway Scotland surveillance programme includes 95% of all children with CP in Scotland. The pelvic radiographs from 640 children in GMFCS levels III-V were chosen. The most displaced hip was analysed and the radiographs used were those taken at the child's first registration in the database to avoid the potential effects of surveillance on subsequent hip centration. A logistic regression model was used with hip displacement (migration percentage [MP] $\geq 40\%$) as outcome and HSA, GMFCS, age and sex as covariates.

Results: The MP was $\geq 40\%$ in 118 hips with a mean HSA of 164° (range $121\text{--}180^\circ$) and $< 40\%$ in 522 hips with a mean HSA of 160° (range $111\text{--}180^\circ$). The logistic regression analysis showed no significant influence of age and sex on MP in this population but a high GMFCS level was strongly associated with hip displacement. An increased HSA was also associated with hip displacement, a 10° difference in HSA for children adjusted for age, sex, and GMFCS gave an odds ratio of 1.26 for hip displacement equal or above 40% ($p = 0.009$) and hips with HSA above 164.5 degrees had an odds ratio of 1.96 compared with hips with HSA below 164.5 degrees ($p = 0.002$).

Conclusion: These findings confirm that HSA is associated with hip displacement in children in GMFCS levels III-V.

Keywords: Children, Cerebral palsy, Hip displacement, Head shaft angle

Background

For hip surveillance in cerebral palsy (CP) it is important to identify risk factors for hip displacement both for the treatment of the individual child and to optimise the monitoring programme. Young age and severe limitation of gross motor function as measured with the Gross Motor Function Classification System (GMFCS) are known risk factors.

The measurement of the head-shaft angle (HSA, Fig. 1) was popularised by Southwick [1] to measure the degree of slip in slipped upper femoral epiphysis. Foroohar et al. [2] evaluated the relationship between the femoral shaft and head in children with cerebral palsy (CP) and compared the HSA in children with CP with the HSA in

typically developing children. They also evaluated the HSA in children with CP who had significant hip displacement. They concluded that the HSA was greater in children with CP and more pronounced in those at risk for displacement and that the evaluation of HSA may be prudent in children with CP.

Lee et al. [3] analysed HSA, Migration Percentage (MP) and the neck-shaft angle (NSA) in 384 patients with CP. They found a higher correlation between NSA and MP than between HSA and MP and concluded that the NSA appeared to be more clinically relevant than the HSA in evaluating proximal femoral deformity in patients with CP. However, the material included children up to 17 years of age some of whom did not have a well demarcated growth plate, making HSA measurement less reliable. Subsequently, van der List et al. [4] reported from a retrospective study that the HSA decreased over

* Correspondence: gunnar.hagglund@med.lu.se

⁴Lund University Department of Clinical Sciences, Orthopedics, Lund, Sweden

Full list of author information is available at the end of the article






Fig. 1 Measurement of the Head Shaft Angle (HSA)

time in normal hips and children in GMFCS levels II-III but not in GMFCS levels IV-V, the two groups most at risk of developing hip subluxation. In the same year, van der List et al. [5] concluded from a retrospective cohort that, at age two years, the GMFCS and HSA were valuable predictors for hip displacement, but at the age of four years, only the MP should be used in the prediction of hip displacement. These studies seemed to confirm that the HSA was higher in GMFCS IV-V children at least up to the age of two years, but that the effect of age on the HSA as a predictor of hip displacement was less clear.

Hermanson et al. [6] reported on a total population of children with CP in a defined geographical area followed up for five years or until the development of a MP \geq 40% of either hip within five years. The use of MP 40% as cut

off was chosen as hips with MP \geq 40 have a high risk for further displacement, indicating the need for surgical intervention [7]. They concluded that a high HSA was a risk factor for hip displacement in children with CP. However, in 2016 Chougule et al. [8] reported no statistically significant correlation between HSA and hip migration in children with CP aged 3–18 years in GMFCS levels III-V.

These seemingly contradictory reports on the relationship between the HSA, age and GMFCS as a risk factor for hip displacement prompted the present study from another defined population of children with CP. The Cerebral Palsy Integrated Pathway Scotland (CPIPS) surveillance programme includes 95% of all children with CP in Scotland. We have used radiographs from this total population to analyse the effect of HSA on hip displacement defined as MP \geq 40%.

Methods


Radiological data from all children in GMFCS levels III-V was used in the analysis. Children receive a pelvic radiograph on first registration in the programme. This is taken in a standardised position which is used across Scotland [9]. Subsequent radiographs are taken according to a protocol based on age and GMFCS. In this study all radiographs taken at first registration into the programme of children in GMFCS levels III-V were used for analysis to avoid potential effects of surveillance on subsequent hip centration. Radiographs showing femoral osteotomy before inclusion in the programme were excluded. Hips were defined as displaced if the MP was \geq 40%. The most displaced hip from each child was analysed. The HSA for all radiographs were measured by one observer (LF). The MP for the index radiograph had been previously been recorded in the database by clinicians responsible for the individual care of the children.

Two logistic regression models were used with hip displacement (MP \geq 40%) as the outcome. In the first model the following covariates were used: HSA and age (both continuous), GMFCS (categorical, level III as reference), sex (categorical, female as reference). For the second model a ROC curve analysis between HSA and MP \geq 40% was done. The ideal cut-off point was selected as the point that maximizes the product of specificity and sensitivity. HSA was dichotomised according to the ideal cutoff point and in the second model the HSA was analysed as a categorical covariate related to this cutoff value. The other covariates were similar to the first model.

The statistical analyses were performed using STATA 13 software.

Results

There were 640 children with a mean age of 8.2 years (range 0–19 years, 271 female and 369 male) fulfilling the inclusion criteria (Fig. 2). Their distribution by GMFCS



level was as follows: III - 160, IV - 184 and V - 296. One hundred and eighteen hips had a MP $\geq 40\%$ in which the mean HSA was 164° (range $121-180^\circ$). Five hundred and twenty two hips with a MP $< 40\%$ had a mean HSA of 160° (range $111-180^\circ$). The first logistic regression analysis using HSA as a continuous variable showed no influence of age and sex had on MP but a high GMFCS level was strongly associated with hip displacement (Table 1). HSA was also statistically significantly associated with hip displacement. A 10° difference in HSA for patients adjusted for age, sex, and GMFCS gave an odds ratio of 1.26 for hip displacement to a MP $\geq 40\%$ ($p = 0.009$) (Table 1). The ROC curve analysis between HSA and MP $\geq 40\%$ showed that the ideal cutoff point was HSA = $164,5$ degrees. The second logistic regression analysis showed that HSA above $164,5$ degrees had an odds ratio of 1.96 for MP $\geq 40\%$ (Table 2).

Discussion

This study has confirmed Hermanson et al's [6] findings that a high HSA is a risk factor for hip displacement in children with CP. Both studies used defined populations

of children with CP and thus probably avoided selection biases. The present study does not give prospective information on HSA changes over time, unlike Hermanson et al's study, because radiological data was taken from radiographs at the time of first registration of children into the surveillance programme. It does, however, provide information on the HSA without the potentially confounding influence of surveillance on hip centration.

The findings of the present study differ from those of Chougule et al. [8] who reported no statistically significant correlation between HSA and hip migration in children with CP aged 3–18 years in GMFCS levels III-V. There are some differences between the two studies which may explain the respective conclusions. Chougule et al. [8] used a linear regression model in their statistical analysis and randomised hips by laterality in the study design. There is a floor and ceiling effect for MP at 0% and 100% that may limit the usefulness of linear regression analysis which is not seen with logistic regression, where a binary outcome of hip 'displaced' or 'not displaced' was used and set at MP $\geq 40\%$. Randomisation

Table 1 Logistic regression estimates on the effect of HSA, GMFCS-level and age on MP using HAS as continuous variable

	Odds Ratio	95% CI		P
HSA ^a	1.26	1.06	1.50	0.009
GMFCS IV	2.86	1.39	5.90	0.004
GMFCS V	4.17	2.13	8.16	0.000
Age ^b	1.03	0.99	1.08	0.174
Sex ^c	0.94	0.62	1.42	0.763

^adegrees, ^byears, ^cfemale as reference

Table 2 Logistic regression estimates on the effect of HSA, GMFCS-level and age on MP with HSA dichotomized

	Odds Ratio	95% CI		P
HSA ^a	1.96	1.29	3.00	0.002
GMFCS IV	2.78	1.35	5.74	0.006
GMFCS V	3.98	2.03	7.80	0.000
Age ^b	1.03	0.99	1.08	0.134
Sex ^c	0.94	0.62	1.42	0.755

^aHSA < 164,5 degrees as reference ^byears, ^cfemale as reference

by laterality may also dilute a study population of hips at risk for displacement. Children in higher GMFCS levels may have pelvic obliquity and the abducted hip would most probably have a MP within normal limits and the opposite, adducted, hip an abnormal MP. For this reason we analysed the most displaced hip in each child. Van der List et al. [5] used both hips in their analyses and also did a logistic regression analysis for displacement with MP $\geq 40\%$.

The present study does have limitations. A single observer measured the HSA in all 640 hips. Training in the measurement was given and, reassuringly, Hermanson et al. [10] have reported excellent observer reliability for HSA. The present study is based on the radiographs of children at the time of registration into the CPIPS programme and effects of time or intervention on hip centration or HSA were not considered. The results are only valid for the age range and range of HSA-angle in the material.

Conclusion

This study, which was based on a total CP population, has confirmed that HSA is associated with hip displacement in children in GMFCS levels III-V.

Abbreviations

CI: Confidence Interval; CP: Cerebral palsy; GMFCS: Gross Motor Function Classification Scale; HSA: Head Shaft Angle; MP: Migration Percentage; ROC: Receiver operation characteristic

Acknowledgements

We gratefully acknowledge the contributions of all clinicians who have or continue to provide data on their patients for CPIPS.

Funding

No funding was obtained for this study.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

JER, MG and GH conceived the research question, MG the data collection, LF the radiographic measurements, and TC the statistical analyses. JER was the major contributor in writing the manuscript to which all authors also contributed. All authors have read and approved the final manuscript.

Ethics approval and consent to participate

The data reported in this study has not involved any research on human subjects. The data collected on the database is purely observational - there is absolutely no change to their treatment or clinical course as part of having this data recorded. The database is a centrally held clinical patient management record for children with cerebral palsy in Scotland. Institutional ethical approval would not apply to the data base as it is a country-wide, national level programme, under the auspices of NHS Scotland, rather than a single institution. We do however have Scottish national level approval to collect the data in the form of an: "NHS Scotland Caldicott Guardian Application to use Multiple Boards or Scotland-Wide Personal Health Information for Health Research or Audit" document, with Data Protection Registration No: Z6650266 - this document can be supplied if required. With regard to consent, all parents have given consent for their data to be used in this way and this is recorded in writing in their record.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹University of Edinburgh, Edinburgh, Scotland. ²Epidemiology and Register Center South, Lund, Sweden. ³Department of Orthopaedic Surgery, Royal Hospital for Sick Children, Edinburgh, Scotland. ⁴Lund University Department of Clinical Sciences, Orthopedics, Lund, Sweden. ⁵School of Medicine, University of St Andrews, St. Andrews, Scotland.

Received: 10 February 2018 Accepted: 24 September 2018

Published online: 05 October 2018

References

- Southwick WO. Osteotomy of the lesser trochanter for slipped capital femoral epiphysis. *J Bone Joint Surg Am.* 1967;49A:803–35.
- Foroohar A, McCarthy JJ, JJ YD, Clarke S, Brey J. Head-shaft angle measurement in children with cerebral palsy. *J Pediatr Orthop.* 2009;29:248–50.
- Lee KM, Kang JY, Chung CY, Kwon DG, Lee SH, Choi IH, Cho TJ, Yoo WJ, Park MS. Clinical relevance of valgus deformity of proximal femur in cerebral palsy. *J Pediatr Orthop.* 2010;30:720–5.
- van der List JP, Witbreuk MM, Buizer AI, van der Sluijs JA. The head–shaft angle of the hip in early childhood. A comparison of reference values for children with cerebral palsy and normally developing hips. *Bone Joint J.* 2015a;97B:1291–5.
- van der List JP, Witbreuk MM, Buizer AI, van der Sluijs JA. The prognostic value of the head–shaft angle on hip displacement in children with cerebral palsy. *J Child Orthop* 2015b; 9:129–135.
- Hermanson M, Häggglund G, Riad J, Wagner P. Head-shaft angle is a risk factor for hip displacement in children with cerebral palsy. *Acta Orthop.* 2015;86:229–32.
- Häggglund G, Lauge-Pedersen H, Persson M. Radiographic threshold values for hip screening in cerebral palsy. *J Child Orthop.* 2007;1:43–7.
- Chougule S, Dabis J, Petrie A, Daly K, Gelfer Y. Is head–shaft angle a valuable continuous risk factor for hip migration in cerebral palsy? *J Child Orthop.* 2016;10:651–6.
- Kinch K, Campbell DM, Maclean JG, Read HS, Barker SL, Robb JE, Gaston MS. How critical is patient positioning in radiographic assessment of the hip in cerebral palsy when measuring migration percentage? *J Pediatr Orthop.* 2015;35:756–61.
- Hermanson M, Häggglund G, Riad J, Rodby-Bousquest E. Inter- and intrarater reliability of the head–shaft angle in children with cerebral palsy. *J Child Orthop.* 2017;11:256–62.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

