

1 **Detecting green shoots of recovery - the importance of long-term individual**
2 **based monitoring of marine turtles**

3 **K. L. STOKES¹, W. J. FULLER^{1,2,3}, F. GLEN⁴, B. J. GODLEY¹, D. J. HODGSON¹, K. A. RHODES¹, R.**
4 **T. SNAPE^{1,3}, AND A. C. BRODERICK^{1,*}**

5 ¹Centre for Ecology and Conservation, University of Exeter, Cornwall Campus, Penryn TR10 9EZ, UK

6 ²Faculty of Veterinary Medicine, Near East University, Nicosia, North Cyprus, Mersin 10, Turkey

7 ³Society for Protection of Turtles, PK65, Kyrenia, North Cyprus, Mersin 10, Turkey

8 ⁴16 Eshton Terrace, Clitheroe, Lancashire, BB7 1BQ, UK

9 *Author for correspondence (a.c.broderick@exeter.ac.uk)

10

11 **Running head:** green turtle population assessment

12 **Word count:** 4373

1 **Abstract**

2 Population monitoring is an essential part of evaluating the effectiveness of management
3 interventions for conservation. Coastal breeding aggregations of marine vertebrate species that come
4 ashore to pup or nest provide an opportunistic window of observation into otherwise widely dispersed
5 populations. Green turtle (*Chelonia mydas*) nesting on the north and west coasts of northern Cyprus
6 has been monitored consistently and exhaustively since 1993, with an intensive saturation tagging
7 programme running at one key site for the same duration. This historically depleted nesting
8 population is showing signs of recovery, possibly in response to nest protection approaching two
9 decades, with increasing nest numbers and rising levels of recruitment. Strong correlation between
10 year to year magnitude of nesting and the proportion of new breeders in the nesting cohort implies
11 that recruitment of new individuals to the breeding population is an important driver of this recovery
12 trend. Recent changes in fishing activities may be impacting the local juvenile neritic stage, however,
13 which may hinder this potential recovery. Individuals returning to breed after two years laid fewer
14 clutches than those returning after three or four years, demonstrating a trade-off between
15 remigration interval and breeding output. Average clutch frequencies have remained stable around a
16 median of three clutches a year per female despite the demographic shift towards new nesters, which
17 typically lay fewer clutches in their first season. We show that where local fecundity has been
18 adequately assessed, the use of average clutch frequencies can be a reliable method for deriving
19 nester abundance from nest counts. Index sites where individual based monitoring is possible will be
20 important in monitoring long-term climate driven changes in reproductive rates.

21

22 **Keywords:** population monitoring, clutch frequency, neophyte, remigration interval, green turtle,
23 trade-off

24

1 Introduction

2 Population monitoring is integral to conservation biology (Goldsmith 1991), and forms an essential
3 part of evaluating the effectiveness of active conservation management (Nichols & Williams 2006).
4 Present-day conservation monitoring must not only endeavour to detect changes in population status,
5 but also climate change driven alterations to reproductive rates, developmental biology (Milligan et
6 al. 2009) and spatio-temporal displacements (Parmesan & Yohe 2003). For many marine, nocturnal
7 or otherwise cryptic species, detection poses additional challenges, and direct monitoring may be
8 difficult, impractical or impossible. Various indirect survey methods are used as indices of abundance,
9 such as redd (nest) counts for salmonids (eg. Rieman & Myers 1997), egg-mass counts for pond
10 breeding amphibians (eg. Raithel et al. 2011), acoustic monitoring for loquacious species (eg. anurans,
11 Crouch & Paton 2002; whales, Simard et al. 2010), and camera trapping, live trapping, hair detection
12 and road casualty data for terrestrial mammals (eg. George et al. 2011, Swan et al. 2013). For marine
13 vertebrates, breeding aggregations are often monitored as an index of overall population status (eg.
14 whales, Andriolo et al. 2010, Fretwell et al. 2014), and species that come ashore to pup or nest present
15 a logistical opportunity to count individuals with greater accuracy and much reduced cost (e.g. sea
16 lions, Pitcher et al. 2007).

17 The vast majority of marine turtle monitoring research is based at nesting beaches. The accessibility
18 of females during this narrow window has made nester abundance a common response variable for
19 sea turtle population trend monitoring (Heppell et al. 2003). Population assessments based solely on
20 abundance of nesting females have drawn criticism (Bjorndal et al. 2010) and should ideally be
21 combined with in-water foraging ground surveys, which are expensive and labour-intensive (Seminoff
22 et al. 2003). Large discrepancies exist in levels of available funding, however, and nesting beach
23 studies are often the only feasible approach to implement monitoring over long time frames at low
24 expense (Meylan 1995; Gerrodette & Taylor 1999).

1 Studies of sea turtle reproductive ecology rely heavily on the practice of tagging individuals to
2 elucidate breeding frequency and fidelity to nesting areas (Balazs 1999). In the past, tag loss has been
3 a major confounding variable, with reports of 78% documented tag loss and upper retention estimates
4 of just six years in early studies (Mortimer & Carr 1987). Tag retention rates have since been enhanced
5 with improved tag design and the introduction of PIT (Passive Integrated Transponder) tags. These
6 developments have increased the accuracy of neophyte/ remigrant classification, reducing
7 uncertainty in the quantification of neophyte turtles and overall nester abundance (McDonald &
8 Dutton 1996).

9 Most marine turtle populations display obligate skipped breeding behaviour due to the high energy
10 demands of migration and reproduction (Prince & Chaloupka 2012), females laying a variable number
11 of clutches within a breeding season (termed clutch frequency) every few years (the remigration
12 interval). Individuals must attain a threshold body condition before embarking on a breeding
13 migration, and so their remigration interval varies in response to fluctuations in environmental
14 conditions (Solow et al. 2002). The low trophic status of the green turtle (*Chelonia mydas*) makes it
15 particularly susceptible to environmental stochasticity, driving large inter-annual oscillations in
16 numbers of nesting females (Limpus & Nicholls 1988; Broderick et al. 2001). The intrinsic variability
17 characteristic of green turtle nesting makes longevity in monitoring programmes essential for
18 identifying underlying population trends (Broderick et al. 2003; Heppell et al. 2003; Jackson et al.
19 2008). Individual plasticity and inter-population variation in clutch frequency add further uncertainty
20 when deriving nesting population estimates from nest abundance counts (Van Buskirk & Crowder
21 1994; Rivalan et al. 2006). Many studies divide nest counts by an average value of clutch frequency
22 to give estimated annual nester abundance or vice versa (e.g. Seminoff 2004; Troeng & Rankin 2005;
23 Beggs et al. 2007). A simplistic model of stochastic nesting behaviour, applied to a loggerhead turtle
24 nest count series with known nester abundance, indicated that this method has the potential to
25 produce biased estimates of population trends (Mazaris et al. 2008). Thus more studies of individually
26 marked populations are needed.

1 Green turtles in the Mediterranean have a history of severe exploitation (Sella 1995). Contemporary
2 rookeries of modest size remain at a handful of sites in Turkey, Cyprus, Syria and Israel (Kasperek et
3 al. 2001; Broderick et al. 2002; Canbolat 2004; Yalcin-Ozdilek 2007; Rees et al. 2008), with ca. 30% of
4 Mediterranean nesting in Cyprus. Modern threats in the Mediterranean include fisheries bycatch and
5 mass tourism (Casale & Margaritoulis 2010); this population has been highlighted as a conservation
6 priority owing to its 'High Risk-High Threat' status (Wallace et al. 2011).

7 Since 1993, an extensive monitoring programme has conducted comprehensive surveys of the nesting
8 beaches of the north and west coasts of northern Cyprus, located in the Eastern Mediterranean (for
9 beach locations see Fig. 1). Intensive survey effort has been concentrated at Alagadi, where continual
10 night patrols of this 2 km stretch of beach for the duration of the breeding season have allowed
11 exhaustive tagging. Here, we examine the apparent recovery of the population, and reveal the range
12 of insights that long-term individual based monitoring can provide.

13

14 **Materials and methods**

15 Daytime monitoring of marine turtle nesting activity was conducted every 1-3 days on beaches with
16 significant nesting on the north and west coasts of northern Cyprus for the duration of the breeding
17 season (end of May to end of September) each year between 1993 and 2013 (less complete
18 monitoring was undertaken in 1992; see Fig. 1 for beach locations). Daytime monitoring involves
19 thorough examination of all nesting activity during the early morning, location of eggs if present, and
20 protection from depredation by stray dogs and foxes using a wide mesh wire screen secured into the
21 sand above the nest (carried out exhaustively since 1994). An intensive night monitoring and tagging
22 programme has been conducted at Alagadi (comprising two coves 1.2 and 0.8 km in length) over the
23 same time period (see Broderick et al. 2002, 2003 for detailed methods). Patrols are undertaken at
24 sufficient frequency to encounter all females nesting at this beach. Internal PIT tags have been
25 administered in addition to external flipper tags to all turtles nesting at this breeding site since 1997.

1 Neophyte/ remigrant analyses were conducted on a subset of the data from 2000 onwards due to
2 increased accuracy of neophyte classification three years (one full nesting cycle for most females)
3 following the introduction of PIT tagging.

4 Long intervals between observed nesting events within a nesting season are indicative that a female
5 has laid elsewhere on a nearby beach. Thus, the number of clutches laid per season at Alagadi by each
6 marked individual (observed clutch frequency, OCF) is adjusted where turtles have interesting
7 intervals of 20 days and over to give the expected clutch frequency (ECF; Frazer & Richardson 1985,
8 see also Broderick et al. 2002 for bimodal distribution of interesting interval data). The remigration
9 interval (RI) for remigrant turtles is calculated as the number of years since that individual was last
10 recorded nesting at Alagadi.

11 Statistical tests and modelling were carried out using R version 2.14.2 (R Development Core Team
12 2012), and packages “nlme” (Pinheiro et al. 2012) and “lme4” (Bates et al. 2011). Tests of correlation
13 were performed using Spearman’s rank order correlation coefficient. LOESS (locally weighted)
14 regressions were fitted to RI and ECF time series data with degree one (linear) and a span of 0.75.
15 Time series analyses of yearly nest counts were conducted using generalized least squares (GLS)
16 modelling to account for temporal autocorrelation in the data.

17 Clutch frequencies were regressed against explanatory variables using generalized linear mixed
18 modelling (GLMM), fitted using the Laplace approximation, restricted maximum likelihood estimates
19 (REML) and stepwise model simplification. GLMMs allow statistical analysis of non-normal data with
20 random effects, which quantify the variation across units/ grouping factors of the fixed effect
21 parameters (Bolker et al. 2009). In this case, models had Poisson error structure and logarithmic link
22 function, with zero-truncation. Explanatory variables included categorical fixed effects for neophytes
23 (first time nesters; true or false) and remigration interval (two vs. three or four years), a fixed covariate
24 of body size, and random effects for individual (to avoid pseudoreplication where females have
25 returned to nest in subsequent years) and year (to account for interannual variation in magnitude of

1 nesting arising from environmental stochasticity). GLMM was also used to regress body size against
2 neophyte/ remigrant nesters whilst accounting for pseudoreplication of individuals. The significance
3 of removing model terms was assessed by likelihood ratio tests using maximum likelihood estimates
4 (Crawley 2007), in order of least significance and with a threshold of $p=0.05$. Model residuals were
5 checked for overdispersion, normality and homoscedasticity.

6

7 **Results**

8 The annual green turtle nesting abundance for Alagadi and the total across the north and west coasts
9 is shown in Figure 2. The high interannual variation typical of green turtle nesting is evident (combined
10 nesting range: 35-335 nests per season, mean \pm standard deviation (SD): 13 ± 77.1), following a two
11 to three year pseudo-cyclical pattern. The coefficient of variation ($CV=SD/mean$: 0.59) lies within the
12 range previously reported for this species by Broderick et al. (2001; 0.41-1.08). Nesting abundance on
13 the two coasts is significantly correlated ($r_{(19)}=0.72$, $p<0.001$) showing a synchrony in reproductive
14 cycles across this area. Comparison of nest count models at Alagadi and across the two coasts
15 demonstrated significant autocorrelation at a time lag of one year (GLS, Alagadi: $\phi=-0.729$,
16 $\chi^2_{(1)}=11.274$, $p<0.001$; overall: $\phi=-0.449$, $\chi^2_{(1)}=4.224$, $p=0.04$). Having accounted for this
17 autocorrelation, nest counts showed a significant quadratic trajectory through time (Fig. 2; Alagadi
18 linear slope: $\beta=-10.663 \pm 2.790$; Alagadi quadratic slope: $\beta=0.709 \pm 0.135$, $\chi^2_{(1)}=17.471$, $p<0.0001$;
19 overall linear slope: $\beta=-20.878 \pm 5.405$; overall quadratic slope: $\beta=1.259 \pm 4.817$, $\chi^2_{(1)}=14.379$,
20 $p=0.0001$). This indicates that nesting in the region has stabilised and may now be increasing. This
21 trend was also significant for nester abundance at Alagadi (Fig. 3a; autocorrelation at one year: $\phi=-$
22 0.743 , $\chi^2_{(1)}=10.711$, $p=0.001$; linear slope: $\beta=-3.151 \pm 0.929$; quadratic slope: $\beta=0.231 \pm 0.045$,
23 $\chi^2_{(1)}=17.079$, $p<0.0001$). Recruitment (as measured by the proportion of nesters that are neophytes)
24 has followed a similar quadratic trend (linear slope: $\beta=-0.103 \pm 0.016$; quadratic slope: $\beta=0.005 \pm$
25 0.001 , $\chi^2_{(1)}=22.005$, $p<0.0001$), but with no significant autocorrelation. Record numbers of nests,
26 nesters and neophytes were observed at Alagadi in 2013 (236 nests, 85 nesting females, 57

1 neophytes). There has been no trend in survey effort, detection probability (imperfect detection of
2 nests or individuals), or detectability (beach fidelity) over the study period (see Supplementary Figure
3 S1 and Pfaller et al. 2013).

4 We confirmed that the recent trajectory describes a significant increase in nests, nesters and
5 recruitment, by considering the number of nests and nesters post-2000, which corresponds with the
6 local minimum of all our quadratic fitted lines. Since 2001, there has been a significant increase
7 through time in the number of nests across all beaches ($\beta=15.993 \pm 3.063$, $\chi^2_{(1)}=11.938$, $p=0.0006$), the
8 number of nests on Alagadi ($\beta=9.799 \pm 1.605$, $\chi^2_{(1)}=15.516$, $p=0.0001$), the number of nesting females
9 on Alagadi ($\beta=3.493 \pm 0.606$, $\chi^2_{(1)}=14.398$, $p=0.0001$) and rates of recruitment ($\beta=0.029 \pm 0.009$,
10 $\chi^2_{(1)}=8.399$, $p=0.004$).

11 A comparison of observed and estimated nester abundance is shown in Figure 3a. Here, the known
12 number of females nesting each year at Alagadi is used to test the accuracy of estimates derived using
13 nest counts and average values of clutch frequency. Estimated nester abundance is taken as the
14 quotient of annual nest abundance divided by an average clutch frequency of three (Seminoff 2004;
15 also the overall mean and median clutch frequency from the current study). Estimated nester
16 abundance and actual/ observed nester abundance were highly correlated ($r_{(19)}=0.97$, $p<0.0001$).
17 Conclusions drawn from these abundance series about the population trend at this breeding
18 aggregation would be analogous.

19 The tagging programme based at Alagadi has revealed a strong correlation between the number of
20 nests and the proportion of neophytes since 2000 ($r_{(12)}=0.94$, $p<0.0001$; Fig. 3b). This strong
21 correlation between the proportion of neophytes in the nesting cohort and the magnitude of nesting
22 implies that recruitment of new individuals into the breeding population is an important driver of year
23 to year nester abundance, an encouraging sign of a population in recovery. The reduced correlation
24 between the number of nesters and the proportion of neophytes seen prior to 2000 provides evidence
25 that the introduction of PIT tagging has had a significant effect on the accuracy of neophyte/ remigrant

1 identification. First time nesters at Alagadi are significantly smaller than remigrant nesters (GLMM, χ^2
2 $(1)=84.95$, $p<0.0001$; mean CCL 87.7 ± 6.5 cm for neophytes cf. 92.0 ± 5.9 cm for remigrants), re-
3 affirming their classification as true neophytes. The intensity of survey effort at this site has afforded
4 near perfect attribution of nests to known females (98% since 2000, 93% since comprehensive
5 monitoring began in 1993).

6 Figure 4a shows RIs observed for the marked green turtle population at Alagadi between 1994 and
7 2013. The majority of remigrants return after 2, 3 or 4 years [87%; median RI: 3, interquartile (IQ)
8 range: 3-4, $n=212$]. The low incidence of unusually long RIs most likely reflects individuals with lower
9 site fidelity, who may have nested elsewhere in Cyprus, or further afield, undetected. The majority
10 (78%) of remigrant turtles observed over three or more seasons varied their RI from one breeding
11 season to the next ($n=51$, see Fig. 4b), exemplifying the high levels of modulated periodicity green
12 turtles show in response to environmental stochasticity. Despite this, the annual average RI has
13 remained relatively stable over the study period (see Fig. 5a), fluctuating mostly between three and
14 four years. Lower RIs at the beginning of the time series are an artefact of time since tagging began;
15 only those remigrants with lower than average remigration interval can be re-encountered within the
16 first three years of monitoring.

17 Median ECF across all years and nesters was three (IQ range: 1-4, $n=485$). No long-term trend in
18 median clutch frequency is apparent from the data (see LOESS smoother Fig. 5b). Instead, median
19 ECF is correlated with the number of nesters present in a given season ($r_{(19)}=0.52$, $p=0.02$), with three
20 of the four lowest nesting seasons having a low average ECF, indicating that females breeding in poor
21 nesting years may be in suboptimal body condition. Median clutch frequency is more variable in the
22 early part of the time series, stabilising as the number of females increases, effectively increasing the
23 sample size and reducing susceptibility to skewed averages. ECF varies between neophytes and
24 remigrants (see Fig. 6a-b), with neophytes most likely to lay a single clutch (40%, $n=194$), and the
25 majority of remigrants laying three to five clutches (77%, $n=212$). GLMM showed the effect to be

1 significant, with remigrants laying an average of 0.6 clutches more than neophytes, whilst accounting
2 for individual and year to year variation ($\chi^2_{(1)}=37.198$, $p<0.0001$). Female body size had a statistically
3 significant but biologically insignificant effect on clutch frequency ($\chi^2_{(1)}=7.689$, $p<0.01$), with a 10 cm
4 increase in curved carapace length (CCL) increasing ECF by an average of 0.04.

5 RI was found to have a significant effect on clutch frequency, with short RIs of less than three years
6 reducing ECF by almost a quarter (0.23), once variation across individuals and years was accounted
7 for (GLMM, $\chi^2_{(1)}=4.009$, $p<0.01$). Body size did not have significant effect to be included in the model.
8 Female nesters returning after a short interval of two years are most likely to lay three clutches (40%,
9 $n=48$, Fig. 6c), whilst those returning after three or four years are more likely to lay four or five clutches
10 (57%, $n=136$, Fig. 6d-e).

11 A total of 273 nesting females have been tagged at Alagadi since 1992. Forty percent of neophytes
12 nesting between 2000 and 2008 ($n=55$) did not remigrate to this site in subsequent breeding seasons
13 (we do not include 2009-2013 as these neophytes may yet return).

14

15 **Discussion**

16 Evaluation of indirect survey method reliability is essential for accurate population monitoring.
17 Validated indices are the primary tool for tracking changes in abundance of many cryptic species of
18 conservation concern (eg. carnivore track counts and camera trap surveys, Balme et al. 2009). Long-
19 term individual based monitoring of green turtles at Alagadi, northern Cyprus has provided
20 fundamental and applied insights into sea turtle nesting ecology. Our data suggest that estimation of
21 nesting population size from nest abundance data is reliable, provided that fecundity is adequately
22 monitored at relevant localised index sites to provide the 'proportionality' information required to
23 interpret these data (Gerrodette & Taylor 1999). Green turtles nesting at Ascension Island in the South
24 Atlantic are larger in size, migrate further (~2300 km, Luschi et al. 1998) and have a longer period of
25 suitable nesting conditions than those nesting in the Mediterranean, and thus perhaps unsurprisingly

1 have a higher average clutch frequency of around 6 nests per season (Weber et al. 2013) compared
2 to the average of 3 detected in the current study. Clutch frequencies derived through tagging efforts
3 alone where complete survey is not possible or site fidelity is low will be underestimated, leading to
4 inflated population assessments. Studies augmenting capture-mark-recapture methods with the use
5 of tracking (Tucker 2010, Weber et al. 2013) and ultrasonography (Blanco et al. 2012) technologies
6 can improve clutch frequency estimates in such cases. Breeding rates will likely be affected by long-
7 term changes in foraging conditions, highlighting the importance of ongoing monitoring at index sites
8 to ascertain multifaceted responses to climate change.

9 Saturation tagging at Alagadi has revealed clutch frequencies that are significantly different among
10 groups (eg. neophytes vs. remigrants), but that are temporally stable across groups. Reduced clutch
11 frequency in neophyte turtles as seen here has previously been reported in green turtles (Carr et al.
12 1978), as well as in leatherback (Tucker & Frazer 1991), loggerhead (Hawkes et al. 2005) and hawksbill
13 turtles (Beggs et al. 2007). It is likely that this phenomenon is caused by both increasing physiological
14 capacity with age, and changes in nesting behaviour such as site fidelity (Carr et al. 1978). Individual
15 green turtles lay increasingly large clutches across (Bjorndal & Carr 1989) and within (Broderick et al.
16 2003) breeding seasons, indicating an increase in reproductive efficiency or capacity. Low subsequent
17 remigration rates of neophytes tagged at Alagadi (this study) suggests lowered site fidelity in new
18 breeders. Broderick et al. (2002) found that single-clutch neophyte females have a lower probability
19 of remigrating to Alagadi in subsequent years than those with higher clutch frequencies (0.3 cf 0.8).
20 Satellite telemetry of internesting loggerhead turtles in Florida has revealed a higher site fidelity in
21 remigrants compared to new breeders (Tucker 2010). Such 'leaky' female nest site fidelity facilitates
22 genetic mixing of the maternal lineage across nesting sites (Lee et al. 2007), and may promote
23 resilience to loss of breeding sites through behavioural adaptation.

24 Our finding that females remigrating after three or four years lay extra clutches in comparison to those
25 remigrating after two years supports the notion that suboptimal foraging conditions can be

1 compensated for by building up energy reserves over a longer interval. A similar relationship between
2 remigration interval and likely clutch frequency has been observed in leatherback turtles (Rivalan et
3 al. 2005), and Van Buskirk and Crowder (1994) describe a comparable trade-off in interspecific
4 reproductive effort resource allotment. Many iteroparous species may skip a breeding year if
5 conditions are not favourable (eg. fat dormouse, Pilastro et al. 2003); this may partly be compensated
6 for if a higher reproductive output can be attained in the following breeding season (eg. four-toed
7 salamander, Harris & Ludwig 2004). The implications for population assessment are that short term
8 fluctuations in breeding activity may be misinterpreted unless populations are monitored in the long
9 term (Hays 2000), and that breeding frequency should be monitored at the individual level where
10 possible in order to detect long term change in the scaling factors used for conversion of monitoring
11 indices to population estimates.

12 The recent upward trend in nest numbers in northern Cyprus may signal the beginning of a recovery
13 phase for this sub-population following the cessation of a heavy harvest and intensive screening of
14 nests against unnaturally elevated predation levels. Recruitment can be viewed as a measure of
15 cohort strength (Heppell et al. 2003), and rising numbers of neophytes as seen in this population are
16 an early indication of population growth (Richardson et al. 2006). Similar nest protection schemes
17 have had measurable success some 20 years later (Garduño-Andrade et al. 1999; Dutton et al. 2005).
18 There is, however, considerable uncertainty surrounding the time it takes for green turtles to reach
19 breeding age; published age at sexual maturity estimates for wild green turtles range from 27 (Frazer
20 & Ladner 1986) to 40 (Limpus & Chaloupka 1997) years. Evidence from living tags, however, has
21 shown that male and female green turtles released from the Cayman Island Turtle Farm (a
22 conservation facility/ tourist attraction/ turtle meat supplier in the Caribbean) as hatchlings can breed
23 at 19 and 17 years respectively (Bell et al. 2005). If this species can indeed reach sexual maturity at
24 less than 20 years, then it is possible that sustained reduction in nest depredation across two decades
25 has aided in the early stages of recovery of this historically depleted breeding aggregation.

1 Behavioural reproductive mechanisms such as natal philopatry and polyandry contribute to the
2 resilience of sea turtles (Bell et al. 2009), which have shown encouraging recovery potential and
3 rebound capacity in response to long-term protection (Garduño-Andrade et al. 1999; Broderick et al.
4 2006; Richardson et al. 2006; Marcovaldi & Chaloupka 2007). The complex life cycle of this group, and
5 others involving multiple distinct habitats and delayed sexual maturity, makes adequate protection
6 particularly challenging (Heppell et al. 2003); protective measures on the nesting beach will not be
7 effective if threats at sea are not addressed (Dutton et al. 2005), and the potential for trophic
8 uncoupling of resources and ability to shift ranges under changing climatic conditions are important
9 considerations for such species (Møller et al. 2008; Robinson et al. 2009).

10 Nest count series should be used in conjunction with data regarding other life stages wherever
11 possible (Bjorndal et al. 2010). Encouragingly, genetic studies at Alagadi have revealed a greater
12 number of males than females in the breeding population (Wright et al. 2012a; Wright et al. 2012b),
13 suggesting that the population increase observed here has occurred across demographic groups.
14 However, a recent assessment of sea turtle bycatch in northern Cyprus (Snape et al. 2013) has found
15 a high incidence of juvenile green turtle mortality. Potential increased fishing effort in the region
16 following changes in trade regulation between northern Cyprus and southern Cyprus (Snape *pers.*
17 *obs.*) may impede the recovery of this population.

18 Monitoring projects must be cost-effective in the long-term in order to ensure the longevity of data
19 required to make meaningful estimations of population trends (Schroeder & Murphy 1999). Re-
20 sampling assessments of extant data from comprehensively monitored nesting sites have found that
21 temporal sub-sampling within the breeding season could save up to 50% of monitoring costs with
22 little loss of statistical power (Jackson et al. 2008; Sims et al. 2008, Whiting et al. 2013). The efficacy
23 of these more parsimonious sampling regimens is reliant on consistency in the temporal distribution
24 of the nesting season, however, which has been shown to be variable in accordance with both long-
25 (Weishampel et al. 2004) and short- (Hawkes et al. 2007) term fluctuations in sea surface temperature.

1 Furthermore, complete sampling of the breeding season yields additional advantages in localities
2 where remedial conservation measures such as nest protection and surveillance of illegal take are
3 beneficial (eg. Bell et al. 2007).

4 A range of strategies is required to cover the breadth and depth necessary to detect changes in
5 biological parameters and spatio-temporal distributions that are likely to occur in response to climate
6 change. Index sites such as Alagadi, where long-term and consistent individual based monitoring is
7 possible, can offer valuable insights into survival and reproductive rates that other localities can use
8 in converting more basic density indices into population estimates. Long-term datasets are vital in
9 documenting change, but are often difficult to maintain with variable funding stability through time
10 (Hays et al. 2005). Monitoring programmes with a core set of simple, robust and inexpensive
11 measurements may have a greater likelihood of remaining consistent and sustainable in the long-term
12 (Bennun 2001; Lovett et al. 2007).

13

14 **Acknowledgments**

15 K.L.S. is funded by a European Social Fund doctoral studentship. We thank the volunteers and funders
16 of the Marine Turtle Conservation Project in North Cyprus, including British Chelonia Group, the British
17 High Commission, the British Residents Society, Erwin Warth Foundation, Friends of SPOT, Kuzey Kibris
18 Turkcell and MEDASSET UK. We also thank project partners Society for the Protection of Turtles in
19 North Cyprus and the Department of Environmental Protection.

1 **References**

- 2 Andriolo, A., P. Kinas, M. Engel, C. Albuquerque Martins, and A. Rufino. (2010). Humpback whales
3 within the Brazilian breeding ground: distribution and population size estimate. *Endangered*
4 *Species Research* 11, 233–243.
- 5 Balazs, G. H. (1999). Factors to Consider in the Tagging of Sea Turtles. In *Research and Management*
6 *Techniques for the Conservation of Sea Turtles No. 4*: 101–109. Eckert, K. L., Bjorndal, K. A.,
7 Abreu-Grobois, F. A., Donnelly, M. (Ed.). Washington, DC: IUCN/SSC Marine Turtle Specialist
8 Group.
- 9 Balme, G. A., L. T. B. Hunter, and R. Slotow. (2009). Evaluating methods for counting cryptic carnivores.
10 *The Journal of Wildlife Management* 73, 433-441.
- 11 Bates, D., M. Maechler, and B. Bolker. (2011). *lme4: Linear mixed-effects models using S4 classes*. R
12 package version 0.999375-42.
- 13 Beggs, J. A., J. A. Horrocks, and B. H. Krueger. (2007). Increase in hawksbill sea turtle *Eretmochelys*
14 *imbricata* nesting in Barbados, West Indies. *Endangered Species Research* 3, 159–168.
- 15 Bell, C. D., J. M. Blumenthal, A. C. Broderick, and B. J. Godley. (2009). Investigating potential for
16 depensation in marine turtles: how low can you go? *Conserv. Biol.* 24, 226–235.
- 17 Bell, C. D. L., J. Parsons, T. J. Austin, A. C. Broderick, G. Ebanks-Petrie, and B. J. Godley. (2005). Some
18 of them came home: the Cayman Turtle Farm headstarting project for the green turtle *Chelonia*
19 *mydas*. *Oryx* 39, 137–148.
- 20 Bell, C. D., J. L. Solomon, J. M. Blumenthal, T. J. Austin, G. Ebanks-Petrie, A. C. Broderick, and B. J.
21 Godley. (2007). Monitoring and conservation of critically reduced marine turtle nesting
22 populations: lessons from the Cayman Islands. *Animal Conservation* 10, 39–47.

- 1 Bennun, L. A. (2001). Long-term monitoring and the conservation of tropical wetlands: high ideals and
2 harsh realities. *Hydrobiologia* 458, 9–19.
- 3 Bjorndal, K. A., B. W. Bowen, M. Chaloupka, L. B. Crowder, S. S. Heppell, and E. Al. (2010). *Assessment*
4 *of Sea-Turtle Status and Trends: Integrating Demography and Abundance*. Washington, DC:
5 National Academies Press.
- 6 Bjorndal, K. A., and A. Carr. (1989). Variation in clutch size and egg size in the green turtle nesting
7 population at Tortuguero, Costa Rica. *Herpetologica* 45, 181–189.
- 8 Blanco, G. S., S. J. Morreale, E. Vélez, R. Piedra, W. M. Montes, F. V. Paladino, and J. R. Spotila. (2012).
9 Reproductive output and ultrasonography of an endangered population of East Pacific green
10 turtles. *The Journal of Wildlife Management* 76.4, 841-846.
- 11 Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.-S. S. White.
12 (2009). Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in*
13 *Ecology & Evolution* 24, 127–35.
- 14 Broderick, A. C., R. Frauenstein, F. Glen, G. C. Hays, A. L. Jackson, T. Pelembe, G. D. Ruxton, and B. J.
15 Godley. (2006). Are green turtles globally endangered? *Global Ecology and Biogeography* 15, 21–
16 26.
- 17 Broderick, A. C., F. Glen, B. J. Godley, and G. C. Hays. (2002). Estimating the number of green and
18 loggerhead turtles nesting annually in the Mediterranean. *Oryx* 36, 227-235.
- 19 Broderick, A. C., F. Glen, B. J. Godley, and G. C. Hays. (2003). Variation in reproductive output of marine
20 turtles. *Journal of Experimental Marine Biology and Ecology* 288, 95–109.
- 21 Broderick, A. C., B. J. Godley, and G. C. Hays. (2001). Trophic status drives interannual variability in
22 nesting numbers of marine turtles. *Proc. R. Soc. Lond. B* 268, 1481–1487.

- 1 Canbolat, A. F. (2004). A review of sea turtle nesting activity along the Mediterranean coast of Turkey.
2 *Biological Conservation* 116, 81–91.
- 3 Carr, A., M. H. Carr, and A. B. Meylan. (1978). The ecology and migrations of sea turtles, 7. The west
4 Caribbean green turtle colony. *Bulletin of the American Museum of Natural History* 162, 1–46.
- 5 Casale, P., and D. Margaritoulis. (2010). *Sea turtles in the Mediterranean: Distribution, threats and*
6 *conservation priorities*. Gland, Switzerland: IUCN.
- 7 Crawley, M. J. (2007). *The R Book*. Chichester: Wiley Publishing.
- 8 Crouch, W. B., and P. W. C. Paton. (2002). Assessing the use of call surveys to monitor breeding
9 anurans in Rhode Island. *Journal of Herpetology* 36, 185-192.
- 10 Dutton, D. L., P. H. Dutton, M. Chaloupka, and R. H. Boulon. (2005). Increase of a Caribbean
11 leatherback turtle *Dermochelys coriacea* nesting population linked to long-term nest protection.
12 *Biological Conservation* 126, 186–194.
- 13 Frazer, N. B., and R. C. Ladner. (1986). A Growth Curve for Green Sea Turtles, *Chelonia mydas*, in the
14 U. S. Virgin Islands. *Copeia* 1986, 798–802.
- 15 Frazer, N. B., and J. I. Richardson. (1985). Annual Variation in clutch size and frequency for loggerhead
16 turtles, *Caretta caretta*, nesting at Little Cumberland Island, Georgia, USA. *Herpetologica* 41,
17 246–251.
- 18 Fretwell, P. T., I. J. Staniland, and J. Forcada. (2014). Whales from space: Counting southern right
19 whales by satellite. *PLOS ONE* 9, e88655.
- 20 Garduño-Andrade, M., V. Guzmán, E. Miranda, R. Briseño-Dueñas, and F. A. Abreu-Grobois. (1999).
21 Increases in hawksbill turtle (*Eretmochelys imbricata*) nestings in the Yucatán Peninsula, Mexico,

- 1 1977 – 1996: Data in support of successful conservation? *Chelonian Conservation and Biology* 3,
2 286–295.
- 3 George, L., J. L. Macpherson, Z. Balmforth, and P. W. Bright. (2011). Using the dead to monitor the
4 living: Can road kill counts detect trends in mammal abundance? *Applied Ecology and*
5 *Environmental Research* 9, 27-41.
- 6 Gerrodette, T., and B. L. Taylor. (1999). Estimating Population Size. In *Research and Management*
7 *Techniques for the Conservation of Sea Turtles No. 4*: 67-71. Eckert, K. L., Bjorndal, K. A., Abreu-
8 grobois, F. A., Donnelly, M. (Ed.). Washington, DC: IUCN/SSC Marine Turtle Specialist Group.
- 9 Goldsmith, B. (1991). *Monitoring for conservation and ecology*. London: Chapman & Hall Ltd.
- 10 Harris, R. N., and P. M. Ludwig. (2004). Resource level and reproductive frequency in female four-toed
11 salamanders, *Hemidactylium scutatum*. *Ecology* 85, 1585-1590.
- 12 Hawkes, L. A., A. C. Broderick, M. S. Coyne, M. H. Godfrey, and B. J. Godley. (2007). Only some like it
13 hot - quantifying the environmental niche of the loggerhead sea turtle. *Diversity and*
14 *Distributions* 13, 447–457.
- 15 Hawkes, L. A., A. C. Broderick, M. H. Godfrey, and B. J. Godley. (2005). Status of nesting loggerhead
16 turtles *Caretta caretta* at Bald Head Island (North Carolina, USA) after 24 years of intensive
17 monitoring and conservation. *Oryx* 39, 65–72.
- 18 Hays, G. C. (2000). The implications of variable remigration intervals for the assessment of population
19 size in marine turtles. *Journal of Theoretical Biology* 206, 221–7.
- 20 Hays, G. C., A. J. Richardson, and C. Robinson. (2005). Climate change and marine plankton. *Trends in*
21 *Ecology & Evolution* 20, 337–44.

- 1 Heppell, S. S., M. L. Snover, and L. B. Crowder. (2003). Sea Turtle Population Ecology. In *Biology of Sea*
2 *Turtles II*: 279–306. Lutz, P. L., Musick, J., Wyneken, J., (Ed.). Boca Raton, Florida: CRC Press.
- 3 Jackson, A., A. Broderick, W. Fuller, F. Glen, G. Ruxton, and B. Godley. (2008). Sampling design and its
4 effect on population monitoring: How much monitoring do turtles really need? *Biological*
5 *Conservation* 141, 2932–2941.
- 6 Kasperek, M., B. J. Godley, and A. C. Broderick. (2001). Nesting of the green turtle, *Chelonia mydas*, in
7 the Mediterranean: a review of status and conservation needs. *Zoology in the Middle East* 24,
8 45–74.
- 9 Lee, P. L. M., P. Luschi, and G. C. Hays. (2007). Detecting female precise natal philopatry in green turtles
10 using assignment methods. *Molecular Ecology* 16, 61–74.
- 11 Limpus, C., and M. Chaloupka. (1997). Nonparametric regression modelling of green sea turtle growth
12 rates (southern Great Barrier Reef). *Marine Ecology Progress Series* 149, 23–34.
- 13 Limpus, C., and N. Nicholls (1988). The Southern Oscillation regulates the annual numbers of green
14 turtles (*Chelonia mydas*) breeding around Northern Australia. *Aust. J. Wildl. Res.* 15, 157-61.
- 15 Lovett, G. M., D. A. Burns, C. T. Driscoll, J. C. Jenkins, M. J. Mitchell, L. Rustad, J. B. Shanley, G. E. Likens,
16 and R. Haeuber. (2007). Who needs environmental monitoring? *Frontiers in Ecology and the*
17 *Environment* 5, 253–260.
- 18 Luschi, P., G. C. Hays, C. Del Seppia, R. Marsh, and F. Papi (1998). The navigational feats of green sea
19 turtles migrating from Ascension Island investigated by satellite telemetry. *Proc. R Soc. Lond. B*
20 265:2279-2284.
- 21 Marcovaldi, M. Â., and M. Chaloupka. (2007). Conservation status of the loggerhead sea turtle in
22 Brazil: an encouraging outlook. *Endangered Species Research* 3, 133–143.

- 1 Mazaris, A., Y. Matsinos, and J. Pantis. (2008). Evaluating the effect of varying clutch frequency in
2 nesting trend estimation of sea turtles. *Amphibia-Reptilia* 29, 361–369.
- 3 McDonald, D. L., and P. H. Dutton. (1996). Use of PIT tags and photoidentification to revise remigration
4 estimates of leatherback turtles (*Demochelys coriacea*) nesting in St. Croix, U.S. Virgin Islands,
5 1979-1995. *Chelonian Conservation and Biology* 2, 148–152.
- 6 Meylan, A. (1995). Sea turtle migration - evidence from tag returns. In *Biology and Conservation of Sea*
7 *Turtles*: 91–100. Bjorndal, K., (Ed.). Washington and London: Smithsonian Institution Press.
- 8 Milligan, S. R., W. V Holt, and R. Lloyd. (2009). Impacts of climate change and environmental factors
9 on reproduction and development in wildlife. *Philosophical transactions of the Royal Society of*
10 *London. Series B, Biological sciences* 364, 3313–9.
- 11 Møller, A. P., D. Rubolini, and E. Lehikoinen. (2008). Populations of migratory bird species that did not
12 show a phenological response to climate change are declining. *PNAS* 105, 16195-16200.
- 13 Mortimer, J. A., and A. Carr. (1987). Reproduction and migrations of the Ascension Island green turtle
14 (*Chelonia mydas*). *Copeia* 1987, 103–113.
- 15 Nichols, J. D., and B. K. Williams. (2006). Monitoring for conservation. *Trends in Ecology & Evolution*
16 21, 668–73.
- 17 Parmesan, C., and G. Yohe. (2003). A globally coherent fingerprint of climate change impacts across
18 natural systems. *Nature* 421, 37–42.
- 19 Pfaller, J. B., K. A. Bjorndal, M. Chaloupka, K. L. Williams, M. G. Frick, and A. B. Bolten. (2013).
20 Accounting for imperfect detection is critical for inferring marine turtle nesting population
21 trends. *PloS one* 8, e62326.

- 1 Pilastro, A., G. Tavecchia, and G. Marin. (2003). Long living and reproduction skipping in the fat
2 dormouse. *Ecology* 84, 1784-1792.
- 3 Pinheiro, J., D. Bates, S. DebRoy, D. Sarkar, and the R. D. C. Team. (2012). *nlme: Linear and Nonlinear*
4 *Mixed Effects Models*. R package version 3.1-105.
- 5 Pitcher, K. W., P. F. Olesiuk, R. F. Brown, M. S. Lowry, S. J. Jeffries, J. L. Sease, W. L. Perryman, C. E.
6 Stinchcomb, and L. F. Lowry. (2007). Abundance and distribution of the eastern North Pacific
7 Steller sea lion (*Eumetopias jubatus*) population. *Fisheries Bulletin* 107, 102–115.
- 8 Prince, R. I. T., and M. Chaloupka. (2012). Estimating demographic parameters for a critically
9 endangered marine species with frequent reproductive omission: hawksbill turtles nesting at
10 Varanus Island, Western Australia. *Marine Biology* 159, 355–363.
- 11 R Development Core Team. (2012). *R: A language and environment for statistical computing*. Vienna,
12 Austria: R Foundation for Statistical Computing.
- 13 Raithel, C. J., P. W. C. Paton, P. S. Pooler, and F. C. Golet. (2011). Assessing long-term population trends
14 of wood frogs using egg-mass counts. *Journal of Herpetology* 45, 23-27.
- 15 Rees, A. F., A. Saad, and M. Jony. (2008). Discovery of a regionally important green turtle *Chelonia*
16 *mydas* rookery in Syria. *Oryx* 42, 456-459.
- 17 Richardson, J. I., D. B. Hall, P. A. Mason, K. M. Andrews, R. Bjorkland, Y. Cai, and R. Bell. (2006). Eighteen
18 years of saturation tagging data reveal a significant increase in nesting hawksbill sea turtles
19 (*Eretmochelys imbricata*) on Long Island, Antigua. *Animal Conservation* 9, 302–307.
- 20 Rieman, B. E., and D. L. Myers. (1997). Use of redd counts to detect trends in bull trout (*Salvelinus*
21 *confluentus*) populations. *Conservation Biology* 11, 1015-1018.

- 1 Rivalan, P., A. Prévot-Julliard, R. Choquet, R. Pradel, B. Jacquemin, and M. Girondot. (2005). Trade-off
2 between current reproductive effort and delay to next reproduction in the leatherback sea
3 turtle. *Population Ecology* 145, 564-574.
- 4 Rivalan, P., R. Pradel, R. Choquet, M. Girondot, and A. Prévot-Julliard. (2006). Estimating clutch
5 frequency in the sea turtle *Dermochelys coriacea* using stopover duration. *Marine Ecology*
6 *Progress Series* 317, 285–295.
- 7 Robinson, R. A. et al. (2009). Travelling through a warming world: climate change and migratory
8 species. *Endangered Species Research* 7, 87-99.
- 9 Schroeder, B., and S. Murphy. (1999). Population Surveys (ground and aerial) on Nesting Beaches. In
10 *Research and Management Techniques for the Conservation of Sea Turtles. No. 4:* 45-55. Eckert,
11 K. L., Bjorndal, K. A., Abreu-Grobois, F. A., Donnelly, M., (Ed.). Washington, DC: IUCN/SSC Marine
12 Turtle Specialist Group.
- 13 Sella, I. (1995). Sea Turtles in the Eastern Mediterranean and Northern Red Sea. In *Biology and*
14 *Conservation of Sea Turtles:* 417–423. Bjorndal, K. A., (Ed.). London: Smithsonian Institution
15 Press.
- 16 Seminoff, J. (2004). *Chelonia mydas*. In *IUCN Red List of Threatened Species*. IUCN, 2012. Version
17 2012.2. <www.iucnredlist.org>.
- 18 Seminoff, J. A., T. T. Jones, A. Resendiz, W. J. Nichols, and M. Y. Chaloupka. (2003). Monitoring green
19 turtles (*Chelonia mydas*) at a coastal foraging area in Baja California, Mexico: multiple indices
20 describe population status. *Journal of the Marine Biological Association of the United Kingdom*
21 83, 1355–1362.

- 1 Simard, Y., N. Roy, S. Giard, C. Gervaise, M. Conversano, and N. Ménard. (2010). Estimating whale
2 density from their whistling activity: Example with St. Lawrence beluga. *Applied Acoustics* 71,
3 1081-1086.
- 4 Sims, M., R. Bjorkland, P. Mason, and L. B. Crowder. (2008). Statistical power and sea turtle nesting
5 beach surveys: How long and when? *Biological Conservation* 141, 2921–2931.
- 6 Snape, R. T. E., D. Beton, A. C. Broderick, B. A. Çiçek, W. J. Fuller, Ö. Özden, and B. J. Godley. (2013).
7 Strand monitoring and anthropological surveys provide insight into marine turtle bycatch in
8 small-scale fisheries of the Eastern Mediterranean. *Chelonian Conservation and Biology* 12, 44-
9 55.
- 10 Solow, A. R., K. A. Bjorndal, and A. B. Bolten. (2002). Annual variation in nesting numbers of marine
11 turtles: the effect of sea surface temperature on re-migration intervals. *Ecology Letters* 5, 742–
12 746.
- 13 Swan, M., J. D. Stefano, F. Christie, E. Steel, A. York. (2013). Detecting mammals in heterogeneous
14 landscapes: Implications for biodiversity monitoring and management. *Biodiversity and
15 Conservation* 23, 343-355.
- 16 Troeng, S., and E. Rankin. (2005). Long-term conservation efforts contribute to positive green turtle
17 *Chelonia mydas* nesting trend at Tortuguero, Costa Rica. *Biological Conservation* 121, 111–116.
- 18 Tucker, A. D. (2010). Nest site fidelity and clutch frequency of loggerhead turtles are better elucidated
19 by satellite telemetry than by nocturnal tagging efforts: Implications for stock estimation. *Journal
20 of Experimental Marine Biology and Ecology* 383, 48–55.
- 21 Tucker, A. D. and N. B. Frazer. (1991). Reproductive variation in leatherback turtles, *Dermochelys
22 coriacea*, at Culebra National Wildlife Refuge, Puerto Rico. *Herpetologica* 47.1, 115-124.

- 1 Van Buskirk, J. V., and L. B. Crowder. (1994). Life-History Variation in Marine Turtles. *Copeia* 1994, 66–
2 81.
- 3 Wallace, B. P. et al. (2011). Global conservation priorities for marine turtles. *PloS one* 6, e24510.
- 4 Weber, N., S. B. Weber, B. J. Godley, J. Ellick and M. Witt. (2013). Telemetry as a tool for improving
5 estimates of marine turtle abundance. *Biological Conservation* 167, 90-96.
- 6 Weishampel, J. F., D. A. Bagley, and L. M. Ehrhart. (2004). Earlier nesting by loggerhead sea turtles
7 following sea surface warming. *Global Change Biology* 10, 1–4.
- 8 Whiting, A. U., M. Chaloupka, and C. J. Limpus. (2013). Comparing sampling effort and errors in
9 abundance estimates between short and protracted nesting seasons for sea turtles. *Journal of*
10 *Experimental Marine Biology and Ecology* 449, 165-170.
- 11 Wright, L. I., W. J. Fuller, B. J. Godley, A. McGowan, T. Tregenza, and A. C. Broderick. (2012a).
12 Reconstruction of paternal genotypes over multiple breeding seasons reveals male green turtles
13 do not breed annually. *Molecular ecology* 21, 3625-3635.
- 14 Wright, L. I., K. L. Stokes, W. J. Fuller, B. J. Godley, A. McGowan, R. Snape, T. Tregenza, and A. C.
15 Broderick. (2012b). Turtle mating patterns buffer against disruptive effects of climate change.
16 *Proc. R. Soc. Lond. B* 279, 2122-2127.
- 17 Yalcin-Ozdilek, S. (2007). Status of sea turtles (*Chelonia mydas* and *Caretta caretta*) on Samandag
18 Beach, Turkey: a five-year monitoring study. *Annales Zoologici Fennici* 44, 333–347.

19

1 **Figures**




Figure 1. Turtle nesting beaches monitored in the current study.

2

1





Figure 2. Green turtle clutches on a) Alagadi beach and b) across all monitored sites against time, with quadratic trend lines (solid lines). These data build on the data 1993-2000 presented in Broderick et al. (2002).

2



1

2 **Figure 3.** Green turtle nesting at Alagadi from 1993 to 2013. a) Number of females nesting at Alagadi
 3 as observed through intensive tagging effort (black dots; data for 1993-2000 previously presented in
 4 Broderick et al. (2002)), and as predicted by dividing annual nest counts by the grand mean expected
 5 clutch frequency for this population (3; open circles). Nesting population trends estimated using these
 6 two measures are almost identical (solid line: actual data, dashed line: predicted data). b) Correlation
 7 between the number of nesting females (solid line), and the proportion of those that are first time
 8 nesters (dotted/ dashed line) at Alagadi over the same time period. The dotted portion indicates
 9 lower confidence in neophyte/ remigrant identification prior to 2000.

10

1





Figure 4. Remigration interval (RI) of green turtles returning to nest at Alagadi. a) Observed RIs (1994-2013). b) Change in RI for green turtles nesting at Alagadi during three or more seasons, taken as the increase/ decrease in RI compared to the previous RI recorded for each individual.






2




1

2 **Figure 5.** Breeding frequency of green turtles at Alagadi from 1993 to 2013. a) Yearly median and
 3 interquartile range for remigration interval (RI) and b) expected clutch frequency (ECF) for nesting at
 4 Alagadi, each with locally weighted regression line (LOESS smoother).

5



1 **Figure 6.** Expected clutch frequency (ECF) for a) neophyte (2000 - 2013), b) all remigrant, c) two year
 2 remigrant, d) three year remigrant and e) four year remigrant green turtles nesting at Alagadi (1994
 3 to 2013). Dashed lines are median values.



1

2 Supplemental Figure S1. Lack of trend in fidelity and detectability. a) Proportion of all nests recorded
 3 across the north and west coasts that are laid at Alagadi. b) Proportion of nests laid at Alagadi that
 4 are assigned to a particular female (through witnessing of oviposition). c) Yearly median ratio of
 5 OCF:ECF (ratios were calculated for each individual female), with 5th - 95th percentiles displayed as
 6 error bars.