

# Relationship between childhood abuse and substance misuse problems is mediated by substance use coping motives, in school attending South African adolescents

Lee Hogarth<sup>1</sup>, Lindi Martin<sup>2</sup>, Soraya Seedat<sup>2</sup>

*Published in Drug and Alcohol dependence*

1. School of Psychology, University of Exeter, Washington Singer Building, Perry Road, Exeter EX4 4QG, UK.
2. Department of Psychiatry, Stellenbosch University, South Africa

**Corresponding author:** Lee Hogarth, School of Psychology, University of Exeter, Washington Singer Building, Perry Road, Exeter EX4 4QG, UK.

**Running head:** Childhood trauma and dependence

**Declaration of Interests:** The secondary analysis was supported by an MRC Confidence in Global Mental Health pump priming award (MC\_PC\_MR/R019991/1) to Hogarth and Seedat, and by an Alcohol Research UK grant (RS17/03) to Hogarth. The original study that collected the data was supported by the South African Research Chair in PTSD, hosted by Stellenbosch University, funded by the Department of Science and Technology, Republic of South Africa (Grant No. 64811) and administered by the National Research Foundation, to Seedat and Martin.

**Word Count:** 3100 (excluding references, tables and figures)

## Abstract

*Background:* In adults, it has been shown that the relationship between childhood abuse and substance misuse problems is mediated by the belief that substance use helps cope with negative affective states. By contrast, in adolescents, it is unknown whether drug use coping motives play this same mediating role.

*Methods:* Secondary analysis of 1149 school attending adolescents in Cape Town South Africa (average age = 16.24 years, range = 13–23; 60% females). Questionnaire measures obtained during a single test session (among a larger battery) assessed childhood trauma (CTQ), alcohol (AUDIT) and drug problems (DUDIT), and coping orientation (A-COPE) which contained three items assessing drug use to cope with negative affect.

*Results:* The three types of childhood abuse measured by the CTQ – emotional, physical and sexual – were positively associated with greater alcohol/drug problems, and drug use coping motives. Drug use coping motives mediated the relationships between childhood abuse types and alcohol/drug problems, and these mediational pathways remained significant when gender and other subscales of the A-COPE were included as covariates.

*Conclusions:* These data are preliminary insofar as drug use coping motives were assessed with a non-validated subscale of the A-COPE. Nevertheless, drug use to cope with negative affect mediated the relationship between all three types of childhood abuse (emotional, physical, sexual) and alcohol/drug problems in school attending adolescents. The implication is that drug prevention programs for this age group should seek to mitigate drug use coping motives.

**Keywords:** Childhood abuse; Substance use coping motives; Alcohol dependence; Drug use problems; Mediation analysis

## 1. Introduction

Childhood abuse is typically associated with greater risk of alcohol and drug problems, in cross sectional and prospective designs, in adolescent and adults, although not all studies replicate this finding (Brady and Back, 2012; Butt et al., 2011; Konkoly Thege et al., 2017; Simpson and Miller, 2002). Childhood abuse/trauma (assessed in adults) is also associated with stronger drug use coping motives, which statistically mediates the relationship between abuse/trauma and alcohol/drug problems (Goldstein et al., 2010; Grayson and Nolen-Hoeksema, 2005; Schuck and Widom, 2001; Vilhena-Churchill and Goldstein, 2014). Similarly, the relationship between adult traumata (e.g. partner violence, sexual coercion etc.) and drug/alcohol problems is mediated by drug use coping motives (Asberg and Renk, 2012; Fossos et al., 2011; Kaysen et al., 2007; Øverup et al., 2015; Ullman et al., 2005). Finally, the relationship between adult psychiatric states (including depression, anxiety) and alcohol/drug problems is mediated by drug use coping motives (Dvorak et al., 2014; Gonzalez et al., 2011; Holahan et al., 2001; McDevitt-Murphy et al., 2015; Mooney et al., 2008; O'Hare and Sherrer, 2011; Peirce et al., 1994; Reardon et al., 2002; Schuckit et al., 2006; Simpson et al., 2014; Stewart et al., 2001; Yeater et al., 2010; Young-Wolff et al., 2009). These studies suggest drug use coping motives are a crucial psychological mechanism driving substance dependence in vulnerable groups.

It is currently unknown whether in school attending adolescents, drug use coping motives mediate the relationship between childhood abuse and alcohol/drug problems, as the forgoing mediation studies have all recruited adults. Indeed, there is a more general question regarding the relative importance of coping motives (as opposed to other substance use motives) as a risk factor for alcohol/drug in the young (Kuntsche et al., 2005). One study, however, does suggest that coping motives are an important risk factor for bullied adolescents (Topper et al., 2011). This study recruited school attending British children aged 13-15, and found that the

prospective relationship between being bullied and alcohol problems at one year follow-up was mediated by drinking to cope with negative affect at follow-up, even when other drinking motives were statistically controlled. It remains to be seen whether coping motives mediate the relationship between child abuse and substance problems in adolescents, specifically.

South African adolescents in particular, and the South African population in general, report high rates of childhood abuse/trauma and alcohol/drug problems (Kaminer et al., 2013a; Kaminer et al., 2013b; Parry et al., 2004; Stansfeld et al., 2018; Stein et al., 2008; Suliman et al., 2009). Furthermore, childhood abuse/trauma is positively associated with alcohol/drug problems in these samples (Morojele et al., 2016; Saban et al., 2014; Sommer et al., 2017) raising questions as to whether this relationship is mediated by coping motives. To address this question, the current study conducted secondary analysis of an existing data set collected to explore the relationship between childhood abuse and psychiatric symptoms in school attending South African adolescents (Martin et al., 2014). The measures relevant to the current hypothesis were the Childhood Trauma Questionnaire (CTQ, Bernstein et al., 2003), the Alcohol and Drug Use Disorders Identification Tests (AUDIT, Babor et al., 2001, and DUDIT, Berman et al., 2005). Crucially, the Adolescent Coping Orientation for Problem Experiences questionnaire (A-COPE, Patterson and McCubbin, 1987) was also recorded, originally to examine protective factors (the ACOPE has been validated with South African Adolescents (Spangenberg and Henderson, 2001). However, the A-COPE includes three items which assess whether individuals use alcohol, smoking or drugs to cope with problems. Although this composite index of coping motives has not previously been validated, the questions are descriptively similar to existing coping motives measures (e.g. Grant et al., 2007), and single item assays of coping motives have previously served as excellent prospective markers for future alcohol problems (Crum et al., 2013; Menary et al., 2011). In

the current analysis, if drug use coping motives indexed by the A-COPE mediate the relationship between childhood abuse (CTQ) and alcohol/drug problems (AUDIT/DUDIT), these mediational paths would suggest that drug prevention programs for vulnerable adolescents should focus on mitigating coping motives specifically – a therapeutic approach that is becoming more widely utilised, and has demonstrated efficacy, in other vulnerable populations (Anker et al., 2016; Bradizza et al., 2017; Conrod et al., 2013; Stasiewicz et al., 2013; Wurdak et al., 2016).

## **2. Method**

### *2.1 Participants*

Participants were 1149 school attending adolescents from a representative sample of secondary schools (n=29) in Cape Town, South Africa. Mean age was 16.24 years (range = 13–23, SD = 1.95), there were 59.97% girls (689/1149), 68.9% identified themselves as Black (792/1149), and mean education level was grade 9 (range = 8–12, SD = 1.30).

### *2.2 Procedure*

Access to secondary schools was granted by the Western Cape Education Department. Ethical approval was provided by Stellenbosch University Health Research Ethics Committee. Randomly selected Cape Town secondary schools were approached. Schools that participated provided names of learners from grades 8 to 12, from which 20 learners per grade were randomly selected. Written consent was obtained from parents/guardians and written assent was obtained from learners. Questionnaires were completed within schools, on a single occasion.

### *2.2 Questionnaires*

The Childhood Trauma Questionnaire (CTQ, Bernstein et al., 2003) is a 28-item retrospective self-report measure of the frequency of and severity of emotional, physical and sexual abuse. The CTQ also contains emotional and physical neglect, and minimisation/denial subscales. This paper only reported the abuse subscales for simplicity. Participants responded to each item in the context of “when you were growing up” and answered according to a 5-point Likert scale ranging from ‘never true’ (1) to ‘very often true’ (5). The abuse subscales were each calculated from 5 items, and ranged from 5 to 25. Severity groups within each abuse type were defined as ‘none’, ‘low’, ‘moderate’ and ‘severe’, using pre-established score boundaries (Bernstein et al., 2003): Emotional Abuse (5-8, 9-12,13-15 and 16+); Physical Abuse (5-7; 8-9; 10-12; 13+); Sexual Abuse (5; 6-7; 8-12; 13+). The Cronbach’s alpha of the three CTQ subscales were .72, .77 and .80 respectively (see Table 1B) indicating acceptable internal consistency of the items.

The Alcohol Use Disorders Identification Test (AUDIT, Babor et al., 2001) is a 10-item questionnaire measuring alcohol use problems. Items are rated on a 5-point scale (total score range 0–40). Scores between 0-7 indicate low-risk alcohol use, whereas scores of 8+ indicate hazardous drinking. The Cronbach’s alpha of the AUDIT was .87 indicating good internal consistency of the items.

The Drug Use Disorders Identification Test (DUDIT, Berman et al., 2005) is an 11-item questionnaire measuring drug use problems (total score range 0–44). For males, scores of 6+, and for females, scores of 2+ are taken to indicate problematic drug use. The Cronbach’s alpha of the DUDIT was .89 indicating good internal consistency of the items.

Adolescent Coping Orientation for Problem Experiences (A-COPE, Patterson and McCubbin, 1987) is a 54-item questionnaire measuring coping strategies, endorsed in the context of the question ‘When you face difficulties or feel tense, how often do you:’. Each coping strategy

was endorsed on a 5-point Likert scale labelled ‘Never’ (1), ‘Hardly’ (2), ‘Sometimes’ (3), ‘Often’ (4) and ‘Most of the time’ (5). Three items assessed the use of drugs to cope: Item 24: ‘Use drugs (not necessarily prescribed by a doctor)’; Item 42: ‘Smoke’; Item 46: ‘Drink beer, wine, liquor’. These items were averaged to create a bespoke composite index of drug use coping motives, descriptively similar to existing validated assays of coping motives (e.g. Grant et al., 2007) and single item measures of coping motives (Crum et al., 2013; Menary et al., 2011). Scores of 3+ (average across items) were considered clinically meaningful insofar as they indicate that the participant (on average) endorsed using drugs to cope at least ‘Sometimes’. The Cronbach’s alpha of the A-COPE bespoke drug coping subscale was .54 indicating poor internal consistency of the items (see Discussion). The A-COPE has 12 standard subscales (ventilating, diversions, self-reliance, social support, solving family problems, avoiding problems, spiritual, close friends, professional support, demanding activity, being humorous and relaxing), which were treated as covariates apart from the Avoiding Problems subscale, which encompasses the three drug coping items.

### *2.3 Data analysis*

In Figures 1A-I, participants were grouped based on the severity of abuse reported in the CTQ childhood abuse questionnaire, into none, low, moderate and severe groups, separately for each gender. The three columns report emotional, physical, sexual abuse respectively. The three rows report the mean (and SEM) of alcohol problems (AUDIT), drug problems (DUDIT), and drug use coping motives (A-COPE), respectively. For each figure (A-I), a separate ANOVA was conducted with the drug measure (AUDIT, DUDIT or A-COPE) as the dependent variable, and the between-subject variables CTQ severity group and gender. A significant main effect of CTQ severity group would indicate that that form of abuse was related to the drug outcome (irrespective of gender). A significant main effect of gender

would indicate that the drug outcome differed between genders. Finally, a significant interaction between CTQ severity group and gender would indicate that that form of abuse was more strongly related to the drug outcome in one gender. Where the main effect of CTQ severity group was significant for a figure, Games-Howell post hoc tests were used to compare every pair of CTQ groups.

Mediation models were tested with 5000 bias corrected bootstrapped confidence intervals, using Hayes and colleagues (Preacher and Hayes, 2004) software within SPSS (all other statistics were in SPSS). This method was favoured because it arguably produces the least Type I and Type II errors (Preacher and Hayes, 2008), and because it is thought to have greater power to detect mediational effects than alternative approaches (MacKinnon et al., 2002). None of the variables used in the analysis were normally distributed, the CTQ abuse groups did not have homogenous variance in AUDIT, DUDIT or drug use coping scores, and there were unequal N for CTQ abuse severity groups. ANOVA, Hayes' bootstrapping method, and Games-Howell post hoc tests are considered robust against these parameters.

### **3. Results**

Figures 1A-I show the relationship between three childhood abuse types separated into columns (physical, emotional, sexual) and the three drug outcome measures separated into rows (AUDIT, DUDIT and drug use coping motives). Inspection of Figure 1 indicates that adolescents in the severe emotional, physical and sexual abuse categories showed higher rates of alcohol and drug problems, and drug use to cope with negative affect (the one exception is sexual abuse which was not associated with greater drug problems on the DUDIT, Figure 1F), Finally, these relationships did not differ between genders (but see Figure 1D).

#### **3.1 Relationship between childhood abuse and alcohol problems**



Following the analytical plan outlined above, ANOVA on Figure 1A produced significant main effects of emotional abuse severity,  $F(3,1141)=9.09$ ,  $p<.001$ ,  $\eta_p^2=.023$ , indicating that emotional abuse was associated with increased alcohol problems across genders. There was also a significant main effect of gender,  $F(1,1141)=10.35$ ,  $p<.001$ ,  $\eta_p^2=.009$ , indicating that males had more alcohol problems. However, there was no significant interaction,  $F<1$ , indicating that the association between emotional abuse and alcohol problems was comparable between genders. Post hoc tests indicated that the severe group differed from the none,  $p<.001$ , and low group,  $p=.01$ , but not the moderate group,  $p=.14$ . ANOVA on Figure 1B produced significant main effects of physical abuse severity,  $F(3,1141)=4.26$ ,  $p=.005$ ,  $\eta_p^2=.011$ , and gender,  $F(1,1141)=4.71$ ,  $p=.03$ ,  $\eta_p^2=.004$ , but no interaction,  $F(3,1141)=1.38$ ,  $p=.25$ ,  $\eta_p^2=.004$ . Only the severe and none groups differed,  $p=.005$ . ANOVA on Figure 1C produced significant main effects of sexual abuse severity,  $F(3,1141)=6.94$ ,  $p<.001$ ,  $\eta_p^2=.018$ , and gender,  $F(1,1141)=6.69$ ,  $p=.01$ ,  $\eta_p^2=.006$ , but no interaction,  $F<1$ . Only the severe and none groups differed,  $p<.005$ .

### 3.2 Relationship between childhood abuse and drug problems

ANOVA on Figure 1D produced significant main effects of emotional abuse severity,  $F(3,1141)=17.83$ ,  $p<.001$ ,  $\eta_p^2=.045$ , and gender,  $F(1,1141)=39.28$ ,  $p<.001$ ,  $\eta_p^2=.009$ , and a significant interaction,  $F(3,1141)=2.85$ ,  $p=.037$ ,  $\eta_p^2=.007$ . In each gender, only the severe group differed from the none group,  $ps\leq.014$ . ANOVA on Figure 1E produced significant main effects of physical abuse severity,  $F(3,1141)=9.18$ ,  $p<.001$ ,  $\eta_p^2=.024$ , and gender,  $F(1,1141)=24.95$ ,  $p<.001$ ,  $\eta_p^2=.021$ , but no interaction,  $F(3,1141)=1.59$ ,  $p=.19$ ,  $\eta_p^2=.004$ . Only the severe and none groups differed,  $p=.005$ . ANOVA on Figure 1F produced no significant main effects of sexual abuse severity,  $F(3,1141)=1.44$ ,  $p=.23$ ,  $\eta_p^2=.004$ ,

(indicating no relationship between sexual abuse and drug problems), a significant main effect of gender,  $F(1,1141)=16.91, p<.001, \eta_p^2=.015$ , and no interaction,  $F<1$ .

### 3.3 Relationship between childhood abuse and drug use coping motives

ANOVA on Figure 1G produced significant main effects of emotional abuse severity,  $F(3,1141)=19.16, p<.001, \eta_p^2=.048$ , and gender,  $F(1,1141)=18.51, p<.001, \eta_p^2=.016$ , but no interaction,  $F<1$ . The severe group differed from all the remaining groups,  $ps\leq.003$ . ANOVA on Figure 1H produced significant main effects of physical abuse severity,  $F(3,1141)=8.16, p<.001, \eta_p^2=.021$ , and gender,  $F(1,1141)=10.31, p=.001, \eta_p^2=.009$ , but no interaction,  $F<1$ . The severe group differed from the none group,  $p<.001$ , and low group,  $p=.037$ . ANOVA on Figure 1I produced significant main effects of sexual abuse severity,  $F(3,1141)=18.48, p<.001, \eta_p^2=.046$ , and gender,  $F(1,1141)=5.91, p=.015, \eta_p^2=.005$ , but no interaction,  $F(3,1141)=1.85, p=.14, \eta_p^2=.005$ . The severe group differed from all the other groups,  $ps\leq.001$ .

### 3.4 Mediation analysis

Table 1A shows 6 mediation models with CTQ childhood abuse types (emotional, physical, sexual) as predictors (X), alcohol (AUDIT) or drug (DUDIT) problems as the outcome (Y), and drug use coping motives (A-COPE) as the mediator (M). The beta coefficients of the component paths (a, b, c) indicate that there were significant interrelationships between the CTQ abuse types, drug use coping motives and alcohol/drug problems (confirmed by the correlation matrix shown in Table 1B). The one exception was CTQ sexual abuse which was not related to drug use problems on the DUDIT (c path total effect; see also Table 1B and Figure 1F). Most importantly, the indirect mediation paths were all significant (the 95% confidence intervals do not encompass zero). Finally, the c' paths between X→Y remained significant when the mediational paths were controlled for the first five models, indicating

partial mediation, whereas the c' path was not significant for the sixth model indicating full mediation. The mediational paths remained significant when gender and the 11 subscales of the A-COPE were entered as covariates into the models, suggesting the mediational role of drug use coping motives could not be explained by other coping styles.

#### **4. Discussion**

The main finding was childhood abuse types (emotional, physical, sexual) were associated with increased alcohol/drug problems, and drug use coping motives. The exception was the relationship between sexual abuse and drug problems, which was not reliable. These findings corroborate the majority of studies which have demonstrated that childhood abuse is associated with substance problems (Brady and Back, 2012; Butt et al., 2011; Konkoly Thege et al., 2017; Simpson and Miller, 2002). There were no differences between the genders in the strength of these associations, apart from emotional abuse which had a stronger relationship with drug problems in males (Figure 1D). Post hoc contrasts suggested that severe abuse was associated with a jump in alcohol/drug problems and coping motives, and that there was no significant linear increase across the less severe abuse groups despite the numerical trend. The implication is that individuals who fall into the severe abuse group could be selected for prevention interventions. Importantly, the relationships between childhood abuse and alcohol/drug problems were demonstrated in a non-western sample, providing rare cross-cultural corroboration (Konkoly Thege et al., 2017).

The most important finding was that drug use coping motives mediated the relationship between childhood abuse and alcohol/drug problems. These mediational pathways remained significant when gender and other subscales of the A-COPE were entered as covariates suggesting that the mediating role of drug use coping motives could not be explained by other coping styles. The study is the first to confirm this mediational pathway in school attending

adolescents (it has previously only observed in adults), and in South African adolescents, suggesting the mediational path is found cross culturally. Such cross cultural generalisation may have been doubted given that previous studies with South African university students found that coping motives were endorsed less than enhancement motives (Maphisa and Young, 2018; Peltzer, 2003; see also Gire, 2002) suggesting that coping motives might be less pronounced in South Africa (but see Kuntsche et al., 2014). The current study suggests that because of their mediating role, drug use coping motives should be targeted by drug prevention programs designed for traumatised South African Adolescents specifically, but also for other vulnerable groups where coping motives play the same mediating role (Anker et al., 2016; Bradizza et al., 2017; Conrod et al., 2013; Stasiewicz et al., 2013; Wurdak et al., 2016).

The main limitation was the non-validated index of drug use coping motives derived from three items of the A-COPE. Although these items are descriptively similar to validated questionnaires of coping motives (e.g. Grant et al., 2007) and single item assays of coping motives have previously served as excellent prospective markers for future alcohol problems (Crum et al., 2013; Menary et al., 2011), the Cronbach's alpha for these three items was .54, indicating poor internal consistency of the items. However, Cronbach's alpha decreases with fewer items so a lower threshold for acceptability is sometimes applied in such cases (Heo et al., 2015). Furthermore, the corrected item total correlations for the smoke, drugs and drink coping items were .44, .32, and .32 respectively, indicating that each item was correlated with a composite of the remaining items, and all are above the desirable cutoff of .3 (Ferketich, 1991). Finally, if each of the individual items was deleted, the overall Cronbach's alpha of .54 would be reduced to .29, .50, and .51, respectively, indicating that each item contributed to the overall Cronbach's alpha value. From this analysis, it might be concluded that the bespoke index of coping motives is acceptable, but minimally so, and thus evidence for the

mediating role of coping motives in the current models may be deemed preliminary. A key objective of future work should be to overcome this limitation by testing whether the current mediational pathways are replicated, or ideally are stronger, when a validated, multiple-item assay of drug use coping motives is tested as the mediating variable. In addition, future studies might utilise measures of alcohol and drug problems which have been validated with adolescents specifically (Read et al., 2006) rather than the AUDIT/DUDIT which have been validated primarily with adult samples (Rumpf et al., 2013). Finally, future work should replicate the cross sectional mediation design with a longitudinal mediation design to support the claim that drug use coping motives are causal in the growth in alcohol/drug problems in abused adolescents (Jose, 2016; Windle and Windle, 2015). These studies would strengthen calls for the development of drug prevention interventions that target drug use coping motives in abused adolescents.

## References

- Anker, J.J., Kushner, M.G., Thuras, P., Menk, J., Unruh, A.S., 2016. Drinking to cope with negative emotions moderates alcohol use disorder treatment response in patients with co-occurring anxiety disorder. *Drug Alcohol Depend.* 159, 93-100.
- Asberg, K., Renk, K., 2012. Substance use coping as a mediator of the relationship between trauma symptoms and substance use consequences among incarcerated females with childhood sexual abuse histories. *Subst. Use Misuse* 47, 799-808.
- Babor, T.F., Higgins-Biddle, J.C., Saunders, J.B., Monteiro, M.G., 2001. AUDIT: The alcohol use disorders identification test guidelines for use in primary care. Second edition. In: *Dependence, D.o.M.H.a.S. (Ed.). World Health Organization.*
- Berman, A.H., Bergman, H., Palmstierna, T., Schlyter, F., 2005. Evaluation of the Drug Use Disorders Identification Test (DUDIT) in criminal justice and detoxification settings and in a Swedish population sample. *Eur. Addict. Res.* 11, 22-31.
- Bernstein, D.P., Stein, J.A., Newcomb, M.D., Walker, E., Pogge, D., Ahluvalia, T., Stokes, J., Handelsman, L., Medrano, M., Desmond, D., Zule, W., 2003. Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Neglect* 27, 169-190.
- Bradizza, C.M., Stasiewicz, P.R., Zhuo, Y., Ruszczyc, M., Maisto, S.A., Lucke, J.F., Brandon, T.H., Eiden, R.D., Slosman, K.S., Giarratano, P., 2017. Smoking cessation for pregnant smokers: Development and pilot test of an emotion regulation treatment supplement to standard smoking cessation for negative affect smokers. *Nicotine and Tobacco Research* 19, 578-584.
- Brady, K.T., Back, S.E., 2012. Childhood trauma, posttraumatic stress disorder, and alcohol dependence. *Alcohol research : current reviews* 34, 408-413.
- Butt, S., Chou, S., Browne, K., 2011. A rapid systematic review on the association between childhood physical and sexual abuse and illicit drug use among males. *Child Abuse Review* 20, 6-38.
- Conrod, P.J., O'Leary-Barrett, M., Newton, N., Topper, L., Castellanos-Ryan, N., Mackie, C., Girard, A., 2013. Effectiveness of a selective, personality-targeted prevention program for adolescent alcohol use and misuse: a cluster randomized controlled trial. *JAMA Psychiatry* 70, 334-342.
- Crum, R.M., Mojtabai, R., Lazareck, S., Bolton, J.M., Robinson, J., Sareen, J., Green, K.M., Stuart, E.A., La Flair, L., Alvanzo, A.A., Storr, C.L., 2013. A prospective assessment of reports of drinking to self-medicate mood symptoms with the incidence and persistence of alcohol dependence. *JAMA Psychiatry* 70, 718-726.
- Dvorak, R.D., Pearson, M.R., Day, A.M., 2014. Ecological momentary assessment of acute alcohol use disorder symptoms: Associations with mood, motives, and use on planned drinking days. *Exp Clin Psychopharmacol* 22, 285-297.
- Ferketich, S., 1991. Focus on psychometrics. Aspects of item analysis. *Res. Nurs. Health* 14, 165-168.
- Fossos, N., Kaysen, D., Neighbors, C., Lindgren, K.P., Hove, M.C., 2011. Coping motives as a mediator of the relationship between sexual coercion and problem drinking in college students. *Addict. Behav.* 36, 1001-1007.
- Gire, J.T., 2002. A cross-national study of motives for drinking alcohol. *Subst. Use Misuse* 37, 215-223.
- Goldstein, A.L., Flett, G.L., Wekerle, C., 2010. Child maltreatment, alcohol use and drinking consequences among male and female college students: An examination of drinking motives as mediators. *Addict. Behav.* 35, 636-639.

- Gonzalez, V.M., Reynolds, B., Skewes, M.C., 2011. Role of impulsivity in the relationship between depression and alcohol problems among emerging adult college drinkers. *Exp Clin Psychopharmacol* 19, 303-313.
- Grant, V.V., Stewart, S.H., O'Connor, R.M., Blackwell, E., Conrod, P.J., 2007. Psychometric evaluation of the five-factor Modified Drinking Motives Questionnaire — Revised in undergraduates. *Addict. Behav.* 32, 2611-2632.
- Grayson, C.E., Nolen-Hoeksema, S., 2005. Motives to drink as mediators between childhood sexual assault and alcohol problems in adult women. *J. Traum. Stress* 18, 137-145.
- Heo, M., Kim, N., Faith, M.S., 2015. Statistical power as a function of Cronbach alpha of instrument questionnaire items. *BMC Medical Research Methodology* 15, 86.
- Holahan, C.J., Moos, R.H., Holahan, C.K., Cronkite, R.C., Randall, P.K., 2001. Drinking to cope, emotional distress and alcohol use and abuse: A ten-year model. *J. Stud. Alcohol* 62, 190-198.
- Jose, P.E., 2016. The merits of using longitudinal mediation. *Educational Psychologist* 51, 331-341.
- Kaminer, D., du Plessis, B., Hardy, A., Benjamin, A., 2013a. Exposure to violence across multiple sites among young South African adolescents. *Peace and Conflict: Journal of Peace Psychology* 19, 112-124.
- Kaminer, D., Hardy, A., Heath, K., Mosdell, J., Bawa, U., 2013b. Gender patterns in the contribution of different types of violence to posttraumatic stress symptoms among South African urban youth. *Child Abuse Negl.* 37, 320-330.
- Kaysen, D., Dillworth, T.M., Simpson, T., Waldrop, A., Larimer, M.E., Resick, P.A., 2007. Domestic violence and alcohol use: Trauma-related symptoms and motives for drinking. *Addict. Behav.* 32, 1272-1283.
- Konkolý Thege, B., Horwood, L., Slater, L., Tan, M.C., Hodgins, D.C., Wild, T.C., 2017. Relationship between interpersonal trauma exposure and addictive behaviors: a systematic review. *BMC Psychiatry* 17, 164.
- Kuntsche, E., Gabhainn, S.N., Roberts, C., Windlin, B., Vieno, A., Bendtsen, P., Hublet, A., Tynjala, J., Valimaa, R., Dankulinova, Z., Aasvee, K., Demetrovics, Z., Farkas, J., van der Sluijs, W., de Matos, M.G., Mazur, J., Wicki, M., 2014. Drinking motives and links to alcohol use in 13 European countries. *Journal of Studies on Alcohol and Drugs* 75, 428-437.
- Kuntsche, E., Knibbe, R., Gmel, G., Engels, R., 2005. Why do young people drink? A review of drinking motives. *Clinical Psychology Review* 25, 841-861.
- MacKinnon, D.P., Lockwood, C.M., Hoffman, J.M., West, S.G., Sheets, V., 2002. A comparison of methods to test mediation and other intervening variable effects. *Psychol Methods* 7, 83-104.
- Maphisa, J.M., Young, C., 2018. Risk of alcohol use disorder among South African university students: The role of drinking motives. *Addict. Behav.* 82, 44-49.
- Martin, L., Viljoen, M., Kidd, M., Seedat, S., 2014. Are childhood trauma exposures predictive of anxiety sensitivity in school attending youth? *J Affect Disord* 168, 5-12.
- McDevitt-Murphy, M.E., Fields, J.A., Monahan, C.J., Bracken, K.L., 2015. Drinking motives among heavy-drinking veterans with and without posttraumatic stress disorder. *Addiction Research & Theory* 23, 148-155.
- Menary, K.R., Kushner, M.G., Maurer, E., Thuras, P., 2011. The prevalence and clinical implications of self-medication among individuals with anxiety disorders. *Journal of Anxiety Disorders* 25, 335-339.
- Mooney, J.L., Minor, K.I., Wells, J.B., Leukefeld, C., Oser, C.B., Staton Tindall, M., 2008. The relationship of stress, impulsivity, and beliefs to drug use severity in a sample of women prison inmates. *Int. J. Offender Ther. Comp. Criminol.* 52, 686-697.
- Morojele, N.K., Brook, J.S., Brook, D.W., 2016. Tobacco and alcohol use among adolescents in South Africa: shared and unshared risks. *Journal of child and adolescent mental health* 28, 139-152.

- O'Hare, T., Sherrer, M., 2011. Drinking motives as mediators between PTSD symptom severity and alcohol consumption in persons with severe mental illnesses. *Addict. Behav.* 36, 465-469.
- Øverup, C.S., DiBello, A.M., Brunson, J.A., Acitelli, L.K., Neighbors, C., 2015. Drowning the pain: Intimate partner violence and drinking to cope prospectively predict problem drinking. *Addict. Behav.* 41, 152-161.
- Parry, C.D., Myers, B., Morojele, N.K., Flisher, A.J., Bhana, A., Donson, H., Pluddemann, A., 2004. Trends in adolescent alcohol and other drug use: findings from three sentinel sites in South Africa (1997-2001). *J. Adolesc.* 27, 429-440.
- Patterson, J.M., McCubbin, H.I., 1987. Adolescent coping style and behaviors: conceptualization and measurement. *J. Adolesc.* 10, 163-186.
- Peirce, R.S., Frone, M.R., Russell, M., Cooper, M.L., 1994. Relationship of financial strain and psychosocial resources to alcohol use and abuse: The mediating role of negative affect and drinking motives. *J. Health Soc. Behav.* 35, 291-308.
- Peltzer, K., 2003. Drinking motives, behaviour and problems among black South African university students. *African Journal of Drug & Alcohol Studies* 2, 1-10.
- Preacher, K.J., Hayes, A.F., 2004. SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behavior Research Methods, Instruments, & Computers* 36, 717-731.
- Preacher, K.J., Hayes, A.F., 2008. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior research methods* 40, 879-891.
- Read, J.P., Kahler, C.W., Strong, D.R., Colder, C.R., 2006. Development and preliminary validation of the young adult alcohol consequences questionnaire. *J. Stud. Alcohol* 67, 169-177.
- Reardon, M.L., Lang, A.R., Patrick, C.J., 2002. An evaluation of relations among antisocial behavior, psychopathic traits, and alcohol problems in incarcerated men. *Alcoholism: Clinical and Experimental Research* 26, 1188-1197.
- Rumpf, H.J., Wohler, T., Freyer-Adam, J., Grothues, J., Bischof, G., 2013. Screening questionnaires for problem drinking in adolescents: performance of AUDIT, AUDIT-C, CRAFFT and POSIT. *Eur. Addict. Res.* 19, 121-127.
- Saban, A., Flisher, A.J., Grimsrud, A., Morojele, N., London, L., Williams, D.R., Stein, D.J., 2014. The association between substance use and common mental disorders in young adults: results from the South African Stress and Health (SASH) Survey. *The Pan African Medical Journal* 17, 11.
- Schuck, A.M., Widom, C.S., 2001. Childhood victimization and alcohol symptoms in females: causal inferences and hypothesized mediators. *Child Abuse Neglect* 25, 1069-1092.
- Schuckit, M.A., Smith, T.L., Chacko, Y., 2006. Evaluation of a depression-related model of alcohol problems in 430 probands from the San Diego prospective study. *Drug Alcohol Depend.* 82, 194-203.
- Simpson, T.L., Miller, W.R., 2002. Concomitance between childhood sexual and physical abuse and substance use problems: A review. *Clinical Psychology Review* 22, 27-77.
- Simpson, T.L., Stappenbeck, C.A., Luterek, J.A., Lehavot, K., Kaysen, D.L., 2014. Drinking motives moderate daily relationships between PTSD symptoms and alcohol use. *J Abnorm Psychol* 123, 237-247.
- Sommer, J., Hinsberger, M., Elbert, T., Holtzhausen, L., Kaminer, D., Seedat, S., Madikane, S., Weierstall, R., 2017. The interplay between trauma, substance abuse and appetitive aggression and its relation to criminal activity among high-risk males in South Africa. *Addict. Behav.* 64, 29-34.
- Spangenberg, J.J., Henderson, K., 2001. Stress and coping in Black South African adolescents. *Studia Psychologica* 43, 77-90.
- Stansfeld, S.A., Rotheron, C., Das-Munshi, J., Mathews, C., Adams, A., Clark, C., Lund, C., 2018. Exposure to violence and mental health of adolescents: South African Health and Well-being Study. *BJPsych Open* 3, 257-264.



- Stasiewicz, P.R., Bradizza, C.M., Schlauch, R.C., Coffey, S.F., Gulliver, S.B., Gudleski, G.D., Bole, C.W., 2013. Affect regulation training (ART) for alcohol use disorders: Development of a novel intervention for negative affect drinkers. *J. Subst. Abuse Treat.* 45, 433-443.
- Stein, D.J., Seedat, S., Herman, A., Moomal, H., Heeringa, S.G., Kessler, R.C., Williams, D.R., 2008. Lifetime prevalence of psychiatric disorders in South Africa. *Br. J. Psychiatry* 192, 112-117.
- Stewart, S.H., Zvolensky, M.J., Eifert, G.H., 2001. Negative-reinforcement drinking motives mediate the relation between anxiety sensitivity and increased drinking behavior. *Pers. Individ. Differ.* 31, 157-171.
- Suliman, S., Mkabile, S.G., Fincham, D.S., Ahmed, R., Stein, D.J., Seedat, S., 2009. Cumulative effect of multiple trauma on symptoms of posttraumatic stress disorder, anxiety, and depression in adolescents. *Comprehensive Psychiatry* 50, 121-127.
- Topper, L.R., Castellanos-Ryan, N., Mackie, C., Conrod, P.J., 2011. Adolescent bullying victimisation and alcohol-related problem behaviour mediated by coping drinking motives over a 12 month period. *Addict. Behav.* 36, 6-13.
- Ullman, S.E., Filipas, H.H., Townsend, S.M., Starzynski, L.L., 2005. Trauma exposure, posttraumatic stress disorder and problem drinking in sexual assault survivors. *J. Stud. Alcohol* 66, 610-619.
- Vilhena-Churchill, N., Goldstein, A.L., 2014. Child maltreatment and marijuana problems in young adults: Examining the role of motives and emotion dysregulation. *Child Abuse Neglect* 38, 962-972.
- Windle, M., Windle, R.C., 2015. A prospective study of stressful events, coping motives for drinking, and alcohol use among middle-aged adults. *Journal of Studies on Alcohol and Drugs* 76, 465-473.
- Wurdak, M., Wolstein, J., Kuntsche, E., 2016. Effectiveness of a drinking-motive-tailored emergency-room intervention among adolescents admitted to hospital due to acute alcohol intoxication — A randomized controlled trial. *Preventive Medicine Reports* 3, 83-89.
- Yeater, E.A., Austin, J.L., Green, M.J., Smith, J.E., 2010. Coping mediates the relationship between posttraumatic stress disorder (PTSD) symptoms and alcohol use in homeless, ethnically diverse women: A preliminary study. *Psychological Trauma: Theory, Research, Practice, and Policy* 2, 307-310.
- Young-Wolff, K.C., Kendler, K.S., Sintov, N.D., Prescott, C.A., 2009. Mood-related drinking motives mediate the familial association between major depression and alcohol dependence. *Alcoholism: Clinical and Experimental Research* 33, 1476-1486.

Mediation models			Component paths			Mediation	
X (predictor)	M (mediator)	Y (outcome)	a path: X→M	b path: M→Y controlling for X	c path: total effect X→Y	c' path: direct effect X→Y controlling for the indirect mediational path	Indirect mediational path: X→M→Y
CTQ Emotional abuse score	A-COPE drug use coping motives	AUDIT total	$\beta=.04$ , CI=.03-.05***	$\beta=4.09$ , CI=3.67-4.52***	$\beta=.26$ , CI=.17-.35***	$\beta=.09$ , CI=.02-.17*	$\beta=.17$ , CI=.12-.22***
CTQ Physical abuse score			$\beta=.04$ , CI=.02-.05***	$\beta=4.12$ , CI=3.70-4.54***	$\beta=.24$ , CI=.15-.33***	$\beta=.10$ , CI=.02-.18*	$\beta=.14$ , CI=.09-.20***
CTQ Sexual abuse score			$\beta=.05$ , CI=.03-.06***	$\beta=4.17$ , CI=3.74-4.59***	$\beta=.24$ , CI=.13-.34***	$\beta=.04$ , CI=-.05-.13	$\beta=.19$ , CI=.13-.26***
CTQ Emotional abuse score		DUDIT total	$\beta=.04$ , CI=.03-.05***	$\beta=2.56$ , CI=2.25-2.87***	$\beta=.19$ , CI=.13-.25***	$\beta=.09$ , CI=.03-.14*	$\beta=.10$ , CI=.07-.14***
CTQ Physical abuse score			$\beta=.04$ , CI=.02-.05***	$\beta=2.58$ , CI=2.27-2.89***	$\beta=.18$ , CI=.11-.24***	$\beta=.09$ , CI=.03-.14**	$\beta=.09$ , CI=.06-.12***
CTQ Sexual abuse score			$\beta=.05$ , CI=.03-.06***	$\beta=2.72$ , CI=2.41-3.03***	$\beta=.06$ , CI=-.01-.14	$\beta=-.06$ , CI=-.13-.01	$\beta=.13$ , CI=.08-.18***

B. Pearson intercorrelations (r); Cronbach's alpha on the diagonal in brackets						
	1	2	3	4	5	6
1. CTQ emotional abuse score	(.72)					
2. CTQ Physical abuse score	$r=.60$ ***	(.77)				
3. CTQ Sexual abuse score	$r=.38$ ***	$r=.42$ ***	(.80)			
4. A-COPE Drug use coping motives	$r=.22$ ***	$r=.18$ ***	$r=.21$ ***	(.54)		
5. AUDIT total	$r=.17$ ***	$r=.15$ ***	$r=.13$ ***	$r=.51$ ***	(.87)	
6. DUDIT total	$r=.18$ ***	$r=.16$ ***	$r=.05$	$r=.45$ ***	$r=.36$ ***	(.89)

Table 1A. Mediation models. Unstandardised beta coefficients ( $\beta$ ) and 95% confidence intervals (CI) from mediational models calculated using Hayes Process bias corrected bootstrap method (5000 samples). \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$ . CTQ = Childhood Trauma Questionnaire, abuse subscales. A-COPE = Adolescent Coping Orientation for Problem Experiences, bespoke subscale assessing drug, smoking and alcohol use to cope. AUDIT = Alcohol Use Disorders Identification Test. DUDIT = Drug Use Disorders Identification Test. The a, b, and c paths indicate that there were significant interrelationships between CTQ abuse types (emotional, physical, sexual), A-COPE drug use coping motives, and alcohol/drug problems indexed by the AUDIT/DUDIT. The one exception is that CTQ sexual abuse was not related to drug use problems in the DUDIT (c path). All six indirect mediational paths were significant (the 95% confidence intervals do not encompass zero) demonstrating that coping motives mediate the relationship between childhood abuse and alcohol/drug problems. The direct effect (c' paths) between  $X \rightarrow Y$  remained significant when the mediational paths were controlled for the first five models, indicating partial mediation, whereas the direct effect was not significant for the sixth model indicating full mediation. The overall conclusion is that drug use coping motives mediate the relationship between childhood abuse and alcohol/drug problems in school attending South African adolescents.

Table 1B. Pearson correlation coefficients between CTQ abuse types (emotional, physical, sexual), drug use to cope and alcohol/drug problems. The Cronbach's alpha of subscales is shown on the diagonal in brackets.




Figure 1

In Figures 1A-I, participants were grouped based on the severity of abuse reported in the CTQ childhood abuse questionnaire, into none, low, moderate and severe groups, separately for each gender. The three columns report emotional, physical, sexual abuse respectively. The three rows report the mean (and SEM) of alcohol problems (AUDIT), drug problems (DUDIT), and drug use coping motives (A-COPE), respectively. For each figure (A-I), a separate ANOVA was conducted with the drug measure (AUDIT, DUDIT or A-COPE) as the dependent variable, and the between-subject variables CTQ severity group and gender.

Adolescents in the severe emotional, physical and sexual abuse categories showed higher rates of alcohol and drug problem, and drug use coping motives (apart from sexual abuse which was not associated with greater drug problems in the DUDIT, Figure 1F). These relationships between childhood abuse and alcohol/drug problems did not differ between genders (apart from the relationship between emotional abuse and drug problems which was stronger in males - Figure 1D). The percentage value shown above each bar is the proportion of each CTQ abuse severity group that reported a ‘clinically meaningful’ level of the drug outcome: AUDIT = hazardous drinking (8+ scores); DUDIT = drug problems (6+ for males, 2+ for females); A-COPE = drug use coping score of 3+ indicating that the participant (on average across items) endorsed using drugs to cope at least ‘Sometimes’. The numbers at the bottom of each bar is the N of participants falling into each CTQ abuse severity group.