

1

2 **Personalised Ecology**

3

4 Kevin J. Gaston,^{1,2,*} Masashi Soga,³ James P. Duffy,¹ Joanne K. Garrett,⁴ Sian Gaston,¹ and
5 Daniel T.C. Cox^{1,4}

6

7 ¹Environment & Sustainability Institute, University of Exeter, Penryn, Cornwall, TR10 9FE, United
8 Kingdom

9 ²Wissenschaftskolleg zu Berlin, Institute for Advanced Study, Wallotstrasse 19, 14193, Berlin,
10 Germany

11 ³School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1, Yayoi, Bunkyo, Tokyo,
12 113-8656, Japan

13 ⁴European Centre for Environment and Human Health, University of Exeter Medical School, Truro,
14 Cornwall, TR1 3HD, United Kingdom

15

16 *Correspondence: k.j.gaston@exeter.ac.uk (K.J. Gaston)

17

18 Keywords: ecosystem services; extinction of experience; human-nature interactions; nature-health
19 interactions; observer bias; urbanisation

20

21 **The field of ecology has focussed on understanding characteristics of natural systems in a**
22 **manner as free as possible from biases of human observers. However, demand is growing**
23 **for knowledge of human-nature interactions at the level of individual people. This is**
24 **particularly driven by concerns around human health consequences of changes in positive**
25 **and negative such interactions. This requires attention to the biased ways in which people**
26 **encounter and experience other organisms. Here we define such a ‘personalised ecology’**
27 **and discuss its connections to other aspects of the field. We propose a framework of focal**
28 **research topics, shaped by whether the unit of analysis is a single person, a single**
29 **population or multiple populations, and whether a human or nature perspective is foremost.**

30

31 **Human-nature Interactions**

32 Ecology has been defined as the study of the abundance and distribution of organisms and the
33 interactions that determine these [1]. As such, it has been important to measure what those
34 abundances and distributions actually are, or at least to have well behaved and characterised
35 proxies, and to limit the influence of the human observer on these estimations. A vast and rich
36 literature has developed particularly around the form of biases in the human detection of individual
37 organisms, the factors that influence those biases (individual and species characteristics, species
38 richness, habitat, season, weather, observer skills, etc.), and the strengths and weaknesses of
39 approaches to their reduction (e.g., [2-5]). Indeed, a major theme of the history of ecology as a
40 discipline has been progressive improvement in documenting the real abundances and
41 distributions of organisms and their respective dynamics.

42

43 By contrast, there has been little consideration of the converse need to understand the interactions
44 that occur between human observers and nature. Nonetheless, demand arises from several
45 quarters to focus on the very effects that traditionally, ecologists have sought to minimise or control
46 for in their studies. First, and perhaps foremost, it has become apparent that people derive a wide
47 array of health and well-being benefits from their personal interactions with nature (reviewed in [6]).
48 This is particularly so in urban areas, which are epicentres for chronic and non-communicable
49 physical and mental health conditions [7] and where opportunities for nature experiences may be

50 less prevalent. These health and well-being benefits include components of mental, physical and
51 social health [6,8,9]. Key to determining how these benefits are achieved is a better understanding
52 of the form, frequency and duration of people's interactions with nature [10].

53

54 Second, there is growing evidence of a progressive reduction in positive human-nature
55 interactions, particularly in more westernised societies and during childhood [11]. This so-called
56 '**extinction of experience**' (see Glossary) [12] results from a combination of local and regional
57 losses of biodiversity, growth of sedentary pastimes, and perceived safety concerns that limit
58 children's independent activities. This may have profound consequences because the loss of
59 human-nature interactions limits the associated health and well-being benefits. There is also
60 evidence that it results in reductions in emotional affinity toward nature and in pro-environmental
61 attitudes and behaviour [11]. Ongoing extinction of experience could thus imply a cycle of
62 disaffection toward nature, and ultimately constitute one of the greatest challenges to conservation
63 policies and management actions aimed at slowing or halting the biodiversity crisis [13]. Again,
64 better understanding the actual nature experiences that people have and how these compare with
65 those that are available is key to addressing these issues.

66

67 Third, there is much discussion and debate around **human-wildlife conflict**, and hence negative
68 human-nature interactions (e.g., [14,15]). One form of this conflict concerns direct interactions
69 between people and wildlife. In the extreme, for example, attacks on humans by large predators
70 appear to be on the rise [16], likely as a consequence of some combination of reductions in
71 available natural undisturbed habitat, increases in ecotourism to previously remote locations,
72 growing familiarity of these animals with people, and inappropriate behaviour of people toward
73 them (possibly in itself evidence of the growing extinction of experience). Other conflicts resulting
74 from direct interactions are doubtless rife, with consequences that range from severe (e.g.,
75 emerging infectious diseases, snake bites, vector-borne disease transmission; e.g., [17,18]) to
76 inconvenient (e.g., noise nuisance, mess and mild aggression; e.g., [19,20]). Management of these
77 interactions would often be improved by better understanding how they arise and with what
78 regularity.

79

80 To address this demand in a more coherent manner, we propose the need for a 'personalised
81 ecology' that is distinguished by its focus on the direct interactions between individual people and
82 nature. In this opinion article, we offer a definition of personalised ecology, suggest a framework of
83 research topics on which personalised ecology should focus, and highlight the connections of
84 personalised ecology to other aspects of ecology.

85

86 **Personalised Ecology**

87 We define personalised ecology as the investigation of the direct interactions between individual
88 people and nature and their ecological dimensions. We define nature to span individual living
89 organisms to ecosystems, but to exclude organisms that are not self-sustained (e.g., crops, house
90 plants, zoo and domesticated animals); we acknowledge that whilst a broadly understood
91 distinction between these two groups is achievable, a precise and uniformly agreed one is
92 challenging. A human-nature interaction is then a particular instance of an individual person being
93 present in the 'same space' as nature or perceiving a stimulus from nature (through sight, sound,
94 smell, taste or touch – although in practice sight and sound tend to predominate). This might be
95 the ecosystems that they experience, the species that they encounter, or the individual organisms
96 they see or hear. Such an interaction could occur intentionally or unintentionally and consciously or
97 unconsciously. To a greater or lesser extent unconscious experiences are likely to be occurring for
98 much of the time that people are outdoors.

99

100 A definition of this breadth allows inclusion of a wide range of types of human-nature interactions,
101 such as visiting urban greenspaces or national parks, viewing trees through a window, listening to
102 bird song, and being bitten by mosquitoes. It excludes interactions with nature through the media
103 (e.g., through books, television, websites), albeit these interactions can have positive outcomes for
104 humans (e.g., [21,22]).

105

106 The focus of personalised ecology is on the ecological dimensions of human-nature interactions,
107 recognising that other important dimensions are not ecological and more relevant to other fields

108 (e.g., medicine, public health, environmental education). We will also exclude for present purposes
109 consideration of organisms that live on or in people, whilst recognising this can be a legitimate
110 topic of ecological enquiry.

111

112 One can view personalised ecology from two perspectives; first, from that of the person, and
113 second from that of nature. Whilst the fundamental unit of study remains the individual person, one
114 can consider both of these perspectives at the level of a single person, a population of people or
115 across multiple human populations (Figure 1). We will address each of these six combinations in
116 turn.

117

118 ***Single Person, Human Perspective***

119 Arguably at its most reductionist, personalised ecology considers the nature that is experienced by
120 a single individual person over a defined period. The vast majority of studies to date have simply
121 assumed that characterization of the environs in which people live, or of the places that they visit
122 (e.g., public parks, protected areas), captures their experience [23]. In the main, even this has
123 been done quite crudely, typically using measures of the extent of green landcover (e.g., [24,25]),
124 although some studies have sought to characterise the abundance or diversity of taxa in these
125 environs or places (e.g., [26-28]). Undoubtedly, the actual nature interactions of people may be
126 very different from what has typically been measured (e.g., [20,29]).

127

128 A key research focus of personalised ecology will need to be on understanding how (e.g.,
129 passively or actively) and what type of nature people are experiencing, and how these experiences
130 are influenced by personal characteristics (e.g., gender, age, observer knowledge, skills and
131 behavioural preferences) and by the physical/environmental conditions under which these nature
132 interactions occur (e.g., time of day, seasonality, weather). Whilst some of these factors
133 (particularly observer skills) have been investigated in attempts to understand the impacts on
134 biodiversity monitoring schemes, the extension of these studies to a much broader cross-section of
135 people and factors has been limited [27,30]. Nonetheless, it has, for example, been shown that
136 ecological knowledge can be important in shaping people's nature experiences (e.g., [31]). The

137 continuing rapid advancement of personal monitoring devices (e.g., eye-tracking glasses, GPS
138 trackers, electroencephalography (EEG), acoustic recorders) will enable much improved
139 characterisation of the nature that people encounter and how this varies.

140

141 ***Single Person, Nature Perspective***

142 If we know which components of nature an individual person is interacting with, then we can ask
143 how these relate to the nature that is potentially available for such experiences. The occurrence
144 and relative frequency of interactions will almost invariably be a non-random subset of those
145 available. For example, abundances of bird species apparent even to a trained observer will often
146 be far less than those actually present (e.g., Figure 2). The numbers of birds that untrained people
147 see and hear as they move around the landscape is likely to be significantly lower [32]. Such
148 differences can arise for a diverse array of reasons, the unpicking of which may be important.
149 These will include the actual distribution and abundance of species, their appearance and
150 behaviour, their response to people (e.g., flight initiation distances, changes in calls), the timings of
151 activities (e.g., daily and seasonal activity patterns, annual migration), and perceptions of where
152 individuals are. Most obviously, people are more likely to interact with species that are common,
153 diurnal, apparent (e.g., large, active, vocal), accustomed to people, and that can be attracted to
154 their vicinity (e.g., through resources such as bird feeders, nest boxes).

155

156 ***Single Population, Human Perspective***

157 Within a human population, nature experiences will vary between individuals in their composition,
158 frequency and duration. Particularly in towns and cities, those having regular nature experiences,
159 or ones of long duration, tend to be rare. A study in the U.K. found that three-quarters of direct
160 nature interactions (instances where people were present in nature) were experienced by just one
161 third of an urban population [33]. As more detailed data on the nature experiences of individual
162 people become easier to collect then so will comparisons between people. Two major sets of
163 factors have been proposed to influence the frequency and duration of human-nature interactions.
164 The first is the opportunity to experience nature, which is particularly shaped by the ease of access
165 to greenspace within the local environs [34]. This can depend heavily on people's socioeconomic

166 circumstances. These strongly determine the kinds and location of the properties that they inhabit,
167 and hence the availability and biodiversity of associated greenspaces [35-37], whether they can
168 invest in green infrastructure [38] and activities to attract wildlife to those environs [39], and also
169 whether they can engage in ecotourism elsewhere. The second influence on the frequency and
170 duration of human-nature interactions is the orientation (or preferences) of people towards
171 exploiting these opportunities. Although more attention has been paid to opportunity in discussions
172 of the design of urban green infrastructure, there is evidence that orientation may be more
173 important in shaping nature experiences [40]. These two tend to be correlated, with people living in
174 greener areas with increased opportunity to experience everyday nature, also having a greater
175 orientation towards doing so [41].

176

177 ***Single Population, Nature Perspective***

178 Different areas and different individual organisms will contribute very differently to the nature
179 experiences of a given human population. Some areas will be visited by many people, others by
180 few or none. This issue is presently best understood with regards to urban greenspaces and
181 protected areas, where human footfall has been measured and associated with their ecological
182 (e.g., [42]) or geographical (e.g., [40]) features. However, it remains challenging to disentangle the
183 influence of the wide array of possible features that may determine whether areas are visited, how
184 often, for how long, and with what consequences for nature experiences and for the management
185 of sites (e.g., to encourage or direct access both to enhance nature experiences and mitigate
186 impacts on wildlife). These include the sizes of areas, their accessibility, their vegetational
187 complexity (e.g., evidence that people prefer 'savannah-like' natural spaces), the presence or
188 absence of key species (e.g., large mammals), and the occurrence of wildlife spectacles. The
189 numbers of people visiting an area will impact their individual nature experiences, due to an
190 increase in numbers of observers (and hence what wildlife is located) and in the disturbance
191 resulting from their activities.

192

193 Equally, there will be great variation in how species and individual organisms interact with the
194 human population. Some individual organisms will interact with many people, others with few or

195 none (e.g., for many years a single black-winged stilt *Himantopus himantopus*, resident on a
196 protected area in Norfolk, U.K., was held to have been watched by more people than any other
197 bird in the country; [43]). These experiences will be further influenced by interactions between
198 species, which increase the probability that the organisms will encounter people or provide a more
199 interesting spectacle. Improvements in remote sensing data and tracking technology have begun
200 to enable evaluation of how individual organisms contribute to nature experiences [44]. In urban
201 areas in particular, those mobile individuals that move between a greater number of greenspaces,
202 are likely to be seen by more people (Figure 3a). Similarly, individuals of those stationary
203 organisms (e.g., trees) that are readily visible, such as besides roadsides, will be experienced by
204 more people than others of the same or similar species (Figure 3b).

205

206 ***Multiple populations, human perspective***

207 There will inevitably be differences in nature experiences of people in different populations, such
208 as different villages, towns and cities. What will be particularly important to understand is the
209 macroecology of such variation – how the frequency, duration and composition of interactions
210 change over large spatial and temporal scales. As with variation within single populations,
211 opportunity and orientation will be significant, with cultural, socioeconomic and environmental
212 differences likely to play profound roles in shaping how people in different populations use their
213 natural environment (e.g. [45]). However, little is known about these patterns, with the majority of
214 studies limited to westernised countries (e.g. [46]), and so the findings may have limited generality.
215 For example, whilst in these usually temperate zones vegetation around the home is often seen as
216 associated with human well-being benefits and to be encouraged (at least where it does not pose a
217 fire risk), in many tropical areas it can harbour species dangerous to human health and is often
218 cleared.

219

220 Even focussing on quite narrow issues, approaches to nature experiences may be very different
221 across the world. This is well illustrated with regard to attitudes toward providing supplementary
222 food for birds and mammals in urban areas. In some parts of Europe and North America the
223 practice, often to increase the likelihood of viewing them, is widespread, and indeed is the basis for

224 a substantial industry (e.g., [46]). In Australia, it is much less favoured, in part because it is seen to
225 encourage alien or unwelcome species (e.g., [47]). In much of the rest of the world, such feeding
226 activities are virtually unknown [48].

227

228 ***Multiple Populations, Nature Perspective***

229 When contrasting the nature experiences of multiple human populations it seems logical to ask to
230 what extent it is the same or analogous components of nature (e.g., the same species or species
231 that have similar traits and ecologies) that are contributing. Such studies will be akin to those in
232 urban ecology that have attempted to characterise the similarities and dissimilarities of species
233 assemblages found in different towns and cities, albeit in this case without explicit reference to
234 their contribution to human-nature experiences (e.g., [49]). In the main, it seems likely that species
235 or groups of species that occupy similar niches in different cities will provide similar kinds of nature
236 experiences to people. However, there are clear cases where quite different species fulfil the same
237 role, with, for example, urban bird feeding tending to focus in some regions on granivorous species
238 and in others on nectivorous ones [48].

239

240 When looking across human populations one can start to map the spatial distribution of nature
241 interactions, which will often be different from the underlying distributions of the species
242 concerned. The distribution across Britain of the Magpie *Pica pica*, as recorded by citizen scientists
243 is, for example, very different from that documented by formal ornithological mapping schemes
244 (Figure 4). Unsurprisingly, the former highlights encounters along major transport routes and in
245 major centres of population, as these are the places in which the vast majority of nature
246 experiences actually occur, while the latter reveal many areas in which the species occurs but
247 interactions are more limited.

248

249 **Linkages**

250 Obviously, personalised ecology is not divorced from a number of other topics of focal interest in
251 ecology. In addition to those already observed above to motivate the need for such an agenda,
252 these include:

253

254 ***Biodiversity monitoring***

255 While biodiversity monitoring has been focussed principally on understanding the relationship
256 between the actual abundances and distributions of species and what expert observers detect,
257 personalised ecology is less concerned with these actual quantities and more with the abundances
258 and distributions experienced by people, and with a focus on 'ordinary' people (i.e., non-experts,
259 and often with a limited knowledge of ecology), and experiences during everyday activities. The
260 growing use of **citizen science** in biodiversity monitoring makes the concerns of personalised
261 ecology increasingly relevant.

262

263 ***Ecosystem services***

264 Whilst the topic of ecosystem services is explicitly concerned with the benefits that people gain
265 from ecosystems [52] rather than emphasising personal nature interactions in the main this is
266 approached in a generic sense of community or societal benefits (e.g., from agricultural production,
267 pollination, carbon sequestration, waste decomposition). The two approaches are obviously
268 complementary, with the ecosystem benefits to individual people often becoming very apparent in
269 terms of cultural ecosystem services (e.g., recreational, sense of place, aesthetic, educational and
270 therapeutic values).

271

272 ***Urban ecology***

273 The bulk of urban ecology research remains focussed on a quite traditional understanding of the
274 determinants of the abundance and distribution of species and the interactions that determine
275 these, albeit in urban areas [53]. Nonetheless, there have been repeated calls for, and important
276 contributions toward, broader approaches (e.g., [54,55]), and particularly those that address the
277 complex interplay between people and urban ecosystems. Personalised ecology would clearly
278 contribute to such an agenda.

279

280

281 ***Human ecology***

282 The field of human ecology studies the relationship between humans and their environment, and
283 typically has a strong emphasis on the anthropological, social or political dimensions to this
284 interaction [56]. Personalised ecology would again serve to add an important dimension to such
285 investigation, by strengthening the links to more conventionally ecological concerns.

286

287 **Implications**

288 A well-developed understanding of personalised ecology would have major practical
289 consequences in two primary arenas. First, it would improve the ability to design policy and
290 management for people's access to nature in such a way that their benefits, the positive
291 interactions, were enhanced and their costs, the negative interactions, were reduced. Second, and
292 more importantly in the face of a global biodiversity crisis, well developed understanding of
293 personalised ecology would improve the ability to determine policy and management of people's
294 interactions with nature in such a way that the benefits to nature were also increased and the costs
295 minimised. Of course, these two arenas interact, and what is presently lacking is a strongly
296 evidence based approach for encouraging the positive engagement of people with nature, whilst
297 promoting the conservation of populations and ecosystems.

298

299 **Concluding Remarks**

300 The global human population is continuing to grow rapidly and become more urbanised, with
301 people less likely to experience regular positive interactions with nature. At the same time, the
302 importance of those interactions to human well-being is becoming increasingly apparent. It thus
303 seems vitally important that ecologists develop a much more comprehensive and detailed
304 understanding of those interactions, their composition, and temporal and spatial dynamics. Such a
305 'personalised ecology' constitutes a challenging agenda, and one that has thus far lagged far
306 behind others in the field of ecology.

307

308

309 **Acknowledgements**

310 We are grateful to I. Geipel, R. Inger and two anonymous reviewers for comments on the
311 manuscript, and to A. Smalley for input to figure 3. This work was supported by the “Fragments,
312 Functions and Flows in Urban Ecosystems” project (NE/J015237/1) funded under the NERC
313 Biodiversity and Ecosystem Services Sustainability (BESS) thematic program, and the Green
314 Infrastructure for Growth project funded through the European Regional Development Fund.

315

316 **References**

- 317 1. Begon, M. *et al.* (2006) *Ecology: from individuals to ecosystems*, Fourth edition, Blackwell
318 Publishing
- 319 2. Buckland, S.T. *et al.* (2004) *Advanced Distance Sampling: estimating abundance of biological*
320 *populations*, Oxford University Press
- 321 3. Chen, G. *et al.* (2012) Imperfect detection is the rule rather than the exception in plant
322 distribution studies. *J. Ecol.* 101, 183-191
- 323 4. Garrard, G.E. *et al.* (2012) A general model of detectability using species traits. *Methods Ecol.*
324 *Evol.* 4, 45-52.
- 325 5. Kellner, K.F. and Swihart, R.K. (2014) Accounting for imperfect detection in ecology: a
326 quantitative review. *PLoS One* 9, e111436
- 327 6. Keniger, L.E. *et al.* (2013) What are the benefits of interacting with nature? *Int. J. Environ. Res.*
328 *Public Health* 10, 913-935
- 329 7. Dye, C. (2008) Health and urban living. *Science* 319, 766-769
- 330 8. Hartig, T. *et al.* (2014) Nature and health. *Annu. Rev. Public Health* 35, 207-228
- 331 9. Shanahan, D.F. *et al.* (2015) Toward improved public health outcomes from urban nature. *Am.*
332 *J. Public Health* 105, 470-477
- 333 10. Shanahan, D.F. *et al.* (2015) The health benefits of urban nature: how much do we need?
334 *BioScience* 65, 476-485
- 335 11. Soga, M. and Gaston, K.J. (2016) Extinction of experience: the loss of human-nature
336 interactions. *Front. Ecol. Environ.* 14, 94-101
- 337 12. Pyle, R.M. (1993) *The Thunder Tree: lessons from an urban wildland*, Houghton Mifflin

- 338 13. Soga, M. and Gaston, K.J. (2018) Shifting baseline syndrome: causes, consequences and
339 implications. *Front. Ecol. Environ.* 16, 222-230
- 340 14. Barua, M. *et al.* (2013) The hidden dimensions of human-wildlife conflict: health impacts,
341 opportunity and transaction costs. *Biol. Conserv.* 157, 309-316
- 342 15. Soulsbury, C.D. and White, P.C.L. (2015) Human-wildlife interactions in urban areas: a review
343 of conflicts, benefits and opportunities. *Wildlife Res.* 42, 541-553
- 344 16. Penteriani, V. *et al.* (2016) Human behaviour can trigger large carnivore attacks in developed
345 countries. *Sci. Rep.* 6, 20552
- 346 17. Kasturiratne, A. *et al.* (2008) The global burden of snakebite: A literature analysis and
347 modelling based on regional estimates of envenoming and deaths. *PLoS Med.* 5, e218
- 348 18. Allen, T. *et al.* (2017) Global hotspots and correlates of emerging zoonotic diseases. *Nature*
349 *Comm.* 8, 1124
- 350 19. Rock, P. (2005) Urban gulls: problems and solutions. *British Birds* 98, 338-355
- 351 20. Belaire, J.A. *et al.* (2015) Urban residents' perceptions of birds in the neighborhood:
352 biodiversity, cultural ecosystem services, and disservices. *The Condor* 117, 192-202
- 353 21. Depledge *et al.* (2011) Can natural and virtual environments be used to promote improved
354 human health and wellbeing? *Environ. Sci. Technol.* 45, 4660-4665
- 355 22. Nadkarni, N.M. *et al.* (2017) Impacts of nature imagery on people in severely nature-deprived
356 environments. *Front. Ecol. Environ.* 15, 395-403
- 357 23. WHO [World Health Organisation]. (2016) *Urban Green Spaces and Health: a review of*
358 *evidence*, WHO Regional Office in Europe
- 359 24. de Vries, S. *et al.* (2003) Natural environments -- healthy environments? An exploratory
360 analysis of the relationship between greenspace and health. *Environ. Plann. A* 35, 1717-1731
- 361 25. Shanahan, D.F. *et al.* (2014) Socio-economic inequalities in access to nature on public and
362 private lands: A case study from Brisbane, Australia. *Landscape Urban Plann.* 130, 14-23.
- 363 26. Fuller, R.A. *et al.* (2007) Psychological benefits of greenspace increase with biodiversity.
364 *Biology Lett.* 3, 390-394
- 365 27. Dallimer, M. *et al.* (2012) Biodiversity and the feel-good factor: understanding associations
366 between self-reported human well-being and species richness. *BioScience* 62, 46-55

- 367 28. Cox, D.T.C. *et al.* (2017) Doses of neighborhood nature: the benefits for mental health of living
368 with nature. *BioScience* 67, 147-155
- 369 29. Arbieu, U. *et al.* (2017) The importance of vegetation density for tourists' wildlife viewing
370 experience and satisfaction in African savannah ecosystems. *PLoS One* 12, e0185793
- 371 30. Cox, D.T.C. and Gaston, K.J. (2015) Likeability of garden birds: importance of species
372 knowledge and richness in connecting people to nature. *PLoS One* 10, e0141505
- 373 31. Muratet, A. *et al.* (2015) Perception and knowledge of plant diversity among urban park users.
374 *Landscape Urban Plann.* 137, 95-106
- 375 32. Clergeau, P. *et al.* (2001) Human perception and appreciation of birds: A motivation for wildlife
376 conservation in urban environments of France. In *Avian Ecology and Conservation in an*
377 *Urbanizing World* (Marzluff, J.M. *et al.*, eds), pp.69-88. Kluwer Academic, Boston
- 378 33. Cox, D.T.C. *et al.* (2017) The rarity of direct experiences of nature in an urban population.
379 *Landscape Urban Plann.* 160, 79-84
- 380 34. Soga, M. *et al.* (2015) Reducing the extinction of experience: association between urban form
381 and recreational use of public greenspace. *Landscape Urban Plann.* 143, 69-75
- 382 35. Martin, C.A. *et al.* (2004) Neighborhood socioeconomic status is a useful predictor of perennial
383 landscape vegetation in residential neighborhoods and embedded small parks of Phoenix, AZ.
384 *Landscape Urban Plann.* 69, 355-368
- 385 36. Luck, G.W. *et al.* (2009) Socio-economics and vegetation change in urban ecosystems:
386 patterns in space and time. *Ecosystems* 12, 604-620
- 387 37. Ferguson, M. *et al.* (2018) Contrasting distributions of urban green infrastructure across social
388 and ethno-racial groups. *Landscape Urban Plann.* 175, 136-148
- 389 38. Avolio, M.L. *et al.* (2018) Biodiverse cities: the nursery industry, homeowners, and
390 neighborhood differences drive urban tree composition. *Ecol. Monogr.* 88, 259-276
- 391 39. van Heezik, Y. and Hight, S.R. (2018) Socio-economic-driven differences in bird-feeding
392 practices exacerbate existing inequities in opportunities to see native birds in cities. *J. Urban*
393 *Ecol.*, in press
- 394 40. Lin, B.B. *et al.* (2014) Opportunity or orientation?: Who uses urban parks and why. *PLoS One*
395 9, e87422

- 396 41. Shanahan, D.F. *et al.* (2017) Variation in experiences of nature across gradients of tree cover
397 in compact and sprawling cities. *Landscape Urban Plann.* 157, 231-238
- 398 42. Shanahan, D.F. *et al.* (2015) What is the role of trees and remnant vegetation in attracting
399 people to urban parks? *Landscape Ecol.* 30, 153-165
- 400 43. Eele, P. (2015) Gone but never forgotten.
401 [https://ww2.rspb.org.uk/community/placestovisit/titchwellmarsh/b/titchwellmarsh-
402 blog/archive/2015/05/21/gone-but-never-forgotten.aspx](https://ww2.rspb.org.uk/community/placestovisit/titchwellmarsh/b/titchwellmarsh-
402 blog/archive/2015/05/21/gone-but-never-forgotten.aspx)
- 403 44. Cox, D.T.C. *et al.* (2016) Movement of feeder-using songbirds: the influence of urban features.
404 *Sci. Rep.* 6, 37669
- 405 45. Buijs, A.E. *et al.* (2009) No wilderness for immigrants: Cultural differences in images of nature
406 and landscape preferences. *Landscape Urban Plann.* 91, 113-123.
- 407 46. Jones, D. (2011) An appetite for connection: why we need to understand the effect and value
408 of feeding wild birds. *Emu* 111, i-vii
- 409 47. Parsons, H. *et al.* (2006) Species interactions and habitat associations of birds inhabiting urban
410 areas of Sydney, Australia. *Austral. Ecol.* 31, 217-227
- 411 48. Reynolds, S.J. *et al.* (2017) Garden bird feeding: insights and prospects from a North-South
412 comparison of this global urban phenomenon. *Front. Ecol. Evol.* 5, 1-24
- 413 49. Ferenc, M. *et al.* (2014) Are cities different? Patterns of species richness and beta diversity of
414 urban bird communities and regional species assemblages in Europe. *Global Ecol. Biogeogr.* 23,
415 479-489
- 416 50. Balmer, D.E. *et al.* (2013) *Bird Atlas 2007-11: the breeding and wintering birds of Britain and
417 Ireland*, BTO Books
- 418 51. Teacher, A.G.F. *et al.* (2013) Smartphones in ecology and evolution: a guide for the app-
419 rehensive. *Ecol. Evol.* 3, 5268-5278
- 420 52. Millennium Ecosystem Assessment (2005) *Ecosystems and Human Well-Being: Current State
421 and Trends, Vol. I*, Island Press
- 422 53. Gaston, K.J. (ed.) (2010) *Urban Ecology*, Cambridge University Press
- 423 54. McDonnell, M.J. and MacGregor-Fors, I. (2016) The ecological future of cities. *Science* 352,
424 936-938

- 425 55. McPhearson, T. *et al.* (2016) Advancing urban ecology toward a science of cities. *BioScience*
426 66, 198-212
- 427 56. Knapp, G. (2007) Human ecology. In *Encyclopedia of Environment and Society, Volume 3*
428 (Robbins, P., ed.), pp. 880-884, Sage Publications
429

430 **Figure 1. Schematic of the different perspectives of personalised ecology.** Personalised
431 ecology can be considered from the perspectives of the person or of nature (arrows), and at
432 different levels, namely a single person (top), a population of people (middle) or across multiple
433 human populations (bottom). The circles represent the (overlapping) components of nature that an
434 individual person, different people within a population or people within different populations interact
435 with. Note the organisms and combinations are for illustrative purposes only.

436

437 **Figure 2. Example variation in the ratio of estimated actual bird abundance to observed bird**
438 **abundance]**). 420 bird surveys (data from [28]) were conducted across three towns in Southern
439 England, U.K. Each town was divided into 500 x 500m tiles in a grid, with 106 tiles being surveyed.
440 Surveys, conducted by trained observers, comprised two early-morning ten-minute point counts at
441 up to four survey points (mean per tile, 3.91 ± 0.32 SD). Actual abundances, adjusted for detection
442 probability were then estimated from observed abundances using distance sampling (see [28] for
443 detailed description of the methodology). The observed and adjusted abundances presented here
444 are per survey point. Icon provided by *Freepik* via www.flaticon.com.

445

446 **Figure 3. Example of variation in the provision of nature experiences contributed by**
447 **different individual organisms.** By moving between bird feeders in multiple gardens, bird A has
448 the potential to be seen by more households, and thus provide nature experiences to more people
449 than bird B, which visits only one feeder. [44] attached Radio Frequency Identification Receivers to
450 20 bird feeders in an equal number of gardens in three neighbourhoods in southern England ($n =$
451 60). They show the number of domestic gardens that songbirds carrying a Passive Integrated
452 Transponder ($n = 348$) visited over a 12-month period. Icons provided by *Freepik* and
453 *Smashicons* via www.flaticon.com.

454

455 **Figure 4.** Differences between (A) the relative abundance of a common, visible and regionally
456 well-known bird species, the Magpie *Pica pica*, and (B) where people interact with this species. (A)
457 is the breeding abundance map from the Bird Atlas 2007-11 [50], which is a joint project between
458 the British Trust for Ornithology (BTO), Bird Watch Ireland and the Scottish Ornithologists Club

459 (reproduced with permission from the BTO). Data were collected through ornithological volunteers
460 carrying out bird counts in at least eight 2km² areas, within each 10km² square across the U.K. (B)
461 is a record of sightings collected in 2013-2014 by a much wider range of people whilst about their
462 daily lives using the Magpie Mapper App [51]. Eye icon provided by *Freepik* via www.flaticon.com.

463 **Glossary**

464

465 **Biodiversity monitoring:** tracking the changes in the state of biodiversity.

466

467 **Citizen science:** scientific research conducted by those who are not professional scientists.

468

469 **Ecosystem services:** the benefits that people gain from the natural environment.

470

471 **Extinction of experience:** progressive loss of daily interactions between people and nature.

472

473 **Human ecology:** the study of the relationship between humans and their environment.

474

475 **Human-wildlife conflict:** interactions between people and wildlife that result in harm to either.

476

477 **Urban ecology:** the study of the abundance and distribution of organisms in urban environments.

478

479