

CORRECTION

Open Access



Correction to: Residual feed intake phenotype and gender affect the expression of key genes of the lipogenesis pathway in subcutaneous adipose tissue of beef cattle

Clare McKenna^{1,2}, Richard K. Porter², Kate A. Keogh¹, Sinead M. Waters¹, Mark McGee¹ and David A. Kenny^{1*}

Correction

In the original publication of this article [1], some errors in Table 4 need to be corrected as below:

“Creating, umil/L” should be “Creatine, umol/L”

“Glucose, nmlo/L” should be “Glucose, nmol/L”

“Urea, nmo/L” should be “Urea, nmol/L”

The original publication has been corrected.

Author details

¹Animal and Bioscience Research Department, Teagasc Grange, Dunsany, Meath C15 PW93, Ireland. ²School of Biochemistry & Immunology, Trinity College Dublin, Dublin 2 D02 R590, Ireland.

Received: 19 October 2018 Accepted: 22 October 2018

Published online: 07 November 2018

Reference

1. McKenna, et al. Residual feed intake phenotype and gender affect the expression of key genes of the lipogenesis pathway in subcutaneous adipose tissue of beef cattle. *J Anim Sci Biotechnol.* 2018;9:68 <https://doi.org/10.1186/s40104-018-0282-9>.

* Correspondence: david.kenny@teagasc.ie

¹Animal and Bioscience Research Department, Teagasc Grange, Dunsany, Meath C15 PW93, Ireland

Full list of author information is available at the end of the article

