

7-Chloro-4-(2-hydroxyethylamino)-quinolin-1-i um chloride

Ivson L. Gama,^a Marcus V. N. de Souza,^a James L. Wardell^{a,b}‡ and Edward R. T. Tiekkink^{c*}

^aFundação Oswaldo Cruz, Instituto de Tecnologia em Fármacos—Farmanguinhos, R. Sizenando Nabuco, 100, Manguinhos, 21041-250, Rio de Janeiro, RJ, Brazil,

^bChemistry Department, University of Aberdeen, Old Aberdeen, AB24 3UE, Scotland, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: edward.tiekkink@gmail.com


Received 27 February 2014; accepted 27 February 2014

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.030; wR factor = 0.076; data-to-parameter ratio = 16.8.

In the title salt, $\text{C}_{11}\text{H}_{12}\text{ClN}_2\text{O}^+\cdot\text{Cl}^-$, the ten non-H atoms comprising the quinolinium residue are coplanar (r.m.s. deviation = 0.041 Å) and the hydroxyethyl group is approximately perpendicular to this plane [$\text{C}_{\text{ring}}-\text{N}-\text{C}_{\text{methylene}}-\text{C}$ torsion angle = $-74.61(18)^\circ$]. A supramolecular chain aligned along [101] mediated by charge-assisted O/N—H \cdots Cl $^-$ hydrogen bonds features in the crystal packing. Chains are connected into a three-dimensional architecture by C—H \cdots O(hydroxy) interactions.

Related literature

For the wide range of pharmacological activities of synthetic and natural products containing the quinoline nucleus, see: Andrade *et al.* (2007); Cunico *et al.* (2006); Font *et al.* (1997); Kaminsky & Meltzer (1968); Musiol *et al.* (2006); Nakamura *et al.* (1999); Sloboda *et al.*, (1991); de Souza *et al.* (2014); Tanenbaum & Tuffanelli (1980); Warshakoon *et al.* (2006). For the crystal structures of related 4-RN(H)-7-chloroquinolines, see: Kaiser *et al.*, (2009).



‡ Additional correspondence author, e-mail: j.wardell@abdn.ac.uk.

Experimental

Crystal data

$\text{C}_{11}\text{H}_{12}\text{ClN}_2\text{O}^+\cdot\text{Cl}^-$
 $M_r = 259.13$
Monoclinic, $P2_1/c$
 $a = 8.2438(13)\text{ \AA}$
 $b = 16.405(2)\text{ \AA}$
 $c = 8.8561(14)\text{ \AA}$
 $\beta = 110.705(2)^\circ$

$V = 1120.3(3)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.56\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.20 \times 0.07 \times 0.04\text{ mm}$

Data collection

Rigaku R-AXIS conversion diffractometer
Absorption correction: multi-scan (*CrystalClear-SM Expert*; Rigaku, 2013)
 $T_{\min} = 0.831$, $T_{\max} = 1.000$

7784 measured reflections
2581 independent reflections
2162 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.076$
 $S = 1.07$
2581 reflections
154 parameters
3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.37\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.24\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1o \cdots Cl2	0.84 (2)	2.30 (2)	3.1338 (14)	179 (2)
N1—H1n \cdots Cl2 ⁱ	0.88 (1)	2.29 (1)	3.1602 (15)	168 (1)
N2—H2n \cdots Cl2 ⁱⁱ	0.87 (1)	2.49 (2)	3.2949 (14)	154 (2)
C2—H2 \cdots O1 ⁱⁱⁱ	0.95	2.60	3.545 (2)	173

Symmetry codes: (i) $-x + 2, -y + 1, -z + 1$; (ii) $-x + 1, -y + 1, -z$; (iii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *CrystalClear-SM Expert* (Rigaku, 2013); cell refinement: *CrystalClear-SM Expert*; data reduction: *CrystalClear-SM Expert*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The use of the EPSRC X-ray crystallographic service at the University of Southampton, England (Coles & Gale, 2012), and the valuable assistance of the staff there is gratefully acknowledged. JLW acknowledges support from CAPES (Brazil). Structural studies are supported by the Ministry of Higher Education (Malaysia) and the University of Malaya through the High-Impact Research scheme (UM-C/HIR/MOHE/SC/3).

Supporting information for this paper is available from the IUCr electronic archives (Reference: HG5387).

References

- Andrade, A. A., Varotti, M. F. D., de Freitas, I. Q., de Souza, M. V. N., Vasconcelos, T. R. A., Boechat, N. & Krettli, A. U. (2007). *Eur. J. Pharm.* **558**, 194–196.

- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Coles, S. J. & Gale, P. A. (2012). *Chem. Sci.* **3**, 683–689.
- Cunico, W., Cechinel, C. A., Bonacorso, H. G., Martins, M. A. P., Zanetta, N., de Souza, M. V. N., Freitas, I. Q., Soares, R. P. P. & Krettli, A. U. (2006). *Bioorg. Med. Chem. Lett.* **16**, 649–653.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Font, M., Monge, A., Ruiz, I. & Heras, B. (1997). *Drug Des. Disc.* **14**, 259–272.
- Kaiser, C. R., Pais, K. C., de Souza, M. V. N., Wardell, J. L., Wardell, S. M. S. V. & Tieckink, E. R. T. (2009). *CrystEngComm*, **11**, 1133–1140.
- Kaminsky, D. & Meltzer, R. I. (1968). *J. Med. Chem.* **11**, 160–163.
- Musiol, R., Jampilek, J., Buchta, V., Silva, L., Halina, H., Podeszwa, B., Palka, A., Majerz-Maniecka, K., Oleksyn, B. & Polanski, J. (2006). *Bioorg. Med. Chem.* **14**, 3592–3598.
- Nakamura, T., Oka, M., Aizawa, K., Soda, H., Fukuda, M., Terashi, K., Ikeda, K., Mizuta, Y., Noguchi, Y., Kimura, Y., Tsuruo, T. & Kohno, S. (1999). *Biochem. Biophys. Res. Commun.* **255**, 618–624.
- Rigaku (2013). *CrystalClear-SM Expert*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sloboda, A. E., Powell, D., Poletto, J. F., Pickett, W. C., Gibbons, J. J., Bell, D. H., Oronsky, A. L. & Kerwar, S. S. (1991). *J. Rheumatol.* **18**, 855–860.
- Souza, M. V. N. de, Goncalves, R. S. B., Rodrigues, F. A. R., Cavalcanti, B. C., Bomfim, I. S., Pessoa, C. O., Wardell, J. L. & Wardell, S. M. S. V. (2014). *Chem. Biol. Drug Des.* **83**, 126–131.
- Tanenbaum, L. & Tuffanelli, D. L. (1980). *Arch. Dermatol.* **116**, 587–591.
- Warshakoon, N. C., Sheville, J., Bhatt, R. T., Ji, W., Mendez-Andino, J. L., Meyers, K. M., Kim, N., Wos, J. A., Mitchell, C., Paris, J. L., Pinney, B. B., Reizes, O. & Hu, X. E. (2006). *Bioorg. Med. Chem. Lett.* **16**, 5207–5211.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2014). E70, o385–o386 [doi:10.1107/S1600536814004565]

7-Chloro-4-(2-hydroxyethylamino)quinolin-1-i um chloride

Ivson L. Gama, Marcus V. N. de Souza, James L. Wardell and Edward R. T. Tiekink

S1. Experimental

S1.1. Synthesis and crystallization

A solution of 7-chloro-4-(2-hydroxyethylamino)quinoline (1 mmol) and $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (1 mmol) in EtOH (25 ml) was refluxed for 30 min. On leaving the reaction mixture at room temperature, crystals of the title compound were formed, M.pt: 538–541 K (dec.).

S2. Refinement


Intensity data was collected at the National Crystallographic Service, England (Coles & Gale, 2012). The C-bound H atoms were geometrically placed ($\text{C}—\text{H} = 0.95\text{--}0.99 \text{\AA}$) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The O- and N-bound H atoms were located from a difference map and refined with $\text{O}—\text{H} = 0.84\pm0.01 \text{\AA}$ and $\text{N}—\text{H} = 0.88\pm0.01 \text{\AA}$, respectively, and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ and $1.2U_{\text{eq}}(\text{N})$, respectively.

S3. Results and discussion


The quinoline nucleus is found in many synthetic and natural products having a wide range of pharmacological activities, such as anti-viral (Font *et al.*, 1997), anti-cancer (Nakamura *et al.*, 1999; de Souza *et al.*, 2014), anti-bacterial (Kaminsky & Meltzer, 1968), anti-malarial (Tanenbaum & Tuffanelli, 1980; Cunico *et al.*, 2006; Andrade *et al.*, 2007), anti-fungal (Musiol *et al.*, 2006), anti-obesity (Warshakoon *et al.*, 2006) and anti-inflammatory (Sloboda *et al.*, 1991) activities. The crystal structures of a series of 4-RN(H)-7-chloro-quinolines was recently reported (Kaiser *et al.*, 2009). We now wish to report the crystal structure of the HCl salt of 4-(HOCH₂CH₂NH)-7-chloro-quinoline, (I), obtained serendipitously from an attempted reaction to generate an iron complex.

The components of salt (I) are illustrated in Fig. 1. The 10 non-hydrogen atoms comprising the quinolinium residue are co-planar with a r.m.s. deviation of 0.041 Å. The hydroxyethyl group is almost perpendicular to this plane as seen in the C3—N2—C10—C11 torsion angle of -74.61 (18)°.


In the crystal structure, charge-assisted O, N—H···Cl⁻ hydrogen bonds, Table 1, lead to a supramolecular chain aligned along [1 0 1], Fig. 2. These are connected into a three-dimensional architecture by methylene-C—H···O(hydroxyl) interactions, Fig. 3 & Table 1.

**Figure 1**

The molecular structures of the ions in (I) showing the atom-labelling scheme and displacement ellipsoids at the 70% probability level.

**Figure 2**

A view of the supramolecular chain along $[1\ 0\ 1]$ in (I) showing O—H \cdots Cl $^-$ and N—H \cdots Cl $^-$ hydrogen bonds as orange and blue dashed lines, respectively.

**Figure 3**

A view in projection down [1 0 1], the direction of the chain shown in Fig. 2, of the unit-cell contents of (I). The O—H···Cl, N—H···Cl and C—H···O interactions are shown as orange, blue and green dashed lines, respectively.

7-Chloro-4-(2-hydroxyethylamino)quinolin-1-i um chloride

Crystal data

$C_{11}H_{12}ClN_2O^+\cdot Cl^-$
 $M_r = 259.13$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 8.2438 (13)$ Å
 $b = 16.405 (2)$ Å
 $c = 8.8561 (14)$ Å
 $\beta = 110.705 (2)^\circ$

$V = 1120.3 (3)$ Å³
 $Z = 4$
 $F(000) = 536$
 $D_x = 1.536$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71075$ Å
Cell parameters from 14670 reflections
 $\theta = 3.0\text{--}27.5^\circ$
 $\mu = 0.56$ mm⁻¹

$T = 100\text{ K}$
Prism, colourless

$0.20 \times 0.07 \times 0.04\text{ mm}$

Data collection

Rigaku R-AXIS conversion
diffractometer
Radiation source: Sealed Tube
Graphite monochromator
Detector resolution: 10.0000 pixels mm⁻¹
profile data from ω -scans
Absorption correction: multi-scan
(*CrystalClear-SM Expert*; Rigaku, 2013)
 $T_{\min} = 0.831$, $T_{\max} = 1.000$

7784 measured reflections
2581 independent reflections
2162 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -10 \rightarrow 9$
 $k = -20 \rightarrow 21$
 $l = -11 \rightarrow 11$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.076$
 $S = 1.07$
2581 reflections
154 parameters
3 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_{\text{o}}^2) + (0.0353P)^2 + 0.170P]$
where $P = (F_{\text{o}}^2 + 2F_{\text{c}}^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.37\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.24\text{ e \AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	1.02274 (5)	0.76756 (2)	0.47284 (4)	0.01888 (11)
O1	0.53371 (15)	0.30013 (7)	0.03702 (13)	0.0206 (3)
H1O	0.595 (2)	0.3400 (9)	0.033 (2)	0.031*
N1	0.90024 (17)	0.49718 (8)	0.69236 (15)	0.0153 (3)
H1N	0.9879 (17)	0.5083 (11)	0.7812 (14)	0.018*
N2	0.50113 (17)	0.43490 (8)	0.27207 (15)	0.0153 (3)
H2N	0.466 (2)	0.4713 (9)	0.1961 (17)	0.018*
C1	0.8212 (2)	0.42491 (10)	0.67877 (18)	0.0162 (3)
H1	0.8571	0.3886	0.7681	0.019*
C2	0.6904 (2)	0.40138 (9)	0.54089 (18)	0.0152 (3)
H2	0.6387	0.3492	0.5354	0.018*
C3	0.63187 (19)	0.45441 (9)	0.40638 (17)	0.0131 (3)
C4	0.72227 (19)	0.53172 (9)	0.41947 (18)	0.0135 (3)

C5	0.68800 (19)	0.58698 (9)	0.28956 (17)	0.0145 (3)
H5	0.5992	0.5747	0.1895	0.017*
C6	0.7804 (2)	0.65795 (9)	0.30511 (18)	0.0157 (3)
H6	0.7581	0.6940	0.2160	0.019*
C7	0.90831 (19)	0.67665 (9)	0.45450 (18)	0.0154 (3)
C8	0.94774 (19)	0.62514 (9)	0.58444 (18)	0.0153 (3)
H8	1.0349	0.6389	0.6845	0.018*
C9	0.85580 (19)	0.55131 (9)	0.56571 (17)	0.0133 (3)
C10	0.4232 (2)	0.35374 (9)	0.24153 (19)	0.0172 (3)
H10A	0.3101	0.3569	0.1517	0.021*
H10B	0.4014	0.3348	0.3388	0.021*
C11	0.5388 (2)	0.29206 (10)	0.19883 (18)	0.0177 (3)
H11A	0.6598	0.2992	0.2736	0.021*
H11B	0.5014	0.2363	0.2144	0.021*
Cl2	0.75804 (5)	0.45204 (2)	0.02211 (4)	0.01726 (10)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0192 (2)	0.01435 (19)	0.02016 (19)	-0.00294 (15)	0.00339 (15)	0.00041 (15)
O1	0.0244 (6)	0.0209 (6)	0.0164 (5)	-0.0031 (5)	0.0071 (5)	-0.0019 (5)
N1	0.0159 (6)	0.0175 (7)	0.0112 (6)	0.0010 (5)	0.0030 (5)	0.0010 (5)
N2	0.0163 (6)	0.0137 (7)	0.0140 (6)	0.0000 (5)	0.0029 (5)	0.0005 (5)
C1	0.0194 (8)	0.0160 (7)	0.0146 (7)	0.0032 (6)	0.0079 (6)	0.0021 (6)
C2	0.0175 (7)	0.0133 (7)	0.0162 (7)	0.0000 (6)	0.0077 (6)	0.0008 (6)
C3	0.0130 (7)	0.0139 (7)	0.0138 (7)	0.0031 (6)	0.0064 (6)	-0.0010 (6)
C4	0.0133 (7)	0.0139 (7)	0.0141 (7)	0.0014 (6)	0.0058 (6)	-0.0016 (6)
C5	0.0143 (7)	0.0151 (7)	0.0120 (7)	0.0021 (6)	0.0021 (6)	-0.0012 (6)
C6	0.0160 (7)	0.0148 (7)	0.0158 (7)	0.0029 (6)	0.0050 (6)	0.0025 (6)
C7	0.0152 (7)	0.0126 (7)	0.0196 (7)	-0.0003 (6)	0.0078 (6)	-0.0016 (6)
C8	0.0133 (7)	0.0173 (8)	0.0143 (7)	0.0011 (6)	0.0035 (6)	-0.0021 (6)
C9	0.0144 (7)	0.0133 (7)	0.0131 (7)	0.0035 (6)	0.0060 (6)	0.0000 (6)
C10	0.0177 (8)	0.0156 (8)	0.0177 (7)	-0.0043 (6)	0.0053 (6)	-0.0034 (6)
C11	0.0209 (8)	0.0159 (8)	0.0164 (7)	-0.0023 (6)	0.0066 (6)	-0.0002 (6)
Cl2	0.01711 (19)	0.0195 (2)	0.01341 (17)	0.00029 (15)	0.00320 (14)	-0.00020 (14)

Geometric parameters (\AA , $^\circ$)

Cl1—C7	1.7413 (16)	C4—C9	1.409 (2)
O1—C11	1.4250 (18)	C4—C5	1.413 (2)
O1—H1O	0.838 (9)	C5—C6	1.371 (2)
N1—C1	1.338 (2)	C5—H5	0.9500
N1—C9	1.3749 (19)	C6—C7	1.404 (2)
N1—H1N	0.879 (9)	C6—H6	0.9500
N2—C3	1.331 (2)	C7—C8	1.371 (2)
N2—C10	1.4615 (19)	C8—C9	1.407 (2)
N2—H2N	0.869 (9)	C8—H8	0.9500
C1—C2	1.368 (2)	C10—C11	1.526 (2)

C1—H1	0.9500	C10—H10A	0.9900
C2—C3	1.415 (2)	C10—H10B	0.9900
C2—H2	0.9500	C11—H11A	0.9900
C3—C4	1.455 (2)	C11—H11B	0.9900
C11—O1—H1O	108.4 (14)	C5—C6—H6	120.5
C1—N1—C9	121.21 (13)	C7—C6—H6	120.5
C1—N1—H1N	119.0 (12)	C8—C7—C6	122.19 (14)
C9—N1—H1N	119.6 (12)	C8—C7—Cl1	119.32 (12)
C3—N2—C10	123.26 (13)	C6—C7—Cl1	118.48 (12)
C3—N2—H2N	117.9 (12)	C7—C8—C9	118.27 (14)
C10—N2—H2N	118.6 (12)	C7—C8—H8	120.9
N1—C1—C2	122.38 (14)	C9—C8—H8	120.9
N1—C1—H1	118.8	N1—C9—C8	118.87 (13)
C2—C1—H1	118.8	N1—C9—C4	119.93 (14)
C1—C2—C3	120.30 (14)	C8—C9—C4	121.20 (14)
C1—C2—H2	119.8	N2—C10—C11	112.19 (13)
C3—C2—H2	119.8	N2—C10—H10A	109.2
N2—C3—C2	122.13 (14)	C11—C10—H10A	109.2
N2—C3—C4	120.72 (13)	N2—C10—H10B	109.2
C2—C3—C4	117.15 (13)	C11—C10—H10B	109.2
C9—C4—C5	117.95 (14)	H10A—C10—H10B	107.9
C9—C4—C3	118.95 (13)	O1—C11—C10	112.87 (13)
C5—C4—C3	123.05 (14)	O1—C11—H11A	109.0
C6—C5—C4	121.27 (14)	C10—C11—H11A	109.0
C6—C5—H5	119.4	O1—C11—H11B	109.0
C4—C5—H5	119.4	C10—C11—H11B	109.0
C5—C6—C7	119.06 (14)	H11A—C11—H11B	107.8
C9—N1—C1—C2	-1.4 (2)	C5—C6—C7—Cl1	-179.28 (11)
N1—C1—C2—C3	-0.9 (2)	C6—C7—C8—C9	0.0 (2)
C10—N2—C3—C2	-9.1 (2)	Cl1—C7—C8—C9	-178.78 (11)
C10—N2—C3—C4	170.11 (13)	C1—N1—C9—C8	-177.54 (14)
C1—C2—C3—N2	-177.69 (14)	C1—N1—C9—C4	1.4 (2)
C1—C2—C3—C4	3.0 (2)	C7—C8—C9—N1	176.63 (13)
N2—C3—C4—C9	177.74 (13)	C7—C8—C9—C4	-2.3 (2)
C2—C3—C4—C9	-3.0 (2)	C5—C4—C9—N1	-176.31 (13)
N2—C3—C4—C5	-5.2 (2)	C3—C4—C9—N1	0.9 (2)
C2—C3—C4—C5	174.04 (13)	C5—C4—C9—C8	2.6 (2)
C9—C4—C5—C6	-0.6 (2)	C3—C4—C9—C8	179.74 (13)
C3—C4—C5—C6	-177.64 (14)	C3—N2—C10—C11	-74.61 (18)
C4—C5—C6—C7	-1.6 (2)	N2—C10—C11—O1	-77.07 (16)
C5—C6—C7—C8	2.0 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1o···Cl2	0.84 (2)	2.30 (2)	3.1338 (14)	179 (2)

N1—H1n···Cl2 ⁱ	0.88 (1)	2.29 (1)	3.1602 (15)	168 (1)
N2—H2n···Cl2 ⁱⁱ	0.87 (1)	2.49 (2)	3.2949 (14)	154 (2)
C2—H2···O1 ⁱⁱⁱ	0.95	2.60	3.545 (2)	173

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $-x+1, -y+1, -z$; (iii) $x, -y+1/2, z+1/2$.