



Range-wide pattern of genetic variation in *Colobanthus quitensis*

Justyna Koc¹ · Piotr Androsiuk¹  · Katarzyna Joanna Chwedorzewska^{2,3} · Marely Cuba-Díaz⁴ · Ryszard Górecki¹ · Irena Giełwanowska¹

Received: 18 May 2017 / Revised: 18 July 2018 / Accepted: 19 July 2018
© The Author(s) 2018

Abstract

There is only one species representing Magnoliopsida which is considered as native to the Antarctic, i.e., Antarctic pearlwort (*Colobanthus quitensis*). Although it was intensively studied toward the morphophysiological adaptation to extreme environmental conditions of that area, there is still a lack of sufficient data on its genetic variability. Nine *C. quitensis* populations from Chile and the Maritime Antarctic were sampled to estimate the pattern of genetic variation in relation to the geographic distribution of analyzed populations and postglacial history of the species. The retrotransposon-based DNA marker system used in our studies appeared to be effective in revealing genetic polymorphism between individuals and genetic differentiation among populations. Although the level of polymorphism was low (9.57%), the Analysis of Molecular Variance showed that overall population differentiation was high ($F_{ST} = 0.6241$) and revealed significant differentiation between the Northern and Southern Group of populations as well as the population from Conguillio Park. The observed genetic subdivision of *C. quitensis* populations was confirmed by Bayesian clustering and results of Principal Coordinates Analysis. The Southern Group of populations was characterized by generally higher genetic diversity, which was expressed by the values of the effective number of alleles, expected heterozygosity and by the distribution of private alleles. Our results suggest that the species may have survived the Last Glacial Maximum in refugia located both on the South American continent and in geographically isolated islands of the Maritime Antarctic, i.e., they support the concept of the multiregional origin of *C. quitensis* in Antarctica.

Keywords Antarctic pearlwort · Genetic diversity · Genetic structure · Antarctica · iPBS

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00300-018-2383-5>) contains supplementary material, which is available to authorized users.

✉ Piotr Androsiuk
piotr.androsiuk@uwm.edu.pl

- ¹ Department of Plant Physiology, Genetics and Biotechnology, Faculty of Biology and Biotechnology, University of Warmia and Mazury in Olsztyn, ul. M. Oczapowskiego 1A, 10-719 Olsztyn, Poland
- ² Institute of Biochemistry and Biophysics, Polish Academy of Sciences, ul. Pawińskiego 5A, 02-106 Warsaw, Poland
- ³ Department of Agronomy, Warsaw University of Life Sciences-SGGW, Nowoursynowska 159, 02-766 Warsaw, Poland
- ⁴ Univ Concepcion, Escuela Ciencias & Tecnol, Lab Biotecnol & Estudios Ambientales, Campus Los Ángeles, Casilla 341, Juan Antonio Coloma 0201, Los Ángeles, Chile

Introduction

Patterns of population genetic structure revealed in the widely distributed plant species are shaped by the interaction of many factors, including natural processes such as climate fluctuations, natural barriers, life history (e.g., breeding system and life-form) as well as human impacts (Convey et al. 2014). Also, demographic events occurring during range contractions and expansions are responsible for modifications of the genome-wide patterns of population genetic diversity (Hewitt 2000; Excoffier et al. 2009). Genetic drift, founder, and bottleneck effects are expected to be especially intense during range expansion, causing gradients or even abrupt modulations in allele frequencies, genetic diversity and population structure (Klopfstein et al. 2006; Excoffier and Ray 2008; François et al. 2008). Expansions can affect the geographic distribution of both neutral and selectively important genetic variation and thus complicate the inferences drawn from clines in alleles or phenotypic traits

(Vasemagi 2006; Keller et al. 2009) or genetic association mapping (Pritchard et al. 2000; Price et al. 2006). Therefore, a proper understanding of how expansion can shape diversity and population structure is crucial for both inferring from the demographic history of the species but also for studying the molecular basis and evolution of the locally adaptive traits in natural populations (Nielsen 2005; Wright and Gaut 2005).

There are a number of molecular phylogeographic studies in which maternally inherited organellar DNA markers were used for determining the patterns of plant population genetic structure and inferring processes such as the number and locations of glacial refugia, past migration routes, and colonization dynamics (Comes and Kadereit 1998). The reason for this approach is that the organellar genomes are usually characterized by uniparental inheritance and generally does not undergo recombination, and therefore haplotypes remain mostly unchanged when passed to the next generation. Nevertheless, the organellar genome reveals only a single gene genealogy and since the genealogical process is highly variable among genes and is affected by demographic events or unexpected loss of lineages (Knowles and Maddison 2002) such analyses may reveal genealogical processes associated with single genes only, which are not necessarily characteristic for the whole species' history.

On the contrary, nuclear DNA markers which are biparentally inherited allow for recombination, integrating many genealogical processes. Currently SSRs (Simple Sequence Repeats) or SNPs (Single Nucleotide Polymorphisms) are the markers of choice for many phylogeographic studies (e.g., DeFaveri et al. 2013; Gao et al. 2013; Faivre-Rampant et al. 2016; Foote and Morin 2016), however due to the high cost and labor intensity involved in their development, they are available generally for model species and those which are economically important. For genetic studies on underutilized crops or plants for which genomic data is limited, several PCR (Polimerase Chain Reaction) marker systems are available, varying in complexity, reliability, and information generating capacity.

Recently, a versatile method of organism genotyping based on the use of transposon sequences has been developed. The iPBS method (inter Primer Binding Sites) is based on the virtually universal presence of a tRNA complement as a reverse transcriptase primer binding site (PBS) in LTR (Long Terminal Repeat) retrotransposons (Kalendar et al. 2010). The iPBS technique has been introduced as a powerful DNA fingerprinting technology which can be applied without the need of prior sequence knowledge (Kalendar et al. 2010). Moreover, iPBS markers are highly reproducible due to the length of primers and the high stringency achieved by the annealing temperature, and they have already found application in clone identification, genetic diversity analyses and phylogenetic studies (e.g., Smýkal et al. 2011; Mehmood

et al. 2016; Özer et al. 2016). *Colobanthus* is often described as a genus showing southern circum-temperate distribution (e.g., West 1991). 26 species are recognized at present in the genus, with the major diversity, occurring in New Zealand—14 species (Sneddon 1999). It exhibits a Gondwanan distribution, occurring in South America, the sub-Antarctic islands, the Antarctic Peninsula, and the Australasian area including New Zealand, Australia, and Tasmania (West 1991). The most recent taxonomic studies in *Colobanthus* are scatter a descriptions of two new species from Australia: *Colobanthus nivicola* (Gray 1976) and *Colobanthus curtisiae* (West 1991) and a revision of *Colobanthus quitensis* (Kunth) Bartl., which is the most wide-ranging species in the genus, occurring from the Antarctic Peninsula (Greene and Greene 1963) north along the southern Andes mountains up to scattered locations northward to Mexico (Moore 1970; Parnikoza et al. 2011). Antarctic pearlwort—*C. quitensis*. and Antarctic hairgrass—*Deschampsia antarctica* Desv. are the two flowering plant species considered as native to the Maritime Antarctic. Due to the broad distribution range, both species undergo various selection forces which shape both their morphological and genetic variability. However, information on the genetic variation of these species is limited and devoted almost exclusively to *D. antarctica* (e.g., Chwedorzewska and Bednarek 2008; van de Wouw et al. 2008; Volkov et al. 2010; Chwedorzewska and Bednarek 2011; González et al. 2016). As regards *C. quitensis*, results of the genetic variation studies can be found so far in only a few papers, which suffer from limited geographic range or the low number of populations studied (Lee and Postle 1975; Gianoli et al. 2004; Acuña-Rodríguez et al. 2014). More recently, two more papers concerning *C. quitensis* genetics appeared, but they were focused on selected areas of the species genomics like the sequence of chloroplast genome (Kang et al. 2016) or genome size estimated by flow cytometry (Cuba-Díaz et al. 2017b).

Our previous studies on the genetic variation of *C. quitensis* (Androsiuk et al. 2015) encompassed the highest number of populations of the species analyzed so far in one study, i.e., eight populations from King George Island (Southern Shetland Islands). They represented diversified habitats of the Maritime Antarctic, which vary considerably when microclimatic conditions as well as soil moisture and its nutrient content are considered. The applied DNA markers (iPBS) revealed low genetic variability and moderate population differentiation. Moreover, the pattern of population differentiation corresponds well with their geographic location. This study demonstrated the usefulness of iPBS markers for genetic diversity studies in non-model plant species and confirmed our hypothesis that genetic variation revealed by this technique is influenced by the abiotic stress and thus shaped in the response to local environment conditions.

In this study, we used the iPBS markers to assess the range-wide genetic variation of *C. quitensis* in order to infer about the population history of the vicariants of *C. quitensis* in South America and the Maritime Antarctic. We aimed to address the following questions: (1) Do the iPBS markers reveal cryptic genetic structure in *C. quitensis*? (2) Do the levels of genetic diversity in small, isolated marginal populations differ from those in other populations?

Materials and methods

DNA extraction and genotyping

The research material consisted of 174 individuals of *C. quitensis* representing its natural stands (hereinafter referred to as populations) from South America and the Maritime Antarctic (South Shetland Islands) (Table 1; Fig. 1). *C. quitensis* from South America was collected from wet environments of both fresh and brackish water, in river deposits flooded by high tides and less frequent on coastal rocks. In the high mountain ranges, it was found mainly in bogs flooded by meltwater runoff. It is usually associated with other herbaceous species. The pPar and pV populations manifest a high degree of deterioration due to human activity: the presence of pitches for winter sports in pPar and heavy grazing in pV. In case of pPA and pL, the ample influence of human activity is observed: both populations are close to roads, especially the pL population is located very nearby to the sites with high industrial activity. In the Antarctic samples were taken from coastal zones, from areas with minor human influence. Each population was represented by 6–30 individuals. The number of sampled individuals was limited by the size of a given population. The plant material

was dried after collection and stored at $-20\text{ }^{\circ}\text{C}$ until DNA extraction in presence of silica gel. The DNA from each individual was extracted using Syngen Plant DNA Mini Kit. The quality of DNA was verified on 1% agarose gel and visualized by staining with $0.5\text{ }\mu\text{g}\cdot\text{mL}^{-1}$ ethidium bromide, while the amount and purity of DNA samples were assessed spectrophotometrically.

Initially, we screened 20 iPBS primers and their combinations for *C. quitensis* (Kalendar et al. 2010). Six of the 20 iPBS primers which gave polymorphic, clearly identifiable and repeatable bands were selected for further analyses. The PCR was performed with the 6 iPBS primers applied individually or in combination with two primers (Table 2), according to the procedure described in Androsiuk et al. (2015). The amplification products were analyzed by gel electrophoresis in 1.5% agarose gels stained with $0.5\text{ }\mu\text{g}\cdot\text{mL}^{-1}$ ethidium bromide.

Data analysis

All amplification products (bands), that could be reliably read were treated as single dominant loci and scored either present (1) or absent (0) across genotypes (Online Resources 1). On the basis of the obtained binary matrix of bands the following genetic parameters were estimated: effective number of alleles (N_B), the percentage of polymorphic bands (P), Shannon's Information Index (I), and expected heterozygosity (H_e). Pairwise F_{ST} values among all analyzed *C. quitensis* populations were also estimated. The above-mentioned calculations were performed with GenAlEx 6.5 software (Peakall and Smouse 2006, 2012), except pairwise F_{ST} values which were estimated using Arlequin 3.5 software (Excoffier et al. 2005).

Table 1 The origin of *Colobanthus quitensis* populations used in the study

Name	Sampling site	Coordinates (GPS)	Altitude (m a.s.l.)	No. individuals
La Parva (pPar)	La Parva hill, Santiago, Chile	33°19'S; 70°16'W	3600	30
Conguillio Park (pC)	Conguillio National Park, Araucanía, Chile	38°36'S; 71°36'W	2575	29
La Vega (pV)	INIA Kampanaike farm, North of Punta Arenas, Chile	52°41'S; 70°56'W	16	10
Laredo (pL)	Laredo Sector, North of Punta Arenas, Chile	52°58'S; 70°49'W	158	30
La Marisma (pPA)	St. María Point, South of Punta Arenas, Chile	53°22'S; 70°58'W	1–3	30
Omora Park (pOm)	Omora Park, Puerto Williams, Patagonia, Chile	54°56'S; 67°39'W	–	7
Elephant Is. (pEl)	Elephant Is., Antarctica	61°06'S; 55°08'W	–	8
Arctowski (pA)	Arctowski Station, King George Is., Antarctica	62°09'S; 58°28'W	23	24
Byers (pBy)	Byers Peninsula, Livingston Is., Antarctica	62°40'S; 60°55'W	40	6

Populations are ordered by latitude of origin

Fig. 1 Map depicting the location of the studied sampling sites of *Colobanthus quitensis*: **a** Geographic location of the populations from South America, **b** contour map of the Drake passage with **c** the enlargement area of South Shetland Archipelago with Elephant Island, King George Island and Livingston Island. *pPar* La Parva; *pC* Conguillio Park; *pV* La Vega; *pL* Laredo; *pPA* La Marisma; *pOm* Omora Park; *pEl* Elephant Is.; *pA* Arctowski; *pBy* Byers




Table 2 The sequence and specification of primers applied in the study

Primer	Sequence	T _m (°C)	Number of amplified bands
iPBS2076	5'-GCTCCGATGCCA-3	54 ^a	20 ^b
iPBS2085	5'-ATGCCGATACCA-3	50	24
iPBS2224	5'-ATCCTGGCAATGGAA CCA-3	52	22
iPBS2228	5'-CATTGGCTCTTGATA CCA-3'	54	20
iPBS2231	5'-ACTTGGATGCTGATA CCA-3'	52	27
iPBS2378	5'-GGTCCTCATCCA-3'	53	24
Total			137

^aAnnealing temperature applied in Polymerase Chain Reaction with combination of primers iPBS2076 × iPBS2085

^bNumber of bands scored when primer iPBS2076 was used in combination with primer iPBS2085

In order to infer about the geographic pattern of genetic variation among *C. quitensis* populations, two methods were used. The first approach was a model-based clustering method using the Bayesian analysis with the STRUCTURE

ver. 2.3.4 software (Pritchard et al. 2000). The model assigns individual multilocus genotypes probabilistically to a user-defined number of clusters (K), achieving linkage equilibrium within clusters (Pritchard et al. 2000). We conducted 10 replicate runs for each K , ranging from 1 to 9. Each run consisted of a burn-in of 500,000 iterations, followed by data collection over 500,000 iterations. The analysis, with the implemented admixture model, was conducted without any prior information on the original population of each sampled individual. In order to determine the optimal number of clusters (K), an ad hoc statistic ΔK (Evanno et al. 2005) was used. The ΔK was evaluated in Structure Harvester ver. 0.6.94 (Earl and Vonholdt 2012). The second method was a Principal Coordinates Analysis (PCoA), based on the matrix of Euclidean distances between individuals from all analyzed populations, performed in PAST software (Hammer et al. 2001).

Hierarchical Analysis of Molecular Variance (AMOVA; Excoffier et al. 1992) was carried out using Arlequin 3.5. For that analysis, the iPBS data were treated as haplotypic, comprising a combination of alleles at one or several loci (Excoffier et al. 2005). The significance of the fixation indices was tested using a non-parametric permutation approach (Excoffier et al. 1992).

The spatial genetic structure was investigated by testing the significance of isolation by distance (IBD) using a Mantel test with 9999 permutations of the relationship between the matrix of pairwise $F_{ST}/(1 - F_{ST})$ and that of the logarithm of geographical distance between populations (Rousset 1997), using GenAlEx 6.5.

Results

Efficiency of applied PCR primers

Our analysis of *C. quitensis* populations, using six iPBS primers/primers combination, yielded 137 clearly distinguished bands (Table 2). The highest number, 27 bands, was revealed by iPBS2231, whereas the lowest number of bands (20) was scored for iPBS2228 and iPBS2076×2085 primers combination. The average number of bands per primer was 22.83. Out of all identified loci, 53 (38.68%) were polymorphic.

Of a total of 137 scored bands, eight (5.84%) were represented as private bands—i.e., observed only in one population and absent in the others. Three private bands were revealed by each of the following primers: iPBS2085 and iPBS2231; whereas primer iPBS2378 revealed two (Table 3). In this context, pEl (Elephant Island) population appeared as the most abundant in private bands—six of them were scored in individuals representing that population. The remaining two private bands were characteristic for pPA (La Marisma) population. Moreover, one of the bands revealed for pEl population (iPBS2378-12), was characteristic for all individuals representing that sampling site, and thus may be considered as a potentially diagnostic marker for that population. In the case of the other private bands, they were scored only for a few (1–5) individuals from a given population.

Table 3 Private alleles (bands) per locus and their respective frequencies revealed in studied *Colobanthus quitensis* populations

Locus	Frequency	Population
iPBS2085-3	0.006	Elephant Island (pEl)
iPBS2085-10	0.03	La Marisma (pPA)
iPBS2085-15	0.006	La Marisma (pPA)
iPBS2231-2	0.025	Elephant Island (pEl)
iPBS2231-3	0.006	Elephant Island (pEl)
iPBS2231-13	0.012	Elephant Island (pEl)
iPBS2378-12	0.05	Elephant Island (pEl)
iPBS2378-15	0.018	Elephant Island (pEl)

Genetic diversity and differentiation

The iPBS markers revealed both the presence of genetic polymorphism between individuals within populations and genetic variation between populations (Table 4). The number of iPBS bands ranged from 122, for pPar, pC, and pV populations, to 131 for pPA and pEl populations. The highest number of polymorphic bands was scored for pPA population (26.28%) whereas the lowest polymorphism was observed for pV and pBy populations (1.46%). The genetic variation was assessed with two parameters: Shannon's Information Index and expected heterozygosity, and in both cases the highest values were observed for pPA population from La Marisma, whereas the lowest—for pV population (La Vega).

The Bayesian clustering revealed that ΔK , the second-order rate of change of the likelihood function with respect to K , has a maximum at $K=3$ (Fig. 2). Consequently, we chose $K=3$ as the optimal number of clusters of the uppermost hierarchical level of population structure. The first cluster consists of pPA, pOm populations from the southern edges of the South American continent and pA, pEl, pBy populations from the South Shetland Islands, the pPar, pV, and pL populations were gathered in the second cluster, whereas pC population form solitary, third cluster.

The AMOVA results revealed that most of the described genetic variation occurred between populations (59.48%), whereas the remaining 40.52% of variation was attributed to the variation among individuals within populations (Table 5). With the purpose of investigating how the region of origin of each population (affiliation to one of the two clusters revealed by STRUCTURE) influences the partition of revealed genetic variation of *C. quitensis*, further analysis of variance was performed using the region of origin as an

Table 4 Population genetic characteristics for analyzed populations of *Colobanthus quitensis*: effective number of alleles (N_B), percentage of polymorphic bands (P), Shannon's Information Index (I), expected heterozygosity (H_e)

Population	N_B	P (%)	I	H_e
pPar	1.046	6.57	0.039	0.026
pC	1.016	2.92	0.016	0.010
pV	1.002	1.46	0.003	0.001
pL	1.032	6.57	0.031	0.020
pPA	1.175	26.28	0.147	0.099
pOm	1.110	13.14	0.083	0.059
pEl	1.076	13.87	0.069	0.046
pA	1.093	13.87	0.078	0.053
pBy	1.010	1.46	0.009	0.006
Mean over loci and populations	1.062	9.57	0.053	0.036

Populations are ordered by latitude of origin

Fig. 2 The uppermost hierarchical level of genetic structure of *Colobanthus quitensis* from Chile and Maritime Antarctic using STRUCTURE (Pritchard et al. 2000). The values of second-order rate of change of $L(K)$, ΔK , of data between successive K values




Table 5 Partitioning of diversity found in *Colobanthus quitensis* from all analyzed populations using AMOVA

Region	Source of variation	df	Sum of squares	Variance components	Percentage of variation	p
All populations ^a	Among populations	8	536.356	3.4747	59.48	<0.0001
	Within populations	165	390.581	2.3672	40.52	–
Three groups distinguished by STRUC ^b	Among groups	2	306.222	1.6852	26.76	0.0017
	Among populations within groups	6	230.134	2.2448	35.65	<0.0001
	Within populations	165	390.581	2.3671	37.59	–

^a F_{ST} =0.5948 when no genetic structure is considered

^b F_{ST} =0.6241 when gene pool subdivision revealed by STRUCTURE is considered

additional factor. The AMOVA revealed that while the most genetic variation still occurs between populations (35.65%), and among individuals within populations (37.59%), there is a significant variation (26.76%) among geographical regions (Table 5).

PCoA indicated that 50.89% of variation was explained by the first three components (20.86, 19.75, and 10.28%, respectively). Figure 3 illustrates the projection of the analyzed populations on the first two axes. The grouping revealed by PCoA pointed on the close relationship between three populations from Arctowski (pA), Elephant Island (pEl), and Byers (pBy) representing the South Shetland Islands. The individuals representing Populations La Marisma (pPA) and Omora Park (pOm) from South America form two the most dispersed clouds, which overlap with each other and also with populations from South Shetlands Islands. The high dispersion of individuals from

pPA and pOm point on the high level of genetic variation found within that populations. Individuals from populations La Parva (pPar) and Laredo (pL) form two overlapping clouds which departed from the others along the Coordinate 1. The most distinct character is shown by Conguillio Park (pC) which individuals departed from the others along Coordinate 2. The individuals representing La Vega population form the smallest and the densest cloud of individuals in the center of Fig. 3.

A Mantel test showed the highly significant correlation between genetic divergence, denoted by pairwise $F_{ST}/(1 - F_{ST})$, and the logarithm of the geographical distances between populations ($R^2 = 0.1299$, $p < 0.016$) (Fig. 4). Pairwise $F_{ST}/(1 - F_{ST})$ values plotted against the logarithm of geographical distances between populations indicated that the values increased with the distance between them (IBD).





Fig. 3 Plot of Coordinate 1 versus Coordinate 2 obtained by Principal Coordinates Analysis (PCoA) based on Euclidean distances between all individuals from nine *Colobanthus quitensis* populations. *pPar* La Parva (plus sign); *pC* Conguillio Park (open square); *pV* La Vega

(filled square); *pL* Laredo (open circle); *pPA* La Marisma (filled circle); *pOm* Omora Park (asterisk); *pEL* Elephant Is.(times); *pA* Arc-towski (open triangle); *pBy* Byers (open diamond)

Fig. 4 Analysis of isolation by distance. The pairwise $F_{ST}/(1 - F_{ST})$ values are plotted against the logarithm of the geographical distances (km) between studied populations of *Colobanthus quitensis*



Discussion

Genetic diversity and differentiation of *Colobanthus quitensis*

Colobanthus quitensis geographic range spreads over thousands of kilometers from Mexico (17°N) to the Antarctic Peninsula (68°S) and from sea level to 4200 m a.s.l. (Moore 1970). Our studies covered the area of Chile and the Maritime Antarctic from 33°S to 62°S and spread over 3300 km, which is the distance between the most distant pPar and pBy populations. In Chile, *C. quitensis* can be found along the Andes, usually in bogs at high altitudes in the north, but close to sea level in the south (Moore 1970; Hoffmann et al. 1998). In the Antarctic, the sites of *C. quitensis* occurrence are disjunct, situated mainly on the Antarctic islands and the Antarctic Peninsula coastal ice-free areas, often separated from one another by natural barriers (open sea, glacial, or mountains) limiting the exchange of gene pool (Lewis-Smith 2003).

Colobanthus. quitensis became an interesting subject of morphophysiological investigations aiming to identify ecotypic variation (Gianoli et al. 2004), to analyze its reproduction performance (Giełwanowska et al. 2011; Sanhueza et al. 2017) or identify the characteristic endophytic fungal communities (Santiago et al. 2012, 2017). Moreover, as the only representative of Magnoliopsida that grows in the extreme environmental conditions of the Antarctic, *C. quitensis* was extensively studied in terms of the morphological, physiological, and biochemical basis of adaptation to extremely cold climate, ample thermal oscillations, short vegetation season, high UV-B radiation, and salinity (Bravo et al. 2007; Bascunan-Godoy et al. 2012; Navarrete-Gallegos et al. 2012; Cuba-Díaz et al. 2017a). In contrast, the genetic diversity of this species is still poorly studied and deserves more attention.

Our data on the genetic variation of *C. quitensis* showed that on average 9.57% of the bands revealed by iPBS primers were polymorphic. However, the uneven distribution of the revealed polymorphism between the analyzed population needs to be emphasized here—the value of this parameter ranged from 26.28% in pPA to 1.46% in pV and pBy. The level of polymorphism seems to be associated here with a sample size for individual populations: the highest polymorphism was found in pPA population, which was represented by 30 individuals, whereas the lowest number of polymorphic loci was characteristic for *C. quitensis* stands represented by 10 and 6 individuals, pV and pBy respectively. However, our data show also that even populations represented by a relatively low number of individuals, e.g., pEl (6 plants) or pOm (7 plants), can reveal higher polymorphism than populations which

consist of 4–5 times higher number of individuals, i.e., pC and pL. Similarly, uneven distribution of polymorphism can be found in our previous studies on *C. quitensis* from King George Island, where the level of revealed polymorphism was on average 14.25% (Androsiuk et al. 2015). In the studies of other authors, the average level of polymorphism for iPBS markers was much higher and reached 85.7% for guava accessions (Mehmood et al. 2013), 86.3% for grape varieties (Guo et al. 2014) or even 97.4% for *Myrica rubra* (Chen and Liu 2014). So far the lowest level of polymorphism of iPBS markers (4.88%) was reported by Baránek et al. (2012), but their analyses aimed at genetic identification of clones of the apricot cultivar.

The AMOVA revealed that the genetic variation was almost equally partitioned between populations (35.65%) and among individuals within studied populations (37.59%); remaining 26.76% was observed between geographical regions with a high fixation index (F_{ST}) pointing at a low historical gene flow between them. Previously high genetic variability between populations (71%) was revealed by AMOVA in Acuña-Rodríguez et al. (2014), who describe the results of the genetic study of *C. quitensis* from populations separated by the Drake Passage; the remaining 29% of revealed genetic variation was portioned between individuals. The results obtained by van de Wouw et al. (2008) for another flowering plant from the Antarctic region, *D. antarctica*, reveal the opposite situation: 74.6% of detected genetic variation was portioned among studied regions and only 15.1% among populations. It needs to be emphasized that the study area covered an extremely large geographic range which included the Indian Ocean, South Georgia, the Falklands, and Antarctic Zone. However, when only the regions within the Antarctic Zone were considered, the situation changed significantly: 45.6% of the variation was found among populations within Antarctic sites and 33.9% among Antarctic sites, the rest of variation was portioned among individuals within populations. High values of the fixation index also indicated low historical gene flow (van de Wouw et al. 2008).

The results of model-based clustering analysis showed that there was a clear gene pool subdivision within the nine populations of *C. quitensis*. According to STRUCTURE, pPA, pOm, pBy, pA, and pEl populations form one cluster (hereinafter called the Southern Group), *C. quitensis* from pL, pV, and pPar form another one (further called the Northern Group), whereas population pC departed from the others and form solitary third cluster. PCoA revealed a similar pattern of population subdivision, where an individual character of population pC was also observed. The unique character of pC population was previously revealed by Cuba-Díaz et al. (2017b). The authors, using flow cytometry, estimated the genome size of three *C. quitensis* populations and

found that mountainous populations from pC present a smaller genome size (0.84 pg), almost half of that found in the other studied pA and pPA populations originating from lower altitudes (1.95 pg), which may suggest that pA and pPA are tetraploids. This variation in genome size could be attributed to ecographic, adaptive variation to site-specific local environmental conditions, and interpreted as a sign of incipient speciation (Murray 2005). The hypothesis on adaptive character of the genetic variation pattern revealed here can be supported by the results of our previous study on *C. quitensis* with application of iPBS markers, which proved to be a very helpful tool in assessment of genome rearrangements, associated with activation of transposable elements arising in a response to various abiotic stress characteristic for a given population site (Androsiuk et al. 2015). Undoubtedly, *C. quitensis* populations analyzed in the present study also represent a very wide range of environmental conditions and therefore undergo various selecting forces which independently shape their genetic variation.

The presence of genetic structure within the analyzed *C. quitensis* population was accompanied by an interesting geographic pattern of genetic diversity distribution. Although low values of the effective number of alleles and expected heterozygosity were found among the populations studied, populations from the Southern Group had generally higher genetic diversity (on average $N_B = 1.0939$ and $H_e = 0.053$) when compared to those from the Northern Group and population pC all together (on average $N_B = 1.024$ and $H_e = 0.014$) (data not shown). Moreover, the Southern Group was characterized by a higher number of bands scored for each population (on average 127.4 bands per population for the Southern Group and 122.25 for the Northern Group + pC population), among which eight private bands can be found which were absent in *C. quitensis* from pC population as well as in populations from the Northern Group. Six of the revealed private bands were scored in pEl, one of the three populations representing a geographically isolated area of the South Shetland Islands. Population pEl was also characterized by a higher than average value of H_e . The unique character of pEl was not an exception since another population (pA) from this remote geographic area has H_e which was higher than the average value of that parameter for the whole Northern Group of *C. quitensis* populations. Only *C. quitensis* from pBy represents a very low level of genetic diversity, which can be attributed to a limited number of individuals available for that site. Significantly higher levels of allelic richness and higher expected heterozygosity for the Maritime Antarctic populations of *C. quitensis* in comparison to the population from Punta Arenas (South America) was also observed by Acuña-Rodríguez et al. (2014).

Phytogeography

Postglacial migration and gene flow associated with this process had major consequences for the population genetics of both animal and plant species. On the one hand, species repeatedly encountered founder events and experienced more or less drastic bottlenecks (Newton et al. 1999; Hewitt 2000). This could be observed both during retreat and re-immigration, which may lead to a reduction in the number of genotypes with increasing distance from glacial refugium (Hewitt 1996). On the other hand, the merging and expanding population from different glacial refugia, which had evolved genetically isolated, could increase genetic variation in zones of secondary contact (Taberlet et al. 1998).

In case of the Antarctic flora, it is difficult to draw conclusions about its evolutionary history since data on contemporary species distributions are not sufficient and this question requires an approach combining knowledge of Antarctic plant species distribution with studies concerning their genetic diversity and rates of evolution. Unfortunately, in case of the *C. quitensis* studies which can provide insights into previous population fragmentation and bottlenecks, historical patterns of dispersal and gene flow, relatedness between the Antarctic and South American populations are still at a very early stage (Gianoli et al. 2004; Acuña-Rodríguez et al. 2014).

Currently available data for *C. quitensis* point at two possible scenarios describing its history and dispersal in the Antarctic. According to one of them, *C. quitensis* is, most likely, a recent postglacial immigrant in the Antarctic and its presence in this region is estimated on Holocene, which was proved by identification of pollen grains and macro-remains of the species in peat deposits dated on 6000 years BP on King George Island, South Shetlands Islands (Birkenmajer et al. 1985). In this context, the Antarctic pearlwort is not an exception—the recent origin for much of the flowering plants is supported by species richness patterns with the highest number of species occurring in localities closest to neighboring continents and areas where there are more favorable climatic conditions for establishment. It is certainly clear that Antarctica experiences a continuous, but at the low level, the input of propagules from the other Southern Hemisphere continents (Marshall 1996). Some species may have potentially overcome natural colonization barriers of the Antarctic, those transported in a natural manner like anemochory (Lewis-Smith 1984, 1991, 1993, 2014; Bargagli et al. 1996; Marshall 1996; Muñoz et al. 2004), hydrochory (Coulson et al. 2002), zoochory (Barnes et al. 2004), or marine debris (Barnes et al. 2004; Lewis et al. 2005), but recent evidence of this kind of transportation is rather limited (Hughes et al. 2006).

According to the second scenario, *C. quitensis* could have survived in highly isolated refugia south of Drake Passage

and inhabited this region from a Mid-Early Cenozoic or even Late Cretaceous (Convey and Stevens 2007; Parnikoza et al. 2007, 2011). Moreover, although during Last Glacial Maximum (LGM) in Pleistocene, all low altitude, coastal regions in the Maritime Antarctic are believed to have been obliterated by expanding glaciers and ice sheets (Larter and Vanneste 1995) and only re-exposed as ice has retreated over, at most, the last 10,000 years (Maslen and Convey 2006; Pugh and Convey 2008), there are palaeolimnological data for coastal continental Antarctic according to which some oases have hosted terrestrial life for up to 150,000 years (Hodgson et al. 2001; Squier et al. 2005), and hence that at least some of them have a considerably longer continuous history of exposure extending beyond the LGM (Peat et al. 2007). Recently even, when propagules from lower latitudes cross geographical barrier isolating the Antarctic the most important obstacle are harsh abiotic conditions, diminishing the probability of their establishment. There are a lot of species which have ecophysiological features required for survival in the polar environment including those from the *Colobanthus* genus like e.g., *C. lycopodioides*, *C. subulatus* and, *C. lechleri*, that partly cover *C. quitensis* range and tolerate similar edaphic conditions. That support partly fact that only records of successful introduction and establishment of species in Maritime Antarctic from lower latitudes (e.g., Chwedorzewska et al. 2015; Pertierra et al. 2017a) are via human vectors (Lityńska-Zajac et al. 2012).

Our molecular data which revealed the presence of evident genetic structure within analyzed *C. quitensis* populations seems to support the concept of the multiregional origin of *C. quitensis* in the Antarctic. The gene pool subdivision as well as relatively high genetic diversity found in the Southern Group of analyzed *C. quitensis* populations suggest that the species may have survived the LGM in refugia located both on the South American continent and also in geographically isolated islands of the Maritime Antarctic. Moreover, high values of fixation index as well as significant results of IBD point at limited gene flow between populations.

However, the connectivity between populations from the Maritime Antarctic and south edges of the South American continent revealed within the Southern Group of *C. quitensis* populations cannot categorically reject the thesis that the current pattern of genetic variation of the species could have been also modified more recently (i.e., during the Holocene) due to low intense, random dispersal events (Muñoz et al. 2004; Parnikoza et al. 2012). Solving of this complex subject needs further, more detailed investigations using more extensive sampling of additional population and application of DNA markers which can provide a better resolution of molecular study, like SNPs or microsatellites.

In face of expanding the human footprint and climate change, the largest redistribution of species since the Last

Glacial Maximum (Pecl et al. 2017) have been observed even in remote Antarctic terrestrial ecosystems (Pertierra et al. 2017b). Expansion of the distribution ranges of two native Antarctic flowering plant species has been the most studied example of a biological response to the recent environmental change in maritime the Antarctic (e.g., Gerighausen et al. 2003; Znój et al. 2017; Wódkiewicz et al. 2018). Thus, with the growing risk of homogenization of Antarctic biota (Terauds et al. 2012) the distribution, population size and diversity of Antarctic native plants became an important ecological bioindicators of environmental changes.

Acknowledgements This research was partly supported by the Polish Ministry of Scientific Research and Higher Education Grant NN303796240. Samples were also collected on Henryk Arctowski Polish Antarctic Station. We would like to thank Editor and three anonymous Reviewers for constructive advice that has improved our paper.

Compliance with ethical standards

Conflicts of interest Authors declare no conflicts of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Acuña-Rodríguez IS, Oses R, Cortés-Vasquez J, Torres-Díaz C, Molina-Montenegro MM (2014) Genetic diversity of *Colobanthus quitensis* across the Drake Passage. *Plant Genet Resour* 12:147–150. <https://doi.org/10.1017/S1479262113000270>
- Androsiuk P, Chwedorzewska K, Szandar K, Gielwanowska I (2015) Genetic variability of *Colobanthus quitensis* from King George Island (Antarctica). *Pol Polar Res* 36:281–295. <https://doi.org/10.1515/popore-2015-0017>
- Baránek M, Meszáros M, Sochorová J, Čechová J, Raddová J (2012) Utility of retrotransposon-derived marker systems for differentiation of presumed clones of the apricot cultivar Velkopavlovická. *Sci Hortic* 143:1–6. <https://doi.org/10.1016/j.scienta.2012.05.022>
- Bargagli R, Broady PA, Walton DWH (1996) Preliminary investigation of the thermal biosystem of Mount Rittmann fumaroles (northern Victoria Land, Antarctica). *Antarct Sci* 8:121–126. <https://doi.org/10.1017/S0954102096000181>
- Barnes DKA, Warren N, Webb K, Phalan B, Reid K (2004) Polar pedunculate barnacles piggy-back on pycnogona, penguins, pinniped seals and plastics. *Mar Ecol Prog Ser* 284:305–310. <https://doi.org/10.3354/meps284305>
- Bascunan-Godoy L, Sanhueza C, Cuba M, Zuniga GE, Corcuera LJ, Bravo LA (2012) Cold-acclimation limits low temperature induced photoinhibition by promoting a higher photochemical quantum yield and a more effective PSII restoration in darkness in the Antarctic rather than the Andean ecotype of *Colobanthus quitensis* (Kunt) Bartl. (Cariophyllaceae). *BMC Plant Biol* 12:114. <https://doi.org/10.1186/1471-2229-12-114>

- Birkenmajer K, Ochyra R, Olsson IU, Stuchlik L (1985) Mid-Holocene radiocarbon-dated peat at Admiralty Bay King George Island (South Shetlands, West Antarctica). *Bull Pol Acad Sci Earth-Sci* 33:7–12
- Bravo LA, Saavedra-Mella FA, Vera F, Guerra A, Cavieres LA, Ivanov AG, Huner NPA, Corcuera LJ (2007) Effect of cold acclimation on the photosynthetic performance of two ecotypes of *Colobanthis quitensis* (Kunth) Bartl. *J Exp Bot* 58:3581–3590. <https://doi.org/10.1093/jxb/erm206>
- Chen FY, Liu JH (2014) Germplasm genetic diversity of *Myrica rubra* in Zhejiang Province studied using inter-primer binding site and start codon-targeted polymorphism markers. *Sci Hortic* 170:169–175. <https://doi.org/10.1016/j.scienta.2014.03.010>
- Chwedorzewska KJ, Bednarek PT (2008) Genetic variability in the Antarctic hairgrass *Deschampsia antarctica* Desv. from maritime Antarctic and sub-Antarctic sites. *Pol J Ecol* 56:209–216
- Chwedorzewska KJ, Bednarek PT (2011) Genetic and epigenetic studies on populations of *Deschampsia antarctica* Desv. from contrasting environments at King George Island (Antarctic). *Pol Polar Res* 32:15–26. <https://doi.org/10.2478/v10183-011-0005-9>
- Chwedorzewska KJ, Gielwanowska I, Olech M, Molina-Montenegro MA, Wódkiewicz M, Galera H (2015) *Poa annua* L. in the maritime-Antarctic—an overview. *Polar Rec* 51:637–643. <https://doi.org/10.1007/s00300-007-0353-4>
- Comes HP, Kadereit JW (1998) The effect of Quaternary climatic changes on plant distribution and evolution. *Trends Plant Sci* 3:432–438. [https://doi.org/10.1016/S1360-1385\(98\)01327-2](https://doi.org/10.1016/S1360-1385(98)01327-2)
- Convey P, Stevens MI (2007) Antarctic biodiversity. *Science* 317:1877–1878. <https://doi.org/10.1126/science.1147261>
- Convey P, Chown SL, Clarke A, Barnes DKA, Bokhorst S, Cummings V, Ducklow HW, Frati F, Green TGA, Gordon S, Griffiths HJ, Howard-Williams C, Huiskes ADHL, Laybourn-Parry J, Lyons WB, Mcminn A, Morley SA, Peck LS, Quesada A, Robinson SA, Schiaparelli S, Wall DH (2014) The spatial structure of Antarctic biodiversity. *Ecol Monogr* 84:203–244. <https://doi.org/10.1890/12-2216.1>
- Coulson SJ, Hodkinson ID, Webb NR, Harrison JA (2002) Survival of terrestrial soil-dwelling arthropods on and in seawater: implications for trans-oceanic dispersal. *Funct Ecol* 16:353–356. <https://doi.org/10.1046/j.1365-2435.2002.00636.x>
- Cuba-Díaz M, Castel K, Acuña D, Machuca Á, Cid I (2017a) Sodium chloride effect on *Colobanthis quitensis* seedling survival and in vitro propagation. *Antarct Sci* 29:45–46. <https://doi.org/10.1017/S0954102016000432>
- Cuba-Díaz M, Cerda G, Rivera C, Gómez A (2017b) Genome size comparison in *Colobanthis quitensis* populations show differences in species ploidy. *Polar Biol* 40:1475–1480. <https://doi.org/10.1007/s00300-016-2058-z>
- DeFaveri J, Viitaniemi H, Leder E, Merila J (2013) Characterizing genic and nongenic molecular markers: comparison of microsatellites and SNPs. *Mol Ecol Resour* 13:377–392. <https://doi.org/10.1111/1755-0998.12071>
- Earl DA, Vonholdt BM (2012) STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conserv Genet Resour* 4:359–361. <https://doi.org/10.1007/s12686-011-9548-7>
- Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol* 14:2611–2620. <https://doi.org/10.1111/j.1365-294X.2005.02553.x>
- Excoffier L, Ray N (2008) Surfing during population expansions promotes genetic revolutions and structuration. *Trends Ecol Evol* 23:347–351. <https://doi.org/10.1016/j.tree.2008.04.004>
- Excoffier L, Smouse P, Quattro JM (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction sites. *Genetics* 131:479–491
- Excoffier L, Laval G, Schneider S (2005) Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evol Bioinform Online* 1:47–50
- Excoffier L, Foll M, Petit RJ (2009) Genetic consequences of range expansions. *Annu Rev Ecol Evol Syst* 40:481–501. <https://doi.org/10.1146/annurev.ecolsys.39.110707.173414>
- Faivre-Rampant P, Zaina G, Jorge V, Giacomello S, Segura V, Scalabrin S, Guerin V, De Paoli E, Aluome C, Viger M, Cattonaro F, Payne A, Paulstephenraj P, Le Paslier MC, Berard A, Allwright MR, Villar M, Taylor G, Bastien C, Morgante M (2016) New resources for genetic studies in *Populus nigra*: genome-wide SNP discovery and development of a 12 k Infinium array. *Mol Ecol Resour* 16:1023–1036. <https://doi.org/10.1111/1755-0998.12513>
- Foote AD, Morin PA (2016) Genome-wide SNP data suggest complex ancestry of sympatric North Pacific killer whale ecotypes. *Heredity* 117:316–325. <https://doi.org/10.1038/hdy.2016.54>
- François O, Blum MGB, Jakobsson M, Rosenberg NA (2008) Demographic history of European populations of *Arabidopsis thaliana*. *PLoS Genet* 4:e1000075
- Gao CH, Ren XD, Mason AS, Li JN, Wang W, Xiao ML, Fu DH (2013) Revisiting an important component of plant genomes: microsatellites. *Funct Plant Biol* 40:645–661. <https://doi.org/10.1071/FP12325>
- Gerighausen U, Brautigam K, Mustafa O, Peter HU (2003) Expansion of vascular plants on an Antarctic island—a consequence of climate change? In: Huiskes AHL, Gieskes WWC, Rozema J, Schorno RML, van der Vies SM, Wolff WJ (eds) *Antarctic biology in a global context*. Backhuys, Leiden, pp 79–83
- Gianoli E, Inostroza P, Zuniga-Feest A, Reyes-Diaz M, Cavieres LA, Bravo LA, Corcuera LJ (2004) Ecotypic differentiation in morphology and cold resistance in populations of *Colobanthis quitensis* (Caryophyllaceae) from the Andes of Central Chile and the Maritime Antarctic. *Arct Antarct Alp Res* 36:484–489
- Gielwanowska I, Bochenek A, Gojło E, Gorecki R, Kellmann W, Pastorzyc M, Szczuka E (2011) Biology of generative reproduction of *Colobanthis quitensis* (Kunth) Bartl. from King George Island, South Shetland Islands. *Pol Polar Res* 32:139–155. <https://doi.org/10.2478/v10183-011-0008-6>
- González M, Urdampilleta JD, Fasanella M, Premoli AC, Chiappella JO (2016) Distribution of rDNA and polyploidy in *Deschampsia antarctica* Desv. in Antarctic and Patagonic populations. *Polar Biol* 39:1663–1677. <https://doi.org/10.1007/s00300-016-1890-5>
- Greene SW, Greene DM (1963) Check list of the sub-Antarctic and Antarctic Vascular Flora. *Polar Rec* 73:411–418
- Guo DL, Guo MX, Hou XG, Hang GH (2014) Molecular diversity analysis of grape varieties based on iPBS markers. *Biochem Syst Ecol* 52:27–32. <https://doi.org/10.1016/j.bse.2013.10.008>
- Hammer Ø, Harper DAT, Ryan PD (2001) PAST: paleontological statistics software package for education and data analysis. *Palaeontol Electronica* 4:1–9
- Hewitt GM (1996) Some genetic consequences of ice ages, and their role in divergence and speciation. *Biol J Linn Soc Lond* 58:247–276. <https://doi.org/10.1111/j.1095-8312.1996.tb01434.x>
- Hewitt GM (2000) The genetic legacy of the Quaternary ice ages. *Nature* 405:907–913. <https://doi.org/10.1038/35016000>
- Hodgson DA, Noon PE, Vyverman W, Bryant CL, Gore DB, Appleby P, Gilmour M, Verleyen E, Sabbe K, Jones VJ, Ellis-Evans JC, Wood PB (2001) Were the Larsemann Hills ice-free through the last glacial maximum? *Antarct Sci* 13:440–454. <https://doi.org/10.1017/S0954102001000608>
- Hoffmann A, Arroyo MTK, Liberona F, Munoz M, Watson J (1998) Plantas Altoandinas en la Flora Silvestre de Chile. Edición Fundación Claudio Gay, Santiago

- Hughes KA, Ott S, Bölter M, Convey P (2006) Colonization Processes. In: Bergstrom DM, Convey P, Huiskes AHL (eds) Trends in Antarctic terrestrial and limnetic ecosystems: Antarctica as a global indicator. Springer, Dordrecht, pp 35–54
- Kalendar R, Antonius K, Smýkal P, Schulman AH (2010) iPBS: a universal method for DNA fingerprinting and retrotransposon isolation. *Theor Appl Genet* 121:1419–1430. <https://doi.org/10.1007/s00122-010-1398-2>
- Kang Y, Lee H, Kim MK, Shin SC, Park H, Lee J (2016) The complete chloroplast genome of Antarctic pearlwort, *Colobanthus quitensis* (Kunth) Bartl. (*Caryophyllaceae*). *Mitochondrial DNA* 27:4677–4678. <https://doi.org/10.3109/19401736.2015.1106498>
- Keller SR, Sowell DR, Neiman M, Wolfe LM, Taylor DR (2009) Adaptation and colonization history affect the evolution of clines in two introduced species. *New Phytol* 183:678–690. <https://doi.org/10.1111/j.1469-8137.2009.02892.x>
- Klopfstein S, Currat M, Excoffier L (2006) The fate of mutations surfing on the wave of a range expansion. *Mol Biol Evol* 23:482–490. <https://doi.org/10.1093/molbev/msj057>
- Knowles LL, Maddison WP (2002) Statistical phylogeography. *Mol Ecol* 11:2623–2635
- Larter RD, Vanneste LE (1995) Relict subglacial deltas on the Antarctic Peninsula outer shelf. *Geology* 23:33–36
- Lee DW, Postle RL (1975) Isozyme variation in *Colobanthus quitensis* (Kunth) Bartl.: Methods and preliminary analysis. *Br Antarct Surv B* 41:133–137
- Lewis PN, Riddle M, Smith SDA (2005) Assisted passage or passive drift: a comparison of alternative transport mechanisms for non-indigenous coastal species into the Southern Ocean. *Antarct Sci* 17:183–191. <https://doi.org/10.1017/S0954102005002580>
- Lewis-Smith RI (1984) Colonization and recovery by cryptogams following recent volcanic activity on Deception Island, South Shetland I. *Br Antarct Surv B* 62:25–51
- Lewis-Smith RI (1991) Exotic sporomorphs as indicators of potential immigrant colonists in Antarctica. *Grana* 30:313–324. <https://doi.org/10.1080/00173139109431986>
- Lewis-Smith RI (1993) Dry coastal ecosystems of Antarctica. In: Maarel E (ed) *Ecosystems of the world, 2A; dry coastal ecosystems, polar regions and Europe*. Elsevier, Amsterdam, pp 51–57
- Lewis-Smith RI (2003) The enigma of *Colobanthus quitensis* and *Deschampsia antarctica* in Antarctica. In: Huiskes AHL, Gieskes WWC, Rozema J, Schorno RML, van der Vies SM, Wolff WJ (eds) *Antarctic biology in a global context*. Backhuys, Leiden, pp 234–239
- Lewis-Smith RI (2014) A fern cultured from Antarctic glacier detritus. *Antarct Sci* 26:341–344. <https://doi.org/10.1017/S0954102013000606>
- Lityńska-Zajac M, Chwedorzewska KJ, Olech M, Korczak-Abshire M, Augustyniuk-Kram A (2012) Diaspores and phyto-remains accidentally transported to the Antarctic Station during three expeditions. *Biodivers Conserv* 21:3411–3421. <https://doi.org/10.1007/s10531-012-0371-6>
- Marshall WA (1996) Biological particles over Antarctica. *Nature* 383:680. <https://doi.org/10.1038/383680a0>
- Maslen NR, Convey P (2006) Nematode diversity and distribution in the southern maritime Antarctic—clues to history? *Soil Biol Biochem* 38:3141–3151. <https://doi.org/10.1016/j.soilbio.2005.12.007>
- Mehmood A, Jaskani MJ, Ahmad S, Ahmad R (2013) Evaluation of genetic diversity in open pollinated guava by iPBS primers. *Pak J Agric Sci* 50:591–597
- Mehmood A, Luo S, Ahmad NM, Dong C, Mahmood T, Sajjad Y, Jaskani MJ, Sharp P (2016) Molecular variability and phylogenetic relationships of guava (*Psidium guajava* L.) cultivars using inter-primer binding site (iPBS) and microsatellite (SSR) markers. *Genet Resour Crop Evol* 63:1345–1361. <https://doi.org/10.1007/s10722-015-0322-7>
- Moore DM (1970) Studies in *Colobanthus quitensis* (Kunth) Bartl. and *Deschampsia antarctica* Desv.: II. Taxonomy, distribution and relationships. *Br Antarct Surv B* 23:63–80
- Muñoz J, Felicísimo AM, Cabezas F, Burgaz AR, Martínez I (2004) Wind as a long-distance dispersal vehicle in the southern hemisphere. *Science* 304:1144–1147. <https://doi.org/10.1126/science.1095210>
- Murray BG (2005) When does intraspecific C-value variation become taxonomically significant? *Ann Bot* 95:119–125. <https://doi.org/10.1093/aob/mci007>
- Navarrete-Gallegos AA, Bravo LA, Molina-Montenegro MA, Corcuera LJ (2012) Antioxidant responses in two *Colobanthus quitensis* (Caryophyllaceae) ecotypes exposed to high UV-B radiation and low temperature. *Rev Chil Hist Nat* 85:419–433. <https://doi.org/10.4067/S0716-078X2012000400005>
- Newton AC, Allnutt TR, Gillies ACM, Lowe AJ, Ennos RA (1999) Molecular phylogeography, intraspecific variation and the conservation of tree species. *Trends Ecol Evol* 14:140–145. [https://doi.org/10.1016/S0169-5347\(98\)01555-9](https://doi.org/10.1016/S0169-5347(98)01555-9)
- Nielsen R (2005) Molecular signatures of natural selection. *Annu Rev Genet* 39:197–218. <https://doi.org/10.1146/annurev.genet.39.073003.112420>
- Özer G, Bayraktar H, Baloch FS (2016) iPBS retrotransposons ‘A Universal Retrotransposons’ now in molecular phylogeny of fungal pathogens. *Biochem Syst Ecol* 68:142–147. <https://doi.org/10.1016/j.bse.2016.07.006>
- Parnikoza IY, Maidanuk DN, Kozeretska IA (2007) Are *Deschampsia antarctica* Desv. and *Colobanthus quitensis* (Kunth) Bartl. migratory relicts? *Cytol Genet* 41:226–229
- Parnikoza I, Kozeretska I, Kunakh V (2011) Vascular plants of the maritime antarctic: origin and adaptation. *Am J Plant Sci* 2:381–395. <https://doi.org/10.4236/ajps.2011.23044>
- Parnikoza I, Dykyy I, Ivanets V, Kozeretska I, Kunakh V, Rozhok A, Ochyra R, Convey P (2012) Use of *Deschampsia antarctica* for nest building by the kelp gull in the Argentine Islands area (maritime Antarctica) and its possible role in plant dispersal. *Polar Biol* 35:1753–1758. <https://doi.org/10.1007/s00300-012-1212-5>
- Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol Ecol Notes* 6:288–295. <https://doi.org/10.1111/1j.1471-8286.2005.01155.x>
- Peakall R, Smouse PE (2012) GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* 28:2537–2539. <https://doi.org/10.1093/bioinformatics/bts460>
- Peat HJ, Clarke A, Convey P (2007) Diversity and biogeography of the Antarctic flora. *J Biogeogr* 34:132–146. <https://doi.org/10.1111/j.1365-2699.2006.01565.x>
- Pecl GT, Araújo MB, Bell JD, Blanchard J, Bonebrake TC, Chen I-C, Clark TD, Colwell RK, Danielsen F, Evengård B, Falconi L, Ferrier S, Frusher S, Garcia RA, Griffis RB, Hobday AJ, Janion-Scheepers C, Jarzyna MA, Jennings S, Lenoir J, Linnæved HI, Martin VY, McCormack PC, McDonald J, Mitchell NJ, Mustonen T, Pandolfi JM, Pettorelli N, Popova E, Robinson SA, Scheffers BR, Shaw JD, Sorte CJB, Strugnell JM, Sunday JM, Tuanmu M-N, Vergés A, Villanueva C, Wernberg T, Wapstra E, Williams SE (2017) Biodiversity redistribution under climate change: impacts on ecosystems and human well-being. *Science* 355:eaai9214. <https://doi.org/10.1126/science.aai9214>
- Pertierra LR, Aragón P, Shaw JD, Bergstrom DM, Terauds A, Olalla-Tárraga MÁ (2017a) Global thermal niche models of two European grasses show high invasion risks in Antarctica. *Glob Chang Biol* 23:2863–2873. <https://doi.org/10.1111/gcb.13596>

- Pertierra LR, Hughes KA, Vega GC, Olalla-Tárraga MÁ (2017b) Correction: high resolution spatial mapping of human footprint across Antarctica and its implications for the strategic conservation of avifauna. *PLoS ONE* 12:e0173649. <https://doi.org/10.1371/journal.pone.0173649>
- Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D (2006) Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet* 38:904–909. <https://doi.org/10.1038/ng1847>
- Pritchard JK, Stephens P, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* 155:945–959
- Pugh PJA, Convey P (2008) Surviving out in the cold: Antarctic endemic invertebrates and their refugia. *J Biogeogr* 35:2176–2186. <https://doi.org/10.1111/j.1365-2699.2008.01953.x>
- Rousset F (1997) Genetic differentiation and estimation of gene flow from F-statistics under isolation by distance. *Genetics* 145:1219–1228
- Sanhueza C, Vallejos V, Cavieres LA, Saez P, Bravo LA, Corcuera LJ (2017) Growing temperature affects seed germination of the antarctic plant *Colobanthus quitensis* (Kunth) Bartl. (*Caryophyllaceae*). *Polar Biol* 40:449–455. <https://doi.org/10.1007/s00300-0-016-1972-4>
- Santiago IF, Alves TMA, Rabello A, Sales PA, Romanha AJ, Zani CL, Rosa CA, Rosa LH (2012) Leishmanicidal and antitumoral activities of endophytic fungi associated with the Antarctic angiosperms *Deschampsia antarctica* Desv. and *Colobanthus quitensis* (Kunth) Bartl. *Extremophiles* 16:95–103. <https://doi.org/10.1007/s00792-011-0409-9>
- Santiago IF, Rosa CA, Rosa LH (2017) Endophytic symbiont yeasts associated with the Antarctic angiosperms *Deschampsia antarctica* and *Colobanthus quitensis*. *Polar Biol* 40:177–183. <https://doi.org/10.1007/s00300-016-1940-z>
- Smýkal P, Bačová-Kerteszová N, Kalendar R, Corander J, Schulman AH, Pavelek M (2011) Genetic diversity of cultivated flax (*Linum usitatissimum* L.) germplasm assessed by retrotransposon-based markers. *Theor Appl Genet* 122:1385–1397. <https://doi.org/10.1007/s00122-011-1539-2>
- Sneddon BV (1999) The Taxonomy and Breeding System of *Colobanthus squarrosus* (Caryophyllaceae). *New Zeal J Bot* 37:195–204. <https://doi.org/10.1080/0028825X.1999.9512627>
- Squier AH, Hodgson DA, Keely BJ (2005) Evidence of late Quaternary environmental change in a continental East Antarctic lake for lacustrine sedimentary pigment distributions. *Antarct Sci* 17:361–376. <https://doi.org/10.1017/S0954102005002804>
- Taberlet P, Fumagalli L, Wust-Saucy AG, Cosson JF (1998) Comparative phylogeography and postglacial colonization routes in Europe. *Mol Ecol* 7:453–464. <https://doi.org/10.1046/j.1365-294x.1998.00289.x>
- Terauds A, Chown SL, Morgan F, Peat HJ, Watts D, Keys H, Convey P, Bergstrom DM (2012) Conservation biogeography of the Antarctic. *Divers Distrib* 18:726–741. <https://doi.org/10.1111/j.1472-4642.2012.00925.x>
- van de Wouw M, van Dijk P, Huiskes AHL (2008) Regional genetic diversity patterns in Antarctic hairgrass (*Deschampsia antarctica* Desv.). *J Biogeogr* 35:365–376. <https://doi.org/10.1111/j.1365-2699.2007.01784.x>
- Vasemagi A (2006) The adaptive hypothesis of clinal variation revisited: single-locus clines as a result of spatially restricted gene flow. *Genetics* 173:2411–2414. <https://doi.org/10.1534/genetics.106.059881>
- Volkov RA, Kozeretska IA, Kyryachenko SS, Andreev IO, Maidanyuk DN, Parnikoza IY, Kunakh VA (2010) Molecular evolution and variability of ITS1–ITS2 in populations of *Deschampsia antarctica* from two regions of the maritime Antarctic. *Polar Sci* 4:469–478. <https://doi.org/10.1016/j.polar.2010.04.011>
- West JG (1991) *Colobanthus curtisiae* (Caryophyllaceae), a new species from Eastern Australia. In: Banks MR, Smith SJ, Orchard AE, Kantvilas G (eds) *Aspects of Tasmanian botany: a tribute to Winifred Curtis*. Royal Society of Tasmania, Hobart, pp 75–77
- Wódkiewicz M, Chwedorzewska KJ, Bednarek PT, Znój A, Androsiuk P, Galera H (2018) How much of the invader's genetic variability can slip between our fingers? A case study of secondary dispersal of *Poa annua* on King George Island (Antarctica). *Ecol Evol* 8:592–600. <https://doi.org/10.1002/ece3.3675>
- Wright SI, Gaut BS (2005) Molecular population genetics and the search for adaptive evolution in plants. *Mol Biol Evol* 22:506–519. <https://doi.org/10.1093/molbev/msi035>
- Znój A, Chwedorzewska KJ, Androsiuk P, Cuba-Diaz M, Gielwanowska I, Koc J, Korczak-Abshire M, Grzesiak J, Zmarz A (2017) Rapid environmental changes in the Western Antarctic Peninsula region due to climate change and human activity. *Appl Ecol Env Res* 15:525–539. https://doi.org/10.15666/aeer/1504_525539