

1 **COCOA INTAKE ATTENUATES OXIDATIVE STRESS**
2 **ASSOCIATED WITH RAT ADJUVANT ARTHRITIS**

3
4 **Authors:** S. Ramos-Romero¹, F.J. Pérez-Cano^{1,2}, E. Ramiro-Puig¹, A. Franch^{1,2}, M.
5 Castell^{1,2*}.

6
7 **Institutional Address:**

8 ¹ Departament de Fisiologia, Facultat de Farmàcia, Universitat de Barcelona, Spain

9 ² Institut de Recerca en Nutrició i Seguretat Alimentària (INSA-UB), Barcelona, Spain

10

11 **e-Mail Addresses:**

12 sara.ramosromero@ub.edu, franciscoperez@ub.edu, emma.ramiro@gmail.com,
13 angelsfranch@ub.edu

14

15 ***Corresponding author:**

16 Margarida Castell, PhD

17 Department de Fisiologia, Facultat de Farmàcia, Universitat de Barcelona

18 Av Joan XXIII s/n, Edifici B, 3^a planta

19 08028, Barcelona, Spain

20 Phone: +34 93 402 45 05

21 Fax: +34 93 403 59 01

22 e-mail address: margaridacastell@ub.edu

23 **ABSTRACT**

24

25 Cocoa contains flavonoids with antioxidant properties. The aim of this study was to
26 ascertain the effect of cocoa intake on oxidative stress associated with a model of
27 chronic inflammation such as adjuvant arthritis. Female Wistar rats were fed with a 5 or
28 10% cocoa-enriched diet or were given p.o. a quercetin suspension every other day for
29 10 days. Arthritis was induced by a heat-killed *Mycobacterium butyricum* suspension.
30 Reactive oxygen species (ROS) produced by macrophages, and splenic superoxide
31 dismutase (total, cytoplasmic and mitochondrial) and catalase activities were
32 determined. Clinically, joint swelling in arthritic rats was not reduced by antioxidants;
33 however, the 5% cocoa diet and quercetin administration reduced ROS production.
34 Moreover, the 5% cocoa diet normalized the activities of superoxide dismutase and
35 catalase. In conclusion, a cocoa diet reduces the oxidative stress associated with a
36 chronic inflammatory pathology, although it was not enough to attenuate joint swelling.

37

38 **Keywords:** cocoa, oxidative stress, adjuvant arthritis

39

40 **Abbreviations:** AA, arthritis reference animals; AIN, American Institute of Nutrition;
41 C5, arthritic animals fed 5% cocoa; C10, arthritic animals fed 10% cocoa; EGTA,
42 ethylene glycol tetraacetic acid; H₂DCF-DA, dichlorofluorescein diacetate; Q, arthritic
43 animals treated p.o. with quercetin; p.o., 'per os' oral administration; RA, rheumatoid
44 arthritis; REF, healthy reference animals; RNS, reactive nitrogen species; ROS, reactive
45 oxygen species; SOD, superoxide dismutase; TNF- α , tumour necrosis factor- α

46

47 **1. Introduction**

48

49 Flavonoids are polyphenol antioxidant compounds found in vegetables. These
50 benzo- γ -pyrone derivatives can prevent injury caused by free radicals by neutralizing
51 free radicals, chelating metals (principally Fe^{2+} and Cu^+) that enhance reactive oxygen
52 species, inhibiting enzymes, and regulating antioxidant defense [1]. Cocoa (product of
53 *Theobroma cacao*) is a natural flavonoid source, mainly containing flavanols such as
54 epicatechin, catechin and procyanidins, and smaller amounts of other flavonoids such as
55 quercetin (3,3',4',5,7-pentahydroxyflavone) [2]. The antioxidant effects of quercetin
56 and catechin have been extensively studied on a variety of inflammatory processes and
57 immune functions [3]. Moreover, other flavanols present in cocoa, such as
58 epigallocatechin-3-gallate and procyanidin B₂, protect Caco2 cells against oxidative
59 stress by reducing ROS production, preventing caspase-3 activation, and increasing
60 antioxidant enzymes as glutathione peroxidase, glutathione reductase and glutathione-S-
61 transferase [4]. Nevertheless, the antioxidant efficacy of flavonoids *in vivo* is less
62 documented. Previous studies show that cocoa flavonoids reduce the macrophage ability
63 to release inflammatory mediators including ROS and nitric oxide *in vitro* and in
64 healthy animals [5-7]. Moreover, a cocoa diet increases the activity of the antioxidant
65 enzymes superoxide dismutase (SOD) and catalase in healthy rat thymus [8].

66

67 Rheumatoid arthritis (RA) is a chronic, systemic, autoimmune and inflammatory
68 disease characterized by joint inflammation and synovial hyperplasia, involving
69 infiltration of activated T cells and macrophages. Macrophages possess broad
70 proinflammatory, destructive and remodeling capacities, and contribute considerably to
71 inflammation and joint destruction in RA. Activation of macrophages is not restricted to

72 the synovial compartment, but extends to circulating monocytes and other cells of the
73 mononuclear phagocyte system [9]. It is believed that macrophages act as amplifiers of
74 systemic inflammation and it has been suggested that RA severity may be associated
75 with the degree of systemic activation of monocytes/macrophages [9]. Activated
76 macrophages release inflammatory mediators such as proteolytic enzymes, ROS and
77 reactive nitrogen species (RNS) [10]. Among ROS, the superoxide anion ($O_2^{\bullet-}$)
78 increases the vascular permeability and promotes neutrophil migration and can be
79 converted into hydroxyl radical (HO^{\bullet}), becoming more aggressive [11]. The SOD is an
80 endogenous ubiquitous enzyme that catalyzes the $O_2^{\bullet-}$ dismutation into H_2O_2 .
81 Subsequently, the endogenous enzyme catalase transforms it to H_2O and O_2 . SOD
82 shows three isoforms: the cytoplasmic SOD1, the mitochondrial SOD2, and the
83 extracellular SOD3. SOD1 and SOD3 use copper and zinc as cofactors, and manganese
84 is the cofactor of SOD2 [11].

85

86 Oxidative stress must be important in the pathogenesis of RA since oxidation
87 markers and impaired antioxidant status are found in the plasma and synovial fluid of
88 RA patients [12-13]. Moreover, several studies have pointed out the beneficial effect of
89 antioxidants in RA patients [14-15]. Due to the antioxidant properties demonstrated by
90 cocoa, it could be hypothesized that cocoa intake modulates the oxidative stress
91 provoked by an inflammatory disease. The aim of this study was to ascertain the effect
92 of cocoa-enriched diets on the oxidant and antioxidant status of rats with adjuvant
93 arthritis considering oxidative stress of peritoneal macrophages, applied in several
94 studies as systemic inflammatory cells [16-17], and antioxidant status in a secondary
95 lymphoid tissue very rich in lymphocytes and monocytes, as the spleen. At the same

96 time, a group treated with the flavonoid quercetin, with recognized antioxidant and anti-
97 inflammatory activities was added [18].

98 **2. Materials and methods**

99 *2.1. Animals and diets*

100 Nine-week-old female outbred Wistar rats (Harlan, Barcelona, Spain) were fed a
101 standard diet, formulated according to the American Institute of Nutrition (AIN-93M),
102 or a diet containing 5% or 10% of partially defatted *Natural Forastero* cocoa (Nutrexpa,
103 Barcelona, Spain) with 21.2 mg of total phenols/g (Folin-Ciocalteu method). Five and
104 ten percent cocoa diets were prepared from a basal mix (Harlan), in which the
105 proportion of proteins, carbohydrates, lipids and fiber had been modified in such a way
106 that the addition of 5 or 10% cocoa resulted in a final isoenergetic diet with the same
107 macronutrient composition as the AIN-93M diet (Table 1). The diets began 14 days
108 before the arthritis induction and lasted until the end of the study (six weeks later)
109 (Fig. 1).

110

111 Animals were randomly distributed in 5 different experimental groups
112 (11-12 rats/group): REF (healthy reference animals), AA (arthritis reference animals), Q
113 (arthritic animals treated p.o. with quercetin at 200 mg/kg on days 0, 2, 4, 6, 8 and 10
114 after arthritis induction), C5 (arthritic animals fed 5% cocoa) and C10 (arthritic animals
115 fed 10% cocoa) (Fig. 1). Studies were performed in accordance with the institutional
116 guidelines for the care and use of laboratory animals established by the Ethical
117 Committee for Animal Experimentation at the University of Barcelona and approved by
118 the Catalanian Government.

119

120 *2.2. Induction and evaluation of adjuvant arthritis*

121 Arthritis was induced in all the animals with the exception of the REF group.
122 Adjuvant arthritis was induced as in previous studies [19-20] by injecting intradermally

123 a suspension of 0.5 mg of heat-killed *Mycobacterium butyricum* (Difco, Detroit, MI,
124 USA) in 0.1 ml of liquid vaseline into the base of the rat tail. Arthritis was clinically
125 assessed by means of hind-paw volume (water plethysmometer LI 7500 Letica, Spain)
126 and arthritic score. The arthritis score was calculated as the sum of the clinical score of
127 the four paws according to scores established by two observers in a blind manner
128 following the criteria: '0' means no signs of inflammation, '1' means erythema or slight
129 swelling in the paw articulations, '2' means erythema and moderate swelling in the paw
130 articulations, '3' means erythema and deep swelling in the paw articulations, '4' means
131 erythema and severe swelling and immobility.

132

133 Body weight, hind-paw volume and arthritis score were determined weekly in all
134 the studied groups. On day 28 post-induction, animals were anaesthetized by an
135 intramuscular injection of ketamine (90 mg/kg; Merial, Lyon, France) and xylazine (10
136 mg/kg; Bayer HealthCare, Kiel, Germany), and peritoneal macrophages, as immune
137 cells involved in the systemic inflammation, and the spleen, as immune cell storage and
138 secondary lymphoid tissue, were obtained. Spleen fragments were immediately frozen
139 at -80 °C until analysis.

140

141 *2.3. Peritoneal macrophages isolation and ROS production*

142 Peritoneal macrophages were obtained by injecting 40 mL of ice-cold sterile
143 PBS (pH 7.2) into the peritoneal cavity. After 2 min of abdominal massages, cell
144 suspension was aspirated. Macrophages were plated and allowed to attach overnight,
145 then they were washed with warm RPMI medium without phenol red (Sigma-Aldrich)
146 containing 100 IU/mL streptomycin-penicillin. Macrophages were incubated with 20
147 $\mu\text{mol/L}$ of reduced 2',7'-dichlorofluorescein diacetate probe ($\text{H}_2\text{DCF-DA}$; Invitrogen,

148 Paisley, UK) for 30 min at 37 °C. Macrophage ROS oxidized H₂DCF to a fluorescent
149 compound (DCF). Fluorescence was measured at 30, 60 and 90 min by a fluorometer
150 (excitation 538 nm, emission 485 nm).

151

152 *2.4. Spleen superoxide dismutase*

153 A fragment of spleen was homogenized in cold 0.1 g/mL HEPES buffer (20 mM
154 pH 7.2, containing 1 mM ethylene glycol tetraacetic acid EGTA, 210 mM mannitol and
155 70 mM sucrose) and centrifuged (1,500 g, 5 min, 4 °C). The total SOD activity was
156 determined in the supernatant using superoxide dismutase assay kit II (Merck KGaA,
157 Darmstadt, Germany) following the manufacturer's instructions. To determine SOD
158 subtypes, homogenate supernatant was centrifuged at 10,000 g. The supernatant and the
159 pellet were used to evaluate the cytoplasmic and mitochondrial SOD activities,
160 respectively. In the mitochondrial SOD quantification, the pellet was resuspended in
161 HEPES buffer, and potassium cyanide (3 mM, Sigma-Aldrich) was added to the
162 samples to inhibit cytoplasmic SOD.

163

164 One unit of SOD is defined as the amount of enzyme needed to exhibit 50%
165 dismutation of the superoxide radical. SOD activity was expressed as units of SOD per
166 g of protein from spleen homogenates (quantified following the Bradford method using
167 the Bio-Rad Protein Assay of Bio-Rad Laboratories Inc., Hercules, CA).

168

169 *2.5. Spleen catalase activity*

170 A spleen fragment was homogenized on ice-cold phosphate buffer (0.1 g/mL; 50
171 mM K₂HPO₄, 50 mM KH₂PO₄ and 1 mM EDTA, pH 7.0) and centrifuged (10,000 g,
172 15 min, 4 °C). The catalase activity was determined in the supernatant by using a

173 catalase assay kit (Merck) following the manufacturer's instructions. The detection
174 method was based on the reaction of the enzyme with methanol in the presence of an
175 optimal concentration of H₂O₂, and the formaldehyde produced was measured
176 spectrophotometrically with 4-amino-3-hydrazino-5-mercapto-1,2,4-triazole as the
177 chromogen. Catalase activity was expressed as μmol of formaldehyde per min per g of
178 protein from spleen homogenates.

179

180 2.6. *Statistics*

181 The software package PASW Statistics 18.0 (SPSS Inc., Chicago, IL) was used
182 for statistical analysis. Levene's and Kolmogorov-Smirnov's tests were applied to assay
183 variance equality and normal distribution of the studied groups, respectively. The one-
184 way analysis of variance (ANOVA) followed by Scheffé's post hoc significance test
185 was applied when the assumptions of normality and equal variance were met. However,
186 non-parametric tests (Kruskal-Wallis and Mann-Whitney U) were used to assay
187 significance. Chi squared test was applied in the frequency study. Significant
188 differences were accepted when $P < 0.05$ and were labeled in tables and figures with a
189 different letter.

190 **3. Results and discussion**

191 Rheumatoid arthritis, as well as experimental models of arthritis, have been
192 associated with oxidative stress [12-13]. Therefore, foods with antioxidant properties
193 could help in the treatment of RA decreasing this harmful state [21]. The results
194 obtained here show the effects of cocoa-enriched diets and quercetin treatment on some
195 markers of oxidative stress in adjuvant arthritis after 4 weeks of induction.

196

197 The inflammatory status of adjuvant arthritis was measured by means of
198 hind-paw volume, arthritis score and body weight increase. As can be seen in Table 2,
199 maximum hind-paw volume in the AA group was achieved 3 weeks after induction and
200 thereafter tended to reduce. Similar results were observed when considering arthritis
201 score. Quercetin group, which was includes as a positive control, did not prevent
202 arthritis as we expected and even induced a more precocious appearance of arthritis
203 (measured by hind-paw volume or arthritis score, Table 2). These results do not agree
204 with those of Mamani-Matsuda and coworkers [18] who show the curative effect of
205 quercetin (30 mg every 2 days from day 10 after induction) in adjuvant arthritis induced
206 in 6 week-old Lewis rats. The differences could be attributed to events that would
207 change susceptibility to this flavonoid, such as the rat strain (Wistar vs Lewis), age of
208 animals (9-week vs 6-week old animals), the protocol (preventive vs curative) or even
209 the dose (about 40 mg per rat vs 30 mg). Similarly, cocoa diets (groups C5 and C10)
210 were not able to prevent joint swelling, which actually was already detected at day 7
211 post-induction. However, after maximum paw swelling on day 21 after induction, the
212 animals from groups C5 and C10 underwent a faster recuperation than the AA rats.
213 These results correlate with the body weight increase: during the first two weeks, body
214 weight decreased in all groups receiving arthritis induction, thereafter body weight rose

215 and this recovery was faster in the C5 and C10 groups than in the AA and Q groups
216 (Table 2).

217

218 Macrophages, obtained 4 weeks after arthritis induction, allow the potential
219 oxidative stress in adjuvant arthritis to be estimated (Fig. 2). Macrophages from the AA
220 animals synthesized higher amounts of ROS than those from the REF group
221 ($P < 0.001$). Macrophages from the C5 and Q groups, but not from the C10 group,
222 produced lower ROS than the AA rats ($P < 0.05$). These results indicate the attenuation
223 of the oxidative stress by certain doses of cocoa flavonoids. Similarly, the increased
224 ROS synthesis in AA rats and their modulation by antioxidant compounds has been
225 reported [22-24]. Moreover, the attenuation of ROS by a cocoa diet is in line with
226 results derived from macrophages obtained from cocoa-fed rats, both healthy ones [6]
227 and those with collagen-induced arthritis [7]. Likewise, administration of quercetin was
228 able to decrease ROS concentration in a kidney inflammation model [25]. The
229 inhibition of ROS synthesis by flavonoids can be associated with their free-radical
230 quenching activity [26]. However, it should be noted that the richest cocoa diet did not
231 produce antioxidant effects. This result does not agree with *in vitro* studies showing a
232 dose-dependent antioxidant effect of cocoa [5, 27], which could be due to the different
233 compounds reaching cells. Thus, the *in vitro* antioxidant activity could be partially due
234 to large flavonoid polymers found in cocoa [1], but these polymers are not intestinally
235 absorbed [28]. On the other hand, flavonoids can act as pro-oxidants when they are in
236 high concentration *in vivo* [1, 29]. Previous studies have shown that low concentrations
237 of quercetin inhibit lipid peroxidation in rat liver and protect human leucocytes against
238 oxidative DNA damage, but high concentrations of this flavonoid enhance the hydroxyl
239 radical formation and increase the DNA damage [1].

240

241 In order to protect tissues from oxidative injuries, the body possesses enzymatic
242 antioxidant enzymatic systems such as superoxide dismutases and catalase enzymes. It
243 has been reported that AA decreases serum or synovial SOD and catalase activities
244 together with other endogenous antioxidant systems [22-24, 30]. Here we found that, 4
245 weeks after induction, arthritis produced a decrease in splenic catalase activity
246 ($P < 0.05$; Fig. 3) and, paradoxically, an increase in splenic total and mitochondrial
247 SOD ($P < 0.05$; Fig. 4). The decreased catalase activity could be associated with the
248 consumption of catalase in neutralizing the H_2O_2 . On the other hand, increased splenic
249 SOD activities could reflect the response of the body to increased ROS concentrations
250 and/or it could be due to the fact that arthritis was in its recovery phase one month after
251 its induction. Moreover, SOD increase could also be explained by the increase in the
252 oxidative stress found in arthritic rats (Fig. 2), and by the increased tumour necrosis
253 factor- α (TNF- α) secretion present in arthritis [7]. Both oxidative stress and TNF- α are
254 shown to induce SOD synthesis [31-32]. It should be added that a similar increase in
255 SOD activity was found in the plasma of RA patients [33] and in the synovial
256 membrane of mice with collagen-induced arthritis [34].

257

258 Interestingly, the changes in the enzymatic antioxidant systems produced by
259 adjuvant arthritis were totally prevented in the C5 group which was fed with a diet
260 containing 5% cocoa ($P < 0.05$; Fig. 3 and 4). SOD and catalase activities in the C10
261 and Q groups were not significantly different from the AA rats (Fig. 3 and 4). The
262 beneficial effect of a 5% cocoa diet on antioxidant systems in adjuvant arthritis is in line
263 with the effect of other flavonoids or compounds of a botanical origin on this
264 experimental inflammatory model [24, 30], although these studies found a clinical

265 improvement on arthritis. The reduction of oxidative stress by a 5% cocoa diet (Fig. 2),
266 as well as the decrease in TNF- α secretion produced by a similar diet as reported in a
267 previous study [7] could be responsible for the reduction in splenic SOD. The effect of
268 5% cocoa diet on SOD seems to be due to its activity on the mitochondrial SOD
269 (Fig. 4C), a unique and essential isoform for life [35]. On the other hand, cocoa intake
270 normalized the catalase activity (Fig. 3), which is a cytosolic enzyme absent in the
271 mitochondria of most cells [36]. These results suggest that the antioxidant effect of
272 cocoa is exerted in both cellular compartments.

273

274 From the results obtained in this study, it can be concluded that cocoa intake
275 reduces the oxidative stress in macrophages provoked by adjuvant arthritis. This
276 antioxidant effect of cocoa is dependent on the doses employed. Nevertheless, we still
277 need to discover the antioxidant effect of cocoa on joint tissues and to ascertain the
278 relationship between this antioxidant effect and inflammatory markers. Further studies
279 are needed to establish more deeply the role of cocoa, a source of antioxidant
280 flavonoids, as a coadjuvant in the treatment of chronic inflammatory diseases.

281

282 **Acknowledgements**

283 This study was supported by the Ministerio de Educación y Ciencia, Spain (AGL2005-
284 002823), and by the Generalitat de Catalunya, Spain (SGR 2005-0083). S.R. is the
285 recipient of a fellowship from the Ministerio de Educación y Ciencia, Spain (BES-2006-
286 13640).

287

288

289 **References**

- 290 [1] Procházková D, Boušová I, Wilhelmová N. Antioxidant and prooxidant properties
291 of flavonoids. *Fitoterapia* 2011;82:513-23.
- 292 [2] Rusconi M, Conti A. *Theobroma cacao* L., the Food of the Gods: A scientific
293 approach beyond myths and claims. *Pharmacol Res* 2010;61:5-13.
- 294 [3] Boots AW, Haenen GR, Bast A. Health effects of quercetin: from antioxidant to
295 nutraceutical. *Eur J Pharmacol* 2008;585:325-37.
- 296 [4] Rodríguez-Ramiro I, Ramos S, Bravo L, Goya L, Martín MA. Procyanidin B2
297 induces Nrf2 translocation and glutathione S-transferase P1 expression via ERKs
298 and p38-MAPK pathways and protect human colonic cells against oxidative stress.
299 *Eur J Nutr* 2011;in press.
- 300 [5] Ramiro E, Franch A, Castellote C, Pérez-Cano F, Permanyer J, Izquierdo-Pulido M,
301 Castell M. Flavonoids from *Theobroma cacao* down-regulate inflammatory
302 mediators. *J Agric Food Chem* 2005;53:8506-11.
- 303 [6] Castell M, Franch A, Ramos-Romero S, Ramiro-Puig E, Pérez-Cano FJ, Castellote
304 C. Effect of a diet rich in cocoa flavonoids on experimental acute inflammation. In:
305 Keller RB, editors. *Flavonoids: Biosynthesis, Biological Effects and Dietary*
306 *Sources*, New York: Nova Science Publishers Inc; 2009, p. 213-29.
- 307 [7] Ramos-Romero S, Pérez-Cano FJ, Pérez-Berezo T, Castellote C, Franch A, Castell
308 C. Effect of a cocoa flavonoid-enriched diet on experimental autoimmune arthritis.
309 *Br J Nutr* 2012;107:523-32.
- 310 [8] Ramiro-Puig E, Urpí-Sardà M, Pérez-Cano FJ, Franch À, Castellote C, Andrés-
311 Lacueva C, Izquierdo-Pulido M, Castell M. Cocoa-enriched diet enhances
312 antioxidant enzyme activity and modulates lymphocyte composition in thymus from
313 young rats. *J Agric Food Chem* 2007;55:6431-8.

- 314 [9] Kinne RW, Bräuer R, Stuhlmüller B, Palombo-Kinne E, Burmester GR.
315 Macrophages in rheumatoid arthritis. *Arthritis Res.* 2000;2:189-202.
- 316 [10] Phillips DC, Dias HKI, Kitas GD, Griffiths HR. Aberrant reactive oxygen and
317 nitrogen species generation in rheumatoid arthritis (RA): Causes and consequences
318 for immune function, cell survival, and therapeutic intervention. *Antioxid Redox*
319 *Signal* 2010;12:743-85.
- 320 [11] Afonso V, Champy R, Mitrovic D, Collin P, Lomri A. Reactive oxygen species and
321 superoxide dismutases: Role in joint diseases. *Joint Bone Spine* 2007;74:324-9.
- 322 [12] Seven A, Güzel S, Aslan M, Hamuryudan V. Lipid, protein, DNA oxidation and
323 antioxidant status in rheumatoid arthritis. *Clin Biochem* 2008;41:538-43.
- 324 [13] Vasanthi P, Nalini G, Rajasekhar G. Status of oxidative stress in rheumatoid
325 arthritis. *Int J Rheum Dis* 2009;12:29-33.
- 326 [14] Tiku ML, Gupta S, Deshmukh DR. Aggrecan degradation in chondrocytes is
327 mediated by reactive oxygen species and protected by antioxidants. *Free Radic Res*
328 1999;30:395-405.
- 329 [15] Nourmohammadi I, Athari-Nikazm S, Vafa MR, Bidari A, Jazayeri S, Hoshyarrad
330 A, Hoseini F, Fasihi-Radmandi M. Effects of antioxidant supplementations on
331 oxidative stress in rheumatoid arthritis patients. *J Biol Sci* 2010;10:63-6.
- 332 [16] Kauss T, Moynet D, Rambert J, Al-Kharrat A, Brajot S, Thiolat D, Enneman R,
333 Fawaz F, Mossalayi MD. Rutoside decreases human macrophage-derived
334 inflammatory mediators and improves clinical signs in adjuvant-induced arthritis.
335 *Arthritis Res Ther.* 2008;10:R19.
- 336 [17] Li R, Li J, Cai L, Hu CM, Zhang L. Suppression of adjuvant arthritis by hesperidin
337 in rats and its mechanisms. *J Pharm Pharmacol.* 2008;60:221-8.
- 338


- 339 [18] Mamani-Matsuda M, Kauss T, Al-Kharrat A, Rambert J, Fawaz F, Thiolat D,
340 Moynet D, Coves S, Malvy D, Mossalayi MD. Therapeutic and preventive
341 properties of quercetin in experimental arthritis correlate with decreased
342 macrophage inflammatory mediators. *Biochem Pharmacol.* 2006;72:1304-10.
- 343 [19] Franch A, Castellote C, Vilà JL, Vilaró S, Castell M. Anticytoskeletal autoantibody
344 development in adjuvant arthritis. *J Rheumatol.* 1994;21:489-97.
- 345 [20] Pelegrí C, Castell M, Serra M, Rabanal M, Rodríguez-Palmero M, Castellote C,
346 Franch A. Prevention of adjuvant arthritis by the W3/25 anti-CD4 monoclonal
347 antibody is associated with a decrease of blood CD4⁺ CD45RC^{high} T cells. *Clin Exp*
348 *Immunol.* 2001;125:470-7.
- 349 [21] Efthimiou P, Kukar M. Complementary and alternative medicine use in rheumatoid
350 arthritis: proposed mechanism of action and efficacy of commonly used modalities.
351 *Rheumatol Int* 2010;30:571-86.
- 352 [22] Jung HJ, Nam JH, Choi J, Lee KT, Park HJ. Antiinflammatory effects of
353 chiisanoside and chiisanogenin obtained from the leaves of *Acanthopanax*
354 *chiisanensis* in the carrageenan- and Freund's complete adjuvant-induced rats. *J*
355 *Ethnopharmacol* 2005;97:359-67.
- 356 [23] Jawed H, Shah SUA, Jamall S, Simjee SU. N-(2-hydroxy phenyl) acetamide
357 inhibits inflammation-related cytokines and ROS in adjuvant-induced arthritic
358 (AIA) rats. *Int Immunopharmacol* 2010;10:900-5.
- 359 [24] Tastekin N, Aydogdu N, Dokmeci D, Usta U, Birtane M, Erbas H, Ture M.
360 Protective effects of l-carnitine and alpha-lipoic acid in rats with adjuvant arthritis.
361 *Pharmacol Res* 2007;56:303-10.

- 362 [25] Liu CM, Ma JQ, Sun YZ. Quercetin protects the rat kidney against oxidative stress-
363 mediated DNA damage and apoptosis induced by lead. *Environ Toxicol Pharmacol*
364 2010;30:264-71.
- 365 [26] Avila V, Bertolotti SG, Criado S, Pappano N, Debattista N, García NA.
366 Antioxidant properties of natural flavonoids: quenching and generation of singlet
367 molecular oxygen. *Int J Food Sci Technol* 2001;36:25-33.
- 368 [27] Erlejman AG, Fraga CG, Oteiza PI. Procyanidins protect Caco-2 cells from bile
369 acid- and oxidant-induced damage. *Free Radic Biol Med* 2006;41:1247-56.
- 370 [28] Monagas M, Urpi-Sarda M, Sánchez-Patán F, Llorach R, Garrido I, Gómez-
371 Cordovés C, Andres-Lacueva C, Bartolomé B. Insights into the metabolism and
372 microbial biotransformation of dietary flavan-3-ols and the bioactivity of their
373 metabolites. *Food Funct* 2010;1:233-53.
- 374 [29] Halliwell B. Are polyphenols antioxidants or pro-oxidants? What do we learn from
375 cell culture and in vivo studies? *Arch Biochem Biophys* 2008;476:107-12.
- 376 [30] Kripa KG, Chamundeeswari D, Thanka J, Uma Maheswara Reddy C. Modulation
377 of inflammatory markers by the ethanolic extract of *Leucas aspera* in adjuvant
378 arthritis. *J Ethnopharmacol* 2011;134:1024-7.
- 379 [31] Kuo S, Cheswron SE, Mellott JK, Rogers RJ, Hsu JL, Nick HS. In vivo
380 architecture of the manganese superoxide dismutase promoter. *J Biol Chem*
381 1999;274:3345-54.
- 382 [32] Xu Y, Kiningham KK, Devalaraja MN, Yeh CC, Majima H, Kasarskis EJ, St Clair
383 DK. An intronic NF- κ B element is essential for induction of the human manganese
384 superoxide dismutase gene by tumor necrosis factor- α and interleukin-1 β . *DNA Cell*
385 *Biol* 1999;18:709-22.

- 386 [33] Bhowmick K, Chakraborti G, Gudi NS, Kutty Moideen AV, Shetty HV. Free
387 radical and antioxidant status in rheumatoid arthritis. *Indian J Rheumatol* 2008;3:8-
388 12.
- 389 [34] Kasama T, Kobayashi K, Sekine F, Negishi M, Ide H, Takahashi T, Niwa Y.
390 Follow-up study of lipid peroxides, superoxide dismutase and glutathione
391 peroxidase in the synovial membrane, serum and liver of young and old mice with
392 collagen-induced arthritis. *Life Sci* 1988;43:1887-96.
- 393 [35] Lebovitz RM, Zhang H, Vogel H, Cartwright JJr, Dionne L, Lu N, Huang S,
394 Matzuk MM. Neurodegeneration, myocardial injury, and perinatal death in
395 mitochondrial superoxide dismutase-deficient mice. *Proc Natl Acad Sci U.S.A.*
396 1996;93:9782-7.
- 397 [36] Hitchon CA, El-Gabalawy HS. Oxidation in rheumatoid arthritis. *Arthritis Res*
398 *Ther* 2004;6:265-78.

399 **Figure legends**


400 **Fig. 1.** Diagram of the experimental design that began 14 days before arthritis induction
401 and finished on day 28 post-induction. White bars mean the standard diet (Ref, AA and
402 Q groups), pale grey color represents the 5% cocoa diet (C5 group), and dark grey color
403 represents the 10% cocoa diet (C10 group). Striped bars are representative of period
404 after arthritis induction. Animals from Q group were treated with quercetin p.o. on days
405 0, 2, 4, 6, 8, 10 post-induction (squared bar). At the end of the study, peritoneal
406 macrophages and spleen were obtained from all animals.



407

408


409 **Fig. 2.** Time course of ROS production by macrophages, expressed in fluorescence
410 units (FU) and determined by DCF assay. Values are expressed as mean \pm S.E.M. ($n =$
411 8). Means with different letter (a, b, c) differ, $P < 0.05$ (Kruskal-Wallis and Mann-
412 Whitney U tests).



413

414


415 **Fig. 3.** Catalase activity in spleen homogenates expressed as $\mu\text{mol}/\text{min}/\text{g}$ of protein.
416 Each bar represents the mean \pm S.E.M. ($n = 11-12$). Means with different letter (a, b)
417 differ, $P < 0.05$ (ANOVA followed by Scheffé's test).



418

419

420 **Fig. 4.** Activity of SOD subtypes in spleen homogenates: (A) total, (B) cytosolic, and
 421 (C) mitochondrial SOD activities. SOD is expressed as units (U)/g of protein. Each bar
 422 represents the mean \pm S.E.M. ($n = 11-12$). Means with different letter (a, b) differ, $P <$
 423 0.05 (Kruskal-Wallis and Mann-Whitney U tests).



424

425 **Tables**426 **Table 1.** Composition of the experimental diets (g/kg).

Components	Standard diet (AIN-93M, g/kg)	5% Cocoa diet (g/kg)	10% Cocoa diet (g/kg)
Casein	140	129	118
L-Cystine	1.8	1.8	1.8
Corn starch	465.69	457.69	449.69
Maltodextrin	155	155	155
Sucrose	100	100	100
Soybean oil	40	34.5	29
Cellulose	50	37.25	24.5
Mineral mix (TD94046)	35	35	35
Vitamin mix (TD94047)	10	10	10
Choline bitartrate	2.5	2.5	2.5
<i>tert</i> -Butylhydroquinone	0.008	0.008	0.008
Natural cocoa powder	-	50	100
Cocoa macronutrients:			
<i>Protein</i>		11	22
<i>Carbohydrate</i>		8	16
<i>Lipid</i>		5.5	11
<i>Insoluble fiber</i>		12.75	25.5
Total cocoa polyphenols ⁽¹⁾		1.06	2.12
Cocoa flavonoids ⁽²⁾ :			
<i>Epicatechin</i>		0.10995	0.2199
<i>Procyanidin B₂</i>		0.08375	0.1675
<i>Catechin</i>		0.0368	0.0736
<i>Isoquercetin</i>		0.00275	0.0055
<i>Quercetin</i>		0.0015	0.003

427

428 ¹Cocoa polyphenols were determined by Folin-Ciocalteu method.429 ²Main cocoa flavonoids were determined by HPLC.

430 **Table 2.** Effect of flavonoid intake on total body weight increase and arthritis severity
 431 (n = 11-12). Body weight (g) and hind-paw volume (mL) values are summarized as
 432 mean ± S.E.M. Means or frequencies with different letter (a, b, c) differ, P < 0.05
 433 (ANOVA followed by Scheffé's test). Arthritis score is presented as frequency of each
 434 score. Frequencies without a common letter differ, P < 0.05 (Chi square).

		post-induction day				
Group		7	14	21	28	
Body Weight	REF	211.7 ± 2.6 ^a	220.1 ± 3.1 ^a	227.1 ± 3.4 ^a	231.8 ± 3.6 ^a	
	AA	195.6 ± 4.1 ^b	194.8 ± 4.8 ^b	193.8 ± 5.2 ^b	200.2 ± 5.4 ^b	
	Q	192.2 ± 3.1 ^b	189.9 ± 2.7 ^b	195.1 ± 3.5 ^b	202.2 ± 3.7 ^b	
	C5	185.0 ± 2.5 ^c	183.1 ± 2.6 ^c	189.5 ± 2.5 ^b	198.9 ± 2.4 ^b	
	C10	180.0 ± 2.4 ^c	181.3 ± 2.3 ^c	184.2 ± 3.4 ^b	192.9 ± 4.0 ^b	
Hind-paw volume ⁽¹⁾	REF	0.81 ± 0.01 ^a	0.83 ± 0.01 ^a	0.84 ± 0.01 ^a	0.85 ± 0.01 ^a	
	AA	0.82 ± 0.01 ^a	1.27 ± 0.09 ^b	1.74 ± 0.13 ^b	1.43 ± 0.09 ^b	
	Q	0.93 ± 0.03 ^b	1.34 ± 0.16 ^b	2.14 ± 0.23 ^b	1.58 ± 0.18 ^b	
	C5	0.87 ± 0.02 ^b	1.40 ± 0.11 ^b	1.67 ± 0.12 ^b	1.27 ± 0.09 ^b	
	C10	0.86 ± 0.02 ^{ab}	1.31 ± 0.12 ^b	1.73 ± 0.16 ^b	1.27 ± 0.10 ^b	
Arthritis Score ⁽²⁾	REF	Arthritis score range				
		0	11 ^a	11 ^a	11 ^a	11 ^a
		1-4	0	0	0	0
		5-8	0	0	0	0
		9-12	0	0	0	0
	13-16	0	0	0	0	
	AA	0	11 ^a	2 ^b	1 ^b	1 ^b
		1-4	1	2	3	5
		5-8	0	8	4	3
		9-12	0	0	4	3
		13-16	0	0	0	0
	Q	0	5 ^b	0 ^b	1 ^b	1 ^b
		1-4	6	5	3	5
		5-8	0	4	3	4
		9-12	0	2	4	1
		13-16	0	0	0	0
	C5	0	5 ^b	0 ^b	0 ^b	0 ^b
		1-4	7	6	3	8
		5-8	0	5	7	3
		9-12	0	1	2	1
		13-16	0	0	0	0
	C10	0	7 ^{ab}	0 ^b	1 ^b	3 ^b
		1-4	4	6	2	3
		5-8	1	5	6	5
		9-12	0	1	3	1
13-16		0	0	0	0	

435 ¹ Hind-paw volume is shown as the average of the two hind-paw volumes, expressed as mL.

436 ² Arthritis score was established according to: '0' means no signs of inflammation, '1' means
 437 erythema or slight swelling in the paw articulations, '2' means erythema and moderate swelling
 438 in the paw articulations, '3' means erythema and deep swelling in the paw articulations, '4'
 439 means erythema and severe swelling and immobility.