

Sustainable strategies for managing *Brassica napus* (oilseed rape) resistance to *Leptosphaeria maculans* (phoma stem canker)

Sustainable strategies for managing *Brassica napus* (oilseed rape) resistance to *Leptosphaeria maculans* (phoma stem canker)

Under the aegis of a European-Australian workshop held at INRA, Versailles, France, in association with the EU SECURE project (QLK5-CT-2002-01813)

**Edited by
B.D.L. Fitt, N. Evans, B.J. Howlett and B.M. Cooke**

Reprinted from *European Journal of Plant Pathology*, Volume 114 Issue 1, 2006

A C.I.P catalogue record for this book is available from the library of Congress

ISBN 1-4020-4524-7 (HB)

ISBN 1-4020-4525-5 (e-book)

Published by Springer,
P.O. Box 17, 3300 AA, Dordrecht, The Netherlands

Printed on acid-free paper

Cover photos:

Phoma stem canker of oilseed rape (*Brassica napus*) caused by *Leptosphaeria maculans* and *L. biglobosa*. From top to bottom: Symptoms on leaves and stems; world-wide distribution; stem cankers of differing severities; pathogen life cycle in Europe in relation to host resistance. Full details in Fitt *et al.* (pp. 3–15)

Springeronline.com

All Rights Reserved

© 2006 Springer

No part of the material protected by this copyright notice may be reproduced or utilized in any form or by any means, electric or mechanical, including photocopying, recording or by any information storage and retrieval system, without written permission from the copyright owner.

Printed in the Netherlands

Contents

Sustainable strategies for managing <i>Brassica napus</i> (oilseed rape) resistance to <i>Leptosphaeria maculans</i> (phoma stem canker)	1
B.D.L. Fitt, N. Evans, B.J. Howlett and B.M. Cooke	
World-wide importance of phoma stem canker (<i>Leptosphaeria maculans</i> and <i>L. biglobosa</i>) on oilseed rape (<i>Brassica napus</i>)	3–15
B.D.L. Fitt, H. Brun, M.J. Barbetti and S.R. Rimmer	
Genetic linkage maps and genomic organization in <i>Leptosphaeria maculans</i>	17–31
M.-L. Kuhn, L. Gout, B.J. Howlett, D. Melayah, M. Meyer, M.-H. Balesdent and T. Rouxel	
Major gene resistance in <i>Brassica napus</i> (oilseed rape) is overcome by changes in virulence of populations of <i>Leptosphaeria maculans</i> in France and Australia	33–40
S.J. Sprague, M.-H. Balesdent, H. Brun, H.L. Hayden, S.J. Marcroft, X. Pinochet, T. Rouxel and B.J. Howlett	
Major gene and polygenic resistance to <i>Leptosphaeria maculans</i> in oilseed rape (<i>Brassica napus</i>)	41–52
R. Delourme, A.M. Chèvre, H. Brun, T. Rouxel, M.H. Balesdent, J.S. Dias, P. Salisbury, M. Renard and S.R. Rimmer	
A large-scale survey of races of <i>Leptosphaeria maculans</i> occurring on oilseed rape in France	53–65
M.-H. Balesdent, K. Louvard, X. Pinochet and T. Rouxel	
Frequency of avirulence alleles in field populations of <i>Leptosphaeria maculans</i> in Europe	67–75
A. Stachowiak, J. Olechnowicz, M. Jedryczka, T. Rouxel, M.-H. Balesdent, I. Happstadius, P. Gladders, A. Latunde-Dada and N. Evans	
Fitness cost associated with loss of the <i>AvrLm4</i> avirulence function in <i>Leptosphaeria maculans</i> (phoma stem canker of oilseed rape)	77–89
Y.-J. Huang, Z.-Q. Li, N. Evans, T. Rouxel, B.D.L. Fitt and M.-H. Balesdent	
Improved resistance management for durable disease control: A case study of phoma stem canker of oilseed rape (<i>Brassica napus</i>)	91–106
J.N. Aubertot, J.S. West, L. Bousset-Vaslin, M.U. Salam, M.J. Barbetti and A.J. Diggle	
Durability of resistance and cost of virulence	107–116
S. Pietravalle, S. Lemarié and F. van den Bosch	
Dissemination of information about management strategies and changes in farming practices for the exploitation of resistance to <i>Leptosphaeria maculans</i> (phoma stem canker) in oilseed rape cultivars	117–126
P. Gladders, N. Evans, S. Marcroft and X. Pinochet	

Foreword

Sustainable strategies for managing *Brassica napus* (oilseed rape) resistance to *Leptosphaeria maculans* (phoma stem canker)

The interaction between the fungus *Leptosphaeria maculans* and oilseed rape (*Brassica napus*) is becoming an excellent model system for studying genetics of host–pathogen interactions. *Leptosphaeria maculans* causes phoma stem canker (blackleg) on oilseed rape and other Brassica crops worldwide. Recently, application of molecular techniques has led to increased understanding of the genetics of this hemibiotrophic interaction. The complete sequences of the genomes of *L. maculans* and *B. rapa* (comprising the *Brassica* A genome) will be available soon. This will provide new opportunities to investigate basic metabolic pathways in the host and the pathogen, and detailed knowledge of the disease process.

Worldwide, the major strategy for control of phoma stem canker is the use of cultivars with resistance to *L. maculans*. However, serious epidemics have occurred recently in Australia and Europe when *L. maculans* populations changed such that major gene resistance in oilseed rape was overcome. Thus there is an urgent need to find and deploy sources of resistance to *L. maculans* in a manner that enhances their durability.

This topic was addressed at a workshop (phoma stem canker durable resistance workshop, 13 September 2004) attended by plant pathologists from Australia and Europe, which was held at INRA (Versailles, France) immediately before the second annual meeting of the European Union-funded SECURE (QLK5-CT-2002-01813) project. This special issue of the European Journal of Plant Pathology is based on papers presented at the workshop, with additional contributions from scientists in Canada, Australia and Europe who were not able to attend. To provide an international perspective on each topic, authors of most of the papers are from several countries. We hope that these papers will provide researchers with a synthesis of the recent studies relating to strategies for management of resistance genes to provide effective control of *L. maculans*, and will stimulate further research on this important model system.

BRUCE D.L. FITT
NEAL EVANS
BARBARA J. HOWLETT
MIKE COOKE