

ID Design 2012/DOOEL Skopje, Republic of Macedonia  
 Open Access Macedonian Journal of Medical Sciences. 2017 Jun 15; 5(3):395-396.  
<https://doi.org/10.3889/oamjms.2017.098>  
 eISSN: 1857-9655  
 Letter to the Editor



## Congenital Progressive Mutilating Hemangioma

Anastasiya Chokoeva<sup>1</sup>, Radica Sokolova<sup>2</sup>, Torello Lotti<sup>3</sup>, Uwe Wollina<sup>4</sup>, Serena Gianfaldoni<sup>4</sup>, Jacopo Lotti<sup>5</sup>, Katlein França<sup>6</sup>, Georgi Tchernev<sup>1,7</sup>

<sup>1</sup>"Onkoderma" - Polyclinic for Dermatology and Dermatologic Surgery, Sofia, Bulgaria; <sup>2</sup>Department of Dermatology and Venereology, Medical University of Plovdiv, 4002 Plovdiv, Bulgaria; <sup>3</sup>University of Rome, "G. Marconi", Rome, Italy; <sup>4</sup>Department of Dermatology and Allergology, Academic Teaching Hospital Dresden-Friedrichstadt, 01067 Dresden, Germany; <sup>5</sup>Department of Nuclear, Subnuclear and Radiation Physics, University of Rome "G. Marconi", Rome, Italy; <sup>6</sup>Institute for Bioethics & Health Policy; Department of Dermatology & Cutaneous Surgery; Department of Psychiatry & Behavioral Sciences, University of Miami Miller School of Medicine - Miami, FL, USA; <sup>7</sup>Medical Institute of Ministry of Interior (MVR-Sofia), Department of Dermatology, Venereology and Dermatologic Surgery, Sofia, Bulgaria

### Abstract

**Citation:** Chokoeva A, Sokolova R, Lotti T, Wollina U, Gianfaldoni S, Lotti J, França K, Tchernev G. Congenital Progressive Mutilating Hemangioma. Open Access Maced J Med Sci. 2017 Jun 15; 5(3):395-396. <https://doi.org/10.3889/oamjms.2017.098>

**Keywords:** infantile hemangioma; plastic surgery; mutilation; beta blockers; topically approach.

**\*Correspondence:** Prof. Dr. Georgi Tchernev. Specialist for Dermatology, Venereology and Dermatologic surgery. "Onkoderma"- Polyclinic for Dermatology and Dermatologic Surgery, Sofia, Bulgaria; Medical Institute of Ministry of Interior (MVR-Sofia), Department of Dermatology, Venereology and Dermatologic Surgery, Sofia, Bulgaria. E-mail: georgi\_tchernev@yahoo.de

Received: 09-May-2017; Revised: 18-May-2017; Accepted: 19-May-2017; Online first: 11-Jun-2017

**Copyright:** © 2017 Anastasiya Chokoeva, Radica Sokolova, Torello Lotti, Uwe Wollina, Serena Gianfaldoni, Jacopo Lotti, Katlein França, Georgi Tchernev. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

**Funding:** This research did not receive any financial support.

**Competing Interests:** The authors have declared that no competing interests exist.

A 73-year-old male patient was admitted with symptoms of decompensated cardiac and pulmonary insufficiency with long-lasting history. A tumor-like formation was observed within the clinical examination, covering the whole skin of the nose, paranasal region of the left part of the face, as well as the upper and lower left eyelids. The lesion was with yellow to brownish surface and dark-reddish to violet discolored peripheral area, composed of nodular formations, smooth central surface and firm texture on palpation. The histopathological examination verified the diagnosis of hemangioma, which had been congenital, regarding the patient's history, treated surgically about 50 years ago, with signs of recurrence. The presented patient had been treated surgically at the age of 20, without medical evidence of the type of the performed excision. The recurrence occurs almost 50 years later, at the age of 78. To the best of our knowledge, this is the first reported recurrence of infantile hemangioma, treated surgically almost 50 years ago.

A 73-year-old male patient was admitted with symptoms of decompensated cardiac and pulmonary insufficiency with long-lasting history. A tumor-like formation was observed within the clinical examination, covering the whole skin of the nose, paranasal region of the left part of the face, as well as the upper and lower left eyelids. The lesion was with yellow to brownish surface and dark-reddish to violet discolored peripheral area, composed of nodular formations, smooth central surface and firm texture on palpation (Fig.1). According to patient's history, he underwent a surgical removal of congenital lesion, at age of 20. Recently, the formation began to increased its size peripherally, becoming more firm in the central area. No subjective complaints were reported. The histopathological examination verified the diagnosis of hemangioma, which had been congenital, regarding the patient's history, treated surgically about 50 years ago, with signs of recurrence.

Surgical excision is nowadays not recommended as a first-line therapy in infantile hemangiomas, because of the well-established tendency of spontaneous regression in one hand, the risk of recurrences, and the proven effectiveness of numerous other non-surgical regimens in other [1].

Although systemic corticosteroids have been the first line of treatment for many years, it has been recently established that, non-selective beta-blockers, such as oral propranolol and topical timolol, are more promising and safer therapeutic approaches. Furthermore, interferon  $\alpha$  and vincristine are suitable for life-threatening haemangiomas that are unresponsive to conventional therapy [2]. The recurrence rate of surgically treated infantile hemangiomas accounts approximately 22%, as it depends on the patients' age at the time of the surgery and the type of the performed resection

(intralesional, marginal, wide and radical resection) [1].



*Figure 1: Clinical manifestation of tumor-like lesion, located predominantly on the left nasal and paranasal region, composed of nodular formations, smooth central surface and firm texture on palpation*

The presented patient had been treated surgically at the age of 20, without medical evidence of the type of the performed excision. The recurrence occurs almost 50 years later, at the age of 78. To the best of our knowledge, this is the first reported recurrence of infantile hemangioma, treated surgically almost 50 years ago.

## References

1. Canavese F, Soo BC, Chia SK, Krajchich JI. Surgical outcome in patients treated for hemangioma during infancy, childhood, and adolescence: a retrospective review of 44 consecutive patients. *J Pediatr Orthop.* 2008; 28(3):381-6. <https://doi.org/10.1097/BPO.0b013e318168d1a7> PMID:18362808
2. Sethuraman G, Yenamandra VK, Gupta V. Management of Infantile Hemangiomas: Current Trends. *J Cutan Aesthet Surg.* 2014; 7(2): 75–85. <https://doi.org/10.4103/0974-2077.138324> PMID:25136206 PMCid:PMC4134656