

Immunological Evaluation in Patients with Familial Mediterranean fever

Shams Kholoussi¹, Naglaa Kholoussi¹, Moushira Erfan Zaki², Hala T. El-Bassyouni³, Hala Elnady⁴, Botros Morcos¹, Asem Abo-Shanab^{1*}

¹National Research Centre - Immunogenetics Department, Cairo, Egypt; ²National Research Centre - Biological Anthropology Department, Cairo, Egypt; ³National Research Centre - Clinical Genetics Department, Cairo, Egypt; ⁴National Research Centre - Child Health, Cairo, Egypt

Abstract

Citation: Kholoussi S, Kholoussi N, Zaki ME, El-Bassyouni HT, Elnady H, Morcos B, Abo-Shanab A. Immunological Evaluation in Patients with Familial Mediterranean fever. Open Access Maced J Med Sci. <https://doi.org/10.3889/oamjms.2018.079>

Keywords: Flow cytometer; Lymphocytes; NK; Immunoglobulins; Nephelometry; FMF; N/L ratio

***Correspondence:** Abo - Shanab. National Research Centre - Immunogenetics Department, Cairo, Egypt. E-mail: asem_nrc@yahoo.com

Received: 29-Oct-2017; **Revised:** 12-Dec-2017; **Accepted:** 13-Dec-2017; **Online first:** 09-Feb-2018

Copyright: © 2018 Shams Kholoussi, Naglaa Kholoussi, Moushira Erfan Zaki, Hala T. El-Bassyouni, Hala Elnady, Botros Morcos, Asem Abo-Shanab. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This study was supported by National Research Centre, grant number: 010010606

Competing Interests: The authors have declared that no competing interests exist

OBJECTIVE: This study aimed to investigate T & B lymphocyte subsets and Natural Killer (NK) cells patterns in children with FMF versus normal control subjects, to estimate the immunoglobulins IgG, IgM, and IgA levels, and to scrutinize the possible use of Neutrophil / Lymphocyte ratio (NLR) as a marker for subclinical inflammation in FMF patients.

PATIENTS AND METHODS: A group of 42 patients with FMF attending the Genetics Clinic at National Research Centre were included in this study. They were 13 males and 19 females; their age ranged from 2 to 17 years old. Normal healthy subjects within the same age and sex range were included as a control group. Complete blood picture was done for all cases, and neutrophil/ lymphocyte ratio was calculated. Flow cytometer analysis was done for CD3, CD4, CD8, CD19 and CD16 using monoclonal antibodies. Immunoglobulins IgG, IgA and IgM were estimated in serum using nephelometry.

RESULTS: Positive consanguinity was present in 20 patients (47.6%). Abdominal pain was the most common manifestation followed by fever, arthritis, and red rash. CD3, CD4 and CD8 were statistically increased in patients group as compared to normal control group, while CD16 was statistically decreased.

CONCLUSION: The study suggests that quantitative measurement of CD expressions of CD3, CD4 and CD8 as well as NLR might be used as valuable markers for subclinical inflammation in FMF.

Introduction

Familial Mediterranean fever (FMF) is the most common systemic autoinflammatory disorder worldwide [1]. It is an autosomal recessive disease [2] that affects mainly the Mediterranean population [3], is characterised by recurrent self - limited fever and aseptic serosal inflammation, causing abdominal, thoracic and joints pain [4]. Children with FMF are more prone to growth retardation [5], chronic normocytic normochromic anaemia [6], and splenomegaly [7].

Malicious activation of many inflammatory pathways occurs during the attacks of FMF, in which T, B and Natural killer cells play a major role [8][9]. The interaction of these cells with endothelial cells

and several inflammatory and immune mediators produced by these cells leads to the formation of atherosclerotic plaques followed by its destabilisation and vessel rupture [10].

In between attacks, inflammation continues subclinically with the associated processes of endothelial dysfunction, increased atherosclerotic changes and platelets activation. The persistent inflammation in FMF may cause endothelial dysfunction, atherothrombosis and systemic amyloidosis [11]. The Neutrophil / Lymphocyte ratio (N/L ratio) showed significant, rapid, and serial changes as immune system response to different conditions like surgical stress, systemic inflammation or severe infections [12], and can be used as a marker for predicting amyloidosis development [13].

The most distressing complication in FMF is renal amyloidosis, leading to nephrotic syndrome and chronic kidney failure. A macromolecular complex, called inflammasome, plays a major role in the activation of IL - 1 and thus induction of inflammation and when inflammasome activity is abnormally stimulated through a mutation, it may be involved in the pathogenesis of FMF [14][15][16]. Natural killer (NK) cells are cytotoxic lymphocytes that participate in innate immunity. In addition to their cytotoxic response, these cells produce cytokines to assist the adaptive immune response [17][18].

The aim of this study was to investigate T and B lymphocytes and Natural Killer (NK) cells patterns through flow - cytometric analysis of CD3, CD4, CD8, CD16 and CD19 expression, to estimate the immunoglobulin levels IgG, IgM and IgA in FMF patients as compared to normal control subjects, and also to scrutinize the possible use of NLR as a marker for subclinical inflammation in patients with FMF.

Patients and Methods

Forty - two patients with FMF attending the Clinical Genetics Clinic at National Research Centre were included in this study. They were following up patients during the year 2015. They were 13 males and 19 females; their age ranged from 2 to 17 years old. Twenty normal healthy subjects within the same age and sex range were included as a control group.

All subjects were subjected to full clinical examination and history taking. Written consents were taken from the children parents', and the study was approved by Medical Ethics Committee of NRC (13/146). Complete blood picture was done for all cases, and neutrophil/lymphocyte ratio was calculated. Flow - cytometer analysis was done for CD3, CD4, CD8, CD19 and CD16 using monoclonal antibodies [19]. Immunoglobulins IgG, IgA, Ig M was estimated in serum using nephelometry [20].

Statistical analysis was performed using SPSS program version 13. Quantitative variables were presented as a mean and standard deviation. The unpaired t-test was used to evaluate differences between the two groups of continuous variables. Two-tailed $P < 0.05$ was considered statistically significant.

Results

The study group consisted of 23 males (54.8%) and 19 females (45.2%), all were diagnosed as FMF and during the attack; while the control group

were normal subjects matching age and sex. The age range in the FMF study group was 2 - 17 years and 2 - 14 years in the control group respectively (Table 1). This study included 42 patients, with the clinical characteristics of FMF. Positive consanguinity was present in 20 patients (47.6 %).

Abdominal pain was the most common manifestation which usually occurs in repeated attacks and stays from 2 hours to several days followed by fever, arthritis, and erysipelas-like rashes. All the patients were treated with colchicine after being identified by molecular diagnosis.

Table 1: Age and sex of patients and control group

	Patients n = 42	Control n = 20
Age range (years)	2-17	2-14
Sex (male/female)	23/19	11/9

CD3, CD4, and CD8 were statistically increased in patients' group as compared to normal control group; p-value were 0.001, 0.002, 0.004, respectively (Table 2). While CD16 has statistically decreased, the p-value was 0.007 as compared the control group.

Table 2: Cellular expression of CD3, CD4, CD8, CD19 and CD16 in FMF patients as compared to control group

	Patients n = 42	Control n = 20	p - value
CD3 %	51.91 ± 15.19	36.31 ± 10.41	0.001*
CD19 %	18.29 ± 9.07	19.78 ± 5.38	0.81
CD16 %	12.81 ± 5.38	19.06 ± 6.9	0.007*
CD4 %	33.74 ± 6.46	22.88 ± 6.81	0.002*
CD8 %	19.37 ± 5.98	12.2 ± 4.15	0.004*

*statistical significant $p < 0.05$.

Immunoglobulins IgA, IgM, IgG and also TLC and N/L ratio did not show the significant statistical difference between the patients and control groups (Table 3).

Table 3: Immunoglobulins IgA, IgM, IgG, TLC and N/L ratio in FMF patients and control group

parameters	Patients n = 42	Control n = 20	p - value
TLC ($\times 10^9$ /mm ³)	7.54 ± 2.39	9.16 ± 2.23	0.13
N/L ratio (NLR)	1.23 ± 0.62	0.8 ± 0.16	0.10
IgA (g/L)	1.35 ± 0.7	1.31 ± 1.06	0.89
IgM (g/L)	1.26 ± 0.45	1.41 ± 0.60	0.54
IgG (g/L)	10.67 ± 3.8	9.62 ± 3.45	0.49

*statistical significant $p < 0.05$

Discussion

Autoinflammatory diseases originate from inappropriate activation of antigen-independent inflammatory mechanisms [21]. FMF is characterised by fever and serositis which flare up as paroxysmal attacks, with intervening asymptomatic periods. The attacks of fever and diffuse abdominal pain in FMF are characterised by subclinical inflammation and associated endothelial dysfunction [11]. We aimed to

investigate a possible immune regulation imbalance in familial Mediterranean fever (FMF) by measuring levels of peripheral blood lymphocyte subsets using flow - cytometry. In this study, we found the increased percentage of T cells CD3, CD4 and CD8 as compared to the control group. This was in agreement with Musabak et al., [22], who stated that CD3, CD4 and CD8 were increased in FMF patients than in the control group indicating that the T cell system is abnormally activated in patients with FMF. Rimar et al., [23] suggested that regulatory T cells play a role in cutting short FMF acute attacks. This may explain the fact that inflammation in FMF is self - limiting and also the absence of autoantibodies, antigen-specific T cells and tissue damage [24].

NK cells mediate cytotoxicity that is regulated by their inhibitory and activating surface receptors. An increase in the activation of NK cells may result in proliferation and change the immune response [25]. In this study, CD16+ NK cells were decreased in FMF patients group as compared to the control group in contrary to what was reported that NK cell numbers were significantly increased in FMF patients as compared to controls. In autoinflammatory diseases, dysregulation is believed to occur in the innate immune system without primary involvement of T and B lymphocytes. Nevertheless, autoinflammation involves crosstalk between the innate immune system – neutrophils, macrophages, and NK cells – with the adaptive immune system [26].

To investigate the B - cells, CD19 showed no significant difference between patients and control groups. As regards to serum immunoglobulins IgA, IgM and IgG estimated in this study, there was no statistically significant difference in patients when compared to the control group. In another genetic disease, results obtained showed a significant difference between patients and controls, in which the results of the control group is nearly the same [27]. However, Livneh et al., [28] compared the immunoglobulin levels in patients with FMF about patients with hyperimmunoglobulinemia D syndrome (HIDS) and not about healthy control subjects, and that the patients with HIDS showed the significantly higher prevalence of elevated immunoglobulins levels than patients with FMF.

The N/L ratio is a reliable marker for evaluating and monitoring the systemic inflammatory response, and prognosis of clinical outcome of inflammatory diseases [29].

No statistically significant difference was detected in TLC nor NLR in FMF patients as compared to the control group while Özer et al., [30] and Duksal et al., [29], found TLC and N/L ratio were significantly higher in children with FMF than in healthy individuals. Celikbilek et al., [31], reported that the N/L ratio was higher in active FMF than in FMF in remission as well as in control subjects.

In conclusion, dysfunction of the innate immune system is the central matter in FMF as a self - reactive autoinflammatory disease.

References

- Ozen S, Aktay N, Lainka E, Duzova A, Bakkaloglu A, Kallinich T. Disease severity in children and adolescents with familial Mediterranean fever: a comparative study to explore environmental effects on a monogenic disease. *Ann Rheum Dis*. 2009; 68(2):246–248. <https://doi.org/10.1136/ard.2008.092031> PMID:18801759
- Caliskan M, Gullu H, Yilmaz S, Erdogan D, Unler GK, Ciftci O, et al. Impaired coronary microvascular function in familial Mediterranean fever. *Atherosclerosis*. 2007; 195(2):e161–167. <https://doi.org/10.1016/j.atherosclerosis.2007.06.014> PMID:17673217
- Onen F. Familial Mediterranean fever. *Rheumatol Int*. 2006; 26: 489–96. <https://doi.org/10.1007/s00296-005-0074-3> PMID:16283319
- Medlej-Hashim M, Loiselet J, Lefranc G, Mégarbané A. Familial Mediterranean Fever (FMF): from diagnosis to treatment. *Sante*. 2004; 14(4): 261–266. PMID:15745878
- Zung A, Barash G, Zadik Z, Barash J. Familial Mediterranean fever and growth: effect of disease severity and colchicine treatment. *J Pediatr Endocrinol*. 2006; 19:15–18. <https://doi.org/10.1515/JPEM.2006.19.2.155>
- Celkan T, Çelik M, Kasapçopur Ö, Özkan A, Apak H, Ocak S, et al. The anemia of familial mediterranean fever disease. *J Pediatric Hematology and Oncology*. 2005; 22, 8: 657–665. <https://doi.org/10.1080/08880010500278681> PMID:16251171
- Korkmaz C, Ozdogan H, Kasapçopur O and Yazici H. Acute phase response in familial Mediterranean fever. *Ann Rheum Dis*. 2002; 61: 79–81. <https://doi.org/10.1136/ard.61.1.79> PMID:11779767 PMID:PMC1753891
- Chae JJ, Aksentijevich I, Kastner DL. Advances in the understanding of familial Mediterranean fever and possibilities for targeted therapy. *Br J Haematol*, 2009; 146: 467–78. <https://doi.org/10.1111/j.1365-2141.2009.07733.x> PMID:19466978 PMID:PMC2759843
- Gattorno M, La Regina M, Martini A and Manna R. An update on auto-inflammatory diseases. New concepts for new and old diseases. *Clin Exp Rheumatol*. 2009; 27:354–65. PMID:19473583
- Galkina E, Ley K. Immune and inflammatory mechanisms of atherosclerosis. *Annu Rev Immunol*, 2009; 27:165–97. <https://doi.org/10.1146/annurev.immunol.021908.132620> PMID:19302038 PMID:PMC2734407
- Yüksel S, Ayvazyan L, Gasparyan AY. Familial Mediterranean Fever as an Emerging Clinical Model of Atherogenesis Associated with Low-Grade Inflammation. *The Open Cardiovascular Medicine Journal*. 2010; 4:51–56. <https://doi.org/10.2174/1874192401004010051>
- Zahorec R. Ratio of neutrophil to lymphocyte counts—rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratislavske Lekarske Listy*. 2001; 102(1):5–14. PMID:11723675
- Uslu AU, Deveci K, Korkmaz S, Aydin B, Senel S, Sancakdar E, Sencan M. Is Neutrophil/ Lymphocyte Ratio Associated with Subclinical Inflammation and Amyloidosis in Patients with Familial Mediterranean Fever? *BioMed Research International*. 2013;185317:1–5. <https://doi.org/10.1155/2013/185317> PMID:23865042 PMID:PMC3705820
- Fonnesu C, Cerquaglia C, Giovinale M, Curigliano V, Verrecchia E, Giuliana de Sicio, et al. Familial Mediterranean Fever: A review for clinical management. *J Joint Bone Spine*. 2009;

- 76(3):227-233. <https://doi.org/10.1016/j.jbspin.2008.08.004> PMID:19091621
15. Manna R, Cerquaglia C, Curigliano V, Fonnesu C, Giovinale M, Verrecchia E, et al. Clinical features of familial Mediterranean fever: an Italian overview. *Eur Rev Med Pharmacol Sci*. 2009; 13(Suppl 1):51–53. PMID:19530512
16. Schroder K, Tschopp J. The inflammasomes. *Cell*. 2010; 140: 821-832. <https://doi.org/10.1016/j.cell.2010.01.040> PMID:20303873
17. Paust S, Senman B, Von Andrian UH. Adaptive Immune Responses Mediated by Natural Killer Cells. *Immunological reviews*. 2010; 235(1): 286-296. <https://doi.org/10.1111/j.0105-2896.2010.00906.x> PMID:20536570 PMID:PMC2911633
18. Vivier E, Raulet DH, Moretta A, Caligiuri MA, Zitvogel L, Lanier LL, et al. Innate or Adaptive Immunity? The Example of Natural Killer Cells. *Science*. 2011; 331(6013):44-49. <https://doi.org/10.1126/science.1198687> PMID:21212348 PMID:PMC3089969
19. Jaroszeski M J, Gilbert R, Heller R. Detection and quantitation of cell-cell electrofusion products by flow cytometry. *Anal Biochem*. 1994; 216:271–275. <https://doi.org/10.1006/abio.1994.1041> PMID:7513970
20. Aksu G, Genel F, Koturoğlu G, Kurugöl Z and Kütükçüler N. Serum immunoglobulin (IgG, IgM, IgA) and IgG subclass concentrations in healthy children: a study using nephelometric technique. *Turk J Pediatr*. 2006; 48(1):19-24. PMID:16562781
21. Wang DQH, Bonfrate L, de Bari O, Wang TY, Portincasa P. Familial Mediterranean Fever: From Pathogenesis to Treatment. *J Genet Syndr Gene Ther*. 2014; 5:248.
22. Musabak U, Sengul A, Oktenli C, Pay S, Yesilova Z, Kenar L, et al. Does immune activation continue during an attack-free period in familial Mediterranean fever? *Clin Exp Immunol*. 2004; 138: 526-533. <https://doi.org/10.1111/j.1365-2249.2004.02632.x> PMID:15544632 PMID:PMC1809227
23. Rimar D, Rosner I, Slobodin G, Boulman N, Toubi E, Kessel A, et al. The Role of Regulatory T Cells in Familial Mediterranean Fever (FMF). *Clin Rheumatol*. 2012; 31(5):885-8. <https://doi.org/10.1007/s10067-011-1935-7> PMID:22234494
24. Manukyan GP, Ghazaryan KA, KtsoyanZh A, TatyamMV, Khachatryan ZA, Hakobyan GS, et al. Cytokine profile of Armenian patients with Familial Mediterranean fever. *Clin Biochem*. 2008; 41:920-2. <https://doi.org/10.1016/j.clinbiochem.2008.03.017> PMID:18440310
25. Erken E, Ozturk OG, Kudas O, Tas DA, Demirtas A, Kibar F, et al. Killer Cell Immunoglobulin-Like Receptor (KIR) Genotype Distribution in Familial Mediterranean Fever (FMF) Patients. *Med Sci Monit*. 2015; 21:3547-3554. <https://doi.org/10.12659/MSM.895211> PMID:26574972 PMID:PMC4655612
26. Schuetz C, Niehues T, Friedrich W, Schwarz K. Autoimmunity, autoinflammation and lymphoma in combined immunodeficiency (CID). *Autoimmunity Reviews*. 2010; 9:477–482. <https://doi.org/10.1016/j.autrev.2010.02.005> PMID:20146941
27. Shanab AM, El-Desouky MA, Kholoussi N, El-Kamah G, Fahmi AA. Evaluation of neopterin as a prognostic factor in patients with beta-thalassemia, in comparison with cytokines and immunoglobulins. *Archives of Hellenic Medicine/Arheia Ellenikes Iatrikes*. 2015; 32(1).
28. Livneh A, Drenth JP, Klasen IS, Langevitz P, George J, Shelton DA, et al. Familial Mediterranean fever and hyperimmunoglobulinemia D syndrome: two diseases with distinct clinical, serologic, and genetic features. *J Rheumatol*. 1997; 24:1558-1563. PMID:9263151
29. Duksal F, Alaygut D, Güven AS, Ekici M, Oflaz MB, Tuncer R et al. Neutrophil- lymphocyte ratio in children with familial Mediterranean fever. *Eur J Rheumatol*. 2015; 1: 20-23. <https://doi.org/10.5152/eurjrheumatol.2015.0093> PMID:27708915 PMID:PMC5047238
30. Özer S, Yılmaz R, Sönmezgöz E, Karaaslan E, Taşkın S, Bütün İ, Demir O. Simple markers for subclinical inflammation in patients with Familial Mediterranean Fever. *Med Sci Monit*. 2015; 23:298-303.
31. Celikbilek M, Dogan S, Akyol L, Borekci E, Zararsiz G, Kozan M, et al. Neutrophil- Lymphocyte Ratio in Patients With Familial Mediterranean Fever. *J Clin Lab Anal*. 2015; 29:80-83. <https://doi.org/10.1002/jcla.21732> PMID:24687426