

ID Design Press, Skopje, Republic of Macedonia
 Open Access Macedonian Journal of Medical Sciences. 2018 Nov 25; 6(11):2040-2044.
<https://doi.org/10.3889/oamjms.2018.416>
 eISSN: 1857-9655
 Clinical Science



IGF1R Gene Alterations in Children Born Small for Gestational Age (SGA)

Aleksandra Janchevska^{1*}, Marina Krstevska-Konstantinova¹, Heike Pfäffle², Marina Schlicke², Nevenka Laban¹, Velibor Tasic¹, Zoran Gucev¹, Kristina Mironska¹, Aleksandar Dimovski³, Jürgen Kratzsch², Jürgen Klammt², Roland Pfäffle²

¹Medical Faculty, Ss. Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia; ²University of Leipzig, Leipzig, Germany; ³Macedonian Academy of Sciences and Arts, Skopje, Republic of Macedonia

Abstract

Citation: Janchevska A, Krstevska-Konstantinova M, Pfäffle H, Schlicke M, Laban N, Tasic V, Gucev Z, Mironska K, Dimovski A, Kratzsch J, Klammt J, Pfäffle R. IGF1R Gene Alterations in Children Born Small for Gestational Age (SGA). Open Access Maced J Med Sci. 2018 Nov 25; 6(11):2040-2044. <https://doi.org/10.3889/oamjms.2018.416>

Keywords: Small for gestational age (SGA); IGF1 receptor (IGF1R); IGF1R gene; Multiplex Ligation-dependent Probe Amplification (MLPA); direct sequencing

***Correspondence:** Aleksandra Janchevska. Medical Faculty, Ss. Cyril and Methodius University in Skopje, Skopje, Macedonia. E-mail: dr.sasha1969@yahoo.com

Received: 18-Aug-2018; **Revised:** 09-Sep-2018; **Accepted:** 10-Sep-2018; **Online first:** 10-Nov-2018

Copyright: © 2018 Aleksandra Janchevska, Marina Krstevska-Konstantinova, Heike Pfäffle, Marina Schlicke, Nevenka Laban, Velibor Tasic, Zoran Gucev, Kristina Mironska, Aleksandar Dimovski, Jürgen Kratzsch, Jürgen Klammt, Roland Pfäffle. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This research did not receive any financial support

Competing Interests: The authors have declared that no competing interests exist

BACKGROUND: Small for gestational age (SGA)-born children are a heterogeneous group with few genetic causes reported. Genetic alterations in the IGF1 receptor (IGF1R) are found in some SGA children.

AIM: To investigate whether alterations in *IGF1R* gene are present in SGA born children.

PATIENTS AND METHODS: We analysed 64 children born SGA who stayed short (mean -3.25 ± 0.9 SDS) within the first 4 years of age, and 36 SGA children who caught up growth (0.20 ± 1.1 SDS). PCR products of all coding IGF1R exons were screened by dHPLC followed by direct sequencing of conspicuous fragments to identify small nucleotide variants. The presence of IGF1R gene copy number alterations was determined by Multiplex Ligation-dependent Probe Amplification (MLPA).

RESULTS: The cohort of short SGA born children revealed a heterozygous, synonymous variant c.3453C > T in one patient and a novel heterozygous 3 bp in-frame deletion (c.3234_3236delCAT) resulting in one amino acid deletion (p.Ile1078del) in another patient. The first patient had normal serum levels of IGF1. The second patient had unusually low IGF1 serum concentrations (-1.57 SD), which contrasts previously published data where IGF1 levels rarely are found below the age-adjusted mean.

CONCLUSIONS: *IGF1R* gene alterations were present in 2 of 64 short SGA children. The patients did not have any dysmorphic features or developmental delay. It is remarkable that one of them had significantly decreased serum concentrations of IGF1. Growth response to GH treatment in one of the patients was favourable, while the second one discontinued the treatment, but with catch-up growth.

Introduction

Small for gestational age children (SGA; low birth weight and/or birth length) are a heterogeneous group both regarding clinical characteristics and the aetiology (fetal, maternal, placental, and/or genetic factors). Most SGA children normalise their stature by 2 yr. of age. Nevertheless, approximately 10-15% of SGA children do not achieve normal growth and height until adolescence and adulthood and remain short [1] [2]. In addition to the short stature SGA children have a reduced lean body mass, fat mass, skin folds, and body mass index (BMI) [3] [4] [5] [6] [7], as well as a lower calorie, fat, and carbohydrate intake [1] [3]. An impaired IGF1R function may lead to

disturbed glucose homeostasis [8], which may partly explain the increased risk for diabetes in SGA adults.

IGF-I, the hormone ligand that binds to the IGF1R, is fundamental for prenatal and postnatal growth and development. Intrauterine and postnatal growth retardation, deafness, microcephaly, and mental retardation have been reported in homozygous deletion or mutation in the *IGF1* gene [1] [5]. The effects of IGF-I are mediated through the type 1 IGF receptor (IGF1R), which is a tyrosine kinase receptor encoded by the *IGF1R* gene [9]. Growth failure and microcephaly have been reported in patients with IGF1R defects.

We used dHPLC and Sanger sequencing and MLPA to detect small nucleotide variants (SNV) or copy number variants (CNV), respectively, to reveal

possible genetic alterations in the *IGF1R* gene as a cause of the observed phenotype in SGA children with or without catch-up growth.

Patients and Methods

SGA was defined as a birth length and/or weight < 2 standard deviation scores (SDS) for the gestational age. SGA children remaining short at age 4 (height > 2.00 SDS) were included in the study. All children had an uncomplicated perinatal and postnatal period. Exclusion criteria included endocrine disorders, skeletal abnormalities, chronic diseases and chromosomal abnormalities.

The study protocol was approved by the Medical Ethics Committee of the Medical Faculty Skopje, Macedonia.

Birth and growth data before the start of treatment were retrieved from records of nurseries, and general practitioners. Height and head circumference were expressed as SD scores [10]. Body mass index was calculated (weight in kg/height in meters²) and expressed as SD scores for age and sex. Bone age was determined according to Greulich and Pyle [11]. The dysmorphological examination was performed by an experienced clinical geneticist.

GH pituitary reserve was assessed by L-dopa and clonidine GH tests. Serum samples were analysed for IGF-1 and IGFBP-3 by either chemiluminescent immunoassays (Mediagnost, Reutlingen, Germany), or by colourimetric ELISA (Mediagnost, Reutlingen, Germany).

IGF-1 inter- and intra-assay variation coefficients were 6.8 and 6.7%, respectively; IGFBP-3 inter- and intra-assay variation coefficients 6.30 and 4.51%, respectively. Serum GH was measured by a solid-phase, two-site, chemiluminescent immunoassay (ARUP, Salt Lake City, Utah, USA). Cortisol, testosterone and estrogens were measured by colourimetric ELISA (Diagnostic Products Corporation, Los Angeles, Calif., USA).

NA was extracted from peripheral blood samples using the QIAamp DNA Blood Mini Kit (Qiagen, Hilden, Germany). PCR products of all coding exons and adjacent intronic sequences of the *IGF1R* gene were generated and subjected to denaturing HPLC (dHPLC) pre-screening (WAVE System; Transgenomic, Glasgow, UK). PCR products with conspicuous chromatograms were further analysed by Sanger dideoxy-sequencing (ABI PRISM 310 Genetic Analyzer; Thermo Fisher Scientific, Waltham, MA). Primer sequences can be obtained upon request. Sequences were compared to the human reference genome (UCSC, version 19 (GRCh37)) and annotated according to the GenBank

reference coding sequence of the *IGF1R* NM_000875 and UniProtKB protein reference P08069. Multiplex Ligation-dependent Probe Amplification (MLPA) to detect copy-number variants (CNV) in the *IGF1R* gene was performed according to the manufacturer's recommendations (SALSA MLPA P217; MRC Holland, Amsterdam, The Netherlands).

Results

All 64 short SGA children were investigated, and mutations in two patients were identified by dHPLC and direct sequencing.

Patient A is a boy who was born spontaneously after 37 weeks of gestation and after uneventful pregnancy and delivery. He is the second child of young, non-consanguineous parents. His brother had a normal birth size, and postnatal growth. The proband's birth weight was 1700 g (-3.36 SDS score) and birth length 41 cm (-3.66 SDS score). The parents' height was: father 166cm (-0.76 SDS score) and of his mother 154.5 cm (-1.9 SDS score), with a target height of 166.2 cm (-1.27 SDS score). His psychomotor development, sight and hearing were normal.

At 6.3 yr. of age, his height was 101.4 cm (-3.90 SDS score), weight 13.8 kg (-3.24 SDS weight for height), and head circumference was not available. L-dopa and clonidine stimulation tests were performed at age 5 years with a maximal GH response of 6.31 ng/ml and 10.8 ng/ml, respectively. At this age, his bone age was 4 yr. His IGF-I level was 52.3 ng/ml (-1.57 SDS score) and IGFBP-3 level 1.17 mg/liter (-1.63 SDS score). Ultrasound of the heart and kidneys were uneventful, antibodies for gliadin negative. MRI of the hypothalamic and pituitary region revealed normal size pituitary and no anomalies. Morphologic examination showed no anomalies. He attends to a regular primary school. His IQ score was 89. Since the age of 11.1 years, the GH treatment (37 µg/kg/day) was given for 18 months. The treatment resulted in catch-up growth, and he reached 144.6 cm (-0.21 SDS) at 12.7 years when the parents interrupted the treatment.

In patient A a heterozygous synonymous nucleotide transition, c.3453C > T (p.I1151 =), was found. Position 3435 is located 5 nucleotides upstream of the last nucleotide of exon 18. In silico analysis using MutationTaster (<http://www.mutationtaster.org/>; accessed April 2018) predicts pathogenicity due to a potential splice site change. The additional computational analysis suggests the introduction of a new exonic splicing silencer site while a potentially existing splicing enhancer site is broken by the nucleotide substitution (Human Splicing Finder 3.1;

<http://www.umd.be/HSF3/>; accessed April 2018). Aberrant splicing at the intron 18 splice donor site would presumably result in a severely disturbed IGF1R function because the affected amino acid residue(s) are part of the tyrosine kinase domain.




Figure 1: Genetic analysis of IGF 1R; a novel heterozygous variant (synonymous); c.3453C>T; a possible impact on splicing has to be verified

The 8.9 old boys was referred for pediatric endocrine evaluation because of short stature. He was born at 40 wk gestation, after uneventful pregnancy and delivery. The parents are young and non-consanguineous. His brother and sister had normal birth size and normal postnatal growth. His birth weight was 2300 g (-2.76 SDS), his birth length 46 cm (-2.14 SDS), the head circumference at birth was not available. The parents' height was: father 158 cm (-2.0 SDS score) and of his mother 158.7 cm (-1.26 SDS score), with a target height of 165.0 cm (-1.38 SDS score). His psychomotor development, sight and hearing were normal. At referral, the boy had a height of 114.6 cm (-3.08 SDS) and a weight of 22.2 kg (-1.2 SDS). His head circumference was 49.8 cm (normal). The Greulich and Pyle male standards bone age was 10 years. His sight, hearing and development were normal. He had average grades in the primary and secondary school. L- Dopa and clonidine tests of pituitary GH reserve were 16.4 ng/ml and 17.7 ng/ml respectively. Initially, IGF-1 was 468 ng/ml (reference 237-996) and IGFBP-3 levels were not available, but under GH treatment IGF-I was 205 ng/ml (reference 226–903), while IGF binding protein-3 (IGFBP)-3 were not available. At the age of 15.41 years, his height was 138.2 cm (-4.11 SDS), his weight 30.3 kg (-2.71 SDS), head circumference normal. T4, TSH, cortisol, renal function, hepatic analysis were normal. Since the age of 16.25 years (-3.77 SDS) the GH treatment (37 µg/kg/day) was initiated and lasted 3 years. This resulted in a final adult height of 160.5 cm (-2.5 SDS) which was within the parental target height range.

In patient B a 3-bp in-frame deletion, c.3234_3236delCAT, was identified. The deletion probably leads to removal of isoleucine at protein position 1078 (p.I1078del) and is predicted to be disease-causing (MutationTaster). The affected position maps to the tyrosine kinase domain of the IGF1R. Isoleucine 1078 is highly conserved among species and paralogues (insulin receptor, insulin receptor-related receptor) with only isoleucine or valine found at this position. Disturbance of the receptor's kinase activity can be assumed but has to be shown experimentally.




Figure 2: Genetic analysis of IGF 1R of Patient B; a novel heterozygous 3 bp deletion (c.3234_3236delCAT) resulting in a one amino acid deletion (p.Ile1078del)

Discussion

Mutations in the *IGF1R* gene resulting in IGF1 resistance underlie some cases of prenatal and postnatal growth failure [4]. Interestingly, there were three types of phenotypes reported. Some SGA children with IGF1R genetic alterations had microcephaly and short stature, others had only short stature without microcephaly, while some reports describe short stature and/or microcephaly and impaired glucose tolerance. Also, most reports describe elevated circulating levels of IGF1, consistent with the expectations when there is a receptor defect. It is of note that some patients were described with low normal levels of IGF1, as was the case in one of our patients with the novel heterozygous 3 bp deletion (c.3234_3236delCAT) resulting in one amino acid deletion (p.Ile1078del). There is one report that describes hypoglycemia found in a patient with a heterozygous mutation (c.94+1G > A, p.D1105E) affecting the splicing site of the *IGF1R* mRNA [12].

Several reports describe SGA with growth failure: a compound heterozygote for point mutations in exon 2 of the *IGF1R* gene bearing a p.R108Q mutation in 1 allele and a p.K115N mutation in the second [4]. Short stature and intrauterine growth retardation (IUGR) were found to be caused by a heterozygous mutation in the *IGF1R* gene, Arg709 to Gln (p.R709Q) [13], while the p.R481Q mutation was described in two family members with increased serum IGF-I levels and intrauterine and postnatal growth retardation [14]. Normal IGF-I, but short stature was found in a boy and several family members with a 19-nucleotide duplication within exon 18 of the *IGF1R* gene and consequently haploinsufficiency of IGF1R protein [15]. One of our patients had unusually low IGF1 serum concentrations (-1.5 SD), which contrasts with previous reports.

Prenatal and postnatal growth failure was found in patients with p.Y387X mutation [16] where the proband had high IGF-I levels, while he and his two paternal aunts had impaired glucose tolerance. SGA was also reported in a novel IGF1R mutation (p.Alal 40fsX20) [17], and intrauterine and postnatal growth retardation was found in a missense mutation

(p.R431L) [18]. Heterozygous nonsense mutations affecting the C-terminal region (p.Q1250X, p.W1249X) of IGF1R were described in two out of 55 analysed Japanese patients with SGA and growth failure [19].

A heterozygous mutation (p.C1248Y) in the IGF1R gene was found in two brothers with prenatal and postnatal growth retardation and their father [8]. It is of note that OGTT showed progressive impaired glucose tolerance, while the father was already treated for type 2 diabetes mellitus. In a child with a deletion on 15q26.2 intrauterine growth retardation, postnatal growth failure, and recurrent hypoglycemia there was only a single copy of the *IGF1R* gene [20].

Microcephaly is frequently associated with IGF1R genetic alterations. SGA, microcephaly, persistent postnatal growth retardation, and elevated IGF-I levels were reported in a 15-year-old girl with heterozygous deletion of 15q26.2-qter which included the IGF1R gene [21]. The p.R59X mutation was reported in two half-brothers with primary microcephaly, mild mental retardation, and intrauterine as well as postnatal growth deficits [4] [22]. Microcephaly, pre- and postnatal growth retardation were found in patients with heterozygous missense mutations in three unrelated patients, de novo p.R1256S, de novo p.N359Y and p.Y865C [23]. Also the c.1549A > T, the p.Y487F mutation was reported in a patient with microcephaly and prenatal and postnatal growth impairment [24]. GH treatment of a patient with short stature, microcephaly, dysmorphic features, developmental delay and a terminal deletion of 15q26.2q26.3 containing the *IGF1R* gene in addition to a terminal duplication of the 4q35.1q35.2 region resulted in a strong growth response [25]. It is of note that the GH treatment in our patients had mixed effects. The first patient did achieve height within the parental target range, while the second one discontinued the treatment without having a catch-up growth.

The compound heterozygous mutation p.E121K/E234K was reported as the cause of intrauterine growth retardation and severe postnatal growth failure [26]. The homozygous c.119G > T (p.R10L) was shown to be associated with dysmorphic features, severe IUGR, and insulin resistance [27].

In conclusion, *IGF1R* gene alterations are an important and relatively frequent cause for SGA. Microcephaly with prenatal or postnatal growth failure should alert the physician on a possible *IGF1R* defect. Increased IGF-I levels are also a major sign of IGF1R defects. It is of note that low normal serum IGF-I levels have also been reported, and therefore are not an argument not to test the *IGF1R* gene. A precise phenotype-genotype correlation is still lacking. The GH treatment in one of the patients did result in a height gain into the parental range. However, the second patient interrupted the treatment but induced a catch-up growth.

Statement

Written informed consent has been obtained from each patient or subject after full explanation of the purpose and nature of all procedures used.

References

1. Woods KA, Camacho-Hubner C, Savage MO, Clark AJ. Intrauterine growth retardation and postnatal growth failure associated with deletion of the insulin-like growth factor I gene. *N Engl J Med*. 1996; 335:1363–1367. <https://doi.org/10.1056/NEJM199610313351805> PMID:8857020
2. Klammt J, Kiess W, Pfäffle R. IGF1R mutations as a cause of SGA. *Best Pract Res Clin Endocrinol Metab*. 2011; 25(1):191–206. <https://doi.org/10.1016/j.beem.2010.09.012> PMID:21396585
3. Kiess W, Kratzsch J, Keller E, Schneider A, Raile K, Klammt J, et al. Clinical examples of disturbed IGF signaling: intrauterine and postnatal growth retardation due to mutations of the insulin-like growth factor I receptor (IGF-IR) gene. *Rev Endocr Metab Disord*. 2005; 6(3):183–187. <https://doi.org/10.1007/s11154-005-3049-5> PMID:16151622
4. Abuzzahab MJ, Schneider A, Goddard A, Grigorescu F, Lautier C, Keller E, et al. Intrauterine Growth Retardation (IUGR) Study Group. IGF-I receptor mutations resulting in intrauterine and postnatal growth retardation. *N Engl J Med*. 2003; 349:2211–2222. <https://doi.org/10.1056/NEJMoa010107> PMID:14657428
5. Walenkamp MJ, Karperien M, Pereira AM, Hilhorst-Hofstee Y, van Doorn J, Chen JW, et al. Homozygous and heterozygous expression of a novel insulin-like growth factor-I mutation. *J Clin Endocrinol Metab*. 2005; 90:2855–2864. <https://doi.org/10.1210/jc.2004-1254> PMID:15769976
6. Walenkamp MJ, van der Kamp HJ, Pereira AM, Kant SG, van Duyvenvoorde HA, Kruithof MF, et al. A variable degree of intrauterine and postnatal growth retardation in a family with a missense mutation in the insulin-like growth factor I receptor. *J Clin Endocrinol Metab*. 2006; 91:3062–3070. <https://doi.org/10.1210/jc.2005-1597> PMID:16757531
7. Wietske AE, van Duyvenvoorde HA, de Wit CC, Broekman AJ, Ruivenkamp CAL, Govaerts LCP et al. Two Short Children Born Small for Gestational Age with Insulin-Like Growth Factor 1 Receptor Haploinsufficiency Illustrate the Heterogeneity of Its Phenotype. *J Clin Endocrinol Metab*. 2009; 94:4717–4727. <https://doi.org/10.1210/jc.2008-1502> PMID:19864454
8. Burkhardt S, Gesing J, Kapellen TM, Kovacs P, Kratzsch J, Schlicke M, et al. Novel heterozygous IGF1R mutation in two brothers with developing impaired glucose tolerance. *J Pediatr Endocrinol Metab*. 2015; 28(1-2):217–25. <https://doi.org/10.1515/jpem-2014-0132> PMID:25153223
9. Abbott AM, Bueno R, Pedrini MT, Murray JM, Smith RJ. Insulin-like growth factor I receptor gene structure. *J Biol Chem*. 1992; 267:10759–10763. PMID:1316909
10. Kromeyer-Hauschild K, Wabitsch M, Kunze D, Geller F, Geiß HC, Hesse V, von Hippel A, Jaeger U, Johnsen D, Korte W, Menner K. Perzentile für den Body-mass-Index für das Kindes- und Jugendalter unter Heranziehung verschiedener deutscher Stichproben. *Monatsschrift Kinderheilkunde*. 2001; 149(8):807–18. <https://doi.org/10.1007/s001120170107>
11. Greulich WW, Pyle SI, Todd TW. Radiographic atlas of skeletal development of the hand and wrist. Stanford: Stanford university press, 1959.
12. Solomon-Zemler R, Basel-Vanagaite L, Steier D, Yakar S, Mel E, Phillip M, et al. A novel heterozygous IGF-1 receptor mutation

- associated with hypoglycemia. *Endocr Connect.* 2017; 6(6):395-403. <https://doi.org/10.1530/EC-17-0038> PMID:28649085 PMCid:PMC5551424
13. Kawashima Y, Kanzaki S, Yang F, Kinoshita T, Hanaki K, Nagaishi J, et al. Mutation at cleavage site of insulin-like growth factor receptor in a short-stature child born with intrauterine growth retardation. *J Clin Endocrinol Metab.* 2005; 90:4679–4687. <https://doi.org/10.1210/jc.2004-1947> PMID:15928254
14. Inagaki K, Tiulpakov A, Rubtsov P, Sverdlova P, Peterkova V, Yakar S, et al. A familial insulin-like growth factor-1 receptor mutant leads to short stature: clinical and biochemical characterization. *J Clin Endocrinol Metab.* 2007; 92:1542–1548. <https://doi.org/10.1210/jc.2006-2354> PMID:17264177
15. Fang P, Schwartz DI, Johnson BD, Derr MA, Roberts CT, Hwa V, et al. Familial Short Stature Caused by Haploinsufficiency of the Insulin-Like Growth Factor I Receptor due to Nonsense-Mediated Messenger Ribonucleic Acid Decay. *The Journal of Clinical Endocrinology & Metabolism.* 2009; 94 (5):1740-1747. <https://doi.org/10.1210/jc.2008-1903> PMID:19240156
16. Mohn A, Marcovecchio ML, de Giorgis T, Pfaeffle R, Chiarelli F, Kiess W. An insulin-like growth factor-I receptor defect associated with short stature and impaired carbohydrate homeostasis in an Italian pedigree. *Horm Res Paediatr.* 2011; 76(2):136-143. <https://doi.org/10.1159/000324957> PMID:21811077
17. Choi JH, Kang M, Kim GH, Hong M, Jin HY, Lee BH, et al. Clinical and functional characteristics of a novel heterozygous mutation of the IGF1R gene and IGF1R haplo-insufficiency due to terminal 15q26.2->qter deletion in patients with intrauterine growth retardation and postnatal catch-up growth failure. *J Clin Endocrinol Metab.* 2011; 96(1):E130-4. <https://doi.org/10.1210/jc.2010-1789> PMID:20962017
18. Kawashima Y, Higaki K, Fukushima T, Hakuno F, Nagaishi J, Hanaki K, et al. Novel missense mutation in the IGF-I receptor L2 domain results in intra-uterine and postnatal growth retardation. *Clin Endocrinol (Oxf).* 2012; 77(2):246-54. <https://doi.org/10.1111/j.1365-2265.2012.04357.x> PMID:22309212
19. Fujimoto M, Kawashima Sonoyama Y, Hamajima N, Hamajima T, Kumura Y, Miyahara N, Nishimura R, Adachi K, Nanba E, Hanaki K, Kanzaki S. Heterozygous nonsense mutations near the C-terminal region of IGF1R in two patients with small-for-gestational-age-related short stature. *Clinical endocrinology.* 2015; 83(6):834-41. <https://doi.org/10.1111/cen.12791> PMID:25866162
20. Okubo Y, Siddle K, Firth H, O'Rahilly S, Wilson LC, Willatt L, et al. Cell proliferation activities on skin fibro-blasts from a short child with absence of one copy of the type 1 insulin-like growth factor receptor (IGF1R) gene and a tall child with three copies of the IGF1R gene. *J Clin Endocrinol Metab.* 2003; 88:5981–5988. <https://doi.org/10.1210/jc.2002-021080> PMID:14671200
21. Walenkamp MJ, de Muinck Keizer-Schrama SM, de Mos M, Kalf ME, van Duyvenvoorde HA, et al. Successful long-term growth hormone therapy in a girl with haploinsufficiency of the insulin-like growth factor-I receptor due to a terminal 15q26.2->qter deletion detected by multiplex ligation probe amplification. *J Clin Endocrinol Metab.* 2008; 93:2421–2425. <https://doi.org/10.1210/jc.2007-1789> PMID:18349070
22. Raile K, Klammt J, Schneider A, Keller A, Laue S, Smith R, et al. Clinical and functional characteristics of the human Arg59Ter insulin-like growth factor I receptor (IGF1R) mutate-on: implications for a gene dosage effect of the human IGF1R. *J Clin Endocrinol Metab.* 2006; 91:2264–2271. <https://doi.org/10.1210/jc.2005-2146> PMID:16569742
23. Juanes M, Guercio G, Marino R, Berensztein E, Warman DM, Ciaccio M, et al. Three novel IGF1R mutations in microcephalic patients with prenatal and postnatal growth impairment. *Clin Endocrinol (Oxf).* 2015; 82(5):704-711. <https://doi.org/10.1111/cen.12555> PMID:25040157
24. Labarta JI, Barrio E, Audí L, Fernández-Cancio M, Andaluz P, de Arriba A, et al. Familial short stature and intrauterine growth retardation associated with a novel mutation in the IGF-I receptor (IGF1R) gene. *Clin Endocrinol (Oxf).* 2013; 78(2):255-262. <https://doi.org/10.1111/j.1365-2265.2012.04481.x> PMID:22738321
25. Mahmoud R, Naidu A, Risheg H, Kimonis V. Response to Growth Hormone Treatment in a Patient with Insulin-Like Growth Factor 1 Receptor Deletion. *J Clin Res Pediatr Endocrinol.* 2017; 9(4):380-386. <https://doi.org/10.4274/jcrpe.4456> PMID:28720553 PMCid:PMC5785648
26. Fang P, Cho YH, Derr MA, Rosenfeld RG, Hwa V, Cowell CT. Severe short stature caused by novel compound heterozygous mutations of the insulin-like growth factor 1 receptor (IGF1R). *J Clin Endocrinol Metab.* 2012; 97(2):E243-247. <https://doi.org/10.1210/jc.2011-2142> PMID:22130793
27. Gannagé-Yared MH, Klammt J, Chouery E, Corbani S, Mégarbané H, Choucair N, Pfäffle R, Mégarbané A. Homozygous mutation of the IGF1 receptor gene in a patient with severe pre- and postnatal growth failure and congenital malformations. *Eur J Endocrinol.* 2012; 168(1):K1-7. <https://doi.org/10.1530/EJE-12-0701> PMID:23045302