

Quantum Gravity in Relativistic Alpha Field Theory (QG in RAFT)

Branko M. Novakovic

Abstract – Classical quantum gravity approach is presented in the frame work of the new Relativistic Alpha Field Theory (RAFT). It is very important to describe gravity at the Planck's scale and near compact astrophysical objects where quantum effects cannot be ignored. Meanwhile, General Relativity Theory (GRT) cannot be applied to the extremely strong gravitational field at the Planck's scale, because of the related singularity. Recently developed RAF theory extends the application of GRT to the extremely strong fields, including Planck's scale. This is the consequence of determination of field parameters by using condition that they should satisfy both relativistic Hamiltonian and field equations with energy momentum tensor. As the result, we obtain, among the others, the following prediction of RAF theory: there exists a minimal gravitational radius at $r = (GM/2c^2)$ that prevents singularity at $r = 0$. The existence of the minimal gravitational radius opens a new avenue for quantization of a gravitational field. One of the possible approaches is presented in this paper.

Index Terms- Quantum gravity, Relativistic Alpha Field Theory (RAFT), Minimal gravitational radius, Extremely strong gravitational fields

I. INTRODUCTION

It is of the special interest to describe gravity at the Planck's scale and near compact astrophysical objects where quantum effects cannot be ignored [1-5]. The main difficulties in quantum gravity (QG) description appear when one attempts to apply the usual prescriptions of quantum field theory to the force of gravity via graviton bosons [6]. The problem is that the theory one gets in this way is not renormalizable and therefore cannot be used to make meaningful physical predictions. Therefore, theorists have taken up more radical approaches to the problem of QG. The most popular approaches are string theory [7-10] and loop quantum gravity [11-15]. The noncommutative theories of geometry and an overview of many current approaches to QG are presented in [16-22]. The quantum cosmology problems and some collections of philosophers' texts of QG and popularization works are illustrated in [23-32].

As it is well known, the Planck's scale is far smaller in distance than those currently accessible at high energy particle accelerators. Therefore, QG is a mainly theoretical enterprise. Meanwhile, in the field of phenomenological QG, we have studies of the possibility of experimental tests of QG [33-37]. Unfortunately, at present, there is no QG theory which is universally accepted and confirmed by experience. One of the major hopes of observing traces of QG phenomena is in the cosmological context. In that sense, we have detection of an electromagnetic counterpart (GRB 170817A)

to the gravitational-wave signal (GW170817) from the merger of two neutron stars [38-41]. This opens a completely new arena for testing theories of gravity.

Further, General Relativity Theory (GRT) [42-47] cannot be applied to the extremely strong gravitational field at the Planck's scale, because of the related singularity. Here we apply a new theory that is called Relativistic Alpha Field (RAF) theory [48-50]. It has been shown [51] that RAF theory extends the capability of the GRT for the application to the extremely strong fields, including the Planck's scale. This is the consequence of determination of field parameters by using condition that they should satisfy both generalized relativistic Hamiltonian and field equations, with energy momentum tensor and without cosmological constant Λ [48,52]. As the result, we obtain the following predictions of RAF theory: a) no a singularity at the Schwarzschild radius [49], b) there exists a minimal gravitational radius at $r = (GM/2c^2)$ that prevents singularity at $r = 0$, i.e. the nature protects itself [49], c) the gravitational force becomes positive (repulsive) if $(GM/rc^2) > 1$, that could be a source of a dark energy [50,53], and d) unification of electrical and gravitational forces can be done in the standard four dimensions (4D) [50]. The existence of the minimal gravitational radius opens a new avenue for quantization of a gravitational field.

The presented predictions of RAF theory may look like too speculative items. But, it is not because the solution of the field parameters generates energy - momentum tensor (EMT) on the right side of the field equations. Thus, this tensor is not added by the hand, as it is suggested in GRT. Furthermore, this EMT obeys the following required properties: symmetry, non-negative energy density and zero trace of EMT. Without EMT, the solution of the field parameters in RAF theory is reduced to the Schwarzschild's solution for the vacuum, as we expected that should be. This vacuum solution cannot be applied to the extremely strong fields, because of the related singularity. In that sense, it has been discussed the possibility that the singularity and dark energy problems are the consequences of vacuum solution of the field equations [54].

This paper is organized as follows. In Sec. II, we show the theoretical proof of the existence of the minimal radius. The quantum gravity procedures are presented in Sec. III. Some possible influences of the presented quantization are considered in Sec. IV. Finally, the related conclusion and the reference list are presented in Sec. V. and Sec. VI, respectively.

II. EXISTENCE OF THE MINIMAL GRAVITATIONAL RADIUS

Many arguments indicate that the Planck length may appear as a minimal meaningful length. But, in RAF theory [48-50] it has been predicted that there exists a minimal

Branko Novakovic, FSB – University of Zagreb, Luciceva 5, P.O.B. 509, 10000 Zagreb, Croatia

radius in gravitational field equal to $r_m = GM/2c^2$, that eliminates the singularity problem. As we can see the minimal radius is proportional to mass. Thus, the minimal length belongs to the minimal mass. Minimal radius of the Planck's mass is half of the Planck's length $r_{pm} = L_p/2$. This means that the Planck's length is the minimal length only for the Planck's mass. As an example, the minimal radius of the proton mass is equal to $r_m = 0.62 \times 10^{-54} m$, which is too smaller than half of the Planck's length. Unfortunately, current technology is not yet capable of observing physical effects at scales that are so small. Therefore, we cannot confirm any of the current tentative QG theories, until the genuine quantum gravitational phenomena are directly or indirectly observed.

In order to theoretically prove the existence of the minimal radius in gravitational field, we started with the general line element ds^2 in an alpha field, derived in the first part of RAFT theory [48]. In the spherical polar coordinates this line element is presented by the nondiagonal form

$$ds^2 = -\alpha\alpha' c^2 dt^2 - \kappa(\alpha - \alpha') c dt dr + dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2. \quad (1)$$

Generally, field parameters α and α' are scalar functions of the space-time coordinates, or normalized potential energy (U/mc^2) in an alpha field [48]. Here U is potential energy, m is the related particle rest mass in that field, c is the speed of light in vacuum and parameter $\kappa = \pm 1$. Since, the line element (1) belongs to the well-known form of the Riemann's type line element, the contra-variant coordinates and components of the covariant metric tensor are given by the relations:

$$\begin{aligned} dx^0 &= c dt, & dx^1 &= dr, & dx^2 &= d\theta, & dx^3 &= d\phi, \\ g_{00} &= -\alpha\alpha', & g_{01} &= g_{10} = -\kappa(\alpha - \alpha')/2, & & & & \\ g_{11} &= 1, & g_{22} &= r^2, & g_{33} &= r^2 \sin^2 \theta. \end{aligned} \quad (2)$$

Starting with the line element (1) we employ, for the convenient, the following substitutions:

$$\eta = \alpha\alpha', \quad \beta = \kappa(\alpha' - \alpha)/2. \quad (3)$$

In that case the nondiagonal line element (1) is transformed into the new relation

$$ds^2 = -\eta c^2 dt^2 + 2\beta c dt dr + dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2. \quad (4)$$

The determination of the parameters η and β in the gravitational field has been obtained by using condition that field parameters should satisfy both the generalized relativistic Hamiltonian [52] and field equations [48], with energy momentum tensor $T_{\mu\nu}$ and without cosmological constant

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = k T_{\mu\nu}, \quad k = \frac{8\pi G}{c^4}, \quad \mu, \nu = 0, 1, 2, 3. \quad (5)$$

Here $R_{\mu\nu}$ is Ricci tensor, R is Ricci scalar, G is the Newton's gravitational constant and c is the speed of the light in a vacuum. Thus, the solutions of the parameters η and β

for the static gravitational field have been presented in the first and second parts of RAFT theory [48, 49]:

$$\begin{aligned} \eta &= 1 - \frac{2GM}{rc^2} + \left(\frac{GM}{rc^2}\right)^2 = \left(1 - \frac{GM}{rc^2}\right)^2, \\ \beta &= \mp \kappa \sqrt{\frac{2GM}{rc^2} \left(1 - \frac{GM}{2rc^2}\right)}, \quad \kappa = \pm 1. \end{aligned} \quad (6)$$

Here M is gravitational mass of the spherically symmetric not rotating body. The quadratic term $(GM/rc^2)^2$ in (6) generates the energy - momentum tensor on the right side of the Einstein's field equations [48-50]:

$$\begin{aligned} T_{\mu\nu} &= (T_{00}, T_{01}, T_{10}, T_{11}, T_{22}, T_{33}), \\ T_{\mu\nu} &= \left(\eta, -\beta, -\beta, -1, r^2, r^2 \sin^2 \theta\right) \frac{(GM)^2}{8\pi G r^4}. \end{aligned} \quad (7)$$

It is easy to prove that the energy - momentum tensor (EMT) in (7) obeys the following required properties: symmetry, non-negative energy density and zero trace of EMT. In relatively weak field, as is in our solar system, or in vacuum, the quadratic term $(GM/rc^2)^2$ is too small and can be neglected. In that case EMT is vanishing and parameters in (6) are transformed into the solution corresponding to the Schwarzschild's solution for the vacuum:

$$\eta = 1 - \frac{2GM}{rc^2}, \quad \beta = \mp \kappa \sqrt{\frac{2GM}{rc^2}}, \quad \kappa = \pm 1. \quad (6a)$$

Including the parameters solutions from (6) into (4), the line element (4) can be presented in the final form:

$$\begin{aligned} ds^2 &= -\left(1 - \frac{GM}{rc^2}\right)^2 c^2 dt^2 \\ &\mp 2\kappa \sqrt{\frac{2GM}{rc^2} \left(1 - \frac{GM}{2rc^2}\right)} c dt dr \\ &+ dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2. \end{aligned} \quad (8)$$

Thus, it is very important to point out that RAFT theory includes both the full form solution (6) and vacuum solution (6a) of the field equations (5).

From (6) and (8) we can theoretically prove the following predictions of RAFT theory: a) no singularity at the Schwarzschild radius and b) there exist the minimal radius at the position $r_m = (GM/2c^2)$ that prevents singularity at $r = 0$, i.e. the nature protects itself:

$$\begin{aligned} r_{sch} &= \frac{2GM}{c^2}, \rightarrow \eta_{sch} = \frac{1}{4}, \beta_{sch} = \mp \kappa \sqrt{\frac{3}{4}}, \kappa = \pm 1, \\ r_m &= \frac{GM}{2c^2}, \rightarrow \eta = 1, \beta = 0, r < r_m \rightarrow \beta = \beta_{im}, \\ r \rightarrow \infty, \eta &\rightarrow 1, \beta \rightarrow 0, r = r_c = \frac{GM}{c^2}, \rightarrow \eta = 0, \beta = 1. \end{aligned} \quad (9)$$

Following (9) we can see that at the Schwarzschild radius, r_{sch} , parameters η and β are regular. This proves the prediction a): no singularity at the Schwarzschild radius.

Further, from (9), we also can see that at the minimal radius, $r_m = GM/2c^2$, parameters η and β are also regular and for $r < r_m$ parameter β becomes imaginary number, $\beta = \beta_{im}$. This proves the prediction b): there exists a minimal radius at $r_m = (GM/2c^2)$ that prevents singularity at $r = 0$. It seems that the existence of the minimal radius tells us that the nature protect itself from the singularity. Thus, we can say that the metrics of the line element in (4-6) is regular for a gravitational field in the region $r_m \leq r \leq \infty$. On that way the proofs of the propositions a) and b) are finished.

Now we shall consider the prediction c) of the RAF theory, given in [49,50]: the gravitational force becomes positive (repulsive) if $(GM/rc^2) > 1$, that could be a source of a dark energy. In order to theoretically prove this prediction, we can start with the solutions of the radial acceleration and radial force. For the time-invariant (or very slowly changed) gravitational field, the radial acceleration \ddot{a} and force F are presented by the relations [50]:

$$\ddot{a} = -\frac{GM}{r^2} \left(1 - \frac{GM}{rc^2}\right), \quad (10)$$

$$F = m\ddot{a} = -\frac{mGM}{r^2} \left(1 - \frac{GM}{rc^2}\right).$$

Here m is a rest mass of a particle. Generally, the gravitational radial force in (10), describes the interactions in the strong fields. In the case of the relatively weak fields (as we have in our solar system) the radial force relation (10) is reduced to the well-known descriptions of the interactions in those fields:

$$\ddot{a} = -\frac{GM}{r^2}, \quad F = m\ddot{a} = -\frac{mGM}{r^2}. \quad (10a)$$

Further, from (10) we can see that the gravitational force is negative (attractive) for $(GM/rc^2) < 1$. This is the case in the relatively weak gravitational field, where the term (GM/rc^2) can be neglected. On the other hand, the gravitational force becomes positive (repulsive) if $(GM/rc^2) > 1$. This is the case in the extremely strong gravitational field where the term (GM/rc^2) has to be taken into account.

In our solar system the term (GM/rc^2) is too small and can be neglected. For an example, on the surface of our Sun the amount of this term is $(GM/rc^2) \approx 2.1193 \times 10^{-6} \ll 1$. On the surface of our planet Earth the related influence of the Sun to this term is $(GM/rc^2) \approx 0.989 \times 10^{-8} \ll 1$. The gravitational influence of the planet Earth on its surface is $(GM/rc^2) \approx 0.695 \times 10^{-9} \ll 1$. The presented amounts of the term (GM/rc^2) in our solar system could be the answer to the question: why our experience is that gravitational force is only the negative (attractive) force?

If the term $(GM/rc^2) = 1$, then the gravitational force is equal to zero. This is the very important situation that is occurring at the gravitational radius $r_c = (GM/c^2)$. This radius separates the regions of the attractive and repulsive forces in a gravitational field. At the minimal radius r_m gravitational force is positive (repulsive): $F_m = (4mc^4)/GM$. Now, we can say that the nature protects itself from the singularity by producing the positive (repulsive) gravitational force at the minimal radius. Further, from (10) we can see that the gravitational force is positive (repulsive) in the region $r_m \leq r <$

r_c , where $r_m = (GM/2c^2)$ and $r_c = (GM/c^2)$. At the Schwarzschild radius the gravitational force is negative (attractive) $F_{sch} = -(mc^4/8GM)$ and belongs to the negative (attractive) set of gravitation forces in the region $r_c < r \leq \infty$.

Thus, the previous consideration theoretically confirms the prediction c) of the RAF theory: the gravitational force becomes positive (repulsive) if $(GM/rc^2) > 1$ that could be a source of a dark energy [53].

III. QUANTUM GRAVITY PROCEDURES

Following (9) we can see that there are two exclusive points in a gravitational static field. The first one is at the minimal radius, $r_m = (GM/2c^2)$, where the parameters $\eta=1$ and $\beta=0$. The second point is at the radius $r_c = (GM/c^2)$, where $\eta=0$ and $\beta=1$. We also can conclude that $r_c = 2r_m$. These two points are very important items and can be useful in derivation of the quantum gravity procedures. Namely, we can calculate the states of the velocities, accelerations (forces) and energies at the two exclusive points. Thus, it can be proved that at the point $r_m = (GM/2c^2)$, radial velocity is equal to zero and radial acceleration (force) is positive and maximal. This means that a particle at this point will be repulsed to the bigger radius. Here we have repulsive (positive) gravitational force that also could be source of the so called dark energy [53]. On the other side, at the point $r_c = (GM/c^2)$, radial velocity is maximal and equal to the speed of the light in vacuum, and radial acceleration (force) is equal to zero [50]. If a gravitational radius $r > r_c$, then radial velocity is decreasing and radial acceleration (force) becomes negative (attractive). Thus, the radius r_c separates the region of the positive gravitational forces $r_m \leq r < r_c$ from the region of the negative gravitational forces $r_c < r \leq \infty$.

In order to make the quantization of a gravitation field we can start with determination of the potential energies of the particle at the minimal radius $r_m = (GM/2c^2)$, and at the radius $r_c = (GM/c^2)$:

$$E_1 = \frac{mGM}{r_m} = \frac{mGM}{GM/2c^2} = 2mc^2, \quad (11)$$

$$E_2 = \frac{mGM}{r_c} = \frac{mGM}{GM/c^2} = mc^2.$$

Here m is a particle rest mass.

A. Definition of graviton

The transition from the energy state E_1 (at the minimal radius $r_m = r_1$) to the energy state E_2 (at the radius $r_c = r_2$) is resulted with the emission (radiation) of the graviton, as a quant of gravitational field, with the maximal energy:

$$E_1 - E_2 = 2mc^2 - mc^2 = mc^2, \quad (12)$$

$$h\nu_{1,2} = \vec{d}\omega_{1,2} = mc^2.$$

Here h is the Planck's constant, \vec{d} is the reduced Planck's constant, $\nu_{1,2}$ is the related gravitational wave frequency between energy states (levels) 1 and 2 and $\omega_{1,2}$ is the related angular frequency. Thus, we can say that this quant of energy (graviton) is equal to the gravitational potential energy between the minimal radius $r_m = r_1$ and the radius $r_c = r_2$. In the opposite case, the transition from the energy state E_2 (at the

radius $r_c=r_2$) to the energy state E_1 (at the minimal radius $r_m=r_1$) is resulted with the absorption of the graviton, as a quant of gravitational field, with the same maximal energy. This is the physical concept of the quantization of a gravitational field.

B. Quantization of gravitational radius, energy and frequency

Quantization of the gravitational radius can be done by using the known values of the minimal radius r_m and the radius r_c , given by (9):

$$r_1 = r_m = \frac{GM}{2c^2}, \quad r_2 = r_c = 2r_m = \frac{GM}{c^2}, \quad (13)$$

$$r_3 = 3r_m = 3\frac{GM}{2c^2}, \dots, \quad r_n = nr_m = n\frac{GM}{2c^2}.$$

Here n is quantization number of a gravitational field. Quantization of the gravitational potential energy follows from (11) and (13):

$$E_1 = \frac{mGM}{r_1} = 2mc^2, \quad E_2 = \frac{mGM}{r_2} = \frac{E_1}{2} = mc^2, \quad (14)$$

$$E_3 = \frac{mGM}{r_3} = \frac{2mc^2}{3} = \frac{E_1}{3}, \quad E_n = \frac{E_1}{n}, \quad n=1,2,..$$

In order to quantize the frequency ν and the related angular frequency ω between two energy states (levels) in a gravitational field, we can employ (12) and (14):

$$h\nu_{1,2} = \hbar\omega_{1,2} = E_1 - E_2 = E_1 - \frac{E_1}{2} = \frac{E_1}{2},$$

$$h\nu_{2,3} = \hbar\omega_{2,3} = E_2 - E_3 = \frac{E_1}{2} - \frac{E_1}{3} = \frac{E_1}{6}, \quad (15)$$

$$h\nu_{n-1,n} = \hbar\omega_{n-1,n} = \frac{E_1}{(n-1)n}, \quad n=2,3,..$$

Here $\nu_{n-1,n}$ is the gravitational wave frequency between energy states (levels) $(n-1)$ and n , where $n=2,3,..$. In that sense $\omega_{n-1,n}$ is the related angular frequency. Thus, the quantization form of the frequency ν and the related angular frequency ω between two energy states (levels) in a gravitational field are given by the relations:

$$\nu_{n-1,n} = \frac{E_1}{(n-1)nh} = \frac{2mc^2}{(n-1)nh}, \quad (16)$$

$$\omega_{n-1,n} = \frac{2mc^2}{(n-1)n\hbar}, \quad n=2,3,..$$

C. Quantization of line element, energy momentum tensor, radial acceleration and force

Now, we are ready to quantize the line element (8). In that sense, we started with parameters η and β from (6):

$$\eta_1 = \left(1 - \frac{GM}{r_1 c^2}\right)^2, \quad \beta_1 = \mp \kappa \sqrt{\frac{2GM}{r_1 c^2} \left(1 - \frac{GM}{2r_1 c^2}\right)},$$

$$\kappa = \pm 1, \quad r_1 = r_m = \frac{GM}{2c^2}, \quad r_n = n \cdot r_m = \frac{nGM}{2c^2}, \quad (17)$$

$$\eta_n = \left(1 - \frac{2}{n}\right)^2, \quad \beta_n = \mp \kappa \sqrt{\frac{4}{n} \left(1 - \frac{1}{n}\right)}, \quad n=1,2,..$$

Thus, the quantization form of the line element (8) is given by the relation:

$$ds_n^2 = -\left(1 - \frac{2}{n}\right)^2 c^2 dt_n^2 \mp 2\kappa \sqrt{\frac{4}{n} \left(1 - \frac{1}{n}\right)} c dt_n dr_n \quad (18)$$

$$+ dr_n^2 + r_n^2 d\theta_n^2 + r_n^2 \sin^2 \theta_n d\phi_n^2, \quad n=1,2,..$$

In order to quantize the energy-momentum tensor on the right side of the Einstein's field equations [48-50], we can employ the relation (7) and the previous quantization procedure:

$$T_{n_{\mu\nu}} = (T_{n_{00}}, T_{n_{01}}, T_{n_{10}}, T_{n_{11}}, T_{n_{22}}, T_{n_{33}}),$$

$$T_n = \left(\frac{2c^2}{n}\right)^4 \frac{1}{8\pi G (GM)^2}, \quad n=1,2,.., \quad (19)$$

$$T_{n_{00}} = \eta_n T_n, \quad T_{n_{01}} = T_{n_{10}} = -\beta_n T_n,$$

$$T_{n_{11}} = -T_n, \quad T_{n_{22}} = r_n^2 T_n, \quad T_{n_{33}} = \sin^2 \theta_n T_{n_{22}}.$$

Finally, the quantization forms of the radial acceleration \ddot{a} and radial force F , given by (10), are presented by the following relations:

$$\ddot{a}_n = -\frac{4c^4}{n^2 GM} \left(1 - \frac{2}{n}\right), \quad F_n = m\ddot{a}_n,$$

$$F_n = -\frac{4c^4}{n^2 G} \left(\frac{m}{M}\right) \left(1 - \frac{2}{n}\right), \quad n=1,2,.., \quad (20)$$

$$m = M \rightarrow F_n = -\frac{4c^4}{n^2 G} \left(1 - \frac{2}{n}\right).$$

From (20) we can see that the quantized gravitational radial force F_n between two particles (bodies) with equal masses ($m=M$) is not function of the mass. At the minimal radius, $n=1$, the gravitational acceleration and force are positive (repulsive) and maximal. For the quantum number $n=2$, the gravitational acceleration and force are equal to zero. If the quantum number $n \geq 3$, the gravitational acceleration and force are negative (attractive). Since our Universe is expanding at the acceleration rate, we can conclude that the present state of the Universe may be between the quantum states $n=1$ and $n=2$, but closer to the quantum state $n=2$. This could be an experimental test of RAF theory and of the presented quantum procedures of a gravitational field. At the same time, this means that between two quantum states there exists a transition process. At the Planck's scale this process is very, very short (looking from the macro time scale). On the other side, this process is very, very long at the macro time scale and should be taken into account.

D. Quantization of radial density

Now, the quantization of the radial density $\rho=M/r$ is presented. This quantization is the function of the minimal radius:

$$\begin{aligned}\rho_1 &= \rho_M = \frac{M}{r_1} = \frac{M}{r_m} = \frac{2c^2}{G}, \\ \rho_2 &= \frac{M}{r_2} = \frac{M}{r_c} = \frac{M}{2r_m} = \frac{\rho_M}{2} = \frac{c^2}{G}, \\ \rho_n &= \frac{M}{r_n} = \frac{M}{nr_m} = \frac{\rho_M}{n} = \frac{2c^2}{nG}, \quad n=1,2,\dots\end{aligned}\quad (21)$$

Here $\rho_M=2c^2/G$ is the maximal radial density in a gravitational field. From (21) we can see that the radial density is not the explicit function of the mass. This means that all massive particles (bodies) have the same maximal radial density and the same radial density at each quantum number $\rho_n=2c^2/nG$, $n=1,2,\dots$. This is very important discovery which predicts that radial density of any particle (body) cannot be greater than $2c^2/G$.

IV. INFLUENCES OF THE PRESENTED QUANTIZATION

The presented quantization of a gravitational field could have very important influences to the regions like black holes, quantum field theory, high energy physics, Big Bang theory, dark energy, Universe acceleration, supernova explosion and cosmology. The presented quantization of gravitational field may also be useful in determination of the present Universe state and its maximal radius.

V. CONCLUSION

Recently developed Relativistic Alpha Field (RAF) theory [48-50] extends the capability of GRT for applications to the extremely strong fields, including Planck's scale [51]. One of the important predictions of RAF theory is the existence of the minimal gravitational radius equal to $r_m=(GM/2c^2)$. This prediction eliminates the singularity problem in a strong gravitational field and opens a new avenue for the quantization of a gravitational field. One of the possible approaches is presented in this paper.

The quantization of the radial density shows that all massive particles (bodies) have the same maximal radial density $\rho_M=2c^2/G$, and the same radial density at each quantum number $\rho_n=2c^2/nG$. Thus, the same maximal radial density have particles like electron and proton, as well as bodies like planet Earth, Sun and Universe. This is very important discovery which predicts that radial density of any particle (body) cannot be greater than $2c^2/G$.

The presented predictions of RAF theory may look like too speculative items. But, it is not. This is just the consequence of determination of field parameters by using condition that they should satisfy both generalized relativistic Hamiltonian and field equations with energy momentum tensor on the right side of them.

ACKNOWLEDGMENTS

The author wishes to thank to the anonymous reviewers for a variety of helpful comments and suggestions. This work is

supported by grants (120-1201842-3048) from the National Scientific Foundation of Republic of Croatia.

REFERENCES

- [1] Rovelli C 2018 *Quantum gravity* (Scholarpedia www.scholarpedia.org)
- [2] Kiefer C 2012 *Quantum gravity*, (Oxford: Oxford University Press)
- [3] Griffiths D J and David J 2004 *Introduction to Quantum Mechanics* (Pearson Prentice Hall, OCLC 803860989)
- [4] Wald R M 1984 *General Relativity* (Chicago: University of Chicago Press p 382 OCLC 471881415)
- [5] Wald R M 1994 *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics* (Chicago: University of Chicago Pres ISBN 0-226-87027-8)
- [6] Zee A 2010 *Quantum Field Theory in a Nutshell* (Princeton: Princeton University Press p 172 OCLC 659549695)
- [7] Green M, Schwarz J and Witten E 1987 *Superstring Theory* (Cambridge: Cambridge University Press)
- [8] Polchinski J 1990 *String Theory* (Cambridge: Cambridge University Press)
- [9] Blau M and Theisen S 2009 *Gen. Relativ. Gravit.* **41** 743
- [10] Blumenhagen R, Lüst D and Theisen S 2013 *Basic Concepts of String Theory* (Berlin: Springer)
- [11] Rovelli C 2004 *Quantum Gravity* (Cambridge: Cambridge University Press)
- [12] Thiemann T 2007 *Modern Canonical Quantum General Relativity* (Cambridge: Cambridge University Press)
- [13] Baez J C and Muniain J P 1994 *Gauge Fields, Knots, and Gravity* (World Scientific Publishing Company)
- [14] Ashtekar A and Singh P 2011 *Class. Quantum Grav.* **28**, 213001
- [15] Rovelli C 2013 *Covariant loop gravity* (Quantum Gravity and Quantum Cosmology, pp 57–66 Lecture Notes in Physics 863 Berlin: Springer).
- [16] Connes A 1994 *Noncommutative Geometry* (Sand Diego: Academic Press)
- [17] Bastos C, Bertolami O, Costa Dias N and Nuno Prata J 2008 *Phys. Rev. D* **78**, 023516
- [18] Ashtekar A and Stachel J 1991 *Conceptual Problems of Quantum Gravity* (Boston-Basel-Berlin: Birkhauser)
- [19] Oriti D 2008 *Approaches to Quantum Gravity: Towards a New Understanding of Space, Time and Matter* (Cambridge UP, Cambridge UK)
- [20] Hamber H W 2009 *Quantum gravitation – The Feynman path integral approach* (Berlin: Springer)
- [21] Pinto-Neto N, Falciano F T, Pereira R and Santini E S 2012 *Phys. Rev. D* **86** 063504
- [22] Haug E G 2016 *Int. J. Astron. Astrophys.* **6**, 206
- [23] Penrose R 2009 *J. Phys. Conf. Ser.* **174** 012001
- [24] Kamenshchik A Y, Kiefer C and Sandhöfer B 2007 *Phys. Rev. D* **76** 064032
- [25] Bojowald M 2012 *Class. Quantum Grav.* **29**, 213001
- [26] Callender C and Huggett N 2001 *Physics Meets Philosophy at the Planck Scale: Contemporary Theories in Quantum Gravity* (Cambridge UP, Cambridge UK)
- [27] Rickles D, French S and Saatsi J 2007 *The Structural Foundations of Quantum Gravity* (Oxford: Oxford University Press)
- [28] Smolin L 2000 *Three Roads to Quantum Gravity* (Oxford: Oxford University Press)
- [29] Penrose R 2005 *The road to reality* (Oxford: Oxford University Press)
- [30] Penrose R 2007 *The road to reality: a complete guide to the laws of the universe* (Vintage p 1017 OCLC 716437154)
- [31] Green B 1999 *The Elegant Universe* (New York: Vintage)
- [32] Rovelli C 2004 *What is Time? What is Space?* (Rome: Di Renzo editore)
- [33] Strominger A 2009 *Nucl. Phys. Proc. Suppl.* 192–193, 119–25
- [34] Bertl E et al 2015 *Testing general relativity with present and future astrophysical observations* *Class. Quantum Grav.* **32** 243001 IOPscience(<http://iopscience.iop.org/0264-0381/32/24/243001>)
- [35] Carlisle C 2014 *First Direct Evidence of Big Bang Inflation* (*SkyandTelescope.com* Retrieved March 18)
- [36] Marongwel S 2017 *Probing quantum gravity through strong gravitational lensing* (DOI: 10.13140/RG.2.2.23029.88806)
- [37] Dey S, Bhat A, Momeni D, Faizal M, Ali A F, Dey T K and Rehman A 2017 *Nuclear Physics B* [10.1016/j.nuclphysb.2017.09.024](https://doi.org/10.1016/j.nuclphysb.2017.09.024) [arXiv:1710.03920](https://arxiv.org/abs/1710.03920) [quant-ph]

- [38] Baker T, Bellini E, Ferreira P G, Lagos M, Noller J, and Sawicki I 2017 *Phys. Rev. Lett.* **119** 251301
- [39] Creminelli P and Vernizzi F 2017 *Phys. Rev. Lett.* **119** 251302
- [40] Sakstein J and Jain B 2017 *Phys. Rev. Lett.* **119** 251303
- [41] Ezquiaga J M and Zumalacárregui M 2017 *Phys. Rev. Lett.* **119** 251304
- [42] Einstein A 1916 *Ann. Phys.* **49** 769-822
- [43] Einstein A 1995 *The Meaning of Relativity* (Princeton: Princeton Univ. Press)
- [44] Sean C 2003 *Spacetime and Geometry: An Introduction to General Relativity* (Amazon.com Books.htm, Hardcover)
- [45] Weinberg S 1972 *Gravitation and Cosmology: Principles and Application of the General Theory of Relativity* (Gebundene Ausgabe, RelEspWeinberg.pdf.)
- [46] S.W. Hawking S W and Ellis G F R 1973 *The Large Scale Structure of Space-Time* (Cambridge: Cambridge Univ. Press)
- [47] Blau M 2014 *Lecture Notes on General Relativity* (Bern: A. Einstein Center for Fundamental Physics, Univ. Bern)
- [48] Novakovic B M 2015 *Int. J. New Techn. Res.* **1** 5 ID-IJNTR01050015 www.ijntr.org.
- [49] Novakovic B M 2015 *Int. J. New Techn. Res.* **1** 5 ID-IJNTR01050016 www.ijntr.org.
- [50] Novakovic B M 2015 *Int. J. New Techn. Res.* **1** 5 ID-IJNTR01050017 www.ijntr.org.
- [51] Novakovic B M 2017 *Int. J. New Techn. Res.* **3** 11 ID-IJNTR03110009 www.ijntr.org.
- [52] Novakovic B M 2010 **CP** 1303 Computing Anticipatory Systems CASYS'09 Ninth Int. Conf. ad. D M Dubois – 2010 AIP 978-0-7354-0858-6/10/\$30.00
- [53] Novakovic B M 2015 *Int. J. New Techn. Res.* **1** 7 ID-IJNTR01070005 www.ijntr.org.
- [54] Novakovic B M 2017 *Int. J. New Techn. Res.* **3** 11 ID-IJNTR03110008 www.ijntr.org.



Branko Novakovic is a Professor emeritus at FSB – University of Zagreb, Croatia. Prof. Novakovic received his PhD from the University of Zagreb in 1978. His research of interest includes physics, control systems, robotics, neural networks, and fuzzy control. He is author of three books; Relativistic Alpha Field Theory (RAFT, e-book, 2016), Control Methods in Robotics, Flexible Manufacturing Systems and Processes (1990), Control Systems (1985) and the first co-author of a book Artificial Neural Networks (1998). He has published over 230 research papers in his research of interest.