

Abstract Control Panel

Abstract ID: 358489

Primary Section:
Volcanology,
Geochemistry and
Petrology (V)

Signed in as: Olga Girina
([Sign out](#))

8 Steps to submit an
Abstract:

1. Session
2. Abstract Details
3. Author(s)
4. Abstract Text
5. Plain-Language Summary (Optional)
6. Special Programs
7. Payment
8. Submit Abstract

Options:

[-View Submission](#)

Abstract #358489

The VolSatView for Satellite Monitoring and Kamchatkan Volcanoes Study

Olga Girina, Institute of Volcanology and Seismology, Petropavlovsk Kamcha, Russia

Abstract Text:

Girina O.A. (1), Loupian E.A. (2), Efremov V.Yu. (2), Melnikov D.V. (1), Manevich A.G. (1), Gordeev E.I. (1), Sorokin A.A. (3), Kramareva L.S. (4), Uvarov I.A. (2), Kashnitskiy A.V. (2), Bourtsev M.A. (2), Konstantinova A. (2)

(1) Institute of Volcanology and Seismology, FEB RAS, Russia, (2) Space Research Institute, RAS, Russia, (3) Computing Center, FEB RAS, Russia, (4) Far East Planeta Center of Space Hydrometeorology Research, Russia

Annually, from 3 to 6 Kamchatkan volcanoes produce eruptions, during which the explosions eject ash to 10-15 km a.s.l., and ash clouds spread thousands of kilometers from volcanoes. Strenuous volcanic activity could cause ash falls in towns and settlements, destruction of forests and communications. Ash clouds plumes pose a serious threat to the modern jet aviation. Scientists of KVERT have conduct daily monitoring of Kamchatka volcanoes since 1993, to mitigate volcanic hazards to airline operations and population. Since 2014, satellite monitoring of volcanoes is carried out by KVERT scientists using the VolSatView (<http://volcanoes.smislab.ru>) IS. The VolSatView (Remote monitoring of active volcanoes of Kamchatka and the Kuril Islands) was created in 2011 by scientists from Space Research Institute (SRI) of Russian Academy of Sciences (RAS), Institute of Volcanology and Seismology of Far East Branch (FEB) RAS, Computing Center of FEB RAS and Far East Planeta Center of Space Hydrometeorology Research, and the IS continues to developing. The system utilize all the available satellite data, weather and video observations to ensure continues monitoring and study of volcanic activity in Kamchatka. Architecture of the VolSatView IS was developed for the work with distributed information resources and computation systems that are used for the acquisition, processing, storage, analysis, and visualization of various instrumental and scientific data. This work was supported by the Russian Science Foundation, project No. 16-17-00042.

Plain-Language Summary:

Girina Olga (1), Loupian Evgenii (2), Efremov Viktor (2), Melnikov Dmitry (1), Manevich Alexander (1), Gordeev Evgenii (1), Sorokin Alexey (3), Kramareva Lyubov (4), Uvarov Ivan (2), Kashnitskiy Alexander (2), Bourtsev Mikhail (2)

(1) Institute of Volcanology and Seismology, FEB RAS, Russia, (2) Space Research Institute, RAS, Russia, (3) Computing Center, FEB RAS, Russia, (4) Far East Planeta Center of Space Hydrometeorology Research, Russia

Annually, from 3 to 6 Kamchatkan volcanoes produce eruptions, during which the explosions eject ash to 10-15 km a.s.l., and ash clouds spread thousands of kilometers from volcanoes. Ash clouds pose a serious threat to the modern jet aviation. Scientists of KVERT have conduct daily monitoring of Kamchatka volcanoes since 1993, to mitigate volcanic hazards to airline operations and population. Since 2014, satellite monitoring of volcanoes they carried out with the VolSatView (Remote monitoring of active volcanoes of Kamchatka and the Kuril Islands) (<http://volcanoes.smislab.ru>) IS. The system utilize all the available satellite data, weather and video observations to ensure continues monitoring and study of volcanic activity in Kamchatka. The VolSatView work with distributed information resources and computation systems. This work was supported by the Russian Science Foundation, project No. 16-17-00042.

The VolSatView for Satellite Monitoring and Kamchatkan Volcanoes Study


Girina Olga (1), Loupian Evgenii (2), Efremov Viktor (2), Melnikov Dmitry (1), Manevich Alexander (1), Gordeev Evgenii (1), Sorokin Alexey (3), Kramareva Lyubov (4), Uvarov Ivan (2), Kashnitskiy Alexander (2), Bourtsev Mikhail (2)

(1) Institute of Volcanology and Seismology, FEB RAS, Russia, (2) Space Research Institute, RAS, Russia, (3) Computing Center, FEB RAS, Russia, (4) Far East Planeta Center of Space Hydrometeorology Research, Russia




Annually, from 3 to 6 Kamchatkan volcanoes produce eruptions, during which the explosions eject ash to 10-15 km a.s.l., and ash clouds spread thousands of kilometers from volcanoes. Strenuous volcanic activity could cause ash falls in towns and settlements, destruction of forests and communications. Ash clouds and plumes pose a serious threat to the modern jet aviation. Scientists of KVERT have conduct daily monitoring of Kamchatka volcanoes since 1993, to mitigate volcanic hazards to airline operations and population. Since 2014, satellite monitoring of volcanoes is carried out by KVERT scientists using the VolSatView (<http://volcanoes.smislab.ru>). The VolSatView (Remote monitoring of active volcanoes of Kamchatka and the Kuril Islands) was created in 2011 by scientists from Space Research Institute (SRI) of Russian Academy of Sciences (RAS), Institute of Volcanology and Seismology of Far East Branch (FEB) RAS, Computing Center of FEB RAS and Far East Planeta Center of Space Hydrometeorology Research, and the IS continues to developing. The system utilize all the available satellite data, weather and video observations to ensure continues monitoring and study of volcanic activity in Kamchatka. Architecture of the VolSatView IS was developed for the work with distributed information resources and computation systems that are used for the acquisition, processing, storage, analysis, and visualization of various instrumental and scientific data.

This work was supported by the Russian Science Foundation, project No. 16-17-00042.



Joint analysis of simulation result and satellite data





Analysis of thermal anomalies over the volcanoes




Bezymianny volcano, 2017 eruptions

Analysis of ash plumes from volcanoes




Ash plumes from active volcanoes of Kamchatka in 2015-2016


Hyperspectral analysis (Hyperion)




Hyperspectral analysis of ash plume different parts, Klyuchevskoy volcano, 08.07.2007.





Rock composition of Klyuchevskoy (basalt) and Bezymianny (andesite)



Analysis of volcano eruption deposits



Classification of Zhupanovskoy volcano collapse deposits



KVERT data