

Can. J. Phys. **91**: 98–104 (2013) [dx.doi.org/10.1139/cjp-2012-0330](https://doi.org/10.1139/cjp-2012-0330)

## Bound state solutions of the Manning–Rosen potential

B.J. Falaye, K.J. Oyewumi, T.T. Ibrahim, M.A. Punyasena, and C.A. Onate

Theoretical Physics Section, Department of Physics, University of Ilorin, P.M.B. 1515, Ilorin, Nigeria

### Abstract:

Using the asymptotic iteration method (AIM), we have obtained analytical approximations to the  $\ell$ -wave solutions of the Schrödinger equation with the Manning–Rosen potential. The energy eigenvalues equation and the corresponding wavefunctions have been obtained explicitly. Three different Pekeris-type approximation schemes have been used to deal with the centrifugal term. To show the accuracy of our results, we have calculated the eigenvalues numerically for arbitrary quantum numbers  $n$  and  $\ell$  for some diatomic molecules (HCl, CH, LiH, and CO). It is found that the results are in good agreement with other results found in the literature. A straightforward extension to the s-wave case and Hulthén potential case are also presented.

Can. J. Phys. **91**: 98–104 (2013) [dx.doi.org/10.1139/cjp-2012-0330](https://doi.org/10.1139/cjp-2012-0330)

Published at [www.nrcresearchpress.com/cjp](http://www.nrcresearchpress.com/cjp) on 24 January 2013.