



Functional β -Adrenoceptors Are Important for Early Muscle Regeneration in Mice through Effects on Myoblast Proliferation and Differentiation

Jarrold E. Church^{1‡}, Jennifer Trieu¹, Radhika Sheorey¹, Annabel Y. -M. Chee¹, Timur Naim¹, Dale M. Baum¹, James G. Ryall¹, Paul Gregorevic², Gordon S. Lynch^{1*}

1 Basic and Clinical Myology Laboratory, Department of Physiology, The University of Melbourne, Victoria, Australia, **2** Laboratory for Muscle Research & Therapeutics Development, Baker IDI Heart and Diabetes Institute, Victoria, Australia

Abstract

Muscles can be injured in different ways and the trauma and subsequent loss of function and physical capacity can impact significantly on the lives of patients through physical impairments and compromised quality of life. The relative success of muscle repair after injury will largely determine the extent of functional recovery. Unfortunately, regenerative processes are often slow and incomplete, and so developing novel strategies to enhance muscle regeneration is important. While the capacity to enhance muscle repair by stimulating β_2 -adrenoceptors (β -ARs) using β_2 -AR agonists (β_2 -agonists) has been demonstrated previously, the exact role β -ARs play in regulating the regenerative process remains unclear. To investigate β -AR-mediated signaling in muscle regeneration after myotoxic damage, we examined the regenerative capacity of tibialis anterior and extensor digitorum longus muscles from mice lacking either β_1 -AR (β_1 -KO) and/or β_2 -ARs (β_2 -KO), testing the hypothesis that muscles from mice lacking the β_2 -AR would exhibit impaired functional regeneration after damage compared with muscles from β_1 -KO or β_1/β_2 -AR null (β_1/β_2 -KO) KO mice. At 7 days post-injury, regenerating muscles from β_1/β_2 -KO mice produced less force than those of controls but muscles from β_1 -KO or β_2 -KO mice did not exhibit any delay in functional restoration. Compared with controls, β_1/β_2 -KO mice exhibited an enhanced inflammatory response to injury, which delayed early muscle regeneration, but an enhanced myoblast proliferation later during regeneration ensured a similar functional recovery (to controls) by 14 days post-injury. This apparent redundancy in the β -AR signaling pathway was unexpected and may have important implications for manipulating β -AR signaling to improve the rate, extent and efficacy of muscle regeneration to enhance functional recovery after injury.

Citation: Church JE, Trieu J, Sheorey R, Chee AY-M, Naim T, et al. (2014) Functional β -Adrenoceptors Are Important for Early Muscle Regeneration in Mice through Effects on Myoblast Proliferation and Differentiation. PLoS ONE 9(7): e101379. doi:10.1371/journal.pone.0101379

Editor: Stephen E. Alway, West Virginia University School of Medicine, United States of America

Received: January 28, 2014; **Accepted:** June 5, 2014; **Published:** July 7, 2014

Copyright: © 2014 Church et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the National Health and Medical Research Council (NHMRC) of Australia (Project grants: 509313; 1026231). JEC was supported by a postdoctoral fellowship from the Association Française contre les Myopathies (AFM, France). PG was supported by a NHMRC Career Development Fellowship and formerly a Pfizer Senior Research Fellowship. JGR is supported by a NHMRC Overseas Postdoctoral Fellowship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist. PG was the recipient of a Senior Research Fellowship sponsored by Pfizer Australia. This award was created to help emerging scientists develop a program of independent, non-commercial research at an Australian university or non-profit research institute. Sponsorship of this award by Pfizer Australia provided PG with salary support and funds to cover establishment costs. Pfizer did not participate in or contribute to the research undertaken by PG, and did not review the findings arising from this research prior to submission for publication. The financial sponsorship of this independent research fellowship by Pfizer Australia does not constitute a competing interest for the authors of this work, and does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* Email: gsl@unimelb.edu.au

‡ Current address: Department of Human Biosciences, Faculty of Health Sciences, La Trobe University, Bundoora, Australia

Introduction

Skeletal muscle is susceptible to damage associated with trauma, metabolic deficits, heritable and acquired diseases, and mechanical strains generated during contractions [1–4]. Although skeletal muscle has an impressive ability to regenerate after injury, the repair process is often slow and incomplete, compromised by factors such as age and pre-existing health conditions [5]. To develop treatment strategies that enhance muscle regeneration after injury and thereby improve quality of life for patients, a more comprehensive understanding of the events taking place during the regeneration of skeletal muscle is required [6].

Many of the most common and important growth factors and hormones can influence the attributes of mature and regenerating

skeletal muscles. For instance, we and others have shown that stimulating β -adrenoceptors (β -ARs) with β -AR agonists (β -agonists) can enhance skeletal muscle repair after injury [6–9]. Recent studies have also shown that $G_{\alpha_{i2}}$, a component of the β -AR signalling pathway, plays an important role in regeneration after injury and general maintenance of muscle mass [10–12]. However, while the capacity to enhance muscle repair via exogenous stimulation of β -ARs is now recognised, it remains unclear as to the role β -ARs play in regulating the regenerative process. To elucidate the significance of β -AR-mediated signalling in regeneration, we examined the regenerative capacity of limb muscles from mice lacking either β_1 -AR and/or the β_2 -AR. On the basis that exogenous administration of β_2 -agonists can promote regeneration, it was hypothesised that muscles from mice

lacking the β_2 -AR (β_2 -KO) would exhibit impaired functional regeneration after damage compared with muscles from β_1 -KO or β_1/β_2 -KO mice.

Materials and Methods

Animals

All procedures were approved by the Animal Ethics Committee of The University of Melbourne and conformed to the Australian code of practice for the care and use of animals for scientific purposes as stipulated by the National Health and Medical Research Council of Australia. Mice lacking both the β_1 - and β_2 -AR (β_1/β_2 -KO) were purchased from The Jackson Laboratory ($Adrb1_{tm1Bkk}Adrb2_{tm1Bkk/j}$, stock #003810) and maintained by mating double homozygote KO mice. Control mice were from a C57BL/6 background, as employed previously for comparison with β_1/β_2 -KO [13,14] and obtained from the Animal Resources Centre (Canning Vale, WA, Australia). Mice lacking either the β_1 -AR or the β_2 -AR were purchased from The Jackson Laboratory and back crossed with C57BL/6 mice to create a heterozygous F1 generation. The heterozygous offspring of the respective strains were subsequently inbred to produce the respective β_1 -KO and β_2 -KO knockout animals and littermate controls (β_1 -WT and β_2 -WT, respectively). Mice were housed in the Biological Research Facility at The University of Melbourne under a 12 h light-dark cycle, with drinking water and standard chow provided *ad libitum*. Male mice aged 8–9 weeks were used in this study.

Myotoxic injury

Injury was induced in the tibialis anterior (TA) or extensor digitorum (EDL) muscles of mice as described previously [15,16]. For TA injuries, mice were anesthetized using a mixture of ketamine (76 mg/kg, i.p.) and xylazine (10 mg/kg, i.p.) and a small portion of the TA muscle of the right hindlimb was surgically exposed by a single incision through the skin. The muscle was filled to its maximal holding capacity ($\sim 40 \mu\text{l}$) via a single intramuscular injection with Notexin (1 $\mu\text{g}/\text{ml}$, Latoxan) using a 29-gauge needle. The wound was closed with Michel clips (Aesculap, Tuttlingen, Germany) and the mice were allowed to recover from the myotoxic injury for 7, 10 or 14 days before assessments of muscle structure and function were performed. Uninjured mice were used as controls for all experiments using the TA muscle. For EDL injuries, mice were anesthetized as described and the EDL muscle of the right hindlimb was surgically exposed by a single incision through the skin and overlying muscles. With the aid of a dissecting microscope, the EDL was filled to its maximal holding capacity (15–20 μl) via intramuscular injection at several sites with Notexin (1 $\mu\text{g}/\text{ml}$, Latoxan) using a 30-gauge needle. The wound was closed with Michel clips (Aesculap, Tuttlingen, Germany) and the mice were allowed to recover from the myotoxic injury for 2, 5, 7, 14 or 21 days. The corresponding muscle from the left (contralateral) limb served as the non-injured control muscle.

Assessment of muscle function

Mice were anesthetized with sodium pentobarbital (60 mg/kg, i.p.) and muscle function was assessed either *in vitro* (for EDL) or *in situ* (for TA) as described previously [17,18]. For assessment of TA muscle force producing capacity *in situ*, the muscles of anesthetised mice were stimulated by supramaximal 0.2 ms square wave pulses of 350 ms duration, delivered via two wire electrodes placed adjacent to the deep peroneal branch of the sciatic nerve. Force generated was measured via attachment of a force transducer to the distal tendon of the muscle. For assessments of EDL muscle

function *in vitro* the muscles from each limb were surgically excised from anesthetized mice and transferred to a custom-built organ bath filled with Krebs-Ringer solution (in mM: NaCl, 1.37; NaHCO_3 , 24; D-glucose, 11; KCl, 5; CaCl_2 , 2; NaH_2PO_4 , 1; MgSO_4 , 0.487, pH 7.4) supplemented with d-tubocurarine chloride (0.293 mM), bubbled with Carbogen (5% CO_2 in O_2) and maintained at 25°C, where the proximal tendon was attached to a micro-manipulator (to facilitate adjustment of muscle length to optimal), and the distal tendon was attached to a force transducer. The EDL was stimulated by supramaximal (40 V) 0.2 ms square wave pulses of 350 ms duration delivered via platinum plate electrodes that flanked the full length of each muscle.

For both *in situ* and *in vitro* assessments, optimal muscle length for contraction (L_o) was defined as the muscle length at which maximal isometric twitch force (P_i) was attained and maximal isometric tetanic force (P_o) was determined from the plateau of the frequency-force relationship. P_o was normalized to the cross-sectional area of the muscle (CSA; calculated using L_o and muscle mass) in order to determine maximum specific isometric tetanic force (sP_o), a measurement that allows for relative comparisons of force production between muscles of different sizes.

Immediately following the functional assessments, muscles were carefully trimmed of any adherent non-muscle tissue and tendons, and weighed on an analytical balance. Muscles were then mounted in Tissue-Tek OCT embedding medium, frozen rapidly in thawing isopentane, and stored at -80°C for later histological and biochemical analyses. The mice were killed by surgical excision of the heart while still anesthetized deeply.

Muscle morphology

Serial transverse cryosections (5 μm) were cut from the midbelly of each muscle and placed onto glass slides (Superfrost Plus, Menzel-Gläser, Kensington, VIC, Australia). General muscle histology was determined by staining sections with hematoxylin and eosin (H & E) to visualize muscle fibers, and digital images of stained sections were obtained using a microscope equipped with digital camera (Carl Zeiss, Wrek, Göttingen, Germany) supported by associated imaging software (Axiovision V4.7.1.0).

Quantitative RT-PCR

Muscle samples were homogenized individually and mRNA extracted using an RNEasy fibrous tissue RNA extraction kit (Qiagen), according to manufacturer's instructions. The concentration and quality of RNA in each sample was determined using a Nanodrop 2000 (Thermo Scientific) and the extracted mRNA was stored at -80°C . mRNA was transcribed into cDNA using the Superscript VILO cDNA synthesis kit (Invitrogen) according to manufacturer's instructions, and stored at -20°C until use. Quantitative RT-PCR was performed using an iCycler Thermal Cycler (Bio-Rad) with SYBR Green supermix (Quantace). Primers were designed using the Perfect Primer online program (Invitrogen) and are listed in Table 1. Due to the severity of the myotoxic damage caused by Notexin at the earlier time points [15], we chose not to use the ΔCt method of analysis as we could not be certain that any reference gene used would remain unchanged between treatment groups. We instead opted to measure the cDNA concentration of each sample using the Quant-iT OliGreen ssDNA Assay Kit (Molecular Probes) and to normalize our data to cDNA content as described previously [19,20].

Cell culture

Primary mouse myoblasts were isolated using a protocol adapted from Huang and colleagues [21]. Briefly, mice were

Table 1. PCR primer sequences.

	Genbank accession number		Primer Sequences (5'-3')
Myf5	NM_008656	F	5'- AACCAGAGACTCCCAAGGT-3'
		R	5'- AGCTGGACACGGAGCTTTTA-3'
MyoD	NM_010866	F	5'- AGTGAATGAGGCCTTCGAGA-3'
		R	5'- GCATCTGAGTCGCCACTGTA-3'
Myogenin	NM_031189	F	5'- CACTCCCTTACGTCCATCGT-3'
		R	5'- CAGGACAGCCCCACTTAAAA-3'
MRF4	NM_008657	F	5'- GGCTGGATCAGCAAGAGAAG-3'
		R	5'- AAGAAAGCGCTGAAGACTG-3'
TNF-α	NM_013693	F	5'- GGCTTCTACCTTCAGACC-3'
		R	5'- AGCAAAGAGGAGGCAACAA-3'
IL-6	NM_031168	F	5'- CCGAGAGGAGACTTCACAG-3'
		R	5'- TCCACGATTCCAGAGAACC-3'
CD68	NM_009853	F	5'- TCCAAGCCCAAATTCAAATC-3'
		R	5'- ATTGTATTCCACCGCCATGT-3'
F4/80	NM_010130	F	5'- CATCAGCCATGTGGGTACAG-3'
		R	5'- CATCACTGCCTCCACTAGCA-3'

doi:10.1371/journal.pone.0101379.t001

anesthetized and killed by cardiac excision and the quadriceps, TA and gastrocnemius muscles were excised, cleaned of all connective tissue, and rinsed in ice-cold culture media (Ham's Nutrient Mix F-12). The muscles were placed in a digest solution consisting of culture media supplemented with 1% Type II collagenase and 0.5% Dispase (Invitrogen), minced with a clean pair of scissors, and incubated in a shaking incubator at 37°C for 90 min. The suspension was filtered through a 100 μm cell filter (BD Biosciences), pelleted (1500×g, 5 min) and resuspended in complete growth media (Ham's Nutrient Mix F-12 supplemented with 20% FBS, 1% L-glutamine, 1% antibiotic-antimycotic and 5 ng/ml rhFGF). The cell suspension was plated into uncoated cell culture dishes overnight to allow fibroblasts to adhere, after which the non-adherent cells were aspirated and plated into a fresh uncoated dish for a further 24 hrs to remove as many fibroblasts as possible. Following the second overnight incubation, non-adherent cells were aspirated and seeded into a cell culture dish pre-coated with extracellular matrix (ECM; Sigma). Cells were expanded in culture for approximately 4 weeks, after which myoblast proliferation and differentiation were assessed.

For proliferation assays, isolated primary myoblasts were seeded into ECM-coated 96 well plates at a density of 1×10⁴ cells/well and incubated in complete growth media for 72 hrs. Subsequently, the cells were fixed (3.7% formaldehyde, 10 min, 25°C), rinsed twice with PBS, and incubated with DAPI (0.235 μM diluted in PBS, 15 min, 25°C). The number of cell nuclei was quantified using an inverted fluorescence microscope and camera (Axiovert 40, Carl Zeiss) and associated imaging software (Axiovision V4.7.1.0, Carl Zeiss).

For differentiation assays, the isolated primary myoblasts were seeded into ECM-coated 12-well plates at a density of 5×10⁵ cells/well. After incubating for 2 hr to permit cell attachment, growth media was replaced with differentiation media (DMEM supplemented with 2% horse serum and 1% L-glutamine). Cells were cultured in differentiation-inducing conditions for 3 days, with differentiation media replaced daily. To examine attributes of differentiation, cells were imaged with an inverted microscope and camera (Axiovision 40, Carl Zeiss) controlled by associated

software (Axiovision V4.7.1.0). Differentiation was measured as total area (μm²) in the field of view occupied by multinucleated myotubes.

Statistical analyses

All values are reported as mean ±SEM. Groups were compared using an unpaired Student's *t*-test, or a one-way or two-way ANOVA and Bonferroni's post hoc multiple comparison procedure where appropriate. Myofiber cross-sectional area (CSA) is not normally distributed, and so medians were compared using a non-parametric Mann-Whitney test by ranks. All statistical analyses were performed using Prism version 3 software (GraphPad Software, Inc. La Jolla, CA). In all cases significance was defined as P<0.05.

Results

Deletion of specific β-ARs has differential effects on muscle mass and force producing capacity

We first compared the morphology and functional characteristics of TA muscles from the different β-AR KO strains of mice. We found no significant differences in body mass between any of the β-KO mice and their respective controls (Fig. 1A). When we compared the muscle mass between the strains, neither β₁-KO mice nor β₂-KO mice were different from their respective controls, but TA muscles of β₁/β₂-KO mice were significantly smaller (Fig. 1B). When maximal force was expressed as absolute force (P₀), β₁-KO mice did not exhibit differences in force production compared with controls, but both β₂-KO mice and β₁/β₂-KO mice produced less force than their controls (Fig. 1C). When maximal force production was normalized to muscle size (sP₀), β₁-KO mice still exhibited no differences in force production and β₂-KO mice still exhibited deficits in force production, but β₁/β₂-KO mice demonstrated equivalent functional capacity compared with control (Fig. 1D). Examining muscle fiber cross-sectional area (CSA) for the various β-KO mouse strains revealed that, as for TA muscle mass, neither the muscles of β₁-KO mice

nor β_2 -KO mice were different from the muscles of their respective littermate controls, but that muscle fiber CSA in β_1/β_2 -KO mice was significantly smaller than muscle fibres from wild type mice (Fig. 1E). We also analyzed a number of twitch characteristics of TA muscles from the β -KO mice (Table 2) and found that the muscles of β_1 -KO mice did not differ from control mice, but the muscles of both β_2 -KO mice and β_1/β_2 -KO mice elicited reduced peak twitch force (P_1) and a reduced rate of contraction (dP_1/dt) compared with the muscles of controls. An analysis of the frequency-force relationship found no significant differences between β_1 -KO mice and control, but significant differences between β_2 -KO mice and control, and β_1/β_2 -KO mice and control (Figure S2).

Deletion of β_2 -ARs impairs the functional performance of regenerating muscles

We next examined the recovery of force production in the TA muscles of the various β -KO mice at 7, 10 and 14 days after myotoxic damage. When force production was expressed as absolute force (P_o), we found that β_1 -KO mice displayed no differences in maximal force production during regeneration, whereas both β_2 -KO mice and β_1/β_2 -KO mice produced significantly less force compared with controls (Figure S1). When force was normalized to muscle size (sP_o) maximal force production was unchanged in β_1 -KO mice from controls (Fig. 2A). Maximal force in β_2 -KO mice was significantly lower than controls throughout regeneration (Fig. 2C), but force production by β_1/β_2 -KO mice was no longer significantly different from control over the 14 days of regeneration (Fig. 2E). In order to take into account the lower sP_o of uninjured muscles from β_2 -KO mice, we then expressed force production during regeneration as a percentage of that produced by uninjured muscles (%UI). When maximal force was expressed this way, we found that none of the β -KO strains exhibited any deficits in the restoration of force producing capacity compared with controls (Fig. 2B, 2D and 2F). When the data at 7 days post-injury were analyzed separately in order to assess the earliest time point in isolation, β_1/β_2 -KO mice had impaired force production compared with controls whether expressed as P_o , sP_o or %UI.

In order to more comprehensively assess the role of β -ARs in skeletal muscle we also compared morphological characteristics of EDL muscles between β_1/β_2 -KO and C57BL/6 control mice. Our findings for the EDL muscles closely mirrored those for the TA muscle; viz., muscle fiber CSA was not significantly different between β_1/β_2 -KO mice and C57BL/6 controls (Fig. 3A) and uninjured muscles from β_1/β_2 -KO mice were significantly smaller than controls (Fig. 3B) and produced lower maximal absolute forces (Fig. 3c and Table 3). Similarly, when maximal force was normalized to muscle cross-sectional area (sP_o) there were no significant differences between β_1/β_2 -KO mice and controls (Fig. 3D and Table 3). When twitch characteristics of uninjured EDL muscles were compared between the two strains, we found that as for the TA muscles, both twitch force (P_1) and rate of contraction (dP_1/dt) were significantly lower in the β_1/β_2 -KO mice (Table 3). Following myotoxic damage, EDL muscles from β_1/β_2 -KO mice exhibited impaired restoration of force production over 21 days, with a particularly significant deficit at 7 days post-injury (Fig. 3C and 3D). In addition to the delayed restoration of contractile force in recovery EDL muscles, qualitative differences in muscle architecture were observed at both 7 and 14 days post-injury (Fig. 3E).

Expression of inflammatory markers and myogenic regulatory factors during early regeneration is potentiated in muscles lacking β_1 - and β_2 -ARs

As we had detected impaired force production in β_1/β_2 -KO mice during the early stages of regeneration (at 7 days post-injury), we examined the expression of both inflammatory markers and myogenic regulatory factors in TA muscles from β_1/β_2 -KO mice and C57BL/6 control mice immediately after myotoxic injury and during early regeneration. Quantitative RT-PCR for the cytokines interleukin-6 (IL-6) and tumour necrosis factor- α (TNF- α) revealed an increase in cytokine expression in C57BL/6 mice after myotoxic damage, which peaked at 2 days post-injury (Fig. 4C and 4D). Injury-induced expression of TNF- α was significantly higher in β_1/β_2 -KO mice than in C57BL/6 controls (Fig. 4D; 2-way ANOVA, main strain effect), and expression of IL-6 showed a non-significant trend toward higher expression in β_1/β_2 -KO mice (Fig. 4C). Expression of the macrophage markers CD68 and F4/80 revealed an increase in macrophage infiltration in injured muscles from both β_1/β_2 -KO and C57BL/6 mice which peaked at 5 days post-injury (Fig. 4A & B). Injury-induced expression of CD68 and F4/80 was significantly higher in β_1/β_2 -KO mice than in uninjured controls (2-way ANOVA, main strain effect).

An examination of myogenic regulatory factor (MRF) mRNA revealed that Myf5, MyoD and myogenin expression all increased after myotoxic damage, peaking at 5 days post-injury (Fig. 5A, B and C). Expression of all three MRFs was significantly higher in β_1/β_2 -KO mice (2-way ANOVA, main strain effect), and also seemed to be more sustained than the increases in control muscles, with MyoD and myogenin in particular remaining elevated for longer than 5 days post-injury (Fig. 5A, B and 5C).

Deletion of β_1 - and β_2 -ARs increases myoblast proliferation and inhibits differentiation ex vivo

To directly determine the proliferation and differentiation of myoblasts from β_1/β_2 -KO mice without the confounding variables inherent to the whole-body knockout mouse, we isolated and cultured myoblasts from β_1/β_2 -KO mice and C57BL/6 controls. We found that myoblasts from β_1/β_2 -KO mice proliferated more rapidly and differentiated less effectively than those from C57BL/6 controls (Fig. 6).

Discussion

Although exogenous stimulation of β -ARs has been shown to enhance mammalian muscle fiber regeneration [6–9], and recent studies have also implicated components of the β -AR signalling pathway in muscle regeneration and hypertrophy [10–12], the contribution of β -ARs to the muscle regenerative process *per se* is less well understood. The findings from the present study have addressed this gap in existing knowledge by examining the functional capacity of skeletal muscles from β_1 -, β_2 - and β_1/β_2 -KO mice. While many previous studies have examined β -KO mice in a cardiovascular context, the attributes of skeletal muscle are less well described [22]. We found that the mass of TA muscles from either β_1 -KO mice or β_2 -KO mice were similar to respective controls, whereas the TA muscles from β_1/β_2 -KO mice were significantly smaller than control. Muscle fiber CSA in the various strains closely mirrored muscle mass, with neither of the single knockout strains displaying significant differences in CSA while the double knockouts had a significant reduction. Hinkle and colleagues [22] also found no difference in TA mass between β_1 -KO mice and control, but noted that both β_2 -KO mice and β_1/β_2 -KO mice exhibited significantly reduced TA muscle mass than controls, from which they concluded that muscle size was




Figure 1. Basic morphology and function of the TA muscles of different β -KO mouse strains. (A) Body masses for the different β -KO mouse strains (white bars) and their respective controls (black bars). No significant differences were observed across the strains. (B) Comparison of TA muscle mass in β -KO mouse strains (white bars) and their respective controls (black bars). There was no significant difference in TA mass for either β_1 -KO or β_2 -KO mice when compared with littermate controls, but β_1/β_2 -KO mice had a significantly lower TA mass than control ($*P < 0.05$ compared with control, *t*-test, $n = 5-6$ per group). (C) and (D) maximal tetanic force production by TA muscles measured as either absolute force (P_o) or force normalized to muscle size (specific force, or sP_o). β_1 -KO mice had no significant differences in maximal force whether expressed as either P_o or sP_o . In contrast, β_2 -KO mice had significant deficits in maximal force when expressed as either P_o or sP_o . β_1/β_2 -KO mice had significantly lower maximal forces than control when expressed as P_o , but there was no significant deficit in force production when force was normalized to muscle size (sP_o) ($*P < 0.05$ compared with control, *t*-test, $n = 5-6$ per group). (E) Comparison of cross-sectional area (CSA) of TA muscle fibers from β -KO mouse strains (white bars) and their respective controls (grey bars). Neither β_1 -KO nor β_2 -KO mice had significant changes in muscle fiber CSA compared with their wild-type littermates, but fiber CSA was significantly lower in TA muscles from β_1/β_2 -KO mice compared with control. Data for each strain were pooled from >1500 fibers from $n \geq 9$ mice ($*P < 0.05$, Mann-Whitney test by ranks).
doi:10.1371/journal.pone.0101379.g001

positively regulated by β_2 -ARs but not β_1 -ARs. In the present study, we did not observe any significant difference in TA muscle mass in β_2 -KO mice, but did note a significant reduction in TA

mass in β_1/β_2 -KO mice. The reason for the discrepancy between the studies is unclear but may reflect the back-crossing onto a different background strain (C57BL/6) in the present study.

Table 2. Selected isometric twitch contractile properties of uninjured TA muscles from β-KO mice.

	β ₁ -WT	β ₁ -KO	β ₂ -WT	β ₂ -KO	C57BL/6	β ₁ /β ₂ -KO
<i>n</i>	5	5	5	5	6	5
P _t (mN)	472±20	502±38	513.8±30.9	425.6±21.6*	457±17	369±27*
TPT (ms)	17.3±0.8	16.7±0.8	15.5±0.5	16.6±1.3	14.4±0.6	15.6±0.4
½ RT (ms)	17.4±2.0	16.4±1.1	14.5±1.1	13.8±1.9	12.2±1.0	16.1±1.6
dP _t /dt (mN/ms)	67.8±2.8	70.4±1.7	79.7±2.9	67.13±3.7*	73.8±1.4	63.2±4.7*

P_t - peak twitch tension; TPT - time to peak twitch tension; ½ RT - one-half relaxation time; dP_t/dt - maximum rate of force development during a twitch contraction. *P<0.05 vs. vehicle control, Student's unpaired *t*-test. doi:10.1371/journal.pone.0101379.t002




Figure 2. Maximal tetanic force production by regenerating TA muscles of different strains of β-KO mice after injury and measured as either specific force (sP_o; a, c, e) or specific force normalized to uninjured muscles (%UI; b, d, f). β₁-KO mice did not exhibit significant deficits in maximal force during regeneration (A and B). When maximal force was expressed as sP_o, β₂-KO mice had force deficits during all stages of regeneration (C; #P<0.05, strain main effect, 2-way ANOVA), although a Bonferroni post-hoc test did not detect any differences at specific time points. When force production was normalised to uninjured muscles (%UI), there was no difference between β₂-KO mice and controls during regeneration (D). β₁/β₂-KO mice did not exhibit delayed regeneration over the 14 day period when analysed with a 2-way ANOVA, but there was a significant deficit in maximal force by β₁/β₂-KO mice (expressed as either sP_o or %UI) at 7 days post-injury when the time point was analysed in isolation (*P<0.05 compared with control, *t*-test, *n* = 7) (E and F). doi:10.1371/journal.pone.0101379.g002




Figure 3. Morphological properties and functional regeneration after myotoxic damage in the EDL muscles of β_1/β_2 -KO mice and C57BL/6 controls. (A) Comparison of muscle fiber CSA in uninjured EDL muscles. There was no significant difference in muscle fiber CSA between β_1/β_2 -KO mice (white bar) when compared with C57BL/6 (black bar). Data were pooled from >1500 fibers from $n=5$ mice (B). Comparison of EDL muscle mass. β_1/β_2 -KO mice had lower mass than C57BL/6 controls ($*P<0.05$ compared with control, t -test, $n=29$ and 21 , respectively). (C) and (D) Maximal force by regenerating EDL muscles expressed as either absolute force (P_o) or force normalized to muscle size (specific force, or sP_o). β_1/β_2 -KO mice (white bars) had lower maximal forces when expressed as either P_o or sP_o ($*P<0.05$, strain main effect, 2-way ANOVA), with significant deficits at 7 days post-injury ($*P<0.05$, Bonferroni's post hoc multiple comparison procedure). (E) H&E stained sections of fibres from regenerating EDL muscles of C57BL/6 or β_1/β_2 -KO mice. Note the qualitative differences in regeneration at 7 and 14 days post-injury. doi:10.1371/journal.pone.0101379.g003

The muscles of mice lacking β_1 -ARs were identical to controls for every functional parameter measured, but muscles from both β_2 -KO mice and β_1/β_2 -KO mice produced less absolute force than controls (whether twitch force or tetanic force) and exhibited a slower rate of force generation. The only difference we observed in functionality between the muscles of β_2 -KO mice and β_1/β_2 -

KO mice was that of specific force (sP_o). When expressed this way muscles from β_1/β_2 -KO mice produced similar forces to control, whereas force production by β_2 -KO mice remained significantly depressed relative to control muscles. Taken together, our data indicate that the loss of β_2 -ARs has profound effects on skeletal

Table 3. Selected isometric twitch contractile properties of uninjured EDL muscles from β₁/β₂-KO mice and C57BL/6 controls.

	C57BL/6	β ₁ /β ₂ -KO
<i>n</i>	26	23
P _o	327.1±6.5	237.4±26.0*
sP _o	244.6±3.9	245.0±7.5
P _t (mN)	83.2±3.1	65.4±3.4*
TPT (ms)	20.9±0.4	21.5±0.5
½ RT (ms)	26.4±0.7	27.9±1.1
dP _t /dt (mN/ms)	13.4±0.6	11.3±0.5*

P_t - peak twitch tension; TPT - time to peak twitch tension; ½ RT - one-half relaxation time; dP_t/dt - maximum rate of force development during a twitch contraction. *P<0.05 vs. vehicle control, Student's unpaired t-test. doi:10.1371/journal.pone.0101379.t003

muscle morphology and contractile function, but the loss of β₁-ARs does not significantly impact on either parameter. This is perhaps not surprising given that β₁-ARs make up only about 7–10% of the total β-ARs in skeletal muscle [23]. It is worth noting however that the expression of β₁-ARs is relatively higher in slow-twitch muscles such as the soleus, and that the present study examined the TA and EDL that are both predominantly fast-twitch muscles. We therefore cannot exclude the possibility that the loss of β₁-ARs may have an impact on contractile function in slow twitch muscles such as the soleus. These findings also confirm

previous studies demonstrating that β₂-AR signaling has significant effects on force producing capacity [24,25].

In the context of muscle regeneration, we found that the muscles from β₁-KO mice did not show any significant differences in force production after myotoxic injury compared with controls. In contrast, muscles from β₂-KO mice displayed significant deficits in force production during regeneration when measured as either absolute force or specific force. When we accounted for the fact that the muscles of β₂-KO mice were returning to a lower absolute force (i.e. we expressed force production as ‘% uninjured’), we found no significant differences in the restoration of force




Figure 4. mRNA expression of (A) CD68, (B) F4/80, (C) IL-6 and (D) TNF-α in regenerating TA muscles of C57BL/6 and β₁/β₂-KO mice. β₁/β₂-KO mice had higher expression of all four markers of inflammation and macrophage infiltration during regeneration (n=3; #P<0.05, strain main effect, 2-way ANOVA; *P<0.05, Bonferroni's post hoc multiple comparison procedure). doi:10.1371/journal.pone.0101379.g004




Figure 5. mRNA expression of (A) Myf5, (B) MyoD and (C) myogenin in regenerating TA muscles of C57BL/6 and β₁/β₂-KO mice. β₁/β₂-KO mice had higher expression of all three MRFs during regeneration (*n* = 3; #*P* < 0.05, strain main effect, 2-way ANOVA; **P* < 0.05, Bonferroni's post hoc multiple comparison procedure). doi:10.1371/journal.pone.0101379.g005

production in regenerating muscles between β₂-KO mice and controls. Similarly, muscles from β₁/β₂-KO (double knockout) mice exhibited significant differences in absolute force production compared with controls, but when force was normalized to either muscle cross-sectional area (sP₀) or to force production by

uninjured muscles (% uninjured) there was no difference in the restoration of force over the 14 day period of regeneration. If the data are normalized to reflect the intrinsic differences in muscle strength and the entire 14 day period of regeneration is analysed in its entirety, then a lack of β-ARs (either β₁-ARs, β₂-ARs or both β₁- and β₂-ARs) does not appear to affect the restoration of muscle function after myotoxic damage. If, however, we confine our analysis to the earliest time point of the regeneration period (i.e. 7 days post-injury), a different pattern emerges. In this case, early regeneration of the TA muscle in both β₁-KO mice and β₂-KO is unaffected compared with controls, whereas the muscles of β₁/β₂-KO mice exhibit significant force deficits. Our results suggest a level of redundancy between the two subtypes of β-AR. This finding may be confounded by these mice having lifelong knock-out of β-ARs, which may promote possible compensatory adaptations in β-AR signaling that accompany such models. In addition, it seems unlikely that β₁-ARs, which comprise only 7–10% of β-ARs within muscle [23] could compensate for the loss of β₂-ARs. Nonetheless, the possibility of β-AR redundancy requires further investigation, possibly by other experimental models such as inducible gene knock-out, or in vivo knockdown of β-ARs with siRNA, that would be less likely to be confounded by compensatory alterations in signaling.

To confirm our findings in the TA muscle, we also examined the characteristics of the EDL muscles. Since our findings from the TA muscle had indicated only β₁/β₂-KO mice exhibited differences in regeneration after myotoxic injury, we confined our analysis of EDL morphology and function to β₁/β₂-KO mice and C57BL/6 controls. Uninjured EDL muscles from β₁/β₂-KO mice exhibited similar differences in morphology and function to TA muscles; i.e. decreased muscle mass, decreased twitch force and rate of contraction, and decreased absolute tetanic force when compared with C57BL/6 controls. Furthermore, EDL muscles from β₁/β₂-KO mice displayed similar deficits in regeneration at 7 days post-injury, confirming our findings in the TA muscle and further suggesting that the loss of both β₁- and β₂-ARs affects the earliest period of regeneration after myotoxic damage.

Skeletal muscle regeneration is preceded by a well-defined and highly coordinated inflammatory response, involving the infiltration of neutrophils and macrophages at the site of damage and subsequent release of pro- and anti-inflammatory cytokines [26]. Non-cytokine anti-inflammatory mediators are also able to modulate the inflammatory process, including glucocorticoids, adenosine, and endogenous β-agonists [27]. β-ARs are not only present in skeletal muscle [23], but have also been detected on the surface of inflammatory cells such as mast cells, eosinophils, neutrophils and macrophages [28]. To determine the degree of macrophage infiltration in regenerating muscles, we examined the expression of the macrophage-specific markers F4/80 and CD68 [29]. The expression of both macrophage markers was significantly increased in β₁/β₂-KO than in controls. Previous studies have revealed that macrophages incubated with LPS have an increased expression of TNF-α and IL-6, which is inhibited by the β-agonist clenbuterol [30,31] and that administering the β-AR antagonist propranolol can potentiate the release of inflammatory cytokines in vivo [32,33]. Due to the well-characterized increase in TNF-α and IL-6 in response to inflammation, as well as the evidence that their expression is controlled to some degree by β-AR signaling, we examined these two cytokines as a measure of inflammation and found that both cytokines were increased acutely after injury, with maximal expression at 2 days post-injury in control mice. This correlates well with previous studies from our laboratory where we have reported that oedema and immune cell infiltration are maximal at this time [15]. In the present study,




Figure 6. (A) Proliferation of primary myoblasts isolated from either C57BL/6 (black bars) or β_1/β_2 -KO (white bars) mice. Myoblasts from β_1/β_2 -KO mice proliferated more rapidly than those from C57BL/6 controls ($*P < 0.05$ compared with control, *t*-test, $n = 6$). (B) and (C) Differentiation of primary myoblasts isolated from either C57BL/6 (black bars) or β_1/β_2 -KO (white bars) mice. Myoblasts from β_1/β_2 -KO mice had significantly impaired differentiation compared with those from C57BL/6 controls ($*P < 0.05$ compared with control, *t*-test, $n = 6$). doi:10.1371/journal.pone.0101379.g006

TNF- α was significantly more elevated in β_1/β_2 -KO mice, and IL-6 mRNA showed a non-significant trend toward higher expression than in regenerating muscles from control mice, suggesting that the acute immune response to injury in β_1/β_2 -KO mice is exacerbated by the lack of β -ARs. Taken together, these data suggest that the inflammatory response and subsequent macrophage infiltration after injury is acutely higher in the β_1/β_2 -KO mice compared with control.

To ascertain whether the observed force deficits in muscles from β_1/β_2 -KO mice at 7 days post-injury were associated with impaired myofiber regeneration, we assessed the expression of the myogenic regulatory factors (MRFs) responsible for muscle formation. Previous studies have demonstrated that clenbuterol represses the expression of MyoD and myogenin in denervated rat soleus muscles [34], and that clenbuterol administration increased myogenin expression in immobilized rat plantaris muscles [35]. Clenbuterol administration also increased MyoD expression in rat soleus muscles [36]. In the present study, Myf5 (the transcription factor responsible for specification of satellite cells to the myogenic lineage) was upregulated after myotoxin injection, with the increase in expression peaking at 5 days post-injury. Furthermore, expression of Myf5 was significantly increased in the regenerating muscles of β_1/β_2 -KO mice than controls, indicating that myoblast proliferation was not compromised in the β_1/β_2 -KO mice, and may be propagated for longer after muscle injury, than in control mice. The induction of MyoD (which initiates the terminal differentiation program in myoblasts) and myogenin (which is involved in the activation of muscle-specific genes in the immature muscle cells [37]) following injury was also exaggerated in the muscles of β_1/β_2 double-KO mice. Taken together, our MRF expression data suggest that myoblast proliferation and differentiation may be enhanced in β_1/β_2 -KO mice at the expense of moderately delayed differentiation. This observation, consistent

with what we had expected and described in a previous review [38], may explain why force producing capacity is impaired at 7 days post-injury in β_1/β_2 -KO mice, but that muscles are capable of restoring functionality (relative to uninjured muscles) similar to control animals at 10 days post-injury. This rapid ‘catch up’ where the muscles of β_1/β_2 -KO mice seemingly overcome their initial delayed regeneration and function impairment, is supported by our observations of cultured primary myoblasts from β_1/β_2 -KO mice, where proliferation was enhanced and prolonged and differentiation was delayed.

The present study utilized whole body β_1/β_2 -KO mice, since to our knowledge there are no muscle-specific β_1/β_2 -KO mice currently available. One concern with using the whole body β_1/β_2 -KO mouse (a concern that is also valid in any *in vivo* study of β -agonist administration) is that any effects on muscle regeneration may be a consequence of perturbations of non-muscle physiological systems, rather than a direct effect on muscle regeneration *per se*. For example, the altered inflammatory response observed in the present study, while not a direct result of the muscle lacking β -ARs, undoubtedly influenced fiber regeneration.

To obviate these concerns we isolated myoblasts from both β_1/β_2 -KO mice and C57BL/6 controls to examine myoblast proliferation and differentiation in the absence of confounding factors, and found that myoblasts isolated from β_1/β_2 -KO mice proliferated more rapidly and differentiated far less effectively than those from C57BL/6 controls. While initially this may seem to be at odds with our MRF expression data from regenerating muscles, it must be remembered that even if myoblast differentiation was impaired *in vivo*, the vastly greater number of myoblasts present in the muscle due to the increased proliferation would still result in an overall increase in MyoD and myogenin expression in the muscle.

Interestingly, we have previously documented a dramatic increase in the gene expression of *adrb1* and *adrb2* (β_1 - and β_2 -adrenoceptors, respectively) during the switch from proliferation to differentiation [38]. Combined with the findings of the present study, these data support a role for β_1/β_2 -ARs in inhibiting myoblast proliferation and promoting differentiation.

Another concern with whole body β_1/β_2 -KO mice is the potential for cardiovascular disturbances (particularly alterations in local blood flow) to influence muscle regeneration. We do not believe that the muscles from β_1/β_2 -KO mice suffered a significant deficit in perfusion as this would have resulted in a constant inhibition of regeneration, whereas we observed a deficit in regeneration only at 7 days post-injury. In fact, the muscles from β_1/β_2 -KO mice subsequently regenerated faster than control in order to ‘catch up’ to the control muscles at the later time points.

With all our data taken together we propose a model in which β_1/β_2 -KO mice exhibit an enhanced inflammatory response to injury and enhanced myoblast proliferation during regeneration. We believe that β_1/β_2 -KO mice have delayed early regeneration due to the prolonged and enhanced inflammatory response, but once the inflammation subsides, the enhanced myoblast proliferation allows the β_1/β_2 -KO muscles to regenerate rapidly – thus accounting for the rapid ‘catch up’ of muscles between 7 and 10 days post-injury. Further studies using muscle specific knockdown of β -ARs or virally-mediated ‘knock in’ of β -ARs to inflammatory cells of β_1/β_2 -KO mice, or other injury models (such as crush or burn injuries) with a higher inflammatory component would help test this hypothesis.

In summary, our findings indicate that β -ARs play an important role in early muscle regeneration, at least in part via a direct effect

on myoblast proliferation and differentiation. Manipulation of β -AR signaling during these early stages of regeneration may therefore improve the rate, extent and efficacy of the regenerative process, to enhance functional recovery after injury.

Supporting Information

Figure S1 Maximal tetanic force of regenerating TA muscles of different strains of β -KO mice following injury and expressed as absolute force (P_0). Force production by β_1 -KO mice did not differ from controls during regeneration (A), but both β_2 -KO mice (B) and β_1/β_2 -KO mice (C) had significant force deficits during all stages of regeneration ($\#P<0.05$, strain main effect, 2-way ANOVA). (TIF)

Figure S2 Frequency-force relationships for different β -KO mouse strains. β_1 -KO mice did not have altered frequency-force relationship compared with controls (A and B), but β_2 -KO (C and D) and β_1/β_2 -KO mice (E and F) produced significantly lower forces at all frequencies when compared with controls. ($\#P<0.05$, strain main effect, 2-way ANOVA). (TIF)

Author Contributions

Conceived and designed the experiments: JEC JGR PG GSL. Performed the experiments: JEC JT RS AC DB TN. Analyzed the data: JEC RS JGR PG GSL. Contributed reagents/materials/analysis tools: PG GSL. Wrote the paper: JEC JGR PG GSL.

References

- Brooks SV, Faulkner JA (1988) Contractile properties of skeletal muscles from young, adult and aged mice. *J Physiol* 404: 71–82.
- Bunn JR, Canning J, Burke G, Mushipe M, Marsh DR, et al. (2004) Production of consistent crush lesions in murine quadriceps muscle – a biomechanical, histomorphological and immunohistochemical study. *J Orthop Res* 22: 1336–1344.
- Lynch GS, Ryall JG (2008) Role of β -adrenoceptor signaling in skeletal muscle: implications for muscle wasting and disease. *Physiol Rev* 88: 729–767.
- Lynch GS, Schertzer JD, Ryall JG (2007) Therapeutic approaches for muscle wasting disorders. *Pharmacol Ther* 113: 461–487.
- Gregorevic P, Plant DR, Stupka N, Lynch GS (2004) Changes in contractile activation characteristics of rat fast and slow skeletal muscle fibres during regeneration. *J Physiol* 558: 549–560.
- Ryall JG, Schertzer JD, Alabakis TM, Gehrig SM, Plant DR, et al. (2008) Intramuscular β_2 -agonist administration enhances early regeneration and functional repair in rat skeletal muscle after myotoxic injury. *J Appl Physiol* 105: 165–172.
- Beitzel F, Gregorevic P, Ryall JG, Plant DR, Sillence MN, et al. (2004) β_2 -adrenoceptor agonist fenoterol enhances functional repair of regenerating rat skeletal muscle after injury. *J Appl Physiol* 96: 1385–1392.
- Beitzel F, Sillence MN, Lynch GS (2007) β -Adrenoceptor signaling in regenerating skeletal muscle after β -agonist administration. *Am J Physiol Endocrinol Metab* 293: E932–940.
- Emery PW, Rothwell NJ, Stock MJ, Winter PD (1984) Chronic effects of β_2 -adrenergic agonists on body composition and protein synthesis in the rat. *Biosci Rep* 4: 83–91.
- Egerman MA, Glass DJ (2014) Signaling pathways controlling skeletal muscle mass. *Crit Rev Biochem Mol Biol* 49: 59–68.
- Minetti GC, Feige JN, Bombard F, Heier A, Morvan F, et al. (2014) *G α i2* signaling is required for skeletal muscle growth, regeneration, and satellite cell proliferation and differentiation. *Mol Cell Biol* 34: 619–630.
- Minetti GC, Feige JN, Rosentiel A, Bombard F, Meier V, et al. (2011) *G α i2* signaling promotes skeletal muscle hypertrophy, myoblast differentiation, and muscle regeneration. *Sci Signal* 4: ra80.
- Chruscinski AJ, Rohrer DK, Schauble E, Desai KH, Bernstein D, et al. (1999) Targeted disruption of the β_2 adrenergic receptor gene. *J Biol Chem* 274: 16694–16700.
- Kiriazis H, Wang K, Xu Q, Gao XM, Ming Z, et al. (2008) Knockout of β_1 - and β_2 -adrenoceptors attenuates pressure overload-induced cardiac hypertrophy and fibrosis. *Br J Pharmacol* 153: 684–692.
- Plant DR, Colarossi FE, Lynch GS (2006) Notexin causes greater myotoxic damage and slower functional repair in mouse skeletal muscles than bupivacaine. *Muscle Nerve* 34: 577–585.
- Schertzer JD, Lynch GS (2006) Comparative evaluation of IGF-I gene transfer and IGF-I protein administration for enhancing skeletal muscle regeneration after injury. *Gene Ther* 13: 1657–1664.
- Harcourt LJ, Holmes AG, Gregorevic P, Schertzer JD, Stupka N, et al. (2005) Interleukin-15 administration improves diaphragm muscle pathology and function in dystrophic *mdx* mice. *Am J Pathol* 166: 1131–1141.
- Gehrig SM, Koopman R, Naim T, Tjoarkarfa C, Lynch GS (2010) Making fast-twitch dystrophic muscles bigger protects them from contraction injury and attenuates the dystrophic pathology. *Am J Pathol* 176: 29–33.
- Lundby C, Nordborg N, Kusuvara K, Kristensen KM, Neuffer PD, et al. (2005) Gene expression in human skeletal muscle: alternative normalization method and effect of repeated biopsies. *Eur J Appl Physiol* 95: 351–360.
- Murphy KT, Koopman R, Naim T, Leger B, Trieu J, et al. (2010) Antibody-directed myostatin inhibition in 21-month-old mice reveals novel roles for myostatin signaling in skeletal muscle structure and function. *FASEB J* 24: 4433–4442.
- Huang YC, Dennis RG, Larkin L, Baar K (2005) Rapid formation of functional muscle in vitro using fibrin gels. *J Appl Physiol* 98: 706–713.
- Hinkle RT, Hodge KM, Cody DB, Sheldon RJ, Kobilka BK, et al. (2002) Skeletal muscle hypertrophy and anti-atrophy effects of clenbuterol are mediated by the β_2 -adrenergic receptor. *Muscle Nerve* 25: 729–734.
- Kim YS, Sainz RD, Molenaar P, Summers RJ (1991) Characterization of β_1 - and β_2 -adrenoceptors in rat skeletal muscles. *Biochem Pharmacol* 42: 1783–1789.
- Cairns SP, Dulhunty AF (1993) The effects of β -adrenoceptor activation on contraction in isolated fast- and slow-twitch skeletal muscle fibres of the rat. *Br J Pharmacol* 110: 1133–1141.
- Ha TN, Posterino GS, Fryer MW (1999) Effects of terbutaline on force and intracellular calcium in slow-twitch skeletal muscle fibres of the rat. *Br J Pharmacol* 126: 1717–1724.
- Tidball JG (1995) Inflammatory cell response to acute muscle injury. *Med Sci Sports Exerc* 27: 1022–1032.
- Van Miert AS (2002) Present concepts on the inflammatory modulators with special reference to cytokines. *Vet Res Comm* 26: 111–126.
- Barnes PJ (1999) Effect of β -agonists on inflammatory cells. *J Allergy Clin Immunol* 104: S10–17.
- Hong EG, Ko HJ, Cho YR, Kim HJ, Ma Z, et al. (2009) Interleukin-10 prevents diet-induced insulin resistance by attenuating macrophage and cytokine response in skeletal muscle. *Diabetes* 58: 2525–2535.

30. Izeboud CA, Mocking JA, Monshouwer M, van Miert AS, Witkamp RF (1999) Participation of β -adrenergic receptors on macrophages in modulation of LPS-induced cytokine release. *J Recept Signal Transduct Res* 19: 191–202.
31. Izeboud CA, Monshouwer M, van Miert AS, Witkamp RF (1999) The β -adrenoceptor agonist clenbuterol is a potent inhibitor of the LPS-induced production of TNF-alpha and IL-6 in vitro and in vivo. *Inflamm Res* 48: 497–502.
32. Boost KA, Flondor M, Hofstetter C, Platacis I, Stegwerth K, et al. (2007) The β -adrenoceptor antagonist propranolol counteracts anti-inflammatory effects of isoflurane in rat endotoxemia. *Acta Anaesthesiol Scand* 51: 900–908.
33. Lang CH, Nystrom G, Frost RA (2008) β -adrenergic blockade exacerbates sepsis-induced changes in tumor necrosis factor alpha and interleukin-6 in skeletal muscle and is associated with impaired translation initiation. *J Trauma* 64: 477–486.
34. Maltin CA, Delday MI, Campbell GP, Hesketh JE (1993) Clenbuterol mimics effects of innervation on myogenic regulatory factor expression. *Am J Physiol* 265: E176–178.
35. Delday MI, Maltin CA (1997) Clenbuterol increases the expression of myogenin but not myoD in immobilized rat muscles. *Am J Physiol* 272: E941–944.
36. Jones SW, Baker DJ, Gardiner SM, Bennett T, Timmons JA, et al. (2004) The effect of the β_2 -adrenoceptor agonist prodrug BRL-47672 on cardiovascular function, skeletal muscle myosin heavy chain, and MyoD expression in the rat. *J Pharmacol Exp Ther* 311: 1225–1231.
37. Le Grand F, Rudnicki MA (2007) Skeletal muscle satellite cells and adult myogenesis. *Curr Opin Cell Biol* 19: 628–633.
38. Ryall JG, Church JE, Lynch GS (2010) Novel role for β -adrenergic signalling in skeletal muscle growth, development and regeneration. *Clin Exp Pharmacol Physiol* 37: 397–401.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Church, JE; Trieu, J; Sheorey, R; Chee, AY-M; Naim, T; Baum, DM; Ryall, JG; Gregorevic, P; Lynch, GS

Title:

Functional beta-Adrenoceptors Are Important for Early Muscle Regeneration in Mice through Effects on Myoblast Proliferation and Differentiation

Date:

2014-07-07

Citation:

Church, J. E., Trieu, J., Sheorey, R., Chee, A. Y. -M., Naim, T., Baum, D. M., Ryall, J. G., Gregorevic, P. & Lynch, G. S. (2014). Functional beta-Adrenoceptors Are Important for Early Muscle Regeneration in Mice through Effects on Myoblast Proliferation and Differentiation. PLOS ONE, 9 (7), <https://doi.org/10.1371/journal.pone.0101379>.

Persistent Link:

<http://hdl.handle.net/11343/190643>

File Description:

Published version

License:

CC BY