

Title Page

Fluid Resuscitation with 6% Hydroxyethyl Starch (130/0.4 and 130/0.42) in Acutely Ill Patients: Systematic Review of Effects on Mortality and Treatment With Renal Replacement Therapy

David J Gattas^{a,b}, Arina Dan^a, John Myburgh^{a,c}, Laurent Billot^d, Serigne Lo^d, Simon Finfer^{a,e}, CHEST Management Committee**

**Crystalloid vs Hydroxyethyl Starch Trial (CHEST) Management Committee: John Myburgh (chair), Rinaldo Bellomo, Alan Cass, Maryam Correa, Simon Finfer, David Gattas, Parisa Glass, Julie Harland, Joanne Lee, Jeffrey Lipman, Bette Liu, Colin McArthur, Shay McGuinness, Maryanne Ng, Dorrilyn Rajbhandari, Colman Taylor, Steve Webb and Lynsey Willenberg

^a Division of Critical Care & Trauma, The George Institute for Global Health, Sydney, Australia. David J Gattas, Fellow; Arina Dan, Fellow; John Myburgh, Professor; Simon Finfer, Professor.

^b Royal Prince Alfred Hospital, University of Sydney, Sydney, Australia.

^c St George Hospital, University of New South Wales, Sydney, Australia.

^d The George Institute for Global Health, Sydney 2000. Laurent Billot, statistician; Serigne Lo, statistician.

^e Royal North Shore Hospital, University of Sydney, Sydney, Australia.

Correspondence to: David J Gattas, Royal Prince Alfred Hospital, Missenden Rd, Camperdown, NSW 2050, Australia. dgattas@med.usyd.edu.au

Sources of Funding

This study was funded by the Division of Critical Care & Trauma, George Institute for Global Health. JM is supported by a Practitioner Fellowship from the National Health and Medical Research Council.

Abstract

Purpose. To determine whether fluid resuscitation of acutely ill adults with 6% hydroxyethyl starch with a molecular weight of 130 kD and a molar substitution ratio of approximately 0.4 (6% HES130) compared with other resuscitation fluids results in a difference in the relative risk of death or treatment with renal replacement therapy (RRT). **Methods.** Systematic review and meta-analysis of randomized controlled trials comparing intravascular fluids for resuscitation of hospitalised adults that reported mortality or treatment with RRT. The risk of bias was assessed independently by two reviewers and meta-analysis was performed using random effects. **Results.** Thirty five trials enrolling 10391 participants were included. The three largest trials had the lowest risk of bias, were published (or completed) in 2012, and together enrolled 77% of all participants. Death occurred in 928 of 4691 patients (19.8%) in the 6% HES130 group vs 871 of 4720 (18.5%) in the control fluid groups (relative risk (RR) in the 6% HES130 group 1.08, 95% confidence interval (CI) 1.00 to 1.17, $I^2=0\%$). Treatment with RRT occurred in 378 of 4236 patients (8.9%) in the 6% HES130 group vs 306 of 4260 (7.2%) in the control fluid group (RR in the 6% HES130 group 1.25, 95% CI 1.08 to 1.44, $I^2=0\%$). **Conclusions.** The quality and quantity of data evaluating 6% hydroxyethyl starch (130/0.4 and 130/0.42) as a resuscitation fluid has increased in the last 12 months. Patients randomly assigned to resuscitation with 6% HES130 are at significantly increased risk of being treated with RRT.

Keywords [MeSH]

hetastarch, colloids, fluid therapy, resuscitation, critical illness

Introduction

Administration of fluid to increase or maintain intravascular volume (resuscitation fluid) is a common intervention in the Intensive Care Unit (ICU). A cross sectional international study reported that over one third of patients in ICUs receive resuscitation fluid each day. In that study colloids were administered to more patients and during more resuscitation episodes than crystalloids, with hydroxyethyl starch solutions being the most frequently administered colloid solutions[1]. Previous meta-analyses have not suggested that colloids in general offer significant advantages over crystalloids[2]. Recent guidelines from the European Society of Intensive Care Medicine taskforce on colloid volume therapy in critically ill patients recommended against the use of 6% HES130 in patients with severe sepsis or at risk of acute kidney injury[3]. The strength of these recommendations may be limited as previous meta-analyses have relied on the results of trials that were generally poor in quality and which reported few patient-centred outcomes[2-9].

Over the preceding twelve months, a number of randomized controlled trials report the effects of 6% HES130 in critically ill patients. As these additional data have the potential to substantially alter the evidence for and against the use of 6% HES130 in critically ill patients, we updated a systematic review and meta-analysis incorporating all the available evidence to determine whether fluid resuscitation of acutely ill adults with 6% HES130 compared to other resuscitation fluids resulted in a difference in patient-centred outcomes.

The aim of this updated review was to determine if there was a difference in the risk of death and treatment with RRT in acutely ill adult patients receiving 6% HES130 for fluid resuscitation compared with other resuscitation fluids.

Methods

Eligibility criteria and assessment for risk of bias[4]. The protocol for the systematic review, including the inclusion and exclusion criteria, was written before the literature search was conducted. Eligible studies were included if all criteria were met: (1) prospective, randomized controlled trials, (2) patients over 18 years, (3) a hospital or pre-hospital clinical setting, (4) patients who were acutely ill or undergoing major surgery, (5) study fluids were administered for resuscitation (defined as fluid required to increase or maintain intravascular volume), (6) at least one intervention group received 6% hydroxyethyl starch with a molecular weight of 130 kD and a molar substitution ratio of approximately 0.4 in any carrier solution, (7) at least one intervention group received another colloid or any type of crystalloid solution for resuscitation, (8) the study reported at least one of the following five outcomes: (i) mortality, (ii) treatment with RRT, (iii) urine output, (iv) transfusion of red blood cells (RBCs), (v) estimated or measured blood loss.

Studies were excluded if any of the following characteristics were present: (1) studies enrolling only healthy volunteers or blood donors, (2) administration of fluid solely for the purposes of a planned anesthetic procedure including spinal or epidural anesthesia, acute normovolemic hemodilution, hypervolemic hemodilution or priming of a cardiopulmonary bypass circuit without subsequent intra- or post-operative use, (3) administration of fluid solely for volume therapy (hemodilution) following ischemic stroke or subarachnoid hemorrhage.

Internal validity was evaluated using a tool based on “yes/no” responses to the following five domains of trial quality: randomization, allocation concealment, blinding, intention-to-treat analysis, and minimal (<10%) loss to follow-up[10]. Low risk of bias was defined as scoring ‘yes’ to all 5 domains. Intermediate risk of bias was defined as scoring ‘yes’ to 4 out of 5 domains. High risk of bias was defined as scoring ‘yes’ to 3 or less out of 5 domains. For randomization, use of term ‘randomization’ in any form without a clear description of sequence generation was deemed pseudo-randomization and judged ‘no’ to randomization. Allocation concealment was considered to have been ‘yes’ if any method for doing so was described. Blinding was assessed in three areas (patient, clinicians, and outcome assessors) and all three elements had to be blinded in order for the trial to be considered blinded overall. Blinding of the fluid alone without any further description was considered to be ‘no’ to blinding. Intention-to-treat analysis was interpreted strictly, with any patients removed from analysis after randomization and receipt of intervention considered to be ‘no’ for that trial. Loss to follow-up of less than 10% for the primary outcome was considered acceptable.

Search strategy. Five electronic databases were searched on 28 July 2012: Ovid MEDLINE, EMBASE, the Cochrane Central Register of Controlled Trials (CENTRAL), the metaRegister of Controlled Trials (controlled-trials.com), and clinicaltrials.gov. In addition, the reference lists from other published systematic reviews were hand searched for any additional studies that met inclusion criteria. No language restriction was placed on the search. Contact was made with experts in the field for any unpublished trials. The search terms used in MEDLINE, EMBASE and CENTRAL are contained in the appendix.

Study selection and data extraction. Two reviewers (DG, AD) screened the results of the search independently. Full-text manuscripts of potentially eligible articles were obtained and assessed independently against inclusion and exclusion criteria. The same two authors independently extracted the data and appraised the internal validity of each study. Differences were then compared and resolved by agreement or referral to a third reviewer (JM). The variables pertaining to patients and setting were: total number of patients, number of participating centres, clinical setting and diagnostic group. We collected details regarding mean daily volume of 6% HES130.

For the clinical setting, we categorized the data into 3 groups: peri-operative (defined as fluid used intra-operatively and post-operatively), operative (fluid used intra-operatively only), and ICU (patients admitted to an ICU at the time of enrolment for a reason that was not associated with routine post-operative care). For volume of fluid administered, we categorized the data into 3 groups according to the mean daily volume of exposure to 6% HES130 throughout the study period: <1 litre, 1-3 litres, or >3 litres. These groupings were chosen to approximate low (<15ml/kg), medium (15-40ml/kg), and higher (>40ml/kg) dose exposure to 6% HES130 for a typical 70kg individual. All-cause mortality was collected preferentially if reported. When mortality was reported at more than one time point, we used the longest complete follow up time after exposure to study fluids.

Data analysis. The relative risk (RR) and 95% confidence intervals (CI) of death and treatment with RRT for 6% HES130 compared to control group fluid were calculated for each study and then pooled via a meta-analysis with random effects using the metan routine in Stata version 11. I^2 was calculated as a measure of consistency. Trials with no deaths were excluded from the pooled estimate of relative risk. Studies with zero deaths in one group were added to the pooled estimate by adding 0.5 to each cell of the 2-by-2 table[11]. For studies with more than one control group, a single control comparison was selected with preference given to a crystalloid control group, then another class of colloid, and finally another hydroxyethyl starch with a molecular weight greater than 130kD as comparator. If there was more than one crystalloid control group, these were pooled in order to make a single comparison between 6% HES130 and crystalloid. The remaining control groups were not included in the pooled mortality or RRT analyses.

Ethical approval was not required.

Results

The process for screening and assessing reports is shown in Figure 1. The search yielded 3984 potential reports of which 35 trials that recruited a total of 10391 participants were eligible for the systematic review; 25 of the 35 studies reported mortality and 11 of 35 reported treatment with RRT. The characteristics of the 35 studies are summarised in Table 1.

List of included studies. Myburgh 2012[12], Perner 2012[13], Siegemund 2012[14], Gondos 2010[15], Guidet 2012[16], Zhu 2011[17], James 2011[18], Yang 2011[19], Nagpal 2012[20], Mittermayr 2007[21], Schramko 2010[22], Lu 2012[23], Volta 2007[24], Dubin 2010[25], Van der Linden 2005[26], Ooi 2009[27], Zdosek 2011[28], Wu 2010[29], Godet 2008[30], Mahmood 2007[31], Dolecek 2009[32], Schramko 2009[33], Mukhtar

2009[34], Inal 2010[35], Palumbo 2006[36], Kasper 2003[37], Gandhi 2003[38], Langeron 2001[39], Sander 2003[40], Gallandat Huet 2000[41], Jungheinrich 2004[42], Mehta 2007[43], Ellger 2006[44], Neff 2003[45], Boldt 2000[46]

The quality of included reports is detailed in Table 2. Only three studies met the predefined criteria for having a low risk of bias, all of these studies have been published or completed in the preceding twelve months[12-14]. 21 of 35 studies were judged to have a high risk of bias, including 7 of 10 studies comparing 6% HES130 to a preparation of hydroxyethyl starch with a molecular weight greater than 130kD. Studies comparing 6% HES130 to at least one crystalloid control group were generally of higher quality (8 of 14 scored low or intermediate risk of bias).

The event rates of the 25 studies that reported mortality and the results of the meta-analysis are shown in Figure 2. In the random effects analysis, the Crystalloid vs Hydroxyethyl Starch Trial (CHEST)[12] and Scandinavian Starch for Severe Sepsis/Septic Shock (6S)[13] studies contribute most to the pooled estimate (combined 85.7% weight). Studies comparing 6% HES130 to another class of colloid or hydroxyethyl starch with a molecular weight greater than 130kD contributed very little to the mortality findings (1.5% and 0.5% weight respectively). 928 of 4691 patients randomly assigned to receive 6% HES130 (19.8%) died compared with 871 of 4720 (18.5%) of those assigned to receive other fluids, relative risk 1.08 (95% confidence interval 1.00-1.17). There was no significant heterogeneity ($I^2=0%$, $p=0.7$).

The event rates of the 11 studies that reported treatment with RRT and the results of the meta-analysis are shown in Figure 3. In the random effects analysis, the Crystalloid vs Hydroxyethyl Starch Trial (CHEST)[11] and Scandinavian Starch for Severe Sepsis/Septic Shock (6S)[12] studies contribute most to the pooled estimate (combined 84.6% weight). Studies comparing 6% HES130 to another class of colloid or hydroxyethyl starch with a molecular weight greater than 130kD contributed very little to the RRT findings (0.9% and 0.7% weight respectively). 378 of 4236 patients randomly assigned to receive 6% HES130 (8.9%) were treated with RRT compared with 306 of 4260 (7.2%) of those assigned to receive other fluids, relative risk 1.25 (95% confidence interval 1.08-1.44). There was no significant heterogeneity ($I^2=0%$, $p=0.8$).

Urine output data were reported in 21 of 35 studies. Time periods of collection were highly variable, or not reported with sufficient detail to enable valid data extraction. Meta-analysis of urine output was deemed inappropriate and not performed. Similar findings were made in trials reporting transfusion (27 of 35 studies) and bleeding (23 of 35 studies). Transfusion was reported using semi-quantitative units of measurement such as 'units transfused' in some studies. Methods used to collect and report bleeding data were usually not stated.

Discussion

The principal finding of this systematic review is that fluid resuscitation with 6% HES130 as compared to other fluids is associated with an 8% increase in the relative risk of death which is of borderline statistical significance, and a significant 25% increase in the relative risk of being treated with RRT compared to other resuscitation fluids. There was no significant heterogeneity among the included trials, and the magnitude and direction of these associations were similar across the recent trials with larger sample size and lower risk of bias.

The strength of this review is that it includes recent large-scale trials that have focussed on mortality and treatment with RRT. The methods we used were the same as for a previous review[4], which resulted from the retraction of clinical trials of 6%HES130[47-51] during the enrolment period of the CHEST trial[12]. The two reviews demonstrate the increase in the number of patients recently randomized into clinical trials evaluating 6%HES130. The limitations of this review are that we did not extract data for other outcomes, and did not contact authors to try and obtain additional unpublished data. The duration of study follow up was not analysed. The results of this review are predominated by two large-scale trials[12, 13] which may limit the applicability of these results in other patients populations and treatment settings.

Several systematic reviews that have been published concerning colloids for resuscitation[2, 5-8]. Some have focussed on 6%HES130 specifically[8], and others have evaluated all hydroxyethyl starch preparations[6] or colloids in general[2, 5]. The risks associated with exposure to the newer formulations of hydroxyethyl starch are now consistent across several large-scale trials with a low risk of bias. It is unlikely that any clinical benefits of using 6%HES130 not studied in this review would outweigh these risks, at least in the patient populations enrolled in these trials.

Further research may identify which patient subgroups are at greater risk of harm from exposure to 6%HES130, in particular individual patient data meta-analysis of existing trials might identify patient characteristics conferring increased risk. We would recommend the use of other fluids until it is understood if there are any patients who are likely to receive a net benefit when they are fluid resuscitated with 6%HES130.

Conclusion. Fluid resuscitation of acutely ill adults with 6%HES130 is associated with an increase in risk of death and treatment with RRT. These associations are consistent across recent large-scale randomised controlled trials with a low risk of bias.

Table and Figure Legends

Figure 1. Study flow diagram

Table 1. **Characteristics of included studies.** ICU = intensive care unit. ns = not stated. RRT = renal replacement therapy. 6%HES130 = 6% hydroxyethyl starch with a molecular weight of 130 kD and a molar substitution ratio of approximately 0.4. Daily Mean 6%HES130 is the mean daily dose, categorised into litre ranges. HES=other (higher molecular weight) forms of hydroxyethyl starch given as control fluids, further defined by xxx/y.y, describing the molecular weight xxx (kDa) and molar substitution y.y. Transfusion = transfusion of red blood cells reported using any measure. Bleeding = estimated or measured blood loss reported using any measure.

Table 2. **Assessment of quality and risk of bias.** Trials were scored 'yes' or 'no' by two reviewers in 5 domains of quality: randomization, allocation concealment, blinding, intention-to-treat analysis, and minimal loss to follow-up. Low risk of bias was defined as scoring 'yes' to all 5 domains. Intermediate risk of bias was defined as scoring 'yes' to 4 out of 5 domains. High risk of bias was defined as scoring 'yes' to 3 or less out of 5 domains.

Figure 2. **Forest plot of pooled estimates for mortality.** 6%HES130=6% hydroxyethyl starch with a molecular weight of 130 kD and a molar substitution ratio of approximately 0.4. CI=confidence interval. Studies reporting at least one event in each group are arranged in ascending year of publication. Weights are from random effects analysis.

Figure 3. **Forest plot of pooled estimates for need for renal replacement therapy.** 6%HES130=6% hydroxyethyl starch with a molecular weight of 130 kD and a molar substitution ratio of approximately 0.4. CI=confidence interval. Studies reporting at least one event in each group are arranged in ascending year of publication. Weights are from random effects analysis.

Author	Year	N	Number of centres	Population	Diagnostic Group	Daily Mean 6%HES130 (range, litres)	Number of control groups	Control Fluid Class	Control Fluid(s)	Reports mortality	Reports treatment with RRT	Reports urine output	Reports tranfusion	Reports bleeding	
<i>Compares 6% HES (130/0.4 or 130/0.42) to at least 1 crystalloid control group</i>															
Myburgh	2012	7000	32	ICU	general ICU population	< 1	1	crystalloid	normal saline	Yes	Yes	Yes	Yes	No	
Perner	2012	804	26	ICU	severe sepsis	1-3	1	crystalloid	Ringer's acetate	Yes	Yes	Yes	Yes	Yes	
Siegemund	2012	241	1	ICU	sepsis	ns	1	crystalloid	normal saline	Yes	Yes	ns	ns	No	
Gondos	2010	200	11	ICU	Postoperative, hypovolaemia	ns	3	2 colloids, crystalloid	4% gelatin, 5% albumin, Ringer's lactate	Yes	No	No	No	Yes	
Guidet	2012	196	24	ICU	severe sepsis	1-3	1	crystalloid	normal saline	Yes	Yes	Yes	Yes	Yes	
Zhu	2011	135	1	ICU	severe sepsis	< 1	2	crystalloid, colloid	7.5% saline, Ringer's lactate	Yes	No	Yes	No	No	
James	2011	115	1	ICU	trauma	> 3	1	crystalloid	normal saline	Yes	Yes	Yes	Yes	No	
Yang	2011	90	1	Perioperative	liver surgery	1-3	2	colloid, crystalloid	20% albumin, Ringer's lactate	Yes	No	Yes	Yes	Yes	
Nagpal	2012	70	1	Operative	cardiac surgery	1-3	1	crystalloid	normal saline	Yes	Yes	Yes	Yes	Yes	
Mittermayr	2007	66	1	Perioperative	orthopaedic surgery	1-3	2	colloid, crystalloid	4% gelatin, Ringer's lactate	No	No	No	Yes	Yes	
Schramko	2010	45	1	Perioperative	cardiac surgery	ns	2	colloid, crystalloid	4% gelatin, Ringer's acetate	No	No	Yes	Yes	Yes	
Lu	2012	42	1	ICU	sepsis	1-3	1	crystalloid	Ringer's lactate	Yes	No	Yes	No	No	
Volta	2007	36	1	Operative	abdominal surgery	1-3	2	colloid, crystalloid	3.4% polygeline, Ringer's lactate	No	No	Yes	Yes	No	
Dubin	2010	25	2	ICU	septic shock	1-3	1	crystalloid	normal saline	Yes	No	Yes	Yes	No	
<i>Compares 6% HES (130/0.4 or 130/0.42) to at least 1 other class of colloid</i>															
Van der Linden	2005	132	1	Perioperative	cardiac surgery	ns	1	colloid	3% gelatin	Yes	No	No	Yes	Yes	
Ooi	2009	90	1	Perioperative	cardiac surgery	1-3	1	colloid	4% gelatin	Yes	Yes	No	Yes	Yes	
Zdolsek	2011	84	1	Operative	orthopaedic surgery	1-3	3	HES, colloid	130/0.42/6:1, 200/0.5, Dextran 70	No	No	No	Yes	Yes	
Wu	2010	80	1	Operative	kidney transplant	1-3	1	colloid	4% gelatin	No	No	Yes	No	No	
Godet	2008	65	7	Perioperative	vascular surgery, renal impairment	1-3	1	colloid	3% gelatin	Yes	Yes	Yes	Yes	No	
Mahmood	2007	62	1	Perioperative	vascular surgery	> 3	2	HES, colloid	200/0.62, 4% gelatin	Yes	Yes	Yes	Yes	Yes	
Dolecek	2009	56	1	ICU	severe sepsis	1-3	1	colloid	20% albumin	Yes	No	No	No	No	
Schramko	2009	45	1	Perioperative	cardiac surgery	1-3	2	HES, colloid	200/0.5, 4% albumin	No	No	Yes	Yes	Yes	
Mukhtar	2009	40	1	Perioperative	liver transplant surgery	> 3	1	colloid	5% albumin	Yes	Yes	Yes	Yes	Yes	
Inal	2010	30	1	ICU	hypovolaemia	< 1	1	colloid	3.5% gelatin	Yes	No	No	No	No	
Palumbo	2006	20	1	ICU	sepsis	ns	1	colloid	20% albumin	No	No	Yes	Yes	No	
<i>Compares 6% HES (130/0.4 or 130/0.42) to other hydroxyethyl starch</i>															
Kasper	2003	120	1	Perioperative	cardiac surgery	> 3	1	HES	200/0.5	Yes	Yes	No	Yes	Yes	
Gandhi	2007	100	6	Operative	orthopaedic surgery	1-3	1	HES	670/0.75	Yes	No	No	Yes	Yes	
Langeron	2001	100	4	Perioperative	orthopaedic surgery	1-3	1	HES	200/0.5	Yes	No	No	Yes	Yes	
Sander	2003	60	1	Perioperative	gynaecological surgery	1-3	1	HES	200/0.5	Yes	No	No	Yes	Yes	
Gallandat Huet	2000	59	2	Perioperative	cardiac surgery	1-3	1	HES	200/0.5	Yes	No	Yes	Yes	Yes	
Jungheinrich	2004	52	1	Perioperative	orthopaedic surgery	1-3	1	HES	200/0.5	No	No	Yes	Yes	Yes	
Mehta	2007	40	1	Perioperative	cardiac surgery	ns	1	HES	200/0.5	No	No	No	No	Yes	
Ellger	2006	40	1	Perioperative	urological surgery	ns	1	HES	200/0.5	No	No	No	Yes	Yes	
Neff	2003	31	1	ICU	traumatic brain injury	> 3	1	HES	200/0.5	Yes	No	Yes	Yes	Yes	
Boldt	2000	20	1	Operative	cardiac surgery	< 1	1	HES	200/0.5	Yes	No	Yes	Yes	Yes	
		35		Number of included trials					Number of trials reporting		25	11	21	27	23
			9	Number of multicentre trials					Number of events reported		1799	684			
		10391		Total number of participants in 36 Included trials					Number of cases reported		9411	8496			
									Crude rate		19.1%	8.1%			

Author	Year	Randomisation	Allocation concealment	Blinding	Intention to treat analysis	No Loss to follow-up
<i>Low risk of bias</i>						
Myburgh	2012	Yes	Yes	Yes	Yes	Yes
Perner	2012	Yes	Yes	Yes	Yes	Yes
Siegemund	2012	Yes	Yes	Yes	Yes	Yes
<i>Intermediate risk of bias</i>						
Guidet	2012	Yes	Yes	Yes	Yes	No
Nagpal	2012	Yes	Yes	Yes	Yes	No
James	2011	Yes	Yes	Yes	No	Yes
Gondos	2010	Yes	Yes	No	Yes	Yes
Schramko	2010	Yes	Yes	No	Yes	Yes
Mukhtar	2009	Yes	Yes	No	Yes	Yes
Schramko	2009	Yes	Yes	No	Yes	Yes
Gandhi	2007	Yes	Yes	No	Yes	Yes
Mahmood	2007	Yes	Yes	No	Yes	Yes
Neff	2003	Yes	Yes	No	Yes	Yes
Boldt	2000	Yes	Yes	No	Yes	Yes
<i>High risk of bias</i>						
Lu	2012	Yes	No	No	Yes	Yes
Zdolsek	2011	Yes	Yes	No	No	Yes
Wu	2010	Yes	Yes	No	No	Yes
Dolecek	2009	Yes	No	No	Yes	Yes
Godet	2008	Yes	Yes	No	No	Yes
Volta	2007	Yes	No	No	Yes	Yes
Ellger	2006	Yes	Yes	No	Yes	No
Van der Linden	2005	Yes	No	No	Yes	Yes
Kasper	2003	Yes	Yes	No	No	Yes
Zhu	2011	No	No	No	Yes	Yes
Dubin	2010	Yes	Yes	No	No	No
Inal	2010	No	No	No	Yes	Yes
Ooi	2009	No	No	No	Yes	Yes
Mittermayr	2007	Yes	Yes	No	No	No
Palumbo	2006	Yes	No	No	No	Yes
Jungheinrich	2004	Yes	Yes	No	No	No
Sander	2003	Yes	No	No	No	Yes
Langeron	2001	No	No	No	Yes	Yes
Gallandat Huet	2000	No	No	No	Yes	Yes
Yang	2011	Yes	No	No	No	No
Mehta	2007	Yes	No	No	No	No

Appendix

Electronic Search strategy. The intersection of: fluid resuscitation, hydroxyethyl starch, and randomized controlled trials

MEDLINE

1. exp Fluid Therapy
2. ((fluid\$ or volume\$ or plasma\$ or rehydrat\$) adj3 (replace\$ or therap\$ or substitut\$ or restor\$ or resuscitat\$ or rehydrat\$)).ab,ti.
3. or/1-2
4. exp Starch
5. exp Blood Substitutes
6. exp Colloids
7. hetastarch\$.tw.
8. hydroxyethyl starch.tw.
9. hydroxyethylstarch.tw.
10. hydroxy ethyl starch.tw.
11. pentastarch.tw.
12. voluven\$.tw.
13. tetrastarch.tw.
14. or/4-13
15. 3 and 14
16. limit 15 to "therapy (sensitivity)" [from the MEDLINE limit 'Clinical Queries', based on Haynes et al[52]]

EMBASE

- #14. #3 AND #12 AND #13
- #13. random:ti OR 'clinical trial':de,m,ab,ti OR 'health care quality'/exp
- #12. #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11
- #11. tetrastarch
- #10. voluven*
- #9. pentastarch
- #8. 'hydroxy ethyl starch'
- #7. hydroxyethylstarch
- #6. 'hydroxyethyl starch'
- #5. hetastarch*
- #4. 'starch'/exp OR starch
- #3. #1 OR #2
- #2. (fluid* OR volum* OR plasma* OR rehydrat*) NEAR/3 (therap* OR substitut* OR restor* OR resusc* OR replac*)
- #1. 'fluid therapy'/exp OR 'fluid therapy'

CENTRAL

- #1 starch* or *starch or voluven* in Clinical Trials
- #2 MeSH descriptor Fluid Therapy explode all trees
- #3 ((fluid* or volume* or plasma* or rehydrat*) NEAR/3 (replace* or therap* or substitut* or restor* or resuscitat* or rehydrat*)):ab,ti in Clinical Trials
- #4 (#2 OR #3)
- #5 (#1 AND #4)

Acknowledgements

The authors acknowledge Dr Nancy Jiang for assistance with language translation.

References

1. Finfer S, Liu B, Taylor C, Bellomo R, Billot L, Cook D, Du B, McArthur C, Myburgh J, (2010) Resuscitation fluid use in critically ill adults: an international cross-sectional study in 391 intensive care units. *Crit Care* 14: R185
2. Perel P, Roberts I, (2012) Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev* 6: CD000567
3. Reinhart K, Perner A, Sprung CL, Jaeschke R, Schortgen F, Johan Groeneveld AB, Beale R, Hartog CS, (2012) Consensus statement of the ESICM task force on colloid volume therapy in critically ill patients. *Intensive Care Med* 38: 368-383
4. Gattas DJ, Dan A, Myburgh J, Billot L, Lo S, Finfer S, (2012) Fluid resuscitation with 6% hydroxyethyl starch (130/0.4) in acutely ill patients: an updated systematic review and meta-analysis. *Anesth Analg* 114: 159-169
5. Bunn F, Trivedi D, (2012) Colloid solutions for fluid resuscitation. *Cochrane Database Syst Rev* 7: CD001319
6. Dart AB, Mutter TC, Ruth CA, Taback SP, (2010) Hydroxyethyl starch (HES) versus other fluid therapies: effects on kidney function. *Cochrane Database Syst Rev*: CD007594
7. Zarychanski R, Turgeon AF, Fergusson DA, Cook DJ, Hébert PC, Bagshaw SM, Monsour D, McIntyre L, (2009) Renal outcomes and mortality following hydroxyethyl starch resuscitation of critically ill patients: systematic review and meta-analysis of randomized trials. *Open Medicine* 3: E196-209
8. Hartog CS, Kohl M, Reinhart K, (2011) A Systematic Review of Third-Generation Hydroxyethyl Starch (HES 130/0.4) in Resuscitation: Safety Not Adequately Addressed. *Anesth Analg* 112: 635-645
9. Antonelli M, Bonten M, Chastre J, Citerio G, Conti G, Curtis JR, De Backer D, Hedenstierna G, Joannidis M, Macrae D, Mancebo J, Maggiore SM, Mebazaa A, Preiser JC, Rocco P, Timsit JF, Wernerman J, Zhang H, (2012) Year in review in *Intensive Care Medicine* 2011. II. Cardiovascular, infections, pneumonia and sepsis, critical care organization and outcome, education, ultrasonography, metabolism and coagulation. *Intensive Care Med* 38: 345-358
10. Juni P, Altman DG, Egger M, (2001) Systematic reviews in health care: Assessing the quality of controlled clinical trials. *BMJ* 323: 42-46
11. Higgins JPT, Green S, (2011) 16.9.2 Studies with zero-cell counts. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.
12. Myburgh JA, Finfer S, Bellomo R, Billot L, Cass A, Gattas D, Glass P, Lipman J, Liu B, McArthur C, McGuinness S, Rajbhandari D, Taylor CB, Webb SAR, (2012) Hydroxyethyl Starch or Saline for Fluid Resuscitation in Intensive Care. *New England Journal of Medicine* DOI: 10.1056/NEJMoa1209759
13. Perner A, Haase N, Guttormsen AB, Tenhunen J, Klemenzson G, Aneman A, Madsen KR, Moller MH, Elkjaer JM, Poulsen LM, Bendtsen A, Winding R, Steensen M, Berezowicz P, Soe-Jensen P, Bestle M, Strand K, Wiis J, White JO, Thornberg KJ, Quist L, Nielsen J, Andersen LH, Holst LB, Thormar K, Kjaeldgaard AL,


- Fabritius ML, Mondrup F, Pott FC, Moller TP, Winkel P, Wetterslev J, (2012) Hydroxyethyl starch 130/0.42 versus Ringer's acetate in severe sepsis. *N Engl J Med* 367: 124-134
14. Siegemund M (2012-09-12) NCT00273728 BaSES Trial: Basel Starch Evaluation in Sepsis, personal communication
 15. Gondos T, Marjanek Z, Ulakcsai Z, Szabo Z, Bogar L, Karolyi M, Gartner B, Kiss K, Havas A, Futo J, (2010) Short-term effectiveness of different volume replacement therapies in postoperative hypovolaemic patients. *European Journal of Anaesthesiology* 27: 794-800
 16. Guidet B, Martinet O, Boulain T, Philippart F, Poussel JF, Maizel J, Forceville X, Feissel M, Hasselmann M, Heininger A, Van Aken H, (2012) Assessment of hemodynamic efficacy and safety of 6% hydroxyethylstarch 130/0.4 vs. 0.9% NaCl fluid replacement in patients with severe sepsis: The CRYSTMAS study. *Crit Care* 16: R94. Renal replacement results were provided by personal communication, Dr B Guidet, 22 August 2012.
 17. Zhu GC, Quan ZY, Shao YS, Zhao JG, Zhang YT, (2011) [The study of hypertonic saline and hydroxyethyl starch treating severe sepsis]. *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue* 23: 150-153
 18. James MF, Michell WL, Joubert IA, Nicol AJ, Navsaria PH, Gillespie RS, (2011) Resuscitation with hydroxyethyl starch improves renal function and lactate clearance in penetrating trauma in a randomized controlled study: the FIRST trial (Fluids in Resuscitation of Severe Trauma). *Br J Anaesth* 107: 693-702
 19. Yang J, Wang WT, Yan LN, Xu MQ, Yang JY, (2011) Alternatives to albumin administration in hepatocellular carcinoma patients undergoing hepatectomy: an open, randomized clinical trial of efficacy and safety. *Chin Med J (Engl)* 124: 1458-1464
 20. Nagpal D (2012-08-10) NCT00964015 Starch or Saline After Cardiac Surgery (SSACS) trial, personal communication
 21. Mittermayr M, Streif W, Haas T, Fries D, Velik-Salchner C, Klingler A, Oswald E, Bach C, Schnapka-Koepf M, Innerhofer P, Mittermayr M, Streif W, Haas T, Fries D, Velik-Salchner C, Klingler A, Oswald E, Bach C, Schnapka-Koepf M, Innerhofer P, (2007) Hemostatic changes after crystalloid or colloid fluid administration during major orthopedic surgery: the role of fibrinogen administration.[see comment]. *Anesthesia & Analgesia* 105: 905-917
 22. Schramko A, Suojaranta-Ylinen R, Kuitunen A, Raivio P, Kukkonen S, Niemi T, (2010) Hydroxyethylstarch and gelatin solutions impair blood coagulation after cardiac surgery: a prospective randomized trial. *British journal of anaesthesia* 104: 691-697
 23. Lu J, Zhao HY, Liu F, An YZ, (2012) [The influence of lactate Ringer solution versus hydroxyethyl starch on coagulation and fibrinolytic system in patients with septic shock]. *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue* 24: 38-41
 24. Volta CA, Alvisi V, Campi M, Marangoni E, Alvisi R, Castellazzi M, Fainardi E, Manfrinato MC, Dallochio F, Bellini T, (2007) Influence of different strategies of volume replacement on the activity of matrix metalloproteinases: an in vitro and in vivo study. *Anesthesiology* 106: 85-91
 25. Dubin A, Pozo MO, Casabella CA, Murias G, Palizas F, Moseinco MC, Kanoore Edul VS, Estensoro E, Ince C, (2010) Comparison of 6% hydroxyethyl starch 130/0.4 and saline solution for resuscitation of the


microcirculation during the early goal-directed therapy of septic patients. *Journal of Critical Care* 25: 659.e651-659.e658

26. Van Der Linden PJ, De Hert SG, Deraedt D, Cromheecke S, De Decker K, De Paep R, Rodrigus I, Daper A, Trenchant A, (2005) Hydroxyethyl starch 130/0.4 versus modified fluid gelatin for volume expansion in cardiac surgery patients: The effects on perioperative bleeding and transfusion needs. *Anesthesia and Analgesia* 101: 629-634
27. Ooi JSM, Ramzisham ARM, Zamrin MD, (2009) Is 6% hydroxyethyl starch 130/0.4 safe in coronary artery bypass graft surgery? *Asian Cardiovasc Thorac Ann* 17: 368-372
28. Zdolsek HJ, Vegfors M, Lindahl TL, Tornquist T, Bortnik P, Hahn RG, (2011) Hydroxyethyl starches and dextran during hip replacement surgery: effects on blood volume and coagulation. *Acta Anaesthesiol Scand* 55: 677-685
29. Wu Y, Wu AS, Wang J, Tian M, Jia XY, Rui Y, Yue Y, (2010) Effects of the novel 6% hydroxyethyl starch 130/0.4 on renal function of recipients in living-related kidney transplantation. *Chin Med J (Engl)* 123: 3079-3083
30. Godet G, Lehot JJ, Janvier G, Steib A, De Castro V, Coriat P, (2008) Safety of HES 130/0.4 (Voluven(R)) in patients with preoperative renal dysfunction undergoing abdominal aortic surgery: a prospective, randomized, controlled, parallel-group multicentre trial.[erratum appears in *Eur J Anaesthesiol*. 2008 Dec;25(12):1042]. *European Journal of Anaesthesiology* 25: 986-994
31. Mahmood A, Gosling P, Vohra RK, (2007) Randomized clinical trial comparing the effects on renal function of hydroxyethyl starch or gelatine during aortic aneurysm surgery. *British Journal of Surgery* 94: 427-433
32. Dolecek M, Svoboda P, Kantorova I, Scheer P, Sas I, Bibrova J, Radvanova J, Radvan M, (2009) Therapeutic influence of 20 % albumin versus 6% hydroxyethylstarch on extravascular lung water in septic patients: A randomized controlled trial. *Hepato-Gastroenterology* 56: 1622-1628
33. Schramko AA, Suojaranta-Ylinen RT, Kuitunen AH, Kukkonen SI, Niemi TT, (2009) Rapidly degradable hydroxyethyl starch solutions impair blood coagulation after cardiac surgery: A prospective randomized trial. *Anesthesia and Analgesia* 108: 30-36
34. Mukhtar A, Aboulfetouh F, Obayah G, Salah M, Emam M, Khater Y, Akram R, Hoballah A, Bahaa M, Elmeteini M, Hamza A, Mukhtar A, Aboulfetouh F, Obayah G, Salah M, Emam M, Khater Y, Akram R, Hoballah A, Bahaa M, Elmeteini M, Hamza A, (2009) The safety of modern hydroxyethyl starch in living donor liver transplantation: a comparison with human albumin. *Anesthesia & Analgesia* 109: 924-930
35. Inal MT, Memis D, Karamanlioglu B, Sut N, (2010) Effects of polygeline and hydroxyethyl starch solutions on liver functions assessed with LIMON in hypovolemic patients. *Journal of critical care* 25: 361.e361-365
36. Palumbo D, Servillo G, D'Amato L, Volpe ML, Capogrosso G, De Robertis E, Piazza O, Tufano R, (2006) The effects of hydroxyethyl starch solution in critically ill patients.[see comment]. *Minerva Anesthesiol* 72: 655-664


37. Kasper SM, Meinert P, Kampe S, Gorg C, Geisen C, Mehlhorn U, Diefenbach C, (2003) Large-dose hydroxyethyl starch 130/0.4 does not increase blood loss and transfusion requirements in coronary artery bypass surgery compared with hydroxyethyl starch 200/0.5 at recommended doses. *Anesthesiology* 99: 42-47
38. Gandhi SD, Weiskopf RB, Jungheinrich C, Koorn R, Miller D, Shangraw RE, Prough DS, Baus D, Bepperling F, Wartier DC, Gandhi SD, Weiskopf RB, Jungheinrich C, Koorn R, Miller D, Shangraw RE, Prough DS, Baus D, Bepperling F, Wartier DC, (2007) Volume replacement therapy during major orthopedic surgery using Voluven (hydroxyethyl starch 130/0.4) or hetastarch. *Anesthesiology* 106: 1120-1127
39. Langeron O, Doelberg M, Ang ET, Bonnet F, Capdevila X, Coriat P, (2001) Voluven, a lower substituted novel hydroxyethyl starch (HES 130/0.4), causes fewer effects on coagulation in major orthopedic surgery than HES 200/0.5. *Anesthesia & Analgesia* 92: 855-862
40. Sander O, Reinhart K, Meier-Hellmann A, (2003) Equivalence of hydroxyethyl starch HES 130/0.4 and HES 200/0.5 for perioperative volume replacement in major gynaecological surgery. *Acta Anaesthesiologica Scandinavica* 47: 1151-1158
41. Gallandat Huet RC, Siemons AW, Baus D, van Rooyen-Butijn WT, Haagenaars JA, van Oeveren W, Bepperling F, (2000) A novel hydroxyethyl starch (Voluven) for effective perioperative plasma volume substitution in cardiac surgery. *Can J Anaesth* 47: 1207-1215
42. Jungheinrich C, Sauermann W, Bepperling F, Vogt NH, (2004) Volume efficacy and reduced influence on measures of coagulation using hydroxyethyl starch 130/0.4 (6%) with an optimised in vivo molecular weight in orthopaedic surgery: A randomised, double-blind study. *Drugs in R and D* 5: 1-9
43. Mehta Y, Dhar A, Sujatha, Meharwal ZS, Trehan N, (2007) Comparison of new HES (130/0.4) and HES (200/0.5) in OPCAB surgery. *Journal of Anaesthesiology Clinical Pharmacology* 23: 273-278
44. Ellger B, Freyhoff J, Van Aken H, Booke M, Marcus MAE, (2006) High-dose volume replacement using HES 130/0.4 during major surgery: Impact on coagulation and incidence of postoperative itching. *Nederlands Tijdschrift voor Anesthesiologie* 19: 63-68
45. Neff TA, Doelberg M, Jungheinrich C, Sauerland A, Spahn DR, Stocker R, (2003) Repetitive large-dose infusion of the novel hydroxyethyl starch 130/0.4 in Patients with severe head injury. *Anesthesia and Analgesia* 96: 1453-1459
46. Boldt J, Lehmann A, Rompert R, Haisch G, Isgro F, (2000) Volume therapy with a new hydroxyethyl starch solution in cardiac surgical patients before cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 14: 264-268
47. Rasmussen LS YS, Schuttler J, Van Aken H, Shafer SL, Eisenach JC, Reilly CS, Miller DR, Zwissler B, Tramer MR, Antonelli M, (2011) Editors-in-Chief Statement Regarding IRB Approval of Clinical Trials by Joachim Boldt <http://www.aeditor.org/EICJointStatement.pdf> Accessed 20 February 2011
48. Rasmussen LS YS, Van Aken H, Shafer SL, Eisenach JC, Reilly CS, Miller DR, Parrillo JE, Zwissler B, Tramer MR, Antonelli M, Kaplan JA, Wiltfang J, Stefano GB, Chiumello D, Mayr WR, (2011) Editors-in-Chief Statement Regarding Published Clinical Trials Conducted without IRB Approval by Joachim Boldt <http://www.aeditor.org/EIC.Joint.Statement.on.Retractions.pdf> Accessed 7 March 2011.
49. Shafer SL, (2011) Shadow of Doubt. *Anesth Analg* 112: 498-500
50. Shafer SL, Notice of retraction. <http://www.aeditor.org/NoticeofRetraction.pdf> Accessed 20 Feb 2011.

51. Shafer SL, (2011) 25 February 2011 notice. <http://www.anesthesia-analgesia.org/site/misc/25.February.2011.Notice.pdf> Accessed 1 March 2011. Anesthesia & Analgesia
52. Haynes RB, McKibbin KA, Wilczynski NL, Walter SD, Werre SR, (2005) Optimal search strategies for retrieving scientifically strong studies of treatment from Medline: analytical survey. BMJ 330: 1179





NOTE: Weights are from random effects analysis



NOTE: Weights are from random effects analysis



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Gattas, DJ; Dan, A; Myburgh, J; Billot, L; Lo, S; Finfer, S

Title:

Fluid resuscitation with 6 % hydroxyethyl starch (130/0.4 and 130/0.42) in acutely ill patients: systematic review of effects on mortality and treatment with renal replacement therapy

Date:

2013-04-01

Citation:

Gattas, D. J., Dan, A., Myburgh, J., Billot, L., Lo, S. & Finfer, S. (2013). Fluid resuscitation with 6 % hydroxyethyl starch (130/0.4 and 130/0.42) in acutely ill patients: systematic review of effects on mortality and treatment with renal replacement therapy. *INTENSIVE CARE MEDICINE*, 39 (4), pp.558-568. <https://doi.org/10.1007/s00134-013-2840-0>.

Persistent Link:

<http://hdl.handle.net/11343/219260>

File Description:

Accepted version